

Infinite products of random matrices and repeated interaction dynamics

Laurent Bruneau^a, Alain Joye^{b,c} and Marco Merkli^{d,1}

^aLaboratoire AGM, Université de Cergy-Pontoise, Site Saint-Martin, BP 222, 95302 Cergy-Pontoise, France.

E-mail: laurent.bruneau@u-cergy.fr; url: <http://www.u-cergy.fr/bruneau>

^bInstitut Fourier, UMR 5582, CNRS-Université de Grenoble I BP 74, 38402 Saint-Martin d'Hères, France. E-mail: Alain.Joye@ujf-grenoble.fr

^cLaboratoire de Physique et Modélisation des Milieux Condensés, UMR 5493, CNRS-Université de Grenoble I, BP 166, 38042 Grenoble, France

^dDepartment of Mathematics and Statistics, Memorial University, Newfoundland, Canada.

E-mail: merkli@mun.ca; url: <http://www.math.mun.ca/~merkli/>

Received 1 April 2008; accepted 15 April 2009

Abstract. Let Ψ_n be a product of n independent, identically distributed random matrices M , with the properties that Ψ_n is bounded in n , and that M has a deterministic (constant) invariant vector. Assume that the probability of M having only the simple eigenvalue 1 on the unit circle does not vanish. We show that Ψ_n is the sum of a fluctuating and a decaying process. The latter converges to zero almost surely, exponentially fast as $n \rightarrow \infty$. The fluctuating part converges in Cesaro mean to a limit that is characterized explicitly by the deterministic invariant vector and the spectral data of $\mathbb{E}[M]$ associated to 1. No additional assumptions are made on the matrices M ; they may have complex entries and not be invertible.

We apply our general results to two classes of dynamical systems: inhomogeneous Markov chains with random transition matrices (stochastic matrices), and random repeated interaction quantum systems. In both cases, we prove ergodic theorems for the dynamics, and we obtain the limit states.

Résumé. On considère le produit Ψ_n de n matrices aléatoires M i.i.d. ayant les propriétés suivantes: la suite $(\Psi_n)_n$ est bornée et M possède un vecteur invariant déterministe (constant). On suppose que la probabilité pour que, sur le cercle unité, M possède une unique valeur propre simple en 1 soit non nulle. On montre que Ψ_n est la somme d'un processus fluctuant et d'un processus décroissant. Ce dernier tend vers zéro presque sûrement et exponentiellement rapidement lorsque n tend vers l'infini. Le terme fluctuant converge en moyenne de Cesaro vers une limite caractérisée explicitement par le vecteur invariant déterministe et par les données spectrales associées à la valeur propre 1 de la matrice $\mathbb{E}[M]$. Aucune hypothèse supplémentaire n'est faite sur les matrices M ; elles peuvent être à valeurs complexes et pas nécessairement inversibles.

On applique les résultats généraux à deux classes de systèmes dynamiques: les chaînes de Markov inhomogènes avec matrices de transition aléatoires (matrices stochastiques), ainsi que les systèmes quantiques avec interactions répétées et aléatoires. Dans les deux cas, on prouve des résultats d'ergodicité pour la dynamique et on détermine les états limites.

MSC: 60H25; 37A30

Keywords: Products of random matrices; Random dynamical systems; Random stochastic matrix; Ergodic theory

1. Introduction

In this paper we study products of infinitely many independent, identically distributed random matrices. The matrices we consider satisfy two basic properties, reflecting the fact that they describe the dynamics of random quantum or

¹Supported by NSERC Discovery Grant 205247.

classical dynamical systems. The first property is that the norm of any product of such matrices is bounded uniformly in the number of factors. It reflects the fact that the underlying dynamics is in a certain sense norm-preserving. The second property is that there is a deterministic invariant vector. This represents a normalization of the dynamics. Two important examples of systems falling into this category are inhomogeneous Markov chains with *random transition matrices*, i.e. products of *random stochastic matrices*, as well as *repeated interaction open quantum systems*. In this paper we present general results on infinite products of random matrices, and applications of these results to the two classes of dynamical systems mentioned. Our main results are convergence theorems of the infinite random matrix product, Theorems 1.1–1.3. They translate into ergodic theorems for the corresponding dynamical systems, Theorems 1.4 and 1.5.

1.1. General results

Let $M(\omega)$ be a random matrix on \mathbb{C}^d , with probability space (Ω, \mathcal{F}, p) . We say that $M(\omega)$ is a *random reduced dynamics operator* (RRDO) if:

- (1) there exists a norm $\|\cdot\|$ on \mathbb{C}^d such that, for all ω , $M(\omega)$ is a contraction on \mathbb{C}^d endowed with the norm $\|\cdot\|$;
- (2) there is a vector ψ_S , constant in ω , such that $M(\omega)\psi_S = \psi_S$, for all ω .

We shall normalize ψ_S such that $\|\psi_S\| = 1$ where $\|\cdot\|$ denotes the euclidean norm.

To an RRDO $M(\omega)$, we associate the (i.i.d.) *random reduced dynamics process* (RRDP)

$$\Psi_n(\bar{\omega}) := M(\omega_1) \cdots M(\omega_n), \quad \bar{\omega} \in \Omega^{\mathbb{N}^*}.$$

We show that Ψ_n has a decomposition into an exponentially decaying part and a fluctuating part. To identify these parts, we proceed as follows. It follows from (1) and (2) that the spectrum of an RRDO $M(\omega)$ must lie inside the closed complex unit disk, and that 1 is an eigenvalue (with eigenvector ψ_S). Let $P_1(\omega)$ denote the spectral projection of $M(\omega)$ corresponding to the eigenvalue 1 ($\dim P_1(\omega) \geq 1$),² and let $P_1^*(\omega)$ be its adjoint operator. Define

$$\psi(\omega) := P_1(\omega)^* \psi_S \tag{1.1}$$

and set

$$P(\omega) = |\psi_S\rangle\langle\psi(\omega)|.$$

For $\psi, \phi \in \mathbb{C}^d$, we denote by $|\psi\rangle\langle\phi|$ the rank-one operator $\mathbb{C}^d \ni \chi \mapsto |\psi\rangle\langle\phi|\chi = \langle\phi, \chi\rangle\psi$, and our convention is to take the inner products linear in the second factor. We put

$$Q(\omega) = \mathbb{1} - P(\omega).$$

Note that the vector $\psi(\omega)$ is normalized as $\langle\psi_S, \psi(\omega)\rangle = 1$. We decompose $M(\omega)$ as

$$M(\omega) = P(\omega) + Q(\omega)M(\omega)Q(\omega) =: P(\omega) + M_Q(\omega). \tag{1.2}$$

Taking into account this decomposition, we obtain (cf. Proposition 2.1)

$$\Psi_n(\bar{\omega}) := M(\omega_1) \cdots M(\omega_n) = |\psi_S\rangle\langle\theta_n(\bar{\omega})| + M_Q(\omega_1) \cdots M_Q(\omega_n), \tag{1.3}$$

where $\theta_n(\bar{\omega})$ is the Markov process

$$\theta_n(\bar{\omega}) = M^*(\omega_n) \cdots M^*(\omega_2)\psi(\omega_1), \tag{1.4}$$

$M^*(\omega_j)$ being the adjoint operator of $M(\omega_j)$. We analyze the two parts in the r.h.s. of (1.3) separately.

Let $\mathcal{M}_{(E)}$ be the set of RRDOs M whose spectrum on the complex unit circle consists only of a simple eigenvalue $\{1\}$.

²We recall that the spectral projection P of a matrix M associated to an eigenvalue λ is the projection onto $\text{Ker}((M - \lambda)^r)$ along $\text{Ran}((M - \lambda)^r)$ where r denotes the algebraic multiplicity of λ .

On $\Omega^{\mathbb{N}^*}$ we define the probability measure $d\mathbb{P}$ in a standard fashion by

$$d\mathbb{P} = \prod_{j \geq 1} dp_j, \quad \text{where } dp_j \equiv dp, \forall j \in \mathbb{N}^*.$$

Theorem 1.1 (Decaying process). *Let $M(\omega)$ be a random reduced dynamics operator. Suppose that $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then there exist a set $\Omega_1 \subset \Omega^{\mathbb{N}^*}$ and constants $C, \alpha > 0$ s.t. $\mathbb{P}(\Omega_1) = 1$ and s.t. for any $\bar{\omega} \in \Omega_1$, there exists an $n_0(\bar{\omega})$ s.t. for any $n \geq n_0(\bar{\omega})$,*

$$\|M_Q(\omega_1) \cdots M_Q(\omega_n)\| \leq Ce^{-\alpha n}. \quad (1.5)$$

Remarks.

1. In the case where $M(\omega) = M$ is constant, and $M \in \mathcal{M}_{(E)}$, one readily shows that for any $\varepsilon > 0$ there is a C_ε such that $\|(M_Q)^n\| \leq C_\varepsilon e^{-n(\gamma - \varepsilon)}$, for all $n \geq 0$, and where $\gamma = \min_{z \in \text{spec}(M) \setminus \{1\}} |\log |z||$ (see e.g. [8], Proposition 2.2). It is remarkable that in the random case, the mere condition of M having an arbitrarily small, non-vanishing probability to be in $\mathcal{M}_{(E)}$ suffices to guarantee the exponential decay of the product in (1.5).

2. Any stochastic matrix whose entries are all nonzero belongs to $\mathcal{M}_{(E)}$.

3. If $\{1\}$ is a simple eigenvalue of $M(\omega)$ then the decomposition (1.2) is just the spectral decomposition of the matrix $M(\omega)$.

4. The choice (1.1) ensures that $\psi(\omega)$ is an eigenvector of $M^*(\omega)$. Other choices of measurable $\psi(\omega)$ which are bounded in ω lead to different decompositions of $M(\omega)$, and can be useful as well. For instance, if $M(\omega)$ is a bistochastic matrix, then one can take for $\psi(\omega)$ an $M^*(\omega)$ -invariant vector which is constant in ω .

Our next result concerns the asymptotics of the Markov process (1.4). Set $\mathbb{E}[f] = \int_\Omega f(\omega) dp(\omega)$ for a random variable f , and denote by $P_{1, \mathbb{E}[M]}$ the spectral projection of $\mathbb{E}[M]$ onto the eigenvalue $\{1\}$.

Theorem 1.2 (Fluctuating process). *Let $M(\omega)$ be a random reduced dynamics operator. Suppose that $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then we have $\mathbb{E}[M] \in \mathcal{M}_{(E)}$. Moreover, there exists a set $\Omega_2 \subset \Omega^{\mathbb{N}^*}$ s.t. $\mathbb{P}(\Omega_2) = 1$ and, for all $\bar{\omega} \in \Omega_2$,*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \theta_n(\bar{\omega}) = \theta, \quad (1.6)$$

where

$$\theta = (\mathbb{1} - \mathbb{E}[M_Q]^*)^{-1} \mathbb{E}[\psi] = P_{1, \mathbb{E}[M]}^* \mathbb{E}[\psi] = P_{1, \mathbb{E}[M]}^* \psi_S. \quad (1.7)$$

Remarks.

5. In the case where M is constant in ω , we have $\mathbb{E}[M_Q]^* = (M_Q)^*$, $\mathbb{E}[\psi] = \psi$, and under the assumption of Theorem 1.2, that $M \in \mathcal{M}_{(E)}$. Therefore, $P_1 = P = |\psi_S\rangle\langle\psi|$ and hence $Q^*\psi = 0$, and $(M_Q)^*\psi = 0$. Consequently, we have $\theta = \psi$. This coincides with the results of [8]. As we will see in Section 3.4, the latter equality is not satisfied for general, ω -dependent matrices M .

6. The ergodic average limit of $\theta_n(\bar{\omega})$ does not depend on the particular choice of $\psi(\omega)$. This follows from the last equality in (1.7).

7. We show in Proposition 2.5 that for every fixed $\bar{\omega}$, $\theta_n(\bar{\omega})$ converges if and only if $\psi(\omega_n)$ converges, and that the limits coincide if they exist.

Combining Theorems 1.1 and 1.2 we obtain the following result.

Theorem 1.3 (Ergodic theorem for RRDP). *Let $M(\omega)$ be a random reduced dynamics operator. Suppose $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then there exists a set $\Omega_3 \subset \Omega^{\mathbb{N}^*}$ s.t. $\mathbb{P}(\Omega_3) = 1$ and, for all $\bar{\omega} \in \Omega_3$,*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N M(\omega_1) \cdots M(\omega_n) = |\psi_S\rangle\langle\theta| = P_{1, \mathbb{E}[M]}. \quad (1.8)$$

Remark. 8. If one can choose $\psi(\omega) \equiv \psi$ to be independent of ω (see also remark 4 above), then we show (see (2.8)) that $\theta_n(\bar{\omega}) = \psi$, for all $n, \bar{\omega}$. It thus follows from (1.3)–(1.5) that $\lim_{n \rightarrow \infty} M(\omega_1) \cdots M(\omega_n) = |\psi_S\rangle\langle\psi|$, a.s., exponentially fast.

A simple example. Fix $R > 0$. Then for any $C \in \mathbb{C}$, $|C| \leq R$, and any $\alpha \in \mathbb{C}$, $|\alpha| \leq 1$, the matrix $M(C, \alpha) = \begin{pmatrix} 1 & C(1-|\alpha|) \\ 0 & \alpha \end{pmatrix}$ is an RRDO for the norm $\| \begin{pmatrix} x \\ y \end{pmatrix} \| = |x| + R|y|$. Moreover, $M(C, \alpha) \in \mathcal{M}_{(E)}$ iff $|\alpha| < 1$.

1.2. Inhomogeneous random Markov chains

Products of random matrices satisfying conditions (1) and (2) are important in the study of inhomogenous random Markov chains. A finite Markov chain is a system consisting of $d < \infty$ states on which a stochastic process is defined in the following way. At time step n , $n \in \mathbb{N}^*$, the probability to make a transition from state i to state j , where $i, j \in \{1, \dots, d\}$, is given by $p_{ij}^{(n)}$. Hence the relation $\sum_{j=1}^d p_{ij}^{(n)} = 1$. Arranged into a $d \times d$ matrix $(M_n)_{ij} = p_{ij}^{(n)}$, these probabilities yield a stochastic matrix. The probability of jumping from state i to state j between times 0 and n is given by the matrix element $i j$ of the product

$$\Psi_n = M_1 M_2 \cdots M_n. \tag{1.9}$$

The Markov chain (1.9) is called *inhomogeneous* if the transition probabilities depend on the time step n . We have that for all n :

- M_n is a contraction for the norm $\|M\| = \max_{i=1, \dots, d} \sum_{j=1}^d |(M)_{ij}|$, and
- the vector $\psi_S = \frac{1}{\sqrt{d}}(1, 1, \dots, 1)^T$ is a normalized eigenvector of M_n , associated to the eigenvalue 1.

Consequently, if the transition probabilities are chosen randomly at each time step, the corresponding operator $M(\omega)$ is an RRDO, and the product (1.9) is an RRDP. If $p_{ij}^{(n)} > 0$, for all i, j , then the Perron–Frobenius theorem asserts that 1 is a simple eigenvalue of M_n , and that all other eigenvalues lie inside the open complex unit disk. Thus the conclusions of Theorems 1.1–1.3 hold for $M(\omega)$ a random stochastic matrix, which, with positive probability, has strictly positive entries.

Results on convergence of products of random stochastic matrices, or non-negative matrices, are numerous, see e.g. the references in Section 1.4. However, these results mostly concern properties of the limiting distribution, if it exists, in terms of the properties of the distribution of the RRDOs. Those results rely heavily on the positivity of the elements of the considered matrices. When studying convergence in distribution, the order of the factors in the product does not matter, and usually products of the form

$$\Phi_n = M_n M_{n-1} \cdots M_1 \tag{1.10}$$

are studied as well (compare with (1.9)). While products of the form (1.10) are not our main concern (and are easier to study), our techniques still yield results for them that are stronger than those for products (1.9). We show in Section 3.5 a general result on products (1.10) which, applied to products of stochastic matrices, yields the following result.

Theorem 1.4. Suppose that $M(\omega)$ takes values in the set of stochastic matrices, and that $p(M(\omega) \in \mathcal{P}_E) > 0$, where \mathcal{P}_E denotes the set of stochastic matrices with strictly positive entries. Then there exists an $\alpha > 0$ and a set $\Omega_4 \subset \Omega^{\mathbb{N}^*}$, such that $\mathbb{P}(\Omega_4) = 1$, and, for all $\bar{\omega} \in \Omega_4$,

$$\Phi_n(\bar{\omega}) = M(\omega_n)M(\omega_{n-1}) \cdots M(\omega_1) = |\psi_S\rangle\langle\eta_\infty(\omega)| + O_{\bar{\omega}}(e^{-\alpha n}), \tag{1.11}$$

where $\eta_\infty(\omega)$ is a vector valued random variable, where $\psi_S = \frac{1}{\sqrt{d}}(1, 1, \dots, 1)^T$, and where the remainder term depends on $\bar{\omega} \in \Omega_4$.

Remark. 9. As expected, $\mathbb{E}[\eta_\infty] = \theta$, with θ given in (1.7).

1.3. Random repeated interaction open quantum systems

In this section we present the mathematical framework of open quantum systems. We establish the link between the dynamics and infinite products of random matrices, and we present an application of Theorems 1.1–1.3.

1.3.1. Definition of repeated interaction quantum systems

A repeated interaction quantum system consists of a subsystem \mathcal{S} which interacts successively with the elements \mathcal{E}_m of a chain $\mathcal{C} = \mathcal{E}_1 + \mathcal{E}_2 + \dots$ of independent quantum systems. At each moment in time, \mathcal{S} interacts precisely with one \mathcal{E}_m (m increases as time does), while the other elements in the chain evolve freely according to their intrinsic (uncoupled) dynamics. The complete evolution is described by the intrinsic dynamics of \mathcal{S} and of all the \mathcal{E}_m , plus an interaction between \mathcal{S} and \mathcal{E}_m , for each m . The latter consists of an interaction time $\tau_m > 0$, and an interaction operator V_m (acting on \mathcal{S} and \mathcal{E}_m); during the time interval $[\tau_1 + \dots + \tau_{m-1}, \tau_1 + \dots + \tau_m)$, \mathcal{S} is coupled to \mathcal{E}_m via V_m . Systems with this structure are important not only from a mathematical, but also from a physical point of view. They arise in fundamental experiments on the interaction of matter with quantized radiation. We refer to the end of this introduction for more information and references concerning this aspect of our work.

According to the fundamental principles of quantum mechanics, states of the systems \mathcal{S} and \mathcal{E}_m are given by normalized vectors (or density matrices) on Hilbert spaces $\mathcal{H}_{\mathcal{S}}$ and $\mathcal{H}_{\mathcal{E}_m}$, respectively, [2,5].³ We assume that $\dim \mathcal{H}_{\mathcal{S}} < \infty$, while $\dim \mathcal{H}_{\mathcal{E}_m}$ may be infinite. Observables $A_{\mathcal{S}}$ and $A_{\mathcal{E}_m}$ of the systems \mathcal{S} and \mathcal{E}_m are bounded operators forming von Neumann algebras $\mathfrak{M}_{\mathcal{S}} \subset \mathcal{B}(\mathcal{H}_{\mathcal{S}})$ and $\mathfrak{M}_{\mathcal{E}_m} \subset \mathcal{B}(\mathcal{H}_{\mathcal{E}_m})$. They evolve according to the Heisenberg dynamics $\mathbb{R} \ni t \mapsto \alpha_{\mathcal{S}}^t(A_{\mathcal{S}})$ and $\mathbb{R} \ni t \mapsto \alpha_{\mathcal{E}_m}^t(A_{\mathcal{E}_m})$, where $\alpha_{\mathcal{S}}^t$ and $\alpha_{\mathcal{E}_m}^t$ are $*$ -automorphism groups of $\mathfrak{M}_{\mathcal{S}}$ and $\mathfrak{M}_{\mathcal{E}_m}$, respectively, see e.g. [5]. We introduce distinguished reference states, given by vectors $\psi_{\mathcal{S}} \in \mathcal{H}_{\mathcal{S}}$ and $\psi_{\mathcal{E}_m} \in \mathcal{H}_{\mathcal{E}_m}$. Typical choices for $\psi_{\mathcal{S}}$, $\psi_{\mathcal{E}_m}$ are equilibrium (KMS) states for the dynamics $\alpha_{\mathcal{S}}^t$, $\alpha_{\mathcal{E}_m}^t$, at inverse temperatures $\beta_{\mathcal{S}}$, $\beta_{\mathcal{E}_m}$. The Hilbert space of states of the total system is the tensor product

$$\mathcal{H} = \mathcal{H}_{\mathcal{S}} \otimes \mathcal{H}_{\mathcal{C}},$$

where $\mathcal{H}_{\mathcal{C}} = \bigotimes_{m \geq 1} \mathcal{H}_{\mathcal{E}_m}$, and where the infinite product is taken with respect to $\psi_{\mathcal{C}} = \bigotimes_{m \geq 1} \psi_{\mathcal{E}_m}$. The non-interacting dynamics is the product of the individual dynamics, defined on the algebra $\mathfrak{M}_{\mathcal{S}} \otimes_{m \geq 1} \mathfrak{M}_{\mathcal{E}_m}$ by $\alpha_{\mathcal{S}}^t \otimes_{m \geq 1} \alpha_{\mathcal{E}_m}^t$. It is useful to consider the dynamics in the Schrödinger picture, i.e. as acting on vectors in \mathcal{H} . To do this, we first implement the dynamics via unitaries, satisfying

$$\alpha_{\#}^t(A_{\#}) = e^{itL_{\#}} A_{\#} e^{-itL_{\#}}, \quad t \in \mathbb{R}, \quad \text{and} \quad L_{\#} \psi_{\#} = 0 \quad (1.12)$$

for any $A_{\#} \in \mathfrak{M}_{\#}$, where $\#$ stands for either \mathcal{S} or \mathcal{E}_m . The self-adjoint operators $L_{\mathcal{S}}$ and $L_{\mathcal{E}_m}$, called Liouville operators, act on $\mathcal{H}_{\mathcal{S}}$ and $\mathcal{H}_{\mathcal{E}_m}$, respectively. The existence and uniqueness of $L_{\#}$ satisfying (1.12) is well known, under general assumptions on the reference states $\psi_{\#}$ [5]. In particular, (1.12) holds if the reference states are equilibrium states. Let $\tau_m > 0$ and $V_m \in \mathfrak{M}_{\mathcal{S}} \otimes \mathfrak{M}_{\mathcal{E}_m}$ be the interaction time and interaction operator associated to \mathcal{S} and \mathcal{E}_m . We define the (discrete) repeated interaction Schrödinger dynamics of a state vector $\phi \in \mathcal{H}$, for $m \geq 0$, by

$$U(m)\phi = e^{-i\tilde{L}_m} \dots e^{-i\tilde{L}_2} e^{-i\tilde{L}_1} \phi, \quad (1.13)$$

where

$$\tilde{L}_k = \tau_k L_k + \tau_k \sum_{n \neq k} L_{\mathcal{E}_n} \quad (1.14)$$

describes the dynamics of the system during the time interval $[\tau_1 + \dots + \tau_{k-1}, \tau_1 + \dots + \tau_k)$, which corresponds to the time-step k of our discrete process. Here L_k is

$$L_k = L_{\mathcal{S}} + L_{\mathcal{E}_k} + V_k, \quad (1.15)$$

³A normalized vector ψ defines a “pure” state $A \mapsto \langle \psi, A \psi \rangle = \text{Tr}(\varrho_{\psi} A)$, where $\varrho_{\psi} = |\psi\rangle\langle\psi|$. A general “mixed” state is given by a density matrix $\varrho = \sum_{n \geq 1} p_n \varrho_{\psi_n}$, where the probabilities $p_n \geq 0$ sum up to one, and where the ψ_n are normalized vectors.

acting on $\mathcal{H}_S \otimes \mathcal{H}_{\mathcal{E}_k}$. Of course, we understand that the operator $L_{\mathcal{E}_n}$ in (1.14) acts nontrivially only on the n th factor of the Hilbert space \mathcal{H}_C of the chain.

A state $\varrho(\cdot) = \text{Tr}(\rho \cdot)$ given by density matrix ρ on \mathcal{H} is called a *normal state*. (A density matrix is a self-adjoint, non-negative trace-class operator of unit trace.) Our goal is to understand the large-time asymptotics ($m \rightarrow \infty$) of expectations

$$\varrho(U(m)^* O U(m)) = \varrho(\alpha_{RI}^m(O)) \quad (1.16)$$

for normal states ϱ and certain classes of observables O that we specify below. We denote the random repeated interaction dynamics by

$$\alpha_{RI}^m(O) = U(m)^* O U(m). \quad (1.17)$$

1.3.2. Reduced dynamics and random matrix products

Let us explain how we link the dynamics to a product of reduced dynamics operators. In order not to muddle the essence of our argument, we only consider the expectation of an observable $A_S \in \mathfrak{M}_S$, and we take the initial state of the entire system to be given by the vector

$$\psi_0 = \psi_S \otimes \psi_C, \quad (1.18)$$

where the ψ_S and ψ_C are the reference states introduced above. (See Section 4 for the general case.) The expectation of A_S at the time-step m is

$$\langle \psi_0, \alpha_{RI}^m(A_S) \psi_0 \rangle = \langle \psi_0, P e^{i\tilde{L}_1} \dots e^{i\tilde{L}_m} A_S e^{-i\tilde{L}_m} \dots e^{-i\tilde{L}_1} P \psi_0 \rangle. \quad (1.19)$$

We write simply A_S for $A_S \otimes \mathbb{1}_C$, and we have introduced

$$P = \mathbb{1}_{\mathcal{H}_S} \bigotimes_{m \geq 1} P_{\psi_{\mathcal{E}_m}}, \quad (1.20)$$

the orthogonal projection onto $\mathcal{H}_S \otimes \mathbb{C} \psi_C$. A *first important ingredient* of our approach is to construct operators K_k with the properties

$$e^{i\tilde{L}_k} A e^{-i\tilde{L}_k} = e^{iK_k} A e^{-iK_k}, \quad (1.21)$$

$$K_k \psi_S \otimes \psi_C = 0, \quad (1.22)$$

where A in (1.21) is any observable of the total system. Equality (1.21) means that the operators K_k implement the same dynamics as the \tilde{L}_k . Relation (1.22) selects a unique generator of the dynamics among all that satisfy (1.21). The existence of operators K_k satisfying (1.21) and (1.22) is linked to the deep Tomita–Takesaki theory of von Neumann algebras, cf. [8] and references therein. It turns out that the K_k are non-normal operators on \mathcal{H} (while the \tilde{L}_k are self-adjoint). We combine (1.21) with (1.19) to obtain

$$\langle \psi_0, \alpha_{RI}^m(A_S) \psi_0 \rangle = \langle \psi_0, P e^{iK_1} \dots e^{iK_m} P A_S \psi_0 \rangle. \quad (1.23)$$

A *second important ingredient* of our approach is to realize that

$$P e^{iK_1} \dots e^{iK_m} P = P e^{iK_1} P \dots P e^{iK_m} P, \quad (1.24)$$

which follows from the independence of the systems \mathcal{E}_m (see (4.7)). We identify $P e^{iK_k} P$ with an operator M_k on \mathcal{H}_S , and thus obtain, from (1.23) and (1.24),

$$\langle \psi_0, \alpha_{RI}^m(A_S) \psi_0 \rangle = \langle \psi_S, M_1 \dots M_m A_S \psi_S \rangle. \quad (1.25)$$

Because the operators $M_k = P e^{iK_k} P$ implement a dynamics, we can show (Lemma 4.1) that the M_k are contractions for some suitable norm $\|\cdot\|$ of \mathbb{C}^d . Moreover, it follows from (1.22) that $M_k \psi_S = \psi_S$, for all k . The matrices M_k satisfy conditions (1) and (2) above and are thus RRDOs.

Remark. 10. *The process of reduction of the dynamics explained in this section does not involve any approximation. This is in contrast e.g. with master equation techniques, where a reduced dynamics is obtained using Born and Markov approximations, or so-called weak coupling approximations. Our method allows us to use the structure of repeated interaction systems, and to do without all these approximations.*

1.3.3. Ergodicity of repeated interaction systems

Let us denote by $\alpha_{RI}^{n,\bar{\omega}}$, $\bar{\omega} \in \Omega^{\mathbb{N}^*}$, the random reduced operator process obtained from (1.17), (1.25), where the M_j in (1.25) are i.i.d. random matrices. We call $\alpha_{RI}^{n,\bar{\omega}}$ the random repeated interaction dynamics determined by the RRDO $M(\omega) = P e^{iK(\omega)} P$.

Theorem 1.5 (Ergodic theorem for random repeated interaction systems). *Let $\alpha_{RI}^{n,\bar{\omega}}$ be the random repeated interaction dynamics determined by an RRDO $M(\omega)$. Suppose that $\mathbb{p}(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then there exists a set $\Omega_5 \subset \Omega^{\mathbb{N}^*}$, s.t. $\mathbb{P}(\Omega_5) = 1$, and s.t. for any $\bar{\omega} \in \Omega_5$, any normal state ϱ and any $A_S \in \mathfrak{M}_S$,*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \varrho(\alpha_{RI}^{n,\bar{\omega}}(A_S)) = \langle \theta, A_S \psi_S \rangle, \quad (1.26)$$

where θ is given by (1.7).

Remarks.

11. *For the normal state ϱ given by the density matrix $\rho = |\psi_0\rangle\langle\psi_0|$, (1.26) follows readily from Theorem 1.3 and relation (1.25). To prove the result for arbitrary normal states, one uses a property of the reference state ψ_0 called cyclicity and separability [5,8].*

12. *Our setup allows us to treat systems having various sources of randomness. Not only do we allow for random interactions, but also for random characteristics of the systems \mathcal{E}_m and \mathcal{S} (interesting cases are random temperatures and dimensions of the \mathcal{E}_m and of \mathcal{S}).*

13. *In [9], we prove a similar result for a general class of so-called instantaneous observables (involving also operators on the chain \mathcal{C}), and we discuss the physical properties of the asymptotic state. See also [8] for the case where $M(\omega) \equiv M$.*

14. *As mentioned above, $\langle \psi_0, \alpha_{RI}^{n,\bar{\omega}}(A_S) \psi_0 \rangle$ is fluctuating, it does not converge pointwise (in $\bar{\omega} \in \Omega^{\mathbb{N}^*}$), but only in the ergodic average sense.*

1.3.4. An explicit example: Spin–spin systems

We consider the application of our general results to the concrete situation where both \mathcal{S} and \mathcal{E} are two-level “atoms,” and where the interaction induces energy exchange processes. This is a particular case of the third example in [8]. We present the following results without proofs, a complete analysis of this (and more general) examples is given in [9].

Both \mathcal{S} and \mathcal{E} are two-level atoms with Hamiltonians h_S, h_E having ground state energy zero, and excited energies E_S and E_E , respectively, i.e. $h_S = \begin{bmatrix} 0 & 0 \\ 0 & E_S \end{bmatrix}$ and $h_E = \begin{bmatrix} 0 & 0 \\ 0 & E_E \end{bmatrix}$. The Hamiltonian describing the interaction of \mathcal{S} with one \mathcal{E} is given by $h = h_S + h_E + \lambda v$, where λ is a coupling parameter, and v induces energy exchange processes,

$$v := a_S \otimes a_E^* + a_S^* \otimes a_E. \quad (1.27)$$

Here, $a_{\#} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $a_{\#}^* = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ are the annihilation and creation operators respectively, of $\# = \mathcal{S}, \mathcal{E}$. The Gibbs state at inverse temperature β is given by

$$\varrho_{\beta,\#}(A) = \frac{\text{Tr}(e^{-\beta h_{\#}} A)}{Z_{\beta,\#}}, \quad \text{where } Z_{\beta,\#} = \text{Tr}(e^{-\beta h_{\#}}). \quad (1.28)$$

We take the reference state to be $\psi_0 = \psi_S \otimes_{m \geq 1} \psi_{\mathcal{E}_m, \beta_{\mathcal{E}_m}}$, where ψ_S is the tracial state on \mathcal{S} , and $\psi_{\mathcal{E}_m, \beta_{\mathcal{E}_m}}$ is the Gibbs state of \mathcal{E}_m (represented by a single vector in an appropriate “GNS” Hilbert space, see [9]).

The following results deal with three situations: (1) The interaction time τ is random. It is physically reasonable to assume that $\tau(\omega)$ varies within an interval of uncertainty, since it cannot be controlled exactly in experiments. (2) The

excitation energy of \mathcal{E} is random. This situation occurs if various kinds of atoms are injected into the cavity, or if some impurity atoms enter it. (3) The temperature of the incoming atoms is random. This is physically reasonable since the incoming atom beam's temperature cannot be controlled exactly in experiments.

Theorem 1.6 (Random spin–spin system). Set $T := \frac{2\pi}{\sqrt{(E_S - E_{\mathcal{E}})^2 + 4\lambda^2}}$.

1. Random interaction time. Suppose that $\beta_{\mathcal{E}_m} = \beta$ is constant, and that $\tau(\omega) > 0$ is a random variable satisfying $p(\tau \notin T\mathbb{N}) \neq 0$. Then there exists a set $\tilde{\Omega} \subset \Omega^{\mathbb{N}^*}$ of probability one, there are constants $C, \alpha > 0$, and there is a random variable $n_0(\bar{\omega})$ satisfying $\mathbb{E}[e^{\alpha n_0}] < \infty$ such that, for each $\bar{\omega} \in \tilde{\Omega}$, for all $n \geq n_0(\bar{\omega})$, all observables A and all normal initial states ϱ :

$$|\varrho(\alpha_{\bar{\omega}}^n(A)) - \varrho_{\beta', S}(A)| \leq C e^{-\alpha n}, \quad (1.29)$$

with $\beta' = \beta_1 := \beta E_{\mathcal{E}}/E_S$.

2. Random excitation energy of \mathcal{E} . Suppose that τ and $\beta_{\mathcal{E}_m} = \beta$ are constant, and that $E_{\mathcal{E}}(\omega) > 0$ is a random variable satisfying $p(\tau \notin T\mathbb{N}) \neq 0$. (Here, $T = T(\omega)$ is random via $E_{\mathcal{E}}(\omega)$.) Then there exists a set $\tilde{\Omega} \subset \Omega^{\mathbb{N}^*}$ of probability one s.t. for all $\bar{\omega} \in \tilde{\Omega}$, for all normal initial states ϱ on \mathfrak{M} and for all observables A of S ,

$$\lim_{\mu \rightarrow \infty} \frac{1}{\mu} \sum_{m=1}^{\mu} \varrho(\alpha_{\bar{\omega}}^m(A)) = \varrho_{\beta', S}(A), \quad (1.30)$$

with $\beta' = \beta_2 := -E_S^{-1} \log(2\{1 - (1 - \mathbb{E}[e_0])^{-1} \mathbb{E}[(1 - e_0)(1 - 2Z_{\beta E_{\mathcal{E}}/E_S, S}^{-1})]\}^{-1} - 1)$, and where

$$e_0 = \left| \frac{(E_S - E_{\mathcal{E}} - \sqrt{(E_S - E_{\mathcal{E}})^2 + 4\lambda^2})^2 + 4\lambda^2 e^{i\tau \sqrt{(E_S - E_{\mathcal{E}})^2 + 4\lambda^2}}}{(E_S - E_{\mathcal{E}} - \sqrt{(E_S - E_{\mathcal{E}})^2 + 4\lambda^2})^2 + 4\lambda^2} \right|^2. \quad (1.31)$$

3. Random temperature of \mathcal{E} . Suppose that $\beta(\omega)$ is a random variable, and that $\tau > 0$ satisfies $\tau \notin T\mathbb{N}$. Then there exists a set $\tilde{\Omega} \subset \Omega^{\mathbb{N}^*}$ of probability one s.t. for all $\bar{\omega} \in \tilde{\Omega}$, for all normal initial states ϱ on \mathfrak{M} and for all observables A of S , (1.30) holds with $\beta' = \beta_3 := -E_S^{-1} \log(\mathbb{E}[Z_{\beta(\omega)E_{\mathcal{E}}/E_S, S}^{-1}]^{-1} - 1)$.

Remark.

15. The better convergence in (1) is due to the fact that in this case we can choose $\psi(\omega) \equiv \psi$ and hence Proposition 3.3 applies.

16. In (3), in general $\beta_3 \neq \bar{\beta}_1$ which reflects the fact that in Lemma 3.4, we usually have $\theta \neq \bar{\psi}$, as mentioned in Remark 1 after the proof of the lemma.

1.4. Related works

Random matrices

Results on the convergence (of some kind) of products of random stochastic matrices, and non-negative matrices, are numerous, see e.g. [15,19,20] and references therein. These results mostly concern properties of the limiting distribution, if it exists, in terms of the properties of the distribution of the RRDOs. The techniques used to obtain those results rely heavily on the positivity of matrix elements. Random matrix products have been heavily studied also for matrices in $Gl_d(\mathbb{R})$, see e.g. [1,12,16]. Again, the main focus of these works is on the study of the properties of the limiting distribution, if it exists, and on the properties of the Lyapunov exponents. In this case, the group property of the set of invertible matrices $Gl_d(\mathbb{R})$ plays a prominent role in the derivation of the results.

By contrast, besides conditions (1) and (2) defining RRDO, we do not require our matrices to be real valued, to have positive entries, or to be invertible. Moreover, we are concerned here with the limiting properties of the products only, not with the limiting distribution. On the one hand, we get a.s. convergence results for the products Φ_n (Theorem 1.4).

On the other hand, in order to eliminate the unavoidable fluctuations in the products Ψ_n , we resort to a limit in an average sense, the Cesaro limit (Theorems 1.2 and 1.3). Let us also point out that our results show as well that the top Lyapunov exponent of products of RRDOs is zero, and that it is almost surely of multiplicity one, see Theorem 3.6.

Getting informations on the fluctuations of the process around its limiting value is certainly an interesting and important issue. It amounts to getting informations about the law of the vector valued random variable η_∞ of Theorem 1.4, which is quite difficult in general. There are recent partial results about aspects of the law of such random vectors in case they are obtained by means of matrices belonging to some subgroups of $Gl_d(\mathbb{R})$ satisfying certain irreducibility conditions, see e.g. [10]. However, these results do not apply to our situation.

Random ergodic theorems for products $\Psi_n(\bar{\omega})$ have been obtained in a more general framework in [4]. They prove the almost sure existence of a Cesaro limit for these products. However, the identification of the limit is not provided by this general result. This identification relies on the detailed properties of the matrices involved, and in particular in the separation of a fluctuating part from a decaying part in the dynamical process. In this respect, our contribution consists in identifying completely the Cesaro limit of products of RRDO.

Actually, in the case under consideration, Theorem 1.1 yields enough information so that Theorem 1.3 becomes a consequence of it and of the existence result proven in [4]. Since a direct self-contained proof of Theorem 1.3 is easy to derive, given Theorem 1.1, we present it for completeness.

We close this comparison by noting that the paper [6] contains results, in a deterministic setup, which are close to some of the results we prove in Section 2. The authors of [6] consider infinite products of contractions for some norm on $M_d(\mathbb{C})$, in the reverse order with respect to RRDP. They prove convergence of the product under the following assumptions: (i) there exists a subsequence of the matrices appearing in the product which converges to a matrix which is paracontracting; (ii) the set of invariant vectors of this matrix is contained in the set of invariant vectors of all matrices appearing in the product. This last assumption implies our condition (2) in a deterministic setup. However, the products in [6] are in the reverse order with respect to those we address, and they address the deterministic framework only.

Repeated interaction systems

In the experimental setup of a ‘‘One-Atom Maser’’ the system \mathcal{S} represents one or several modes of the quantized electromagnetic field in a cavity, while the elements \mathcal{E}_m describe atoms injected into the cavity, one by one, interacting with the radiation while passing through the cavity. After interaction, the atoms encode certain properties of the field that can be measured after they exit the cavity [17,22]. In an idealized model for this process, one assumes that all the elements \mathcal{E}_m represent a copy of the same, fixed quantum system, and that the interaction is given by a fixed interaction time and a fixed interaction operator (the same for all m). Such *idealized repeated interaction systems* have been analyzed mathematically in [8,23]. However, it is clear that in actual experiments, neither the interaction time τ_m (or V_m) nor the elements \mathcal{E}_m can be exactly the same for all m ! Typically, the interaction time will be *random* (because of the random velocities of the atoms in the beam, see e.g. [7,11]), given e.g. by a Gaussian or by a uniform distribution around a mean value, and the state of the incoming atoms will be random as well, for instance determined by a temperature that fluctuates slightly around a mean temperature. (In experiments, the atoms are ejected from an atom oven, then they are cooled down to a wanted temperature before entering the cavity.) It is therefore important to develop a theory that allows for *random repeated interactions*, as we do in the present paper. Our approach is based on the techniques of non-equilibrium quantum statistical mechanics developed in [8]. We are not aware of any other work dealing with variable and random interactions in the quantum context.

1.5. *Organization of this paper*

Our paper is organized as follows. In Section 2, we prove several deterministic results, including the fact that Ψ_n converges (for fixed $\bar{\omega}$) if and only if there are no fluctuations in the matrix product (Proposition 2.5). In Section 3 we prove our main results, Theorems 1.1 and 1.2. In Section 4 we explain the link between repeated interaction quantum dynamics and matrix products.

2. Deterministic results

In this section, we derive some algebraic formulae and some uniform bounds which will play a crucial role in our analysis. However, there is no probabilistic issue involved here and we shall therefore *freeze* the random variable. We will thus simply denote $M_j = M(\omega_j)$, and

$$\Psi_n := M_1 \cdots M_n. \tag{2.1}$$

2.1. Decomposition of the M_j

Let $P_{1,j}$ denote the spectral projection of M_j for the eigenvalue 1 and define

$$\psi_j := P_{1,j}^* \psi_S, \quad P_j := |\psi_S\rangle\langle\psi_j|. \tag{2.2}$$

Note that $\langle\psi_j|\psi_S\rangle = 1$ so that P_j is a projection and, moreover, $M_j^* \psi_j = \psi_j$. We introduce the following decomposition of M_j

$$M_j := P_j + Q_j M_j Q_j, \quad \text{with } Q_j = \mathbb{1} - P_j. \tag{2.3}$$

We denote the part of M_j in $Q_j \mathbb{C}^d$, by $M_{Q_j} := Q_j M_j Q_j$. It easily follows from these definitions that

$$P_j P_k = P_k, \quad Q_j Q_k = Q_j, \tag{2.4}$$

$$Q_j P_k = 0, \quad P_k Q_j = P_k - P_j = Q_j - Q_k. \tag{2.5}$$

Proposition 2.1. *For any n ,*

$$\Psi_n = |\psi_S\rangle\langle\theta_n| + M_{Q_1} \cdots M_{Q_n}, \tag{2.6}$$

where

$$\theta_n = \psi_n + M_{Q_n}^* \psi_{n-1} + \cdots + M_{Q_n}^* \cdots M_{Q_2}^* \psi_1 \tag{2.7}$$

$$= M_n^* \cdots M_2^* \psi_1 \tag{2.8}$$

and where $\langle\psi_S, \theta_n\rangle = 1$.

Proof. Inserting the decomposition (2.3) into (2.1), and using (2.4), (2.5), we have

$$\Psi_n = \sum_{j=1}^n P_j M_{Q_{j+1}} \cdots M_{Q_n} + M_{Q_1} \cdots M_{Q_n}.$$

Since $P_j = |\psi_S\rangle\langle\psi_j|$, this proves (2.6) and (2.7). From (2.5), we obtain for any j, k ,

$$M_{Q_j} M_{Q_k} = M_{Q_j} M_k = Q_j M_j M_k. \tag{2.9}$$

Hence, $\Psi_n = P_1 M_1 \cdots M_n + Q_1 M_1 \cdots M_n = |\psi_S\rangle\langle M_n^* \cdots M_2^* \psi_1| + M_{Q_1} \cdots M_{Q_n}$, which proves (2.8). \square

The following lemma is useful in the study of the asymptotic behaviour of Ψ_n . It is proven as the passage from (2.7) to (2.8) is, simply by considering the sequence $(M_{j_n})_n$ instead of $(M_n)_n$.

Lemma 2.2. *We have $\sum_{k=1}^n M_{Q_{j_1}}^* \cdots M_{Q_{j_k}}^* \psi_{j_{k+1}} = M_{j_1}^* \cdots M_{j_n}^* \psi_{j_{n+1}}$, for any sequence of indices $(j_n)_n$.*

2.2. Uniform bounds

The operators M_j , and hence the product Ψ_n , are contractions on \mathbb{C}^d for the norm $\|\cdot\|$. In order to study their asymptotic behaviour, we need some uniform bounds on the P_j, Q_j, \dots . We denote by $\|\cdot\|$ the Euclidean norm on \mathbb{C}^d . The operator norm of a rank one projection $P = |\phi\rangle\langle\chi|$ is simply $\|P\| = \|\phi\|\|\chi\|$ (see also after (1.1)). Recall also that $\|\psi_S\| = 1$.

Lemma 2.3. *We have $\sup\{\|M_{j_n}M_{j_{n-1}}\cdots M_{j_1}\|, n \in \mathbb{N}^*, j_k \in \mathbb{N}^*\} = C_0 < \infty$. In particular, $\|\Psi_n\| \leq C_0$.*

Proof. This follows from $\|M_j\| \leq 1, j \in \mathbb{N}^*$, and from the equivalence of the norms $\|\cdot\|$ and $\|\cdot\|$. \square

As a consequence, we get the following fundamental uniform estimates.

Proposition 2.4. *Let C_0 be as in Lemma 2.3. Then, the following bounds hold:*

1. *For any $j \in \mathbb{N}^*$, $\|P_j\| = \|\psi_j\| \leq C_0$ and $\|Q_j\| \leq 1 + C_0$.*
2. *$\sup\{\|M_{Q_{j_n}}M_{Q_{j_{n-1}}}\cdots M_{Q_{j_1}}\|, n \in \mathbb{N}^*, j_k \in \mathbb{N}^*\} \leq C_0(1 + C_0)$.*
3. *For any $n \in \mathbb{N}^*$, $\|\theta_n\| \leq C_0^2$.*

Proof.

1. Let $j \in \mathbb{N}^*$. By means of Von Neumann's ergodic theorem, we have

$$P_{1,j} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=0}^{N-1} M_j^k.$$

Hence, we get from Lemma 2.3 that $\|P_{1,j}\| \leq C_0$. Since $\|\psi_S\| = 1$ we also have $\|\psi_j\| \leq C_0$, and hence $\|P_j\| = \|\psi_j\| \leq C_0$ and $\|Q_j\| \leq 1 + C_0$.

2. Using (2.9), we have $M_{Q_{j_n}}M_{Q_{j_{n-1}}}\cdots M_{Q_{j_1}} = Q_{j_n}M_{j_n}M_{j_{n-1}}\cdots M_{j_1}$, so that

$$\|M_{Q_{j_n}}M_{Q_{j_{n-1}}}\cdots M_{Q_{j_1}}\| \leq \|Q_{j_n}\|C_0 \leq C_0(1 + C_0).$$

3. From (2.8) and the above estimates, we obtain $\|\theta_n\| \leq C_0\|\psi_1\| \leq C_0^2$. \square

2.3. Asymptotic behaviour

We now turn to the study of the asymptotic behaviour of Ψ_n . If $M_n \equiv M \in \mathcal{M}(E)$ (recall the definition of $\mathcal{M}(E)$ given before Theorem 1.1), then M_Q^n converges to zero, and Ψ_n converges to the rank one projection $P = |\psi_S\rangle\langle\psi|$ (exponentially fast). The following result shows that in the general case, Ψ_n converges if and only if ψ_n does.

Proposition 2.5. *Suppose that $\lim_{n \rightarrow \infty} \sup\{\|M_{Q_{j_n}}\cdots M_{Q_{j_1}}\|, j_k \in \mathbb{N}^*\} = 0$. Then θ_n converges if and only if ψ_n does. If they exist, these two limits coincide, and thus*

$$\lim_{n \rightarrow \infty} \Psi_n = |\psi_S\rangle\langle\psi_\infty|,$$

where $\psi_\infty = \lim_{n \rightarrow \infty} \psi_n$. Moreover, $|\psi_S\rangle\langle\psi_\infty|$ is a projection.

In general thus, we cannot expect to obtain pointwise convergence of the θ_n , but we have to consider the ergodic average of $\theta_n(\bar{\omega})$ in Theorem 1.2. This is natural in view of the interpretation of our results in terms of dynamical systems (a fluctuating system does not converge). For instance, if, in the random setting, $M(\omega) = P(\omega) = |\psi_S\rangle\langle\psi(\omega)|$, then $\Psi_n(\bar{\omega}) = P(\omega_n)$ converges iff $\psi(\omega_n)$ does (this is actually a particular case of Proposition 2.5). For a general

random vector $\psi(\omega)$, there is no reason to expect a limit, due to the fluctuations of the $\psi(\omega_j)$. Nevertheless, Birkhoff's ergodic theorem asserts that the ergodic limit exists almost surely [1], and that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N P(\omega_j) = \mathbb{E}(P) = |\psi_S\rangle \langle \mathbb{E}(\psi) | = |\psi_S\rangle \langle \bar{\psi} |, \quad \text{a.s.}, \quad (2.10)$$

where the projection $|\psi_S\rangle \langle \bar{\psi} |$ is constant in ω . Theorem 1.2 gives an analogous result in the general case.

Proof of Proposition 2.5. Suppose $\lim \theta_n = \theta_\infty$ exists, and let $\phi_n := \theta_n - \theta_\infty$. We have $M_n^* \theta_\infty = M_n^* \theta_{n-1} - M_n^* \phi_{n-1} = \theta_n - M_n^* \phi_{n-1} = \theta_\infty + \phi_n - M_n^* \phi_{n-1}$. Hence, for any n , $M_{Q_n}^* \theta_\infty = Q_n^* \theta_\infty + Q_n^* \phi_n - M_{Q_n}^* \phi_{n-1} =: Q_n^* \theta_\infty + \chi_n$, where, by Proposition 2.4, $\lim_{n \rightarrow \infty} \chi_n = 0$. For any $p \in \mathbb{N}$, we can write $(M_{Q_n}^*)^p \theta_\infty = (M_{Q_n}^*)^{p-1} \theta_\infty + (M_{Q_n}^*)^{p-1} \chi_n$, and thus we obtain, by induction, $(M_{Q_n}^*)^p \theta_\infty = Q_n^* \theta_\infty + \sum_{k=0}^{p-1} (M_{Q_n}^*)^k \chi_n$. By assumption, given any $\varepsilon > 0$, we can chose p_0 large enough so that, for any n , $\|(M_{Q_n}^*)^{p_0}\| \leq \varepsilon$. Using Proposition 2.4, we thus get that for any n ,

$$\|Q_n^* \theta_\infty\| \leq \varepsilon \|\theta_\infty\| + p_0 C_0 (C_0 + 1) \|\chi_n\|.$$

Since χ_n goes to zero and ε is arbitrary, this proves that $\lim_{n \rightarrow \infty} Q_n^* \theta_\infty = 0$. Now, $\langle \psi_S, \theta_\infty \rangle = 1$ because $\langle \psi_S, \theta_n \rangle = 1$ for all n . Thus we have $\psi_n = P_n^* \theta_\infty = \theta_\infty - Q_n^* \theta_\infty$. Therefore, $\lim \psi_n$ exists and equals θ_∞ .

Conversely, suppose $\lim \psi_n = \psi_\infty$ exists. Since $M_n^* \psi_n = \psi_n$ for all n , we have

$$M_n^* \psi_\infty = M_n^* (\psi_\infty - \psi_n) + \psi_n.$$

Combining this with Lemma 2.3 shows that $\lim M_n^* \psi_\infty = \psi_\infty$. Furthermore, we have for any $n, p \in \mathbb{N}$,

$$\begin{aligned} \theta_{n+p} &= \sum_{j=1}^{n+p} M_{Q_{n+p}}^* \cdots M_{Q_{j+1}}^* \psi_j \\ &= M_{Q_{n+p}}^* \cdots M_{Q_{n+1}}^* \sum_{j=1}^n M_{Q_n}^* \cdots M_{Q_{j+1}}^* \psi_j + \sum_{j=n+1}^{n+p} M_{Q_{n+p}}^* \cdots M_{Q_{j+1}}^* \psi_j \\ &= M_{Q_{n+p}}^* \cdots M_{Q_{n+1}}^* \theta_n + M_{n+p}^* \cdots M_{n+1}^* \psi_{n+1}, \end{aligned}$$

where we used Lemma 2.2 to obtain the last line. It follows that

$$\begin{aligned} \|\theta_{n+p} - \psi_\infty\| &\leq \|M_{Q_{n+p}}^* \cdots M_{Q_{n+1}}^*\| \|\theta_n\| + \|M_{n+p}^* \cdots M_{n+1}^* (\psi_{n+1} - \psi_\infty)\| \\ &\quad + \|M_{n+p}^* \cdots M_{n+1}^* \psi_\infty - \psi_\infty\|. \end{aligned}$$

Let $\varepsilon > 0$. By assumption, there exists p_0 such that for any n , $\|M_{Q_{n+p_0}}^* \cdots M_{Q_{n+1}}^*\| < \varepsilon$. Thus we have, for all n ,

$$\begin{aligned} \|\theta_{n+p_0} - \psi_\infty\| &\leq \varepsilon \|\theta_n\| + C_0 \|\psi_{n+1} - \psi_\infty\| \\ &\quad + \left\| M_{n+p_0}^* \psi_\infty - \psi_\infty + \sum_{j=n+1}^{n+p_0-1} M_{n+p_0}^* \cdots M_{j+1}^* (M_j^* \psi_\infty - \psi_\infty) \right\| \\ &\leq \varepsilon C_0^2 + C_0 \|\psi_{n+1} - \psi_\infty\| + \|M_{n+p_0}^* \psi_\infty - \psi_\infty\| \\ &\quad + C_0 (p_0 - 1) \sup_{n+1 \leq j \leq n+p_0-1} \|M_j^* \psi_\infty - \psi_\infty\|. \end{aligned}$$

Since $\lim M_n^* \psi_\infty = \psi_\infty$, and since ε is arbitrary, we obtain $\lim \theta_n = \psi_\infty$. To prove that $|\psi_S\rangle \langle \psi_\infty |$ is a projection, it suffices to note that $\langle \psi_n, \psi_S \rangle = 1$ for all n . \square

The previous convergence result relies on the (exponential) decay of the product of operators M_{Q_j} . One can show that such a decay holds, in a deterministic situation, provided there exists $\delta > 0$ such that $\|M_{Q_j}\| \leq e^{-\delta}$, $\forall j \in \mathbb{N}$. One may obtain this bound for specific models (e.g. in the model of Section 1.3.4, if the sequence $(\tau_n)_n$ of interaction times is deterministic and satisfies $d(\tau_n, T\mathbb{Z}) > \varepsilon$ for some $\varepsilon > 0$ and where T is defined as in Theorem 1.6). To satisfy this bound, it is not only required that the M_j belong to $\mathcal{M}_{(E)}$ for all j , but also that:

1. we have some uniformity in the spectrum of the M_{Q_j} , i.e. there exists a $\gamma > 0$, such that

$$\sigma(M_j) \setminus \{1\} \subset \{z \mid |z| \leq e^{-\gamma}\} \quad \forall j \in \mathbb{N}, \quad (2.11)$$

2. we can prove that the corresponding spectral projectors do not behave too badly so that we can still bound M_{Q_j} efficiently, i.e. $\|Q_j\|e^{-\gamma} \leq e^{-\delta}$, for all j .

Such conditions look quite stringent. However, Theorem 1.1 shows that in the random setting, a similar exponential decay holds under much weaker assumptions.

3. Proofs of Theorems 1.1 and 1.2

3.1. The probabilistic setting

Let $(\Omega, \mathcal{F}, \mathbb{p})$ be a probability space underlying the random process. Given a random variable $f(\omega)$, we denote by $\bar{f} := \mathbb{E}(f) = \int f(\omega) d\mathbb{p}(\omega)$ its expectation. We define the probability measure $d\mathbb{P}$ on $\Omega_{\text{ext}} := \Omega^{\mathbb{N}^*}$ in a standard fashion by

$$d\mathbb{P} = \prod_{j \geq 1} dp_j, \quad \text{where } dp_j \equiv dp, \quad \forall j \in \mathbb{N}^*.$$

We denote points in Ω_{ext} by $\bar{\omega}$. For any $r \in \mathbb{N}^*$, and for any collection of measurable sets in Ω , $\{A_j\}_{1 \leq j \leq r}$, we have

$$\mathbb{P}((\omega_1, \omega_2, \dots, \omega_r) \in (A_1, A_2, \dots, A_r)) = \prod_{1 \leq j \leq r} \int_{A_j} dp(\omega_j).$$

We also define the shift $T : \Omega_{\text{ext}} \rightarrow \Omega_{\text{ext}}$ by

$$(T\bar{\omega})_j = \omega_{j+1} \quad \forall \bar{\omega} = (\omega_j)_{j \in \mathbb{N}^*} \in \Omega_{\text{ext}}. \quad (3.1)$$

T is an ergodic transformation of Ω_{ext} . Further, we assume that the distribution of the matrices M is characterized by a measurable map

$$\Omega \ni \omega_1 \mapsto M(\omega_1) \in M_d(\mathbb{C}). \quad (3.2)$$

We write sometimes $M(\omega)$ instead of $M(\omega_1)$. Hence, for any set $B \in M_d(\mathbb{C})$,

$$p(M(\omega) \in B) = p(M^{-1}(B)) = \int_{M^{-1}(B)} dp(\omega),$$

and similarly for other random variables. With these notations we have for the i.i.d. process

$$\begin{aligned} \Psi_n(\bar{\omega}) &:= M(\omega_1)M(\omega_2) \cdots M(\omega_n) \\ &= M(T^0\omega)M(T^1\omega) \cdots M(T^{n-1}\omega), \end{aligned} \quad (3.3)$$

where $\bar{\omega} \in \Omega^{\mathbb{N}^*}$.

In the same way as in (2.2), we introduce the random variable $\psi(\omega_1) \in \mathbb{C}^d$ defined as

$$\psi(\omega_1) := P_1(\omega_1)^* \psi_S, \quad (3.4)$$

where $P_1(\omega_1)$ denotes the spectral projection of $M(\omega_1)$ for the eigenvalue 1, and we decompose

$$M(\omega_1) := |\psi_S\rangle\langle\psi(\omega_1)| + M_Q(\omega_1) = P(\omega_1) + M_Q(\omega_1) \tag{3.5}$$

as in (2.3). Note that $\psi(\omega_1)$ and $M_Q(\omega_1)$ define *bona fide* random variables: $\omega_1 \mapsto P_1(\omega_1)$ is measurable since $\omega_1 \mapsto M(\omega_1)$ is [3]. Consider finally the process

$$\begin{aligned} \theta_n(\bar{\omega}) &= M^*(T^{n-1}\omega)M^*(T^{n-2}\omega)\cdots M^*(T\omega)\psi(\omega) \\ &= \sum_{j=1}^n M_Q^*(\omega_n)M_Q^*(\omega_{n-1})\cdots M_Q^*(\omega_{j+1})\psi(\omega_j). \end{aligned} \tag{3.6}$$

Note that θ_n is a Markov process, since $\theta_{n+1}(\bar{\omega}) = M_{n+1}^*(\omega_{n+1})\theta_n(\bar{\omega})$.

3.2. Decay estimates

In a first step, we derive a bound similar to the one in (1.5), but only for some “good ω .” Then we will prove (1.5) via the Borel–Cantelli lemma.

Lemma 3.1. *Suppose $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then there exists an $M_0 \in \mathcal{M}_{(E)}$ such that for any $\varepsilon > 0$, $p(\|M(\omega) - M_0\| < \varepsilon) > 0$.*

Proof. Let M be such that $\|M\| \leq 1$ and $M\psi_S = \psi_S$. We decompose M as in (2.3), and then $M \in \mathcal{M}_{(E)}$ if and only if $\sigma(M_Q) \subset \{z \in \mathbb{C} \text{ s.t. } |z| < 1\}$. Note that it is equivalent to $\text{sr}(M_Q) < 1$, where sr denotes the spectral radius. For $n \in \mathbb{N}$, set

$$\mathcal{M}_{(n)} := \{M \in M_d(\mathbb{C}) \text{ s.t. } \|M\| \leq 1, M\psi_S = \psi_S, \sigma(M_Q) \subset \{z \in \mathbb{C} \text{ s.t. } |z| \leq 1 - 1/n\}\}.$$

The sets $\mathcal{M}_{(n)} \subset M_d(\mathbb{C})$ are compact, and we have $\mathcal{M}_{(E)} = \bigcup_n \mathcal{M}_{(n)}$. Thus, since $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$, there exists n_0 such that $p(M(\omega) \in \mathcal{M}_{(n_0)}) > 0$. The lemma then follows with $M_0 \in \mathcal{M}_{(n_0)}$ by the following standard argument using the compactness of $\mathcal{M}_{(n_0)}$.

We construct a sequence $M_k \in \mathcal{M}_{(n_0)}$ as follows. For any $k \in \mathbb{N}^*$ we cover $\mathcal{M}_{(n_0)}$ with a finite number of balls with centres in $\mathcal{M}_{(n_0)}$ and radius $1/k$. For each fixed k , since $p(M(\omega) \in \mathcal{M}_{(n_0)}) > 0$, there is at least one of these balls which has a non-zero probability. Pick one of these balls and denote its centre by M_k . Thus, $\forall k \in \mathbb{N}^*$, $p(\|M(\omega) - M_k\| \leq \frac{1}{k}) > 0$. Since $\mathcal{M}_{(n_0)}$ is compact, the sequence M_k converges to some $M_0 \in \mathcal{M}_{(n_0)}$ (up to taking a subsequence). Now, for any given $\varepsilon > 0$, take k large enough so that $1/k < \varepsilon/2$ and $\|M_k - M_0\| < \varepsilon/2$. Then

$$p(\|M(\omega) - M_0\| < \varepsilon) \geq p(\|M(\omega) - M_k\| < \varepsilon/2) > 0. \quad \square$$

Lemma 3.2. *Suppose $p(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Then, there exists $\tilde{\Omega}_1 \subset \Omega$ such that $p(\tilde{\Omega}_1) > 0$, and there exist constants $\delta > 0$ and $C > 0$ such that $\forall k \in \mathbb{N}^*$,*

$$(\omega_1, \omega_2, \dots, \omega_k) \in \tilde{\Omega}_1^k \implies \|M_Q(\omega_1)M_Q(\omega_2)\cdots M_Q(\omega_k)\| \leq Ce^{-\delta k}.$$

Proof. Let M_0 be as in Lemma 3.1 and decompose it as $M_0 = |\psi_S\rangle\langle\psi_0| + M_{Q_0}$. Using the Riesz representation,

$$P_1 = \frac{-1}{2\pi i} \int_{\Gamma} (M - z)^{-1} dz$$

for a small contour Γ encircling the eigenvalue one, one gets that the map $M \mapsto M_Q$ is continuous on $\{M \in M_d(\mathbb{C}) \text{ s.t. } \|M - M_0\| < \eta\}$, for η small enough. Thus Lemma 3.1 shows that $p(\|M_Q(\omega) - M_{Q_0}\| < \varepsilon) > 0$.

Hence, for any $\varepsilon > 0$, there exists $\Omega_\varepsilon \subset \Omega$ such that $p(\Omega_\varepsilon) > 0$, and such that for any $\omega_j \in \Omega_\varepsilon$, we can write

$$M_Q(\omega_j) = M_{Q_0} + \Delta(\omega_j) =: M_{Q_0} + \Delta_j$$

with $\|\Delta_j\| < \varepsilon$. With these notations, we have

$$\begin{aligned} & M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_k) \\ &= \sum_{j=0}^k \sum_{1 \leq k_1 < k_2 < \cdots < k_j \leq k} M_{Q_0}^{k_1-1} \Delta_{k_1} M_{Q_0}^{k_2-k_1-1} \Delta_{k_2} \cdots M_{Q_0}^{k_j-k_{j-1}-1} \Delta_{k_j} M_{Q_0}^{k-k_j}. \end{aligned} \quad (3.7)$$

Since $M_0 \in \mathcal{M}(E)$, we know that there exist $\gamma > 0$ and $C > 0$ such that $\|M_{Q_0}^n\| \leq Ce^{-\gamma n}$, for any n . Inserting this estimate in (3.7), we get for any k and $(\omega_1, \dots, \omega_k) \in \Omega_\varepsilon^k$,

$$\begin{aligned} \|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_k)\| &\leq \sum_{j=0}^k \sum_{1 \leq k_1 < k_2 < \cdots < k_j \leq k} C^{j+1} e^{-\gamma(k-j)} \varepsilon^j \\ &= Ce^{-\gamma k} \sum_{j=0}^k \binom{k}{j} (Ce^\gamma \varepsilon)^j = Ce^{-\gamma k} (1 + Ce^\gamma \varepsilon)^k. \end{aligned}$$

We now choose ε small enough so that $e^{-\gamma}(1 + Ce^\gamma \varepsilon) = e^{-\delta}$ for some $\delta > 0$, and the proof is complete. \square

3.3. Proof of Theorem 1.1

Let $\tilde{\Omega}_1$ be as in Lemma 3.2, and consider a product of L matrices

$$M_Q(\omega_{j+1})M_Q(\omega_{j+2}) \cdots M_Q(\omega_{j+L}).$$

The L matrices belong to the set $M_Q(\tilde{\Omega}_1)$ with positive probability, and by Lemma 3.2, we have $\|M_Q(\omega_{j+1})M_Q(\omega_{j+2}) \cdots M_Q(\omega_{j+L})\| \leq Ce^{-\delta L}$. In other words, for any $j \in \mathbb{N}$,

$$\mathbb{P}(\|M_Q(\omega_{j+1})M_Q(\omega_{j+2}) \cdots M_Q(\omega_{j+L})\| \leq Ce^{-\delta L}) =: \mathcal{P} \geq p(\tilde{\Omega}_1)^L > 0. \quad (3.8)$$

We consider now a sequence of nL matrices and estimate the probability that

$$\|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_{Ln})\| \leq (Ce^{-\delta L})^m C^{m+1}. \quad (3.9)$$

We decompose the above product into n blocks of length L . Estimate (3.9) holds in particular if among these n blocks, there are at least m blocks in which (3.8) holds. Indeed, let $j_1, \dots, j_m \in \{0, \dots, n-1\}$ be such that

$$\|M_Q(\omega_{j_k L+1})M_Q(\omega_{j_k L+2}) \cdots M_Q(\omega_{j_k L+L})\| \leq Ce^{-\delta L}, \quad k = 1, \dots, m. \quad (3.10)$$

We then isolate the m ‘‘good blocks’’ and write

$$\begin{aligned} & \|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_{Ln})\| \\ &\leq \prod_{k=1}^m \|M_Q(\omega_{j_k L+1}) \cdots M_Q(\omega_{j_k L+L})\| \|M_Q(\omega_1) \cdots M_Q(\omega_{j_1 L})\| \\ &\quad \times \prod_{k=1}^{m-1} \|M_Q(\omega_{j_k L+L+1}) \cdots M_Q(\omega_{j_{k+1} L})\| \|M_Q(\omega_{j_m L+L+1}) \cdots M_Q(\omega_{Ln})\|. \end{aligned}$$

Using Proposition 2.4 and (3.10), we obtain (3.9) with $C \geq C_0(1 + C_0)$. Hence,

$$\mathbb{P}(\|M_Q(\omega_1) \cdots M_Q(\omega_{Ln})\| \leq (Ce^{-\delta L})^m C^{m+1}) \geq \sum_{j=m}^n \binom{n}{j} \mathcal{P}^j (1 - \mathcal{P})^{n-j}. \quad (3.11)$$

We now choose $m = \lfloor \beta n \rfloor$, for some $\beta > 0$ to be chosen later, and where $\lfloor \cdot \rfloor$ denotes the integer part. For L large enough we have the following bound, uniform in β ,

$$(C e^{-\delta L})^{\lfloor \beta n \rfloor} C^{\lfloor \beta n \rfloor + 1} \leq C e^{\delta L} (C^{2\beta} e^{-\delta \beta L})^n = \tilde{C} e^{-\gamma \beta n} \tag{3.12}$$

for some $\gamma > 0$. For $0 < \beta < \mathcal{P}$, the following classical tail estimate follows from Hoeffding’s inequality [13]

$$\begin{aligned} \sum_{j=\lfloor \beta n \rfloor}^n \binom{n}{j} \mathcal{P}^j (1 - \mathcal{P})^{n-j} &= 1 - \sum_{j=0}^{\lfloor \beta n \rfloor - 1} \binom{n}{j} \mathcal{P}^j (1 - \mathcal{P})^{n-j} \\ &\geq 1 - e^{-2(n\mathcal{P} - \lfloor \beta n \rfloor)^2/n} \geq 1 - e^{-\sigma n} \end{aligned} \tag{3.13}$$

for $\sigma = 2(\mathcal{P} - \beta)^2$. Let us denote

$$G_n := \{ \bar{\omega} = \{\omega_j\}_{j \geq 1} \text{ s.t. } \|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_{Ln})\| \leq \tilde{C} e^{-\gamma \beta n} \}.$$

Then, (3.11)–(3.13) show that, for all n , we have $\mathbb{P}(G_n) \geq 1 - e^{-\sigma n}$. Let G_n^c be the complement of G_n in Ω_{ext} . The first Borel–Cantelli lemma asserts that

$$\sum_{n \geq 0} \mathbb{P}(G_n^c) < \infty \implies \mathbb{P}\left(\bigcap_n \bigcup_{m \geq n} G_m^c\right) = 0.$$

Since

$$\mathbb{P}(G_n^c) \leq e^{-\sigma n}, \tag{3.14}$$

we get that G_n^c occurs infinitely often with probability zero. Therefore, there exists $\Omega_1 \subset \Omega_{\text{ext}}$, of measure one, such that for any $\bar{\omega} \in \Omega_1$, there exists $n(\bar{\omega})$ s.t.

$$\|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_{Ln})\| \leq \tilde{C} e^{-\alpha Ln} \quad \text{if } n \geq n(\bar{\omega}), \tag{3.15}$$

where $\alpha = \gamma \beta / L$. The result of Theorem 1.1 now follows: fix L large enough and write $n = mL + p$ (with $0 \leq p < L$). We conclude from Proposition 2.4 and (3.15) that if $\bar{\omega} \in \Omega_1$ (of measure one) and if $n \geq n_0(\bar{\omega}) := Ln(\bar{\omega}) + L$ (so that $m \geq n(\bar{\omega})$), then

$$\|M_Q(\omega_1) \cdots M_Q(\omega_n)\| \leq C_0 \tilde{C} e^{-\alpha Lm} \leq C_0 \tilde{C} e^{\alpha p} e^{-\alpha n} \leq C_0 \tilde{C} e^{\alpha L} e^{-\alpha n}.$$

This completes the proof of Theorem 1.1.

Remarks.

1. We can take $n_0(\bar{\omega}) := \min\{m \geq 0 \mid \forall k \geq m, \|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_k)\| \leq \tilde{C} e^{-\alpha k}\}$, so that, defining $g_n = \{\bar{\omega} \text{ s.t. } \|M_Q(\omega_1)M_Q(\omega_2) \cdots M_Q(\omega_n)\| \leq \tilde{C} e^{-\alpha n}\}$, we have $\{\bar{\omega} \mid n_0(\bar{\omega}) = m\} = \bigcap_{k \geq m} g_k \cup g_{m-1}^c$. Hence, by the estimate (3.14) and our definition of α, β and σ , we get

$$\mathbb{E}(e^{\alpha n_0}) \leq \sum_{m \geq 0} e^{\alpha m} \mathbb{P}(g_{m-1}^c) \leq e^{\sigma/L} \sum_{m \geq 0} e^{(\gamma \beta - \sigma)m/L} = e^{\sigma/L} \sum_{m \geq 0} e^{(\gamma \beta - 2(\mathcal{P} - \beta)^2)m/L} < \infty \tag{3.16}$$

for β small enough.

2. By Proposition 2.4, (1.5) holds for all $\bar{\omega} \in \Omega_1$ and for all $n \in \mathbb{N}^*$, up to replacing \tilde{C} by $C_{\bar{\omega}} = \tilde{C} e^{\alpha n_0(\bar{\omega})}$.

As we mentioned in the Introduction, if one can chose $\psi(\omega) \equiv \psi$ one may expect a convergence without any ergodic averaging. Indeed, the following result is a direct consequence of (2.8) and Theorem 1.1.

Proposition 3.3. *Let $M(\omega)$ be a random reduced dynamics operator. Suppose that $\mathbb{p}(M(\omega) \in \mathcal{M}_{(E)}) > 0$ and there exists $\psi \in \mathbb{C}^d$ such that $M(\omega)^*\psi = \psi$ for all ω . Then there exists a set $\Omega_1 \subset \Omega^{\mathbb{N}^*}$, and a constant $\alpha > 0$, s.t. $\mathbb{P}(\Omega_1) = 1$ and s.t. for any $\bar{\omega} \in \Omega_1$, there exists $C_{\bar{\omega}}$ so that for any $n \in \mathbb{N}^*$,*

$$\left\| M_1(\omega) \cdots M_n(\omega) - \frac{|\psi_S\rangle\langle\psi|}{\langle\psi, \psi_S\rangle} \right\| \leq C_{\bar{\omega}} e^{-\alpha n}.$$

This result applies in particular to bistochastic matrices with $\psi_S = \psi = \frac{1}{\sqrt{d}}(1, \dots, 1)$. Infinite product of doubly stochastic matrices have been considered in e.g. [21], where the author looks for sufficient conditions which ensure that such a product converges to the rank one projection $|\psi_S\rangle\langle\psi_S|$. However, only deterministic products are considered.

3.4. Proof of Theorem 1.2

The proof of Theorem 1.2 relies on the estimate (1.5), but not directly on the more stringent assumption $\mathbb{p}(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Consider the condition

(RE) There exist $\tilde{\Omega} \subset \Omega^{\mathbb{N}^*}$, $C, \alpha > 0$ such that $\mathbb{P}(\tilde{\Omega}) = 1$, and such that for any $\bar{\omega} \in \tilde{\Omega}$, there exists $n_0(\bar{\omega}) \in \mathbb{N}$ so that $\|M_Q(\omega_1) \cdots M_Q(\omega_n)\| \leq C e^{-\alpha n}$ if $n \geq n_0(\bar{\omega})$, and, $\|M_Q(\omega_1) \cdots M_Q(\omega_n)\| \leq C$, otherwise.

Theorem 1.1 and Proposition 2.4 show that (RE) is a consequence of $\mathbb{p}(M(\omega) \in \mathcal{M}_{(E)}) > 0$. Consequently, Theorem 1.2 follows from Theorem 3.5 below. We denote by $P_{1, \bar{M}}$ the spectral projection of $\bar{M} := \mathbb{E}(M(\omega))$ for the eigenvalue 1.

Remark. *Note that $M(\omega)\psi_S = \psi_S$ for all ω implies $\bar{M}\psi_S = \psi_S$. In general, it may happen that 1 is a degenerate eigenvalue for \bar{M} even though it is non-degenerate for any $M(\omega)$. This will however not be the case here as the following lemma shows.*

Lemma 3.4. *Suppose assumption (RE) holds, then $\bar{M} \in \mathcal{M}_{(E)}$. Moreover, we have $P_{1, \bar{M}} = |\psi_S\rangle\langle\theta|$, where θ satisfies*

$$\theta = (\mathbb{1} - \bar{M}_Q^*)^{-1} \bar{\psi} = \sum_{k=0}^{\infty} \bar{M}_Q^{*k} \bar{\psi} = P_{1, \bar{M}}^* \bar{\psi} = P_{1, \bar{M}}^* \psi_S. \tag{3.17}$$

Proof. We have $\bar{M} = |\psi_S\rangle\langle\bar{\psi}| + \bar{M}_Q$. Since $M(\omega)\psi_S = \psi_S$ and $\langle\psi(\omega)|\psi_S\rangle = 1$ for any ω , we get $\bar{M}_Q\psi_S = 0$ and $\langle\bar{\psi}|\psi_S\rangle = 1$. Thus, for any $n \in \mathbb{N}$ we have (see also (2.6) and (2.7))

$$\bar{M}^n = |\psi_S\rangle\langle\bar{\psi} + \bar{M}_Q^* \bar{\psi} + \cdots + \bar{M}_Q^{*n-1} \bar{\psi}| + \bar{M}_Q^n. \tag{3.18}$$

By independence, we have, for any $k \in \mathbb{N}$, $\bar{M}_Q^k = \int d\mathbb{P} M_Q(\omega_1) \cdots M_Q(\omega_k)$ so that

$$\|\bar{M}_Q^k\| \leq \int d\mathbb{P} \|M_Q(\omega_1) \cdots M_Q(\omega_k)\|. \tag{3.19}$$

Due to assumption (RE) and considerations similar to (3.16), we can bound the right-hand side by

$$\mathbb{P}\{n_0(\bar{\omega}) > k\} C + \mathbb{P}\{n_0(\bar{\omega}) \leq k\} C e^{-\alpha k} \leq \bar{C} e^{-\alpha k} \tag{3.20}$$

for some constant \bar{C} . Hence we have

$$\|\bar{M}_Q^k\| \leq C e^{-\alpha k} \quad \forall k \in \mathbb{N}. \tag{3.21}$$

In particular, the spectrum of \bar{M}_Q lies inside the open complex unit disk.

This estimate and (3.18) prove $\lim_{n \rightarrow \infty} \overline{M}^n = |\psi_S\rangle\langle\theta|$ where

$$\theta = \sum_{k=0}^{\infty} \overline{M}_Q^{*k} \overline{\psi} = (\mathbb{1} - \overline{M}_Q^*)^{-1} \overline{\psi}.$$

Hence $\overline{M} \in \mathcal{M}(E)$ and $P_{1, \overline{M}} = |\psi_S\rangle\langle\theta|$. The last identities in (3.17) are immediate since $\langle\psi_S, \overline{\psi}\rangle = \langle\psi_S, \psi_S\rangle = 1$. \square

Remarks.

1. In general $\overline{\psi}$ is not an eigenvector of \overline{M}^* , and $\theta \neq \overline{\psi}$; see Theorem 1.6 and the remark thereafter.
2. Without assumption (RE), instead of (3.21), we can only prove that $\|\overline{M}_Q^k\| \leq C_0(1 + C_0)$ and that $\|\overline{M}^k\| \leq C_0$, $\forall k \in \mathbb{N}$, which is not enough to exclude that $1 \in \sigma(\overline{M}_Q)$.

Theorem 3.5. Suppose assumption (RE) holds. Then, there exists a set $\Omega_0 \subset \Omega^{\mathbb{N}^*}$ such that $\mathbb{P}(\Omega_0) = 1$ and, for all $\overline{\omega} \in \Omega_0$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \theta_n(\overline{\omega}) = \theta.$$

Proof. Recall that T denotes the shift map on Ω_{ext} (see (3.1)). Using (2.7), we get

$$\begin{aligned} \sum_{n=1}^N \theta_n(\overline{\omega}) &= \sum_{n=1}^N \sum_{j=0}^{n-1} M_Q^*(T^{n-1}\omega) \cdots M_Q^*(T^{j+1}\omega) \psi(T^j\omega) \\ &= \sum_{k=1}^N \sum_{j=0}^{N-k} M_Q^*(T^{k+j-1}\omega) \cdots M_Q^*(T^{j+1}\omega) \psi(T^j\omega). \end{aligned} \tag{3.22}$$

Let us introduce random vectors $\theta^{(k)}$, $k = 1, 2, \dots$, by

$$\begin{aligned} \theta^{(k)}(\overline{\omega}) &= \theta^{(k)}(\omega_1, \dots, \omega_k) \\ &= M_Q^*(T^{k-1}\omega) M_Q^*(T^{k-2}\omega) \cdots M_Q^*(T\omega) \psi(T^0\omega) \\ &= M_Q^*(\omega_k) \theta^{(k-1)}(\omega_1, \dots, \omega_{k-1}), \end{aligned} \tag{3.23}$$

so that, by (3.22),

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^N \theta_n(\overline{\omega}) &= \sum_{k=1}^N \frac{1}{N} \sum_{j=0}^{N-k} \theta^{(k)}(T^j\overline{\omega}) \\ &= \sum_{k=1}^{\infty} \chi_{\{k \leq N\}} \sum_{j=0}^{N-k} \theta^{(k)}(T^j\overline{\omega}) \frac{1}{N} =: \sum_{k=1}^{\infty} g(k, N, \overline{\omega}). \end{aligned}$$

For each fixed k , by ergodicity, there exists a set $\Omega_{(k)} \subset \Omega^{\mathbb{N}^*}$ of probability one, such that, for all $\omega \in \Omega_{(k)}$, the following limit exists

$$\begin{aligned} \lim_{N \rightarrow \infty} g(k, N, \overline{\omega}) &= \lim_{N \rightarrow \infty} \frac{1}{N-k+1} \sum_{j=0}^{N-k} \theta^{(k)}(T^j\overline{\omega}) \frac{N-k+1}{N} \\ &= \lim_{M \rightarrow \infty} \frac{1}{M+1} \sum_{j=0}^M \theta^{(k)}(T^j\overline{\omega}) = \mathbb{E}(\theta^{(k)}). \end{aligned}$$

Therefore, on the set $\Omega_\infty := \bigcap_{k \in \mathbb{N}} \Omega(k)$ of probability one, for any $k \in \mathbb{N}$, we have by independence of the $M(\omega_j)$,

$$\lim_{N \rightarrow \infty} g(k, N, \bar{\omega}) = \mathbb{E}(\theta^{(k)}) = (\overline{M_Q^*})^{k-1} \overline{\psi}. \quad (3.24)$$

Also, by assumption (RE) and Proposition 2.4, on $\Omega_0 = \tilde{\Omega} \cap \Omega_\infty$, a set of probability 1, we have the bound $\|\theta^{(k)}(T^j \bar{\omega})\| \leq C_0 C e^{\alpha n_0(T^j \bar{\omega})} e^{-\alpha(k-1)}$. Hence, for all N large enough, and for all $1 \leq k \leq N$, by ergodicity

$$\|g(k, N, \bar{\omega})\| \leq C_0 C \frac{1}{N} \sum_{j=0}^{N-1} e^{\alpha n_0(T^j \bar{\omega})} e^{-\alpha(k-1)} \leq 2C_0 C \mathbb{E}(e^{\alpha n_0}) e^{-\alpha(k-1)},$$

which is finite by (3.16). For $k > N$, $\|g(k, N, \bar{\omega})\| = 0 \leq 2C_0 C \mathbb{E}(e^{\alpha n_0}) e^{-\alpha(k-1)}$. Since $e^{-\alpha(k-1)} \in l^1(\mathbb{N})$, we can apply the Lebesgue dominated convergence theorem in (3.24) to conclude that $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \theta_n(\bar{\omega}) = \theta$ a.s. (recall Lemma 3.4). \square

3.5. The mean ergodic theorem and Lyapunov exponents

We present in this section some results for products “in reverse order” of the form $\Phi_n(\bar{\omega}) := M(\omega_n) \cdots M(\omega_1)$, as well as results related to the multiplicative ergodic theorem. The following results are standard, see e.g. [1], Theorem 3.4.1 and Remark 3.4.10(ii). The limits

$$\Lambda_\Phi(\bar{\omega}) = \lim_{n \rightarrow \infty} (\Phi_n(\bar{\omega})^* \Phi_n(\bar{\omega}))^{1/2n} \quad \text{and} \quad \Lambda_\Psi(\bar{\omega}) = \lim_{n \rightarrow \infty} (\Psi_n(\bar{\omega})^* \Psi_n(\bar{\omega}))^{1/2n}$$

exist almost surely, the top Lyapunov exponent of $\Lambda_\Phi(\bar{\omega})$ and of $\Lambda_\Psi(\bar{\omega})$ are the same, called $\gamma_1(\bar{\omega})$; $\gamma_1(\bar{\omega})$ is constant a.s., and so is its multiplicity. It is usually difficult to prove that the multiplicity of $\gamma_1(\bar{\omega})$ is 1. In the present case however, we can get this result.

Theorem 3.6. *Suppose $p(M(\omega) \in \mathcal{M}(E)) > 0$. Then there exist $\alpha > 0$, a random vector $\eta_\infty(\bar{\omega})$ and $\Omega_0 \subset \Omega^{\mathbb{N}^*}$ with $\mathbb{P}(\Omega_0) = 1$ such that for any $\bar{\omega} \in \Omega_0$ and $n \in \mathbb{N}^*$*

$$\|\Phi_n(\bar{\omega}) - |\psi_S\rangle \langle \eta_\infty(\bar{\omega})|\| \leq C_{\bar{\omega}} e^{-\alpha n} \quad \text{for some } C_{\bar{\omega}}. \quad (3.25)$$

As a consequence, for any $\bar{\omega} \in \Omega_0$, $\gamma_1(\bar{\omega})$ is of multiplicity one.

Proof. We first decompose $\Phi_n(\bar{\omega})$ in the same way as $\Psi_n(\bar{\omega})$. We have,

$$\Phi_n(\bar{\omega}) = |\psi_S\rangle \langle \eta_n(\bar{\omega})| + M_Q(\omega_n) \cdots M_Q(\omega_1), \quad (3.26)$$

where

$$\eta_n(\bar{\omega}) = \psi(\omega_1) + M_Q(\omega_1)^* \psi(\omega_2) + \cdots + M_Q(\omega_1)^* \cdots M_Q(\omega_{n-1})^* \psi(\omega_n) \quad (3.27)$$

$$= M(\omega_1)^* \cdots M(\omega_n)^* \psi(\omega_n) \quad (3.28)$$

and $\langle \psi_S, \eta_n(\bar{\omega}) \rangle = 1$. Using (3.27) and Lemma 2.2, we have for any $n, p \in \mathbb{N}$

$$\begin{aligned} \eta_{n+1+p}(\bar{\omega}) - \eta_{n+1}(\bar{\omega}) &= M_Q(\omega_1)^* \cdots M_Q(\omega_n)^* \sum_{j=1}^p M_Q(\omega_{n+1})^* \cdots M_Q(\omega_{n+j})^* \psi(\omega_{n+j+1}) \\ &= M_Q(\omega_1)^* \cdots M_Q(\omega_n)^* M(\omega_{n+1})^* \cdots M(\omega_{n+p})^* \psi(\omega_{n+p+1}). \end{aligned}$$

Thus, Lemma 2.3 and Proposition 2.4 imply

$$\|\eta_{n+p+1}(\bar{\omega}) - \eta_{n+1}(\bar{\omega})\| \leq C_0^2 \|M_Q(\omega_n)M_Q(\omega_{n-1}) \cdots M_Q(\omega_1)\|,$$

uniformly in $p \geq 0$. Using Theorem 1.1, there exists Ω_0 with $\mathbb{P}(\Omega_0) = 1$ and $\alpha > 0$ such that for any $n, p \in \mathbb{N}$, $\bar{\omega} \in \Omega_0$, there exists $C_{\bar{\omega}}$ s.t. $\|\eta_{n+p+1}(\bar{\omega}) - \eta_{n+1}(\bar{\omega})\| \leq C_0^2 C_{\bar{\omega}} e^{-n\alpha}$. Therefore, $\eta_n(\bar{\omega}) = \eta_\infty(\bar{\omega}) + O_{\bar{\omega}}(e^{-n\alpha})$. Together with (3.26) and Theorem 1.1, this proves (3.25).

Now, we can write

$$\begin{aligned} \Phi_n(\bar{\omega})^* \Phi_n(\bar{\omega}) &= |\eta_\infty(\bar{\omega})\rangle\langle\eta_\infty(\bar{\omega})| + O_{\bar{\omega}}(e^{-n\alpha}) \\ &= \|\eta_\infty(\bar{\omega})\|^2 |\hat{\eta}(\bar{\omega})\rangle\langle\hat{\eta}(\bar{\omega})| + O_{\bar{\omega}}(e^{-n\alpha}), \end{aligned}$$

where $\hat{\eta}(\bar{\omega}) = \eta_\infty(\bar{\omega})/\|\eta_\infty(\bar{\omega})\|$, so that $|\hat{\eta}(\bar{\omega})\rangle\langle\hat{\eta}(\bar{\omega})|$ is an orthogonal projector. Hence, by perturbation theory, the non-negative self-adjoint operator $\Phi_n(\bar{\omega})^* \Phi_n(\bar{\omega})$ has a simple eigenvalue $\|\eta_\infty(\bar{\omega})\|^2 + O_{\bar{\omega}}(e^{-n\alpha})$ and all its other eigenvalues are non-negative and bounded from above by $C_{\bar{\omega}} e^{-n\alpha}$, for some $C_{\bar{\omega}} > 0$. Thus the $2n$ th root of $\Phi_n^*(\bar{\omega})\Phi_n(\bar{\omega})$ has in the limit $n \rightarrow \infty$, a simple eigenvalue 1, and all its other eigenvalues are bounded from above by $e^{-\alpha/2}$. \square

4. From repeated interactions to matrix products

In this section, we show how to reduce the evolution of states of a R.I. quantum system (for observables on the small system \mathcal{S}) to the study of a product of non-identical matrices given by a RDO process.

4.1. The repeated interaction model

We call the *time step* m the index $m \geq 1$ that labels the interaction step during which the system \mathcal{S} interacts with the element \mathcal{E}_m of the chain $\mathcal{C} = \mathcal{E}_1 + \mathcal{E}_2 + \mathcal{E}_3 + \cdots$, even though the duration τ_m of this interaction may vary with m , with the understanding that $\inf_{m \in \mathbb{N}^*} \tau_m > 0$.

The Hilbert space $\mathcal{H}_{\mathcal{S}}$ is finite-dimensional. We denote by $L_{\mathcal{E}_k}$ the free Liouvillean of \mathcal{E}_k , acting on $\mathcal{H}_{\mathcal{E}_k}$. The invariant reference vector $\psi_{\mathcal{E}_k} \in \mathcal{H}_{\mathcal{E}_k}$ is cyclic and separating for $\mathfrak{M}_{\mathcal{E}_k}$, it is independent of time and satisfies $L_{\mathcal{E}_k} \psi_{\mathcal{E}_k} = 0$. The interaction operator between the m th element of the chain and the reference system at time m is a self-adjoint operator $V_m \in \mathfrak{M}_{\mathcal{S}} \otimes \mathfrak{M}_{\mathcal{E}_m}$. The coupled dynamics on $\mathcal{H}_{\mathcal{S}} \otimes \mathcal{H}_{\mathcal{E}_m}$ is then generated by the interaction Liouvillean $L_m = L_{\mathcal{S}} + L_{\mathcal{E}_m} + V_m$. Moreover, if J_m and Δ_m denote the modular data of the pair $(\mathfrak{M}_{\mathcal{S}} \otimes \mathfrak{M}_{\mathcal{E}_m}, \psi_{\mathcal{S}} \otimes \psi_{\mathcal{E}_m})$ (see e.g. [5]), we further require

$$\Delta_m^{1/2} V_m \Delta_m^{-1/2} \in \mathfrak{M}_{\mathcal{S}} \otimes \mathfrak{M}_{\mathcal{E}_m} \quad \forall m \geq 1.$$

The evolution operator acting on $\mathcal{H} = \mathcal{H}_{\mathcal{S}} \otimes \mathcal{H}_{\mathcal{C}}$ in the Schrödinger picture at time step m is given by the unitary $U(m) = e^{-i\tilde{L}_m} e^{-i\tilde{L}_{m-1}} \cdots e^{-i\tilde{L}_1}$, where \tilde{L}_m is given in (1.14).

Let us finally define the Liouville operator K_m at time m , see [8,14,18], by

$$K_m = \tau_m [L_{\mathcal{S}} + L_{\mathcal{E}_m} + V_m - J_m \Delta_m^{1/2} V_m \Delta_m^{-1/2} J_m].$$

Its main dynamical features,

$$e^{i\tau_m L_m} A e^{-i\tau_m L_m} = e^{iK_m} A e^{-iK_m} \quad \text{for } A \in \mathfrak{M}_{\mathcal{S}} \otimes \mathfrak{M}_{\mathcal{E}}, t \in \mathbb{R}, \quad (4.1)$$

$$K_m \psi_{\mathcal{S}} \otimes \psi_{\mathcal{E}_m} = 0, \quad (4.2)$$

(see also (1.21) and (1.22)), are proven to hold by using standard relations of the modular data Δ_m, J_m , see e.g. [8]. Note also the bound $\|e^{\pm iK_m}\| \leq e^{\tau_m \|\Delta_m^{1/2} V_m \Delta_m^{-1/2}\|}$.

We proceed along the same lines as in [8], taking into account that in the present setup, V_m may vary with m . Let $\mathcal{H}_C := \bigotimes_{k \geq 1} \mathcal{H}_{\mathcal{E}_k}$, where the infinite tensor product is understood with respect to the reference vector $\psi_C := \bigotimes_{k \geq 1} \psi_{\mathcal{E}_k}$. Since L_m and $L_{\mathcal{E}_k}$ commute for $k \neq m$, we can write successively

$$\begin{aligned} e^{-i\tilde{L}_1} &= e^{-i\tau_1 L_1} e^{-i\tau_1 \sum_{k>1} L_{1k}}, \\ e^{-i\tilde{L}_2} &= e^{-i\tau_2 L_{\mathcal{E}_1}} e^{-i\tau_2 L_2} e^{-i\tau_2 \sum_{k>2} L_{2k}}, \\ &\vdots \\ e^{-i\tilde{L}_m} &= e^{-i\tau_m \sum_{k<m} L_{\mathcal{E}_k}} e^{-i\tau_m L_m} e^{-i\tau_m \sum_{k>m} L_{\mathcal{E}_k}}, \end{aligned}$$

so that we obtain

$$\begin{aligned} U(m) &= \exp \left[-i \sum_{j=1}^m \sum_{k=1}^{j-1} \tau_j L_{\mathcal{E}_k} \right] e^{-i\tau_m L_m} e^{-i\tau_{m-1} L_{m-1}} \dots e^{-i\tau_1 L_1} \exp \left[-i \sum_{j=1}^m \sum_{k>j} \tau_j L_{\mathcal{E}_k} \right] \\ &\equiv U_m^- e^{-i\tau_m L_m} e^{-i\tau_{m-1} L_{m-1}} \dots e^{-i\tau_1 L_1} U_m^+. \end{aligned} \quad (4.3)$$

The unitaries U_m^\pm act trivially on \mathcal{H}_S and satisfy $U_m^\pm \psi_C = \psi_C$, $\forall m \in \mathbb{N}^*$. The corresponding Heisenberg evolution of an observable $O \in \mathfrak{M} = \mathfrak{M}_S \otimes \mathfrak{M}_C$ at time m is (1.17). Given a normal state ϱ on \mathfrak{M} , we want to understand the limit of large m of the expectation value $\varrho(\alpha_{RI}^m(O)) = \varrho(U(m)^* O U(m))$, for observables on the small system S , i.e. $O = A_S \otimes \mathbb{1}_C$ (more general observables, such as the so-called instantaneous observables following the terminology of [8], are considered in [9]). In other words, we wish to address the long time behaviour of the evolution of the state ϱ defined by $\varrho \circ \alpha_{RI}^m$ on \mathfrak{M} .

Any normal state is a convex combination of vector states, i.e., its density matrix is given by a convex combination of rank-one projections, $\rho = \sum_n p_n |\phi_n\rangle\langle\phi_n|$, where $\sum_n p_n = 1$, and where the ϕ_n are normalized vectors in \mathcal{H} . It is then clear that convergence of the vector states $|\phi_n\rangle\langle\phi_n|$ implies convergence of ρ (see [8] for more details). Therefore, we will limit our attention to vector states $\langle\phi, \cdot\rangle$ from now on. Moreover, because $\psi := \psi_S \otimes \psi_C$ is cyclic for the commutant \mathfrak{M}' , it is enough to consider ϕ of the form

$$\phi = B' \psi_S \otimes \psi_C =: B' \psi \quad (4.4)$$

for some

$$B' = B'_S \bigotimes_{n=1}^N B'_n \bigotimes_{n>N} \mathbb{1}_{\mathcal{E}_n} \in \mathfrak{M}', \quad (4.5)$$

with $B'_S \in \mathfrak{M}'_S$, $B'_n \in \mathfrak{M}'_{\mathcal{E}_n}$ (up to a vanishing error as $N \rightarrow \infty$, see [8]). Hence, we are lead to consider

$$\langle\psi, (B')^* \alpha_{RI}^m(O) B' \psi\rangle = \langle\psi, (B')^* B' \alpha_{RI}^m(O) \psi\rangle \quad (4.6)$$

for observables $O = A_S \otimes \mathbb{1}_C$. In (4.6) we use that B' commutes with $\alpha_{RI}^m(O)$.

4.2. Observables on the small system

In this paper we restrict our attention to the Heisenberg evolution of observables $A_S \in \mathfrak{M}_S$. That is, we consider

$$\begin{aligned} \alpha_{RI}^m(A_S \otimes \mathbb{1}) &= U(m)^*(A_S \otimes \mathbb{1})U(m) \\ &= (U_m^+)^* e^{i\tau_1 L_1} e^{i\tau_2 L_2} \dots e^{i\tau_m L_m} (A_S \otimes \mathbb{1}) e^{-i\tau_m L_m} e^{-i\tau_{m-1} L_{m-1}} \dots e^{-i\tau_1 L_1} U_m^+, \end{aligned}$$

where we made use of the fact that U_m^- acts trivially on \mathcal{H}_S . Considering (4.6), we note that due to the properties of the unitary $U^+(m)$ and those of the Liouvillean, (4.1) and (4.2), we have

$$\begin{aligned} & \alpha_{RI}^m(A_S \otimes \mathbb{1})\psi \\ &= U^+(m)^* e^{i\tau_1 L_1} e^{i\tau_2 L_2} \dots e^{i\tau_m L_m} (A_S \otimes \mathbb{1}) e^{-i\tau_m L_m} e^{-i\tau_{m-1} L_{m-1}} \dots e^{-i\tau_1 L_1} \psi \\ &= U^+(m)^* e^{iK_1} e^{iK_2} \dots e^{iK_m} (A_S \otimes \mathbb{1})\psi. \end{aligned}$$

Let us introduce $P_N = \mathbb{1}_S \otimes \mathbb{1}_{\mathcal{E}_1} \otimes \dots \otimes \mathbb{1}_{\mathcal{E}_N} \otimes P_{\psi_{\mathcal{E}_{N+1}}} \otimes P_{\psi_{\mathcal{E}_{N+2}}} \otimes \dots$, where $P_{\psi_{\mathcal{E}_k}} = |\psi_{\mathcal{E}_k}\rangle\langle\psi_{\mathcal{E}_k}|$. Due to (4.5), we have $\langle\psi|(B')^* B' = \langle\psi|(B')^* B' P_N$. Moreover, introducing the unitary operator

$$\tilde{U}^+(N) = \exp \left[-i \sum_{j=1}^{N-1} \sum_{k=j+1}^N \tau_j L_{\mathcal{E}_k} \right],$$

which is *independent* of m , we can write for $m > N$,

$$\begin{aligned} & \langle\psi, (B')^* B' \alpha_{RI}^m(A_S \otimes \mathbb{1})\psi\rangle \\ &= \langle\psi, (B')^* B' \tilde{U}^+(N)^* P_N e^{iK_1} e^{iK_2} \dots e^{iK_m} (A_S \otimes \mathbb{1})\psi\rangle \\ &= \langle\psi, (B')^* B' \tilde{U}^+(N)^* e^{iK_1} \dots e^{iK_N} P_N e^{iK_{N+1}} \dots e^{iK_m} (A_S \otimes \mathbb{1})\psi\rangle. \end{aligned}$$

At this level, introducing the projector $P = \mathbb{1}_S \otimes |\psi_C\rangle\langle\psi_C|$, satisfying $P\psi = \psi$, we observe that $P_N e^{iK_{N+1}} \dots e^{iK_m} (A_S \otimes \mathbb{1})\psi = P e^{iK_{N+1}} \dots e^{iK_m} P (A_S \otimes \mathbb{1})\psi$. Thus, we can invoke Proposition 4.1 of [8] which states that for any $q \geq 1$ and any distinct integers n_1, \dots, n_q ,

$$P e^{iK_{n_1}} e^{iK_{n_2}} \dots e^{iK_{n_q}} P = P e^{iK_{n_1}} P e^{iK_{n_2}} P \dots P e^{iK_{n_q}} P. \quad (4.7)$$

We note that the proof of this statement does not require that the K_n 's be identical as operators on $\mathcal{H}_S \otimes \mathcal{H}_E$. Therefore, introducing operators M_j acting on \mathcal{H}_S by

$$P e^{iK_j} P = M_j \otimes |\psi_C\rangle\langle\psi_C| \quad \text{or} \quad M_j \simeq P e^{iK_j} P, \quad (4.8)$$

we arrive at

$$\begin{aligned} & \langle\psi|(B')^* B' \alpha_{RI}^m(A_S \otimes \mathbb{1})\psi\rangle \\ &= \langle\psi|(B')^* B' \tilde{U}^+(N)^* e^{iK_1} \dots e^{iK_N} P M_{N+1} M_{N+2} \dots M_m (A_S \otimes \mathbb{1})\psi\rangle. \end{aligned} \quad (4.9)$$

Hence, given the repeated interactions dynamics, in order to understand the long time behaviour of the evolution of any states $\varrho \circ \alpha_{RI}^m$ on the set of observables on the reference system \mathcal{S} , it is necessary (and sufficient) to address the asymptotic behaviour of the product of matrices $M_j \simeq P e^{iK_j} P$ of the type $\Psi_m = M_1 M_2 \dots M_m$ on \mathcal{H}_S .

Finally, we recall the main features of these matrices M_j inherited from those of e^{iK_j} which shows that they indeed satisfy properties (1) and (2) of an RRDO. The next lemma is an easy generalization of Proposition 2.1 in [8].

Lemma 4.1 ([8]). *With the definition (4.8), we have for any $j \in \mathbb{N}^*$, $M_j \psi_S = \psi_S$. Moreover, to any $\phi \in \mathcal{H}_S$ corresponds a unique $A \in \mathfrak{M}_S$ such that $\phi = A \psi_S$, and $\|\phi\| := \|A\|_{\mathcal{B}(\mathcal{H}_S)}$ defines a norm on \mathcal{H}_S . As an operator on \mathcal{H}_S endowed with the norm $\|\cdot\|$, M_j is a contraction for any $j \in \mathbb{N}^*$.*

4.3. Proof of Theorem 1.5

For random repeated interactions, Theorem 1.3 shows that there exists a vector $\theta \in \mathcal{H}_S$ satisfying $\langle\theta, \psi_S\rangle = 1$, such that, with probability 1,

$$\lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M \Psi_m(\omega) = |\psi_S\rangle\langle\theta| = \pi, \quad (4.10)$$

where $\pi = |\psi_S\rangle\langle\theta|$ is a rank one projector in \mathcal{H}_S . Therefore, (4.9) yields

$$\begin{aligned} & \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M \langle \psi | (B')^* B' \alpha_{RI}^m(A_S \otimes \mathbb{1}) \psi \rangle \\ &= \langle \psi, (B')^* B' \tilde{U}^+(N)^* e^{iK_1} \dots e^{iK_N} P \pi(A_S \otimes \mathbb{1}) \psi \rangle \\ &= \langle \psi, (B')^* B' \tilde{U}^+(N)^* e^{iK_1} \dots e^{iK_N} |\psi\rangle\langle\theta, A_S \psi_S\rangle_{\mathcal{H}_S} \rangle \\ &= \langle \psi | (B')^* B' \psi \rangle_{\langle\theta | A_S \psi_S\rangle_{\mathcal{H}_S}} \\ &= \|B' \psi\|^2_{\langle\theta | A_S \psi_S\rangle_{\mathcal{H}_S}}, \end{aligned}$$

where $\|B' \psi\|$ is arbitrary close to $\|\psi\| = 1$, for N large enough, see (4.4).

In other words, (4.10) implies the existence of an asymptotic ergodic state ϱ_E on \mathfrak{M}_S , defined by $\varrho_E(A_S) := \langle\theta, A_S \psi_S\rangle_{\mathcal{H}_S}$, $\forall A_S \in \mathfrak{M}_S$, such that for any normal state ϱ on \mathfrak{M} , $\lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M \varrho(\alpha_{RI}^m(A_S \otimes \mathbb{1})) = \varrho_E(A_S)$, which is the statement of Theorem 1.5.

Acknowledgement

We thank the ACI Jeunes Chercheurs ‘‘Stochastic modeling of non-equilibrium systems’’ for financial support.

References

- [1] L. Arnold. *Random Dynamical Systems*. Springer, Berlin, 2003. [MR1723992](#)
- [2] S. Attal, A. Joye and C.-A. Pillet (Eds). *Open Quantum Systems I-III. Lecture Notes in Mathematics 1880–1882*. Springer, Berlin, 2006. [MR2261249](#)
- [3] E. A. Azoff. Borel measurability in linear algebra. *Proc. Amer. Math. Soc.* **42** (1974) 346–350. [MR0327799](#)
- [4] A. Beck and J. T. Schwartz. A vector-valued random ergodic theorem. *Proc. Amer. Math. Soc.* **8** (1957) 1049–1059. [MR0098162](#)
- [5] O. Bratteli and D. W. Robinson. *Operator Algebras and Quantum Statistical Mechanics. Texts and Monographs in Physics 1,2*, 2nd edition. Springer, New York, 1996.
- [6] R. Bru, L. Elsner and M. Neumann. Convergence of infinite products of matrices and inner–outer iteration schemes. *Electron. Trans. Numer. Anal.* **2** (1994) 183–193. [MR1308895](#)
- [7] M. Brune, J. M. Raimond and S. Haroche. Theory of the Rydberg-atom two-photon micromaser. *Phys. Rev. A* **35** (1987) 154–163.
- [8] L. Bruneau, A. Joye and M. Merkli. Asymptotics of repeated interaction quantum systems. *J. Funct. Anal.* **239** (2006) 310–344. [MR2258226](#)
- [9] L. Bruneau, A. Joye and M. Merkli. Random repeated interaction quantum systems. *Comm. Math. Phys.* **284** (2008) 553–581. [MR2448141](#)
- [10] B. De Saporta, Y. Guivarc’h and E. LePage. On the multidimensional stochastic equation $Y(n+1) = a(n)Y(n) + b(n)$. *C. R. Math. Acad. Sci. Paris* **339** (2004) 499–502. [MR2099549](#)
- [11] P. Filipowicz, J. Javanainen and P. Meystre. Theory of a microscopic maser. *Phys. Rev. A* **34** (1986) 3077–3087.
- [12] Y. Guivarc’h. Limit theorem for random walks and products of random matrices. In *Proceedings of the CIMPA-TIFR School on Probability Measures on Groups, Recent Directions and Trends, September 2002* 255–330. TIFR, Mumbai, 2006.
- [13] W. Hoeffding. Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.* **58** (1963) 13–30. [MR0144363](#)
- [14] V. Jaksic and C.-A. Pillet. Non-equilibrium steady states of finite quantum systems coupled to thermal reservoirs. *Comm. Math. Phys.* **226** (2002) 131–162. [MR1889995](#)
- [15] H. Kesten and F. Spitzer. Convergence in distribution of products of random matrices. *Z. Wahrsch. Verw. Gebiete* **67** (1984) 363–386. [MR0761563](#)
- [16] Y. Kifer and P.-D. Liu. Random dynamics. In *Handbook of Dynamical Systems 1B* 379–499. B. Hasselblatt and A. Katok (Eds). North-Holland, Amsterdam, 2006. [MR2186245](#)
- [17] D. Meschede, H. Walther and G. Müller. One-atom maser. *Phys. Rev. Lett.* **54** (1985) 551–554.
- [18] M. Merkli, M. Mück and I. M. Sigal. Instability of equilibrium states for coupled heat reservoirs at different temperatures. *J. Funct. Anal.* **243** (2007) 87–120. [MR2291433](#)
- [19] A. Mukherjea. *Topics in Products of Random Matrices*. TIFR, Mumbai, 2000. [MR1759920](#)
- [20] E. Seneta. *Non-negative Matrices and Markov Chains. Springer Series in Statistics*. Springer, New York, 2000. [MR2209438](#)
- [21] S. Schwarz. Infinite product of doubly stochastic matrices. *Acta Math. Univ. Comenian.* **39** (1980) 131–150. [MR0619269](#)
- [22] M. Weidinger, B. T. H. Varcoe, R. Heerlein and H. Walther. Trapping states in the micromaser. *Phys. Rev. Lett.* **82** (1999) 3795–3798.
- [23] T. Wellens, A. Buchleitner, B. Kümmerer and H. Maassen. Quantum state preparation via asymptotic completeness. *Phys. Rev. Lett.* **85** (2000) 3361–3364.