

Decay of geometry for Fibonacci critical covering maps of the circle [☆]

Eduardo Colli ^a, Marcio L. do Nascimento ^{b,1}, Edson Vargas ^{a,*,2}

^a Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, Cep 05508-090, São Paulo, SP, Brazil

^b Instituto de Ciências Exatas e Naturais, Universidade Federal do Pará, Rua Augusto Corrêa 01, Cep 66075-110, Belém, PA, Brazil

Received 19 February 2008; received in revised form 9 March 2009; accepted 16 March 2009

Available online 7 April 2009

Abstract

We study the growth of $Df^n(f(c))$ when f is a Fibonacci critical covering map of the circle with negative Schwarzian derivative, degree $d \geq 2$ and critical point c of order $\ell > 1$. As an application we prove that f exhibits exponential decay of geometry if and only if $\ell \leq 2$, and in this case it has an absolutely continuous invariant probability measure, although not satisfying the so-called Collet–Eckmann condition.

© 2009 Elsevier Masson SAS. All rights reserved.

Résumé

Nous étudions la croissance de $Df^n(f(c))$ lorsque f est un revêtement critique de Fibonacci du cercle avec dérivée Schwarzienne négative, degré $d \geq 2$ et point critique c d'ordre $\ell > 1$. Comme application nous démontrons que f exhibe une décroissance exponentielle de géométrie si et seulement si $\ell \leq 2$, et dans ce cas f a une mesure de probabilité invariante absolument continue, sans satisfaire la condition de Collet–Eckmann.

© 2009 Elsevier Masson SAS. All rights reserved.

MSC: 37E10; 37C40; 37L40

Keywords: Circle maps; Covering maps; Fibonacci combinatorics; Decay of geometry; Invariant measures

1. Introduction

A critical covering map of the circle $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$ is a C^r ($r \geq 1$) covering map $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ with just one critical point, say c , which must be of inflection type. In the present work, we will consider critical covering maps of degree $d \geq 2$. This kind of map has been considered before, for example in [20–22,30]. Under the point of view of dynamical systems such critical coverings with neither wandering intervals nor periodic attractors are all topologically conju-

[☆] This work is part of the “Projeto Temático – Fapesp # 2006/03829-2”.

* Corresponding author. Tel.: +55 (11) 30916156; fax: +55 (11) 30916183.

E-mail addresses: colli@ime.usp.br (E. Colli), marcion@ufpa.br (M.L. do Nascimento), vargas@ime.usp.br (E. Vargas).

¹ Partially supported by CNPq, grant # 154594/2006-7.

² Partially supported by CNPq, grant # 304517/2005-4.

gate to the map of the circle induced by the map $x \mapsto dx$ of the real line \mathbb{R} . This implies in particular that the set $\mathcal{B}(f) := \{x \in \mathbb{S}^1 : \omega(x) = \mathbb{S}^1\}$ is a countable intersection of open and dense subsets of \mathbb{S}^1 . But metric properties of the underlying dynamics as the Lebesgue measure of $\mathcal{B}(f)$ or the growth of $Df^n(f(c))$, as $n \rightarrow \infty$, depend on the order and combinatorial behavior of the forward critical orbit. Here we will consider the case of critical points of order $\ell > 1$ with the *Fibonacci combinatorics*, which will be defined below.

The Fibonacci combinatorics appeared before in the context of unimodal maps related to a question posed by J. Milnor [28] about the classification of the measure-theoretical attractors in dynamical systems on compact spaces. Among the quadratic real polynomials $Q_\alpha(x) = \alpha x(1-x)$ there is one whose turning point has this combinatorics and, as it implies a strong recurrence of the critical point, it was considered in [14] as a candidate to exhibit a *wild attractor*. A wild attractor for Q_α (also called *absorbing Cantor attractor* in [13]) is a compact invariant Cantor set whose basin of attraction is a meager subset of $[0, 1]$ with full Lebesgue measure [28]. It was proved in [24] that a quadratic real polynomial with the Fibonacci combinatorics has no wild attractor. This was generalized later for any combinatorics [23] (see also [7,10–13,16,26]). On the other hand, in [1] it was proved that if a unimodal real polynomial of degree big enough has the Fibonacci combinatorics then it has a wild attractor.

Under mild hypotheses a critical covering map f as above is ergodic with respect to the Lebesgue measure [35], that is, every totally invariant set has Lebesgue measure zero or one. Then, if it has an absolutely continuous invariant probability measure it has no wild attractor and the topological conjugacy between f and a map of the circle induced by $x \mapsto dx$ is absolutely continuous.

In order to prove that there is an absolutely continuous invariant probability measure one examines the derivative on the orbit of the critical value and tries to show that it undergoes a relative expansion, for example by showing that the map satisfies the so-called *Collet–Eckmann condition*, stating that the derivative on the critical orbit grows exponentially, or weaker conditions as the summability condition in [2,31] and also [3]. This may be done either by directly examining the derivative for some specially chosen moments of the orbit (at closest returns, for example) or by studying the geometric aspects of sequences of inductively defined return maps around the critical point. In the latter case, if some metric relations between successive stages decay exponentially fast then one says that f exhibits *exponential decay of geometry* (see below for a precise definition), the opposite situation being *bounded geometry*, where scales do not decay at all. Exponential decay of geometry and topological additional hypotheses (as finiteness of central returns) are related to the expansion of the derivative on the critical orbit and have also been a way of proving existence of absolutely continuous invariant probability measures (see [27] for the unimodal case). We also mention that in the proof of denseness of hyperbolicity in the logistic family Q_α (see [8,9,25]), a central problem in one-dimensional dynamics, the exponential decay of geometry played an important role.

Several questions about dynamics on the real line were treated with tools from complex analysis. But these tools can be applied just to a somewhat narrow set of cases which do not include situations where the order of the critical point is a general real number greater than one. It is also a natural aim that real analysis is enough to solve the questions from real dynamics. In this line of thought the non-existence of wild attractors for negative Schwarzian derivative Fibonacci unimodal maps with critical point of order $1 < \ell \leq 2$ was proved by G. Keller and T. Nowicki in [17] with just real analysis (in fact the result extends a little above $\ell = 2$). Moreover, the exponential decay of geometry was proved by W. Shen in [33] for non-renormalizable unimodal maps with any combinatorics with critical point of order $1 < \ell \leq 2$. Shen strongly uses the natural symmetry around the critical point that arises in the unimodal case, contrary to what happens with the inflection critical point of critical covering maps.

Here the decay of geometry for a map with an inflection critical point is considered for the first time. We prove that Fibonacci critical covering maps of the circle of degree $d \geq 2$ having critical point of order $1 < \ell \leq 2$ display decay of geometry. Moreover, our result is sharp since in [22] it was proved that Fibonacci critical coverings have bounded geometry for every $\ell > 2$. This is similar to the unimodal case where bounded geometry for the Fibonacci combinatorics follows for every $\ell > 2$ from a result in [17].

We also mention that the study of the growth of $|Df^n(f(c))|$ connected to the question of the existence of an absolutely continuous invariant probability measure and the regularity of topological conjugacies appeared at least three decades ago, see for example [5,15,18,19,29,32,34] and others references in the books [4] and [6].

2. Basic concepts and main results

We look at the circle \mathbb{S}^1 as the quotient space \mathbb{R}/\mathbb{Z} with the orientation and metric induced from the real line \mathbb{R} . Given a C^1 critical covering map $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ of degree $d \geq 2$ we choose one of its fixed points, say p , and set $I_1 := \mathbb{S}^1 \setminus \{p\}$. The distance between 2 points x and y in I_1 will be denoted by $|x - y|$ and the length of an interval $I \subset I_1$ will be denoted by $|I|$.

Note that f has d branches mapping onto I_1 (below we will identify I_1 with $(0, 1) \subset \mathbb{R}$ in order to consider homeomorphic extensions of these branches and prove a technical but essential lemma). From now on we will assume that the critical point c of f is recurrent and define the sequence $I_1 \supset I_2 \supset I_3 \supset \dots \{c\}$ such that, for $n \geq 1$, the interval I_{n+1} is a component of the domain of the first return map ϕ_n to I_n . If $\phi_n(c) \in I_{n+1}$, for some $n \geq 1$, we say that n is a *central return moment*. The *critical return time* s_n is defined by $\phi_n(c) = f^{s_n}(c)$. If the sequence $s_1, s_2, s_3, s_4, \dots$ coincides with the Fibonacci sequence $1, 2, 3, 5, \dots$ the critical covering map f is called a *Fibonacci critical covering map*. It is not difficult to show that Fibonacci critical covering maps have no central returns.

One of the main geometric aspects of one-dimensional dynamics with critical points is the evolution of the *scaling factor* $\mu_n = |I_{n+1}|/|I_n|$, particularly in the subsequence $(n_i)_i$ of non-central return moments (which is the sequence itself in the case of Fibonacci critical covering maps). If $\mu_{n_i} \rightarrow 0$ exponentially fast as $i \rightarrow \infty$ we say that f has *exponential decay of geometry*.

Let \mathcal{C}_d denote the set of C^1 critical coverings f of degree $d \geq 2$ which also have the following properties:

- The critical point c is recurrent and not periodic.
- There exist a C^1 map $\psi : I_1 \rightarrow I_1$ satisfying $\lim_{x \rightarrow c} \psi(x) = 0$ and real constants $\vartheta > 0$ and $\ell > 1$ such that

$$f(x) = f(c) + \vartheta \operatorname{sgn}(x - c)|x - c|^\ell(1 + \psi(x)) \tag{1}$$

for every x in a neighborhood of c .

- Restricted to $\mathbb{S}^1 \setminus \{c\}$, the map f is C^3 and has negative Schwarzian derivative.

Remark that a map $f \in \mathcal{C}_d$ cannot have neither a wandering interval nor a periodic attractor, see [6] and also [35]. Our main result claims the exponential growth of $Df^{s_n}(f(c))$ as $n \rightarrow \infty$ for Fibonacci critical coverings.

Theorem 1. *There are constants $\rho > 1$ and $C > 0$ such that if $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell \in (1, 2]$ then $Df^{s_n}(f(c)) \geq C\rho^n$, for all $n \geq 1$.*

Three consequences are the following.

Theorem 2. *If $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell \in (1, 2]$ then f has an absolutely continuous invariant probability measure.*

Theorem 3. *If $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell \in (1, 2]$ then f exhibits exponential decay of geometry.*

Theorem 4. *If $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell \in (1, 2]$ then the ω -limit set $\omega(c)$ of its critical point is a minimal invariant Cantor set with zero Hausdorff dimension.*

In order to prove Theorem 1 we find a recursive difference equation (12) involving $Df^{s_n}(f(c))$ and the parameter $\lambda_n = |f^{s_n}(c) - c|/|f^{s_{n+2}}(c) - c|$. Then we proceed as in [17] to get the exponential growth of $Df^{s_n}(f(c))$ with n . Using a classical argument, we show that the sequence of derivatives $Df^k(f(c))$ goes to infinity as $k \rightarrow \infty$ and this is exactly the criterion in [3] assuring the existence of an absolutely continuous invariant probability measure, as stated in Theorem 2. Finally, Theorem 1 together with distortion control imply the third and the fourth theorems.

A technical difference with respect to the work of Keller and Nowicki [17] is that we do not get an *a priori* numeric lower bound for λ_n , and in fact we do not need it for the proof. It suffices to use that λ_n is uniformly bounded away from one, which is a consequence of the real bounds (see Theorem 9 below). Another difference is a cross-ratio argument where an asymmetric huge extendibility allowed by topological properties plays a central role. For this

argument, the assumption of negative Schwarzian derivative is essential, since the range of extendibility goes much beyond the neighborhood of the critical point.

We also derive some more precise statements about the geometry and the derivative growth in the Fibonacci combinatorics, as a consequence of these theorems and a technical result proved in [22] for any $\ell > 1$. This result states, in particular, that the two components of $I_n \setminus I_{n+1}$ are comparable with I_n . Hence I_{n+1} is not only exponentially small with respect to I_n but also with respect to these two adjacent components.

The derivative growth presents three distinct pictures, depending on the value of ℓ , as the following theorems tell.

Theorem 5. *Let $f \in \mathcal{C}_d$ with the Fibonacci combinatorics and $\ell \in (1, 2)$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \log Df^{s_n}(f(c)) = \mu,$$

where

$$\mu = \frac{1 + \sqrt{1 + 4\ell}}{2\ell}.$$

In particular, the geometry decays super exponentially.

This theorem says that $Df^{s_n}(f(c))$ grows super exponentially with n , for $\ell < 2$. But as s_n grows with ω^n , where $\omega = \frac{1+\sqrt{5}}{2}$ is the golden number, and $\mu = \mu(\ell)$ decreases from ω , when $\ell = 1$, to 1, when $\ell = 2$, then we can conclude that $Df^{s_n}(f(c))$ does not grow exponentially with s_n . Therefore, as a consequence, f is not Collet–Eckmann. But we can extend this conclusion to all iterates, by the following theorem.

Theorem 6. *Let $f \in \mathcal{C}_d$ with the Fibonacci combinatorics and $\ell \in (1, 2)$. Then*

$$\lim_{k \rightarrow \infty} \frac{\log \log Df^k(f(c))}{\log k} = \frac{\log \mu}{\log \omega} < 1,$$

where $\mu = \mu(\ell)$ is given as in Theorem 5 and ω is the golden number. *In particular, f does not satisfy the Collet–Eckmann condition.*

For $\ell = 2$ the exponential growth of $Df^{s_n}(f(c))$ with n is the best that we can hope.

Theorem 7. *Let $f \in \mathcal{C}_d$ with the Fibonacci combinatorics and $\ell = 2$. Then the growth of $Df^{s_n}(f(c))$ is not more than exponential. In particular, the geometry decays with a ratio which is not more than exponential.*

Finally, for $\ell > 2$ we conclude that $Df^{s_n}(f(c))$ and the geometry are bounded. This last assertion, in particular, is also proved in [22].

Theorem 8. *Let $f \in \mathcal{C}_d$ with the Fibonacci combinatorics and $\ell > 2$. Then $Df^{s_n}(f(c))$ is bounded, and in particular the geometry is bounded.*

This paper is organized as follows. In Section 3 we review the technical tools related to the uniform distortion control of powers of f which are widely used in one-dimensional dynamics. The so-called real bounds play a central role: they state an *a priori* property on the geometry of the induced sequence of first return maps. In Section 4 we discuss in more details the Fibonacci combinatorics, which is explicitly used in the proofs. The facts stated in this section are very similar to the Fibonacci combinatorics of unimodal maps. In Section 5 we state the extendibility properties of a first entry map from the critical value to a neighborhood of the critical point which will be essential to get lower and upper bounds for its derivative, through a cross-ratio argument. These bounds are obtained in Section 6 and used to show Theorem 1, in Section 7. In this section we also prove the existence of an absolutely continuous invariant probability measure, the exponential decay of geometry and the zero Hausdorff dimension of $\omega(c)$. Finally, in Section 8 we derive, using a result in [22], the more precise consequences of these findings, which are stated in Theorems 5, 6, 7 and 8.

3. Cross-ratio, distortion control

Koebe Principles are the most important tools to controlling distortion and finding lower bounds for derivatives that will ultimately prove Theorem 1. We list some important definitions and results that may be found, for example, in [6].

Let $J, T \subset I_1 \subset \mathbb{S}^1$ be a pair of intervals such that $J \subset T$ and $T \setminus J$ has 2 non-empty connected components L and R . We say that J is α -well inside T if $\min\{|L|, |R|\} \geq \alpha|J|$. The *cross-ratio* of J and T is the ratio

$$C(T, J) = \frac{|T| |L|}{|J| |R|}.$$

Let $h : T \rightarrow \mathbb{S}^1$ be a diffeomorphism onto its image and define

$$B(h, T, J) = \frac{C(h(T), h(J))}{C(T, J)}.$$

If h is C^3 and has negative Schwarzian derivative then $B(h, T, J) > 1$. As a consequence, if $J \subset T$ is degenerated to a point x we have

$$|Dh(x)| \geq \frac{|h(L)||h(R)|}{|h(T)|} \frac{|T|}{|L||R|}. \tag{2}$$

On the other hand, if we degenerate R to a point x then

$$|Dh(x)| \leq \frac{|h(T)||h(J)|}{|h(L)|} \frac{|L|}{|T||J|}. \tag{3}$$

There is also the Macroscopic Koebe Principle saying that if $h(J)$ is α -well inside $h(T)$ then J is also α -well inside T . Under the same condition the distortion is controlled by

$$\frac{Dh(x)}{Dh(y)} \leq \left(\frac{1 + \alpha}{\alpha} \right)^2, \tag{4}$$

for all $x, y \in J$.

In [20] and [35] one may find the following important result, that will be used everywhere throughout this work.

Theorem 9 (Real bounds). *There is $\alpha = \alpha(\ell) > 0$ (which can be uniform for $\ell \leq \ell_0$, for every $\ell_0 > 1$), such that if $f \in \mathcal{C}_d$ has order ℓ then I_{n+1} is α -well inside I_n for every non-central return moment n after two non-central return moments. In particular, if f has the Fibonacci combinatorics, this is true for every $n \geq 3$.*

Again we consider ϕ_n , the first return map to the interval I_n associated to the critical covering $f \in \mathcal{C}_d$. All branches of ϕ_n map their domains diffeomorphically onto I_n except by the *critical branch* $\phi_n|_{I_{n+1}}$, which is a C^1 homeomorphism onto I_n . In the case of the critical branch, even if the distortion cannot be bounded because of the presence of the critical point, the corollary of the next lemma guarantees that it can be written as the composition of f with a map (a power of f) having uniformly bounded distortion.

Lemma 10. *If $f \in \mathcal{C}_d$ and $x \in I_1 \setminus I_{n+1}$ is a point such that there exists the smallest $k \geq 1$ such that $f^k(x) \in I_{n+1}$ then there is an interval T containing x which is mapped by f^k diffeomorphically onto I_n .*

Proof. The statement follows from the fact that I_n is a nice interval in the sense of Martens [26]. \square

Lemma 11. *There is $K = K(\ell) > 0$ such that if $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell > 1$ then for every $x \in I_1 \setminus I_{n+1}$, $n \geq 3$, and k the first positive integer such that $f^k(x) \in I_{n+1}$ there is an interval J to which x belongs such that $f^k(J) = I_{n+1}$ and*

$$\frac{Df^j(y)}{Df^j(z)} \leq K,$$

for all $y, z \in J$ and $j = 0, 1, 2, \dots, k$. The constant K is given by

$$K = \left(\frac{1 + \alpha}{\alpha} \right)^2,$$

where $\alpha = \alpha(\ell) > 0$ is the constant given in Theorem 9.

Proof. By Lemma 10 there is an interval T such that $x \in T$ and $f^k|_T$ is a diffeomorphism onto I_n . Let J be the pre-image of I_{n+1} under $f^k|_T$. By Theorem 9, I_{n+1} is α -well inside I_n . Now we apply the Macroscopic Koebe Principle to $h = f^{k-j}|_f^j(T)$ and conclude that $f^j(J)$ is α -well inside $f^j(T)$, for every $j = 0, 1, \dots, k$. Finally the distortion control of (4) applied to $h = f^j|_T$ implies the lemma. \square

4. The Fibonacci combinatorics

The domain of ϕ_n has a connected component M_{n+1} , called *post critical domain*, which contains $\phi_n(c)$. Following the terminology in [9], a branch of ϕ_n which is a restriction of a branch of ϕ_{n-1} is called an *immediate branch*.

Lemma 12. *If $f \in \mathcal{C}_d$ has the Fibonacci combinatorics then the following properties hold:*

- (1) *The post critical domain of ϕ_n is an immediate branch. In particular $\phi_{n+1}(c) = \phi_{n-1} \circ \phi_n(c)$, for all $n \geq 2$.*
- (2) *Two consecutive post critical domains M_n and M_{n+1} (and in particular $\phi_{n-1}(c)$ and $\phi_n(c)$), $n \geq 1$, are on opposite sides of c inside I_1 .*

Proof. Due to the fact that the critical branch of ϕ_n is a homeomorphism from I_{n+1} onto I_n , for all $n \geq 1$, ϕ_n has exactly one immediate branch, for every $n \geq 2$. As $d \geq 2$ the first return map ϕ_1 has exactly d branches and all these branches are restrictions of f . Now ϕ_2 has its critical branch with return time equal to 2 (since by the hypothesis $s_2 = 2$), its immediate branch with return time equal to 1 and any of its branches which is not an immediate branch with return time at least equal to 2. Then, as $s_3 = 3$, $\phi_3(c) = \phi_1 \circ \phi_2(c)$. By induction, assume, for $n \geq 2$, that the immediate branch of ϕ_n has return time s_{n-1} and that the return time of its others branches (including the critical branch) are at least s_n . So, as $s_{n+1} = s_{n-1} + s_n$, the post critical branch of ϕ_n must be its immediate branch. Also, except for its immediate and critical branches, all the other branches of ϕ_{n+1} will have return time greater or equal than $2s_n$, which is greater than s_{n+1} , and the induction follows.

The second statement follows from the fact that the branches of ϕ_n preserve orientation and the fact that its post critical branch is an immediate branch. \square

The following lemma illustrates the strong recurrence property of Fibonacci critical coverings.

Proposition 13. *If $g \in \mathcal{C}_d$ has a sequence of return times $s_1, s_2, s_3, s_4, \dots$ smaller (with respect to the lexicographic order) than the Fibonacci sequence $1, 2, 3, 5, \dots$ then g has central returns.*

Proof. We argue that if $g \in \mathcal{C}_d$ has a sequence of return times $s_1, s_2, s_3, s_4, \dots$ smaller (with respect to the lexicographic order) than the Fibonacci sequence $1, 2, 3, 5, \dots$ then g has central returns. Among the branches of the first return map of a critical covering the immediate branch has the smallest possible return time. So, if a critical covering has no central returns its sequence of critical returns is at least as big as the Fibonacci sequence $1, 2, 3, 5, \dots$. \square

The following lemma is one of the applications of Lemma 11 in the Fibonacci case.

Lemma 14. *Let $K = K(\ell) > 0$ be the constant given in Lemma 11. If $f \in \mathcal{C}_d$ has the Fibonacci combinatorics and order $\ell > 1$ then*

$$K^{-1} < \frac{Df^j(f^{s_{n+1}}(f(c)))}{Df^j(f(c))} < K,$$

for all $n \geq 4$ and for all $j = 0, 1, \dots, s_n - 1$.

Proof. As $\phi_n|_{I_{n+1}}$ has c as its only critical point and c is not periodic, then the interval $J = f(I_{n+1})$ is diffeomorphically mapped onto I_n by f^{s_n-1} , it is disjoint from I_n , it is the first entry map into I_n and it contains the points $f(c)$ and $f(f^{s_{n+1}}(c))$, since c and $f^{s_{n+1}}(c)$ are in I_{n+1} . Therefore the lemma follows directly from Lemma 11 (applied to I_n). \square

5. Extensibility around the critical value

From now on it will be always assumed in the statements that we are considering a function $f \in \mathcal{C}_d$ with the Fibonacci combinatorics.

We define $f^m(c) = c_m$, for $m \geq 1$, the points of the critical orbit. Among them there are the points c_{s_n} , that belong to $M_{n+1} \subset I_n$ and are in alternate sides of c , accordingly to the second statement of Lemma 12. We also define, for each $n \geq 1$, the point $z_n = (\phi_n|_{I_{n+1}})^{-1}(c)$. The points z_n will help us to find the right cross-ratios in order to evaluate $Df^{s_n-1}(f(c))$.

Lemma 15 (Definition and placement of z_n). *Let $z_n = (\phi_n|_{I_{n+1}})^{-1}(c)$, for every $n \geq 1$. Then z_n is between $c_{s_{n-1}}$ and $c_{s_{n+1}}$.*

Proof. As $\phi_n(c) = c_{s_n} \in M_{n+1}$ it is easy to see that z_n and c_{s_n} are in opposite sides of c . That z_n is between c and $c_{s_{n-1}}$ follows from the fact that $c_{s_{n-1}} \in M_n$ and $z_n \in I_n$, and that they are both in the same side of c . That $c_{s_{n+1}}$ is between z_n and c follows from the monotonicity of $\phi_n|_{I_{n+1}}$, that send the points $z_n, c_{s_{n+1}}, c$ to $c, c_{s_{n+2}}, c_{s_n}$, in this order, and $c_{s_{n+2}}$ is between c and c_{s_n} . \square

We now identify I_1 with the interval $(0, 1)$ in the real line, in order to consider extendibility properties of the return map $\phi_n|_{I_{n+1}}$ not restricted by the circle topology. Regarding as an interval map, f has d branches in $(0, 1)$, two of them being $\phi_1|_{M_2}$ and $\phi_1|_{I_2}$.

This identification allows that *all results will be valid also in the context of interval maps with d branches*. These maps are in correspondence with circle maps with (in general) d points of discontinuity of the derivative. But these discontinuity points will make no difference in the arguments.

Let g_0 and g_1 be the lifts of f that extend $\phi_1|_{M_2}$ and $\phi_1|_{I_2}$, respectively, to the whole line. Their difference is constant and equal to some integer τ with absolute value smaller than d . More precisely, defining the unitary vector $e_n = \frac{c_{s_n}-c}{|c_{s_n}-c|}$ then

$$g_1 = g_0 + \tau e_1, \tag{5}$$

for some $\tau \in \{1, \dots, d-1\}$.

Therefore if we define by induction $g_n = g_{n-2} \circ g_{n-1}$, for $n \geq 2$, then g_n is the homeomorphic extension of $\phi_n|_{I_{n+1}}$ to the whole line. We claim that

$$g_n \circ g_{n-1} = g_{n-1} \circ g_n + (d-1)\tau e_{n+1}. \tag{6}$$

For $n = 1$ the assertion follows after explicitly developing the equality

$$g_1 \circ g_0 = (g_0 + \tau e_1) \circ (g_1 - \tau e_1).$$

If it is true for n then

$$g_{n+1} \circ g_n = g_{n-1} \circ g_n \circ g_n = (g_n \circ g_{n-1} - (d-1)\tau e_{n+1}) \circ g_n = g_n \circ g_{n+1} + (d-1)\tau e_{n+2}.$$

Another auxiliary point is x_n , a pre-image of one of the boundary points of $(0, 1)$.

Lemma 16 (Definition and placement of x_n). *For $n \geq 1$ let ∂_n be the boundary point of $(0, 1)$ which is opposite to c_{s_n} with respect to c and let $x_n = g_n^{-1}(\partial_n)$. Then*

- (1) x_n is between $c_{s_{n-1}}$ and z_n , for every $n \geq 2$;
- (2) $x_n = g_{n-1}^{-1}(x_{n-2})$, for every $n \geq 3$.

Proof. (1) It is enough to prove that $g_n(x_n)$ is between $g_n(c_{s_{n-1}})$ and $g_n(z_n) = c$. But by (6)

$$g_n(c_{s_{n-1}}) = c_{s_{n+1}} - (d - 1)\tau e_n,$$

that is, $g_n(c_{s_{n-1}})$ is outside $(0, 1)$ and in the same side of ∂_n with respect to c . The first claim follows.

(2) By definition $g_n(x_n) = \partial_n = \partial_{n-2}$, hence $g_{n-2}(g_{n-1}(x_n)) = \partial_{n-2}$. This implies that $g_{n-1}(x_n) = x_{n-2}$, proving the second claim. \square

Lemma 17. For $n \geq 3$, the restriction of g_n to (x_n, z_{n-1}) is a C^1 homeomorphism onto $(\partial_n, c_{s_{n-2}})$ having c as its unique critical point.

Proof. Although the statement is for $n \geq 3$, we first look at g_1 and g_2 . Let $U_1 = I_2$ and $U_2 = g_1^{-1}(M_2)$. Then $g_1|_{U_1}$ is a C^1 homeomorphism onto $(0, 1)$ having c as its unique critical point. The same is true for $g_2|_{U_2}$, since $g_2 = g_0 \circ g_1$ and $g_0|_{M_2}$ is a diffeomorphism onto $(0, 1)$. Notice that $x_1 \in \partial U_1$, $x_2 \in \partial U_2$, and $U_2 \subset U_1$.

Now we let $U_n = (x_n, z_{n-1})$, for every $n \geq 3$. We claim that z_n and x_{n+1} belong to \bar{U}_n , for all $n \geq 1$, hence $U_{n+1} \subset U_n$ for every $n \geq 1$. This assertion immediately follows in the case $n \geq 3$ from the localization Lemmas 15 and 16. Moreover, $z_1 \in U_1 = I_2$ since $z_1 = (\phi_1|_{I_2})^{-1}(c)$ and $z_2 \in U_2 = g_1^{-1}(M_2)$, since $I_3 \subset g_1^{-1}(M_2)$ and $z_2 = (\phi_2|_{I_3})^{-1}(c)$. That $x_2 \in \bar{U}_1$ follows from $x_2 \in \partial U_2$ and $U_2 \subset U_1$. That $x_3 \in \bar{U}_2$ follows from $x_3 = g_2^{-1}(x_1)$ (by the second assertion of Lemma 16), $g_2(U_2) = (0, 1)$ and $x_1 \in (0, 1)$.

We claim that $g_n|_{U_n}$ is a C^1 homeomorphism with unique critical point c for all $n \geq 1$. As for $n = 1, 2$ it is already proved, assume by induction that $n \geq 3$ and that the claim is valid for $n - 1$ and $n - 2$. First $U_n \subset U_{n-1}$ implies that $g_{n-1}|_{U_n}$ is a C^1 homeomorphism with unique critical point c . Moreover, as $z_{n-1} \in \partial U_n$ and $g_{n-1}(z_{n-1}) = c$ then $g_{n-1}(U_n)$ intersects just one component of $U_{n-2} \setminus \{c\}$. But in fact $g_{n-1}(U_n)$ is exactly this component, since $g_{n-1}(x_n) = x_{n-2} \in \partial U_{n-2}$, by the second assertion of Lemma 16. Then g_{n-2} restricted to $g_{n-1}(U_n)$ is a diffeomorphism onto its image and the composition $g_n = g_{n-2} \circ g_{n-1}$, restricted to U_n , is a C^1 homeomorphism onto its image having c as its unique critical point.

As $g_n(x_n) = \partial_n$ and $g_n(z_{n-1}) = g_{n-2}(c) = c_{s_{n-2}}$ the image of $g_n|_{U_n}$ is the interval given in the assertion. \square

It is the following corollary of Lemma 17 that will be explicitly used when estimating the derivative growth using cross-ratio expansion. The notation introduced in the lemma closely follows previous works on this subject, for example [1].

Lemma 18. For $n \geq 2$ let $x_n^f = f(x_n)$, $z_n^f = f(z_n)$, $c_{s_n}^f = f(c_{s_n})$, $c^f = f(c)$ (in this case, x_n, z_n, c_{s_n} and c are all in the closure of I_2 , and f means the continuous extension of the branch $\phi_1|_{I_2}$ to \bar{I}_2). Then, for every $n \geq 3$, f^{s_n-1} diffeomorphically sends (x_n^f, z_{n-1}^f) onto $(\partial_n, c_{s_{n-2}})$. Moreover, the points c^f and $c_{s_{n+1}}^f$ are sent to c_{s_n} and $c_{s_{n+2}}$.

Proof. By Lemma 17, f^{s_n} sends (x_n, z_{n-1}) homeomorphically onto $(\partial_n, c_{s_{n-2}})$ with c as its unique critical point. As c is a critical point for the first iterate, if f^{s_n-1} restricted to (x_n^f, z_{n-1}^f) had a critical point then there would exist $j < s_n$ such that $f^j(c) = c$. This would be a contradiction, since c is not periodic.

Moreover, $f^{s_n-1}(c^f) = f^{s_n}(c) = c_{s_n}$ and

$$f^{s_n-1}(c_{s_{n+1}}^f) = f^{s_n}(c_{s_{n+1}}) = f^{s_n+2}(c) = c_{s_{n+2}}. \quad \square$$

6. Derivative on the critical orbit

We will control the growth of $Df^{s_n}(f(c))$ stated in Theorem 1 using the expansion of suitably chosen (degenerated) cross-ratios, that will allow us to construct difference equations, similarly as was done in [17].

We use the notation introduced in Lemma 18 (following [1]), where $w^f = f(w)$, for every point $w \in I_2$, and define also the distances $d_n := |c_{s_n} - c|$, $d_n^f := |c_{s_n}^f - c^f|$, but just once we use x_n^f to denote $|x_n^f - c|$. We also introduce the parameter

$$\lambda_n = \frac{d_n}{d_{n+2}},$$

that will play an important role in the proof. Since c_{s_n} and $c_{s_{n+2}}$ are in the same side of $\{c\}$, λ_n is always greater than one. But the real bounds of Theorem 9 give more.

Lemma 19. *Let $\alpha > 0$ be as in Theorem 9, $f \in C_d$ with the Fibonacci combinatorics and λ_n defined as above. Then*

$$\lambda_n \geq 1 + \alpha$$

for every $n \geq 2$.

Proof. There is a component R of $I_{n+1} \setminus I_{n+2}$ between $c_{s_{n+2}} \in I_{n+2}$ and $c_{s_n} \in I_n \setminus I_{n+1}$. Theorem 9 implies that $|R| \geq \alpha |I_{n+2}|$. Then

$$d_n \geq R + d_{n+2} \geq \alpha |I_{n+2}| + d_{n+2} \geq (\alpha + 1)d_{n+2}. \quad \square$$

The local approximation given in (1) gives useful estimates involving the d_n^f 's and also the derivative of f at the c_{s_n} 's. One of the estimates also uses Lemma 19.

Lemma 20. *For sufficiently high n , there are numbers $\theta_n^{(1)}$, $\theta_n^{(2)}$ and $\theta_n^{(3)}$ going to 1 as $n \rightarrow \infty$, such that*

$$d_n^f = \theta_n^{(1)} \vartheta d_n^\ell, \tag{7}$$

$$Df(c_{s_n}) = \theta_n^{(2)} \vartheta \ell d_n^{\ell-1}, \tag{8}$$

$$\frac{d_n^f}{d_n^f - d_{n+2}^f} = \theta_n^{(3)} \frac{d_n^\ell}{d_n^\ell - d_{n+2}^\ell} = \theta_n^{(3)} \frac{1}{1 - \lambda_n^{-\ell}}, \tag{9}$$

where ϑ is the constant given in (1).

Proof. Eq. (7) follows directly from (1). From the same equation we get that

$$Df(x) = \vartheta \ell |x - c|^{\ell-1} (1 + \psi_1(x)) \tag{10}$$

for x in a neighborhood of the critical point, where ψ_1 is a continuous function such that $\lim_{x \rightarrow c} \psi_1(x) = 0$, and this proves (8).

Let $\xi(a, x) = \frac{1-x}{1-ax}$, for $x \in [0, (1 + \alpha)^{-1}]$ and $a < 1 + \alpha$, where $\alpha > 0$ is the constant given in Lemma 19. We have

$$\frac{d_n^f}{d_n^f - d_{n+2}^f} = \frac{1}{1 - \frac{d_{n+2}^f}{d_n^f}} = \frac{1}{1 - \frac{\theta_n^{(1)}}{\theta_n^{(1)}} \lambda_n^{-\ell}} = \theta_n^{(3)} \frac{1}{1 - \lambda_n^{-\ell}},$$

where $\theta_n^{(3)} = \xi(\frac{\theta_n^{(1)}}{\theta_n^{(1)}}, \lambda_n^{-\ell})$, which is defined for n sufficiently high, since $\theta_n^{(1)} \rightarrow 1$. As $\xi(a, \cdot)$ uniformly tends to 1 as $a \rightarrow 1$ then $\theta_n^{(3)} \rightarrow 1$. \square

Now taking into account Lemma 18 we apply (2) to the diffeomorphism f^{s_n-1} in the interval (x_n^f, c^f) and obtain a fundamental lower bound for its derivative.

Lemma 21. *For sufficiently high n , there is $K_n^{(1)}$ going to 1 as $n \rightarrow \infty$ such that*

$$Df^{s_n-1}(c_{s_{n+1}}^f) \geq K_n^{(1)} \frac{d_n - d_{n+2}}{d_{n+1}^f} \frac{d_{n-1}^f}{d_{n-1}^f - d_{n+1}^f}.$$

Proof. Let $T_n = (x_n^f, c^f)$ with subintervals $L_n = (x_n^f, c_{s_{n+1}}^f)$, $R_n = (c_{s_{n+1}}^f, c^f)$ and $J_n = \{c_{s_{n+1}}^f\}$ (degenerated), which are sent, under f^{s_n-1} , to the intervals (∂_n, c_{s_n}) , $(\partial_n, c_{s_{n+2}})$, $(c_{s_{n+2}}, c_{s_n})$ and $\{c_{s_{n+2}}\}$, respectively. Eq. (2) yields

$$Df^{s_n-1}(c_{s_{n+1}}^f) \geq \frac{|f^{s_n-1}(L_n)|}{|f^{s_n-1}(T_n)|} \frac{d_n - d_{n+2}}{d_{n+1}^f} \frac{x_n^f}{x_n^f - d_{n+1}^f}.$$

As $|\partial_n - c|$ alternates between two positive values and $c_{s_n} \rightarrow 0$ (since c is recurrent), then it follows that $K_n^{(1)} \equiv |f^{s_n-1}(L_n)|/|f^{s_n-1}(T_n)| \rightarrow 1$ as $n \rightarrow \infty$. Moreover, as $x_n^f < d_{n-1}^f$, by Lemma 16,

$$\frac{x_n^f}{x_n^f - d_{n+1}^f} = \frac{1}{1 - \frac{d_{n+1}^f}{x_n^f}} > \frac{1}{1 - \frac{d_{n+1}^f}{d_{n-1}^f}} = \frac{d_{n-1}^f}{d_{n-1}^f - d_{n+1}^f},$$

and the assertion follows. \square

At this point but only for later use we also give, in the following lemma, an upper bound for $Df^{s_n-1}(c^f)$, using (3).

Lemma 22. For sufficiently high $n \geq 1$,

$$Df^{s_n-1}(c^f) \leq (K_n^{(1)})^{-1} \frac{d_n - d_{n+2}}{d_{n+1}^f} \frac{d_{n-1}^f - d_{n+1}^f}{d_{n-1}^f},$$

where $K_n^{(1)}$ is the same as in Lemma 21.

Proof. Let $T_n = (x_n^f, c^f)$ with subintervals $L_n = (x_n^f, c_{s_{n+1}}^f)$, $J_n = (c_{s_{n+1}}^f, c^f)$ and $R_n = \{c^f\}$ (degenerated), which are sent, under f^{s_n-1} , to the intervals (∂_n, c_{s_n}) , $(\partial_n, c_{s_{n+2}})$, $(c_{s_{n+2}}, c_{s_n})$ and $\{c_{s_n}\}$, respectively. Eq. (3) yields

$$Df^{s_n-1}(c^f) \leq \frac{|f^{s_n-1}(T_n)|}{|f^{s_n-1}(L_n)|} \frac{d_n - d_{n+2}}{d_{n+1}^f} \frac{x_n^f - d_{n+1}^f}{x_n^f}$$

and the result follows similarly as in the proof of Lemma 21. \square

We define the auxiliary parameter

$$q_n \equiv \frac{Df^{s_n-1}(c_{s_{n+1}}^f)}{Df^{s_n-1}(c^f)},$$

which is bounded away from zero and from infinity, by Lemma 14. The next lemma will be the starting point to obtain the exponential growth of $Df^{s_n-1}(c^f)$ with n for $1 < \ell \leq 2$.

Lemma 23. For sufficiently high n , there is $K_n^{(2)} \rightarrow 1$ as $n \rightarrow \infty$ such that

$$\frac{Df^{s_{n+1}}(c^f)^2}{Df^{s_n}(c^f)Df^{s_{n-1}}(c^f)} \geq K_n^{(2)} \ell^2 \lambda_{n-1}^{2-\ell} \frac{q_{n-1}}{q_n} \frac{1 - \lambda_n^{-1}}{1 - \lambda_{n-2}^{-\ell}} \frac{1 - \lambda_{n-1}^{-1}}{1 - \lambda_{n-1}^{-\ell}}. \tag{11}$$

Proof. A straightforward calculation using the chain rule and $s_{n+1} = s_n + s_{n-1}$ shows that the left-hand side of (11) is equal to

$$\frac{q_{n-1}}{q_n} Df^{s_{n-1}-1}(c_{s_n}^f) Df^{s_n-1}(c_{s_{n+1}}^f) \frac{Df(c_{s_{n+1}})^2 Df(c_{s_n})}{Df(c_{s_{n-1}})}.$$

Eq. (8) implies that

$$\frac{Df(c_{s_{n+1}})^2 Df(c_{s_n})}{Df(c_{s_{n-1}})} = \theta_n^{(4)} \vartheta^2 \ell^2 \frac{d_{n+1}^{2\ell-2} d_n^{\ell-1}}{d_{n-1}^{\ell-1}},$$

where $\theta_n^{(4)} \rightarrow 1$ as $n \rightarrow \infty$. Using Lemma 21 for the two remaining factors of this product and also Lemma 20 the Inequality follows, with

$$K_n^{(2)} = \frac{\theta_n^{(4)} \theta_{n-1}^{(3)} \theta_{n-2}^{(3)}}{\theta_{n+1}^{(1)} \theta_n^{(1)}} K_n^{(1)} K_{n-1}^{(1)}. \quad \square$$

Finally, the next lemma will be useful at obtaining finer results on the growth of the derivative along the orbit of the critical value.

Lemma 24. *There is $C > 0$ such that, for sufficiently high n ,*

$$C^{-1}\lambda_{n-1}^{2-\ell} \leq \frac{Df^{s_{n+1}}(c^f)^2}{Df^{s_n}(c^f)Df^{s_{n-1}}(c^f)} \leq C\lambda_{n-1}^{2-\ell}.$$

Proof. The first inequality is a consequence of Lemmas 14 and 19 applied to (11) of Lemma 23. The second inequality can be obtained in two stages: first by following the same arguments as in the proof of Lemma 23, using Lemma 22. And then using Lemmas 14 and 19 again. \square

7. Exponential growth of derivative

7.1. Exponential growth

We aim at proving the exponential growth of $Df^{s_n}(c^f)$ with n , which is the same as finding a positive \liminf for $\frac{1}{n} \log Df^{s_n}(c^f)$. From Lemma 23, there is n_0 such that $n \geq n_0$ implies

$$\left[\frac{Df^{s_{n+1}}(c^f)}{1 - \lambda_n^{-1}} \right]^2 \left[\frac{Df^{s_n}(c^f)}{1 - \lambda_{n-1}^{-1}} \right]^{-1} \left[\frac{Df^{s_{n-1}}(c^f)}{1 - \lambda_{n-2}^{-1}} \right]^{-1} \geq \frac{Q_{n-1}}{Q_n} \sigma_n, \tag{12}$$

where

$$Q_n = q_n(1 - \lambda_n^{-1})$$

and

$$\sigma_n = K_n^{(2)} \ell^2 \lambda_{n-1}^{2-\ell} \frac{1 - \lambda_{n-1}^{-1}}{1 - \lambda_{n-1}^{-\ell}} \frac{1 - \lambda_{n-2}^{-1}}{1 - \lambda_{n-2}^{-\ell}} \tag{13}$$

(for the sake of precision, as $K_n^{(2)}$ is defined only for high n in Lemma 23, let σ_n be the minimum (positive) value such that (12) is true for the finite set of values of n where $K_n^{(2)}$ is not defined). Taking logarithms on both sides and defining $\delta_n = \frac{1}{2} \log Q_n$, $\epsilon_n = \frac{1}{2} \log \sigma_n$ and

$$X_n = \log \left[\frac{Df^{s_n}(c^f)}{1 - \lambda_{n-1}^{-1}} \right],$$

we obtain

$$X_{n+1} - \frac{1}{2}X_n - \frac{1}{2}X_{n-1} \geq \delta_{n-1} - \delta_n + \epsilon_n. \tag{14}$$

The following lemma gives an explicit “solution” of this inequation and gives an inferior limit for $\frac{X_n}{n}$ when the δ_n ’s are bounded.

Lemma 25. *Let $(X_n)_{n \geq 0}$ be a sequence satisfying (14), for arbitrary X_0 and X_1 . Then*

$$X_n \geq \alpha_n X_0 + (1 - \alpha_n) X_1 + \sum_{j=1}^{n-1} \alpha_{n-j} (\delta_{j-1} - \delta_j) + \sum_{j=1}^{n-1} \alpha_{n-j} \epsilon_j, \tag{15}$$

for $n \geq 2$, where $\alpha_0 = 0$, $\alpha_1 = 1$ and, by induction, $\alpha_n = \frac{1}{2}(\alpha_{n-1} + \alpha_{n-2})$, for every $n \geq 2$. If moreover $|\delta_n| \leq c$ for every $n \geq 0$ and $\liminf_n \epsilon_n > -\infty$ then

$$\liminf_n \frac{X_n}{n} \geq \frac{2}{3} \liminf_n \epsilon_n.$$

Proof. The proof of (15) can be done by straightforward induction. Remark that $\alpha_n \rightarrow \frac{2}{3}$ when $n \rightarrow \infty$ and in fact

$$\alpha_n - \frac{2}{3} = (-1)^{n+1} \frac{2}{3} 2^{-n}.$$

In particular the sequence α_n is bounded, implying that the first two terms of the right-hand side of this inequation go to zero when divided by n . The third term can be written as

$$\sum_{j=1}^{n-1} \left(\alpha_{n-j} - \frac{2}{3} \right) (\delta_{j-1} - \delta_j) + \frac{2}{3} \sum_{j=1}^{n-1} \delta_{j-1} - \delta_j.$$

The first sum is bounded by a sum of a finite geometric progression (with ratio $\frac{1}{2}$) and the second is equal to $\frac{2}{3}(\delta_0 - \delta_{n-1})$, hence it is also bounded. Therefore the third term also goes to zero when divided by n . The last term is treated in the same way: it is equal to

$$\sum_{j=1}^{n-1} \left(\alpha_{n-j} - \frac{2}{3} \right) \epsilon_j + \frac{2}{3} \sum_{j=1}^{n-1} \epsilon_j.$$

The \liminf of the first sum divided by n is greater or equal than 0 since the sum is uniformly bounded away from $-\infty$, and the \liminf of the second sum divided by n is greater or equal $\liminf_n \epsilon_n$. \square

Now we use this lemma to obtain a positive lower bound for $\liminf_n \frac{X_n}{n}$. We know that the δ_n 's are bounded, since $\delta_n = \frac{1}{2} \log q_n (1 - \lambda_n^{-1})$, q_n is uniformly bounded away from 0 and from ∞ , by Lemma 14, and $1 - \lambda_n^{-1}$ is smaller than one and also bounded away from 0, by Lemma 19. The next step is to prove that $\liminf_n \epsilon_n > 0$ whenever $\ell \in (1, 2]$.

Lemma 26. *Let $\ell \in (1, 2]$ and take $\epsilon_n = \frac{1}{2} \log \sigma_n$, where σ_n is defined as in (13). Then there is $\epsilon > 0$ such that*

$$\liminf_n \epsilon_n \geq \epsilon.$$

Proof. Let $\psi_\ell : [1, +\infty) \rightarrow \mathbb{R}$ be defined by $\psi_\ell(1) = 1$ and

$$\psi_\ell(t) = \ell \frac{1 - t^{-1}}{1 - t^{-\ell}},$$

for $t > 1$. It is a Calculus' exercise to show that ψ_ℓ is continuous, strictly increasing with $\lim_{t \rightarrow \infty} \psi_\ell(t) = \ell$, and also that $\ell \mapsto \psi_\ell(t_0)$ is strictly increasing for every $t_0 > 1$. With this definition, $\sigma_n = K_n^{(2)} \lambda_{n-1}^{2-\ell} \psi_\ell(\lambda_{n-1}) \psi_\ell(\lambda_{n-2})$ and, as $K_n^{(2)} \rightarrow 1$ it suffices to show that $\lambda_{n-1}^{2-\ell} \psi_\ell(\lambda_{n-1}) \psi_\ell(\lambda_{n-2})$ is uniformly greater than one. We divide the proof into two cases: (i) $1 < \ell < \frac{3}{2}$ and (ii) $\frac{3}{2} \leq \ell \leq 2$. In the first case, we use $\psi_\ell(x) \geq 1$, so that the product is greater or equal than $\lambda_{n-1}^{2-\ell}$, which in turn is greater or equal than $\sqrt{1 + \alpha}$, by Lemma 19. In the second case, the first factor is just assumed to be greater or equal than one (it is equal to one if $\ell = 2$), but $\psi_{\frac{3}{2}}(1 + \alpha)^2$ is a lower bound for the rest of the product, using the properties of ψ_ℓ stated above and also Lemma 19. This proves the lemma with $\epsilon = \min\{\sqrt{1 + \alpha}, \psi_{\frac{3}{2}}(1 + \alpha)^2\}$. \square

Therefore we conclude that

$$\liminf_n \frac{X_n}{n} \geq \frac{2\epsilon}{3},$$

for $\ell \in (1, 2]$, where $\epsilon > 0$ is given by Lemma 26. But once again we use Lemma 19: as $1 - \lambda_{n-1}^{-1}$ is greater or equal than $\frac{\alpha}{1 + \alpha}$ this implies

$$\liminf_n \frac{1}{n} \log Df^{s_n}(c^f) \geq \frac{2\epsilon}{3}$$

and the proof of Theorem 1 is complete.

7.2. The invariant measure

Lemma 27 below shows a classic summability condition on the critical orbit which has been used as a criterion to prove the existence of an absolutely continuous invariant probability measure. We refer to the work [3] and references therein, since it includes the case of maps with inflection critical points. There, in fact, the authors prove the existence with the weaker condition that $Df^j(c^f)$ goes to infinity as $j \rightarrow \infty$.

Lemma 27. *If $f \in \mathcal{C}_d$ is a Fibonacci critical covering with order $\ell \in (1, 2]$ then $\sum_{j=0}^{\infty} Df^j(f(c))^{-\frac{1}{\ell}}$ converges.*

Proof. Define the monotone sequence $H_n = \sum_{j=0}^{s_n-1} Df^j(c^f)^{-\frac{1}{\ell}}$, which is a subsequence of the partial sums of the series in the statement, and note that

$$H_{n+1} = H_n + Df^{s_n}(c^f)^{-\frac{1}{\ell}} \sum_{j=0}^{s_{n-1}-1} Df^j(c_{s_n}^f)^{-\frac{1}{\ell}}.$$

By Lemma 14,

$$H_{n+1} \leq H_n + K Df^{s_n}(c^f)^{-\frac{1}{\ell}} \sum_{j=0}^{s_{n-1}-1} Df^j(c^f)^{-\frac{1}{\ell}} = H_n + K Df^{s_n}(c^f)^{-\frac{1}{\ell}} H_{n-1},$$

and, as $H_{n-1} \leq H_n$, we conclude that

$$H_{n+1} \leq \prod_{k=1}^n (1 + K Df^{s_k}(c^f)^{-\frac{1}{\ell}}).$$

The exponential growth of Theorem 1 shows that H_n is uniformly bounded, hence the sum of the statement converges. \square

7.3. Exponential decay of geometry

We decompose the proof into three parts, since the first two will be used in Section 8.

Lemma 28. *There is n_0 such that*

$$2^{-\ell-1} \vartheta |I_n|^\ell \leq |f(I_n)| \leq 4\vartheta |I_n|^\ell$$

for all $n \geq n_0$, where $\vartheta > 0$ is the constant of (1). Moreover, for every interval $J \subset I_n \setminus \{c\}$,

$$|f(J)| \geq \frac{\vartheta}{2} |J|^\ell.$$

Proof. Write $I_n = (c - a, c + b)$ for positive a and b . As in the proof of (7),

$$\frac{\vartheta}{2} (a^\ell + b^\ell) \leq |f(I_n)| \leq 2\vartheta (a^\ell + b^\ell)$$

for large n . But a or b is greater or equal than $\frac{|I_n|}{2}$, and from this it follows the first inequality of the first expression. On the other hand, both a and b are smaller than $|I_n|$, and this implies the second inequality.

Now write $J = (c + a, c + b)$, with $b > a > 0$ (without loss of generality). Then, using (10) and n big,

$$|f(J)| \geq \frac{\vartheta \ell}{2} \int_a^b |x|^{\ell-1} dx \geq \frac{\vartheta \ell}{2} \int_0^{b-a} |u|^{\ell-1} du = \frac{\vartheta}{2} |J|^\ell. \quad \square$$

Lemma 29. Let $K > 0$ be the constant of Lemma 11, ϑ the constant of (1). There is n_0 such that

$$\frac{K^{-1}\vartheta^{-1}}{4} \frac{|I_n|}{|I_{n+1}|^\ell} \leq Df^{s_n-1}(c^f) \leq 2^{\ell+1} K \vartheta^{-1} \frac{|I_n|}{|I_{n+1}|^\ell}$$

for all $n \geq n_0$.

Proof. As f^{s_n-1} sends the interval $f(I_{n+1})$ diffeomorphically onto I_n with bounded distortion, by Lemmas 10 and 11, it suffices to combine this information with Lemma 28. \square

Proof of Theorem 3. Let n_0 be given by Lemma 29 and such that $Df(c_{s_n}) \leq 2\vartheta \ell d_n^{\ell-1}$, given by (8). By the Chain Rule

$$|Df^{s_n}(c^f)| \leq 2^{\ell+2} K \ell \frac{|I_n|}{|I_{n+1}|^\ell} d_n^{\ell-1},$$

where $K > 0$ is the constant of Lemma 11. As $f^{s_n}(c) = c_{s_n} \in I_n$, then $d_n^{\ell-1}|I_n| \leq |I_n|^\ell$. Then we get

$$Df^{s_n}(c^f) \leq 2^{\ell+2} K \ell \left(\frac{|I_n|}{|I_{n+1}|} \right)^\ell. \tag{16}$$

As the left-hand side grows exponentially, by Theorem 1, then the exponential decay of geometry of Theorem 3 follows. \square

We can also prove the exponential decay of the ratio $\frac{|M_{n+1}|}{|I_n|}$, which follows from the following lemma combined with Theorem 3.

Lemma 30. If n is big enough then

$$\frac{|I_n|}{|I_{n+1}|} \leq 8K \left(\frac{|I_{n+1}|}{|M_{n+2}|} \right)^\ell, \tag{17}$$

where $K > 0$ is the constant of Lemma 11.

Proof. Observe that the intervals $f(M_{n+2}) \subset f(I_{n+1})$ are mapped onto the intervals $I_{n+1} \subset I_n$ by f^{s_n-1} , with bounded distortion, see the proof of Theorem 3. This shows that

$$\frac{|I_n|}{|I_{n+1}|} \leq K \frac{|f(I_{n+1})|}{|f(M_{n+2})|},$$

K is the constant of Lemma 11. By Lemma 28, if n is big then

$$|f(I_{n+1})| \leq 4\vartheta |I_{n+1}|^\ell,$$

where ϑ is the constant of (1). On the other hand $|f(M_{n+2})| \geq \frac{\vartheta}{2} |M_{n+2}|^\ell$, by Lemma 28, and the result follows. \square

7.4. Hausdorff dimension

The critical ω -limit set $\omega(c)$ is a subset of

$$\Lambda = \bigcap_{n=1}^{\infty} \Lambda_n,$$

where

$$\Lambda_n = \left(\bigcup_{i=0}^{s_n-1} f^i(I_{n+1}) \right) \cup \left(\bigcup_{j=0}^{s_n-1} f^j(M_{n+1}) \right).$$

This, together with the fact that a Fibonacci critical covering $f \in \mathcal{C}_d$ has no wandering intervals imply that Λ is a minimal invariant Cantor set and $\omega(c) = \Lambda$.

The components of Λ_n are a covering of $\omega(c)$, for each $n \geq 1$. On the other hand, the intervals $f^i(I_{n+1})$ with $1 \leq i < s_n$ and $f^j(M_{n+1})$ with $1 \leq j < s_{n-1}$ are connected components of the domain of the first entry map \mathcal{E}_n to I_n . Moreover, $f^i(I_{n+1})$ contains $f^i(I_{n+2})$ and $f^i(M_{n+2})$, while $f^j(M_{n+1})$ contains $f^{j+s_n}(I_{n+2})$. The intervals $f^i(M_{n+2})$ and $f^{j+s_n}(I_{n+2})$ are mapped by \mathcal{E}_n onto I_{n+1} while the interval $f^i(I_{n+2})$ is mapped by \mathcal{E}_n into M_{n+1} .

Since the branches of \mathcal{E}_n have uniformly bounded distortion, by Lemmas 10 and 11, and using Theorem 3, we get that the size of the intervals in the union Λ_n decay at least as a^{n^2} , for all $n \geq 1$ and some $a < 1$. The number of these intervals grows at most as 2^n and then we conclude that the Hausdorff dimension of $\omega(c)$ is zero.

8. Further results

In [22] it was proved a stronger result on *a priori* bounds involved in the Fibonacci geometry which will be the basis to finer estimates for the derivative growth and the decay of geometry. Let G_{n+1}^M, G_{n+1}^0 and G_{n+1}^I be the three connected components of $I_n \setminus (I_{n+1} \cup M_{n+1})$, where G_{n+1}^0 is the component between I_{n+1} and M_{n+1} and G_{n+1}^M is the lateral component adjacent to M_{n+1} . Let

$$r_n = \max \left\{ \frac{|I_n|}{|G_{n+1}^M|}, \frac{|I_n|}{|G_{n+1}^0|}, \frac{|I_n|}{|G_{n+1}^I|} \right\}.$$

Lemma 31 (Levin–Świątek). *For every $\ell > 1$ there is $r = r(\ell) > 0$ such that*

$$\limsup_n r_n \leq r,$$

for every $f \in \mathcal{C}_d$ with the Fibonacci combinatorics and order ℓ . In particular, for every such f , there is $R = R(f)$ such that $r_n \leq R$ for all $n \geq 1$.

In particular, Theorem 3 and Lemma 30 imply that I_{n+1} and M_{n+1} exponentially decrease not only with respect to I_n but also with respect to these components.

A basic consequence is that the distance of c_{s_n} to c , that we called d_n , is comparable to $|I_n|$. As $c_{s_n} \in I_n$ we already know that $d_n \leq |I_n|$. The following lemma states the opposite inequality, up to a constant.

Lemma 32. *Let $f \in \mathcal{C}_d$ have the Fibonacci combinatorics and order $\ell > 1$. Then there is n_0 such that*

$$d_n \geq \frac{|I_n|}{2r},$$

for every $n \geq n_0$, where $r = r(\ell)$ is given by Lemma 31.

Proof. As $c_{s_n} \in M_{n+1}$ and $c \in |I_{n+1}|$ then $d_n > |G_{n+1}^0|$. By Lemma 31, there is n_0 such that $r_n \leq 2r$ for every $n \geq n_0$, in particular $|I_n| \leq 2r|G_{n+1}^0|$. \square

This lemma, in turn, immediately allows one to relate ratios between d_n 's and ratios between $|I_n|$'s. For example,

$$(2r)^{-1} \frac{d_n}{d_{n+1}} \leq \frac{|I_n|}{|I_{n+1}|} \leq 2r \frac{d_n}{d_{n+1}} \tag{18}$$

for large n (or with another f -dependent constant for any $n \geq 1$).

Moreover, Lemma 32 gives, for large n ,

$$Df(c_{s_n}) \geq \frac{\vartheta}{2} \ell \left(\frac{|I_n|}{2r} \right)^{\ell-1},$$

implying, by Lemma 29, that there is a constant $C_1 > 0$ such that

$$Df^{s_n}(c^f) \geq C_1^{-1} \left(\frac{|I_n|}{|I_{n+1}|} \right)^\ell, \tag{19}$$

which is the counterpart of (16). This inequation, together with (16) and (18) gives a constant $C_2 > 0$ such that for large n

$$C_2^{-1} \left(\frac{d_n}{d_{n+1}} \right)^\ell \leq Df^{s_n}(c^f) \leq C_2 \left(\frac{d_n}{d_{n+1}} \right)^\ell. \quad (20)$$

This, in turn, implies the existence of a constant $C_3 > 0$ such that

$$-\frac{C_3}{\ell} \leq \log \lambda_n - \frac{1}{\ell} \log Df^{s_n}(c^f) - \frac{1}{\ell} \log Df^{s_{n+1}}(c^f) \leq \frac{C_3}{\ell}. \quad (21)$$

If we define $X_n = \log Df^{s_n}(c^f)$ this, together with Lemma 24, gives a constant $C_4 > 0$ such that

$$-C_4 \leq X_{n+1} - \frac{1}{\ell}(X_{n-1} + X_n) \leq C_4, \quad (22)$$

for large n . With this inequation we will derive precise estimates on the growth of the derivative on the critical orbit. We start by the stating the following lemma, which can be easily proved by induction.

Lemma 33. *Let $(X_n)_n$ be a sequence satisfying*

$$X_{n+1} \leq \frac{1}{\ell}(X_{n-1} + X_n) + \Delta, \quad (23)$$

for $\Delta \in \mathbb{R}$, and let $(Y_n)_n$ be another sequence satisfying $Y_0 = X_0$, $Y_1 = X_1$ and $Y_{n+1} = \frac{1}{\ell}(Y_n + Y_{n-1}) + \Delta$. Then

$$X_n \leq Y_n,$$

for all $n \geq 0$. Similarly, if the inequality in (23) is inverted then $X_n \geq Y_n$.

Hence, estimating the growth of Y_n gives estimates about the growth of X_n . The sequence Y_n , in turn, can be studied by the two-dimensional affine iteration $(u_n, v_n) \mapsto (v_n, \frac{1}{\ell}(u_n + v_n) + \Delta)$, by making $u_n = Y_{n-1}$ and $v_n = Y_n$. If $\ell \neq 2$ this iteration has the unique fixed point (Y_*, Y_*) , where

$$Y_* = \frac{-\Delta}{\frac{2}{\ell} - 1},$$

and eigenvalues

$$\mu_{\pm} = \mu_{\pm}(\ell) = \frac{1 \pm \sqrt{1 + 4\ell}}{2\ell},$$

where the eigenvectors associated to μ_{\pm} are multiples of $(1, \mu_{\pm})$. The eigenvalue μ_- is negative with absolute value smaller than one, for every $\ell > 1$, giving a contractive direction. On the other hand, μ_+ decreases with ℓ from the golden number $\omega = \frac{1+\sqrt{5}}{2}$, when $\ell = 1$, to zero, as $\ell \rightarrow \infty$, being greater than 1 for $\ell < 2$ and smaller than one for $\ell > 2$. This gives an expanding or contracting direction, for $\ell \in (1, 2)$ and $\ell > 2$, respectively. From this it is easy to conclude that:

- (1) if $\ell \in (1, 2)$ and if Y_0 and Y_1 are both greater than Y_* then $Y_n > 0$ for large n and $\lim_n \frac{1}{n} \log Y_n = \mu_+$;
- (2) if $\ell > 2$ then $|Y_n|$ is bounded.

In the case $\ell = 2$ the recurrence $Y_{n+1} = \frac{1}{2}(Y_n + Y_{n-1}) + \Delta$ has the explicit solution $Y_n = \frac{2\Delta}{3}(n-1) +$ (a converging term), arguing as in the proof of Lemma 25. From these considerations, we derive the conclusions in the three cases: $1 < \ell < 2$, $\ell = 2$, and $\ell > 2$.

Proof of Theorem 5. If $X_n = \log Df^{s_n}(c^f)$ then the sequence $(X_n)_n$ satisfies (22), for large n and some (universal) $C_4 > 0$. Let $Y_* = \frac{C_4}{\frac{2}{\ell} - 1}$, which is positive, by the hypothesis. By Theorem 1, X_n grows at least linearly, then there is some (perhaps) larger n_0 such that X_{n_0} and X_{n_0+1} are both greater than Y_* . Now let $(Y_n)_n$ be a sequence satisfying

$Y_0 = X_{n_0}$, $Y_1 = X_{n_0+1}$ and $Y_{n+1} = \frac{1}{\ell}(Y_{n-1} + Y_n) - C_4$. Then, by Lemma 33, $X_n \geq Y_{n-n_0}$ for every $n \geq n_0$. By the considerations preceding this proof, $\liminf_n \frac{1}{n} \log X_n \geq \mu_+(\ell) = \mu$.

On the other hand, if we redefine $(Y_n)_n$ in such a way that $Y_{n+1} = \frac{1}{\ell}(Y_{n-1} + Y_n) + C_4$ then we conclude that $\limsup_n \frac{1}{n} \log X_n \leq \mu_+(\ell) = \mu$.

That the geometry decays super exponentially immediately follows from (16). \square

Proof of Theorem 6. Let $k \geq 1$ and n_0 be such that $s_{n_0} \leq k < s_{n_0+1}$. If $k = s_{n_0}$ then $Df^k(c^f) = Df^{s_{n_0}}(c^f)$, otherwise

$$Df^k(c^f) = Df^{s_{n_0}}(c^f) Df^{k-s_{n_0}}(c_{s_{n_0}}^f).$$

Let $k_1 = k - s_{n_0}$. As $k_1 < s_{n_0+1} - s_{n_0} = s_{n_0-1}$ then we can apply Lemma 11 to obtain

$$\left(1 + \frac{1}{\alpha_{n_0}}\right)^{-2} \leq \frac{Df^{k_1}(c_{s_{n_0}}^f)}{Df^{k_1}(c^f)} \leq \left(1 + \frac{1}{\alpha_{n_0}}\right)^2,$$

where α_{n_0} is such that I_{n_0-1} is α_{n_0} -well inside I_{n_0-2} . Now let n_1 such that $s_{n_1} \leq k_1 < s_{n_1+1}$ (certainly $n_1 \leq n_0 - 2$, since $k_1 < s_{n_0-1}$, see above). Then either $k_1 = s_{n_1}$ and then we stop or

$$Df^{k_1}(c^f) = Df^{s_{n_1}}(c^f) Df^{k_2}(c_{s_{n_1}}^f)$$

and

$$\left(1 + \frac{1}{\alpha_{n_0}}\right)^{-2} \leq \frac{Df^{k_1}(c_{s_{n_0}}^f)}{Df^{k_1}(c^f)} \leq \left(1 + \frac{1}{\alpha_{n_0}}\right)^2,$$

where α_{n_1} is such that I_{n_1-1} is α_{n_1} -well inside I_{n_1-2} . Following this procedure, by induction, there will be some $j \geq 0$ such that $k_j = s_{n_j}$. Consequently, $Df^k(c^f)$ and

$$Df^{s_{n_0}}(c^f) Df^{s_{n_1}}(c^f) \dots Df^{s_{n_j}}(c^f)$$

will differ by a multiplicative constant

$$\prod_{i=0}^j \left(1 + \frac{1}{\alpha_{n_i}}\right)^2.$$

This constant is uniformly bounded (independently of k), since the α_{n_i} 's exponentially grow with i , by Theorem 3 and Lemma 31. If $M > 0$ is such a constant, then

$$-\log M \leq \log Df^k(c^f) - \log Df^{s_{n_0}}(c^f) \sum_{i=0}^j \frac{\log Df^{s_{n_i}}(c^f)}{\log Df^{s_{n_0}}(c^f)} \leq \log M.$$

By Theorem 5, $\log Df^{s_n}(c^f)$ is exponential with n , then for large k the sum in the last expression is positive and uniformly bounded from zero and infinity.

One of the inequations then gives

$$\limsup_k \frac{\log \log Df^k(c^f)}{\log k} \leq \limsup_k \frac{\log \log Df^{s_{n_0}}(c^f)}{\log k}.$$

But

$$1 \leq \frac{k}{s_{n_0}} < \frac{s_{n_0+1}}{s_{n_0}}$$

and this last quotient is bounded (it approaches the golden number ω), then k and s_{n_0} are comparable up to a multiplicative constant. Moreover, s_{n_0} is comparable to ω^{n_0} . These considerations imply that

$$\limsup_k \frac{\log \log Df^k(c^f)}{\log k} \leq \limsup_{n_0} \frac{\log \log Df^{s_{n_0}}(c^f)}{n_0 \log \omega} = \frac{\log \mu}{\log \omega}.$$

By a similar reasoning, we obtain $\liminf_k \frac{\log \log Df^k(c^f)}{\log k} \geq \frac{\log \mu}{\log \omega}$ and the theorem is proved. \square

Proof of Theorem 8. We know that $X_n = \log Df^{s_n}(c^f)$ satisfies (22), and by Lemma 33 that X_n is bounded from above by Y_n , where the sequence $(Y_n)_n$ satisfies the equality $Y_{n+1} = \frac{1}{\ell}(Y_n + Y_{n-1}) + C_4$. As we concluded that Y_n is bounded then X_n is bounded, and the conclusion follows.

That the geometry is bounded follows from (19). \square

Proof of Theorem 7. If $X_n = \log Df^{s_n}(c^f)$ then $X_{n+1} \leq \frac{1}{2}(X_n + X_{n-1}) + C_4$, accordingly to (22). Once again we conclude that X_n is bounded by Y_n satisfying the corresponding equality and, as remarked above, Y_n grows linearly with n . Then X_n cannot grow faster than linearly and the theorem follows. \square

Acknowledgement

The second author would like to thank the Institute of Mathematics and Statistics of the University of São Paulo for the hospitality during his visit to execute this work.

References

- [1] H. Bruin, G. Keller, T. Nowicki, S. van Strien, Wild Cantor attractors exist, *Ann. of Math. (2)* 143 (1996) 97–130.
- [2] H. Bruin, S. van Strien, Existence of absolutely continuous invariant probability measures for multimodal maps, in: *Global Analysis of Dynamical Systems*, Inst. Phys., Bristol, 2001, pp. 433–447, 17 (4) (2004) 749–782 (electronic).
- [3] H. Bruin, J.R. Letelier, W. Shen, S. van Strien, Large derivatives, backward contraction and invariant densities for interval maps, *Invent. Math.* 172 (3) (2008) 509–533.
- [4] P. Collet, J.P. Eckmann, *Iterated Maps on the Interval as Dynamical Systems*, Prog. Phys., vol. 1, Birkhäuser Boston, Massachusetts, 1980.
- [5] P. Collet, J.P. Eckmann, Positive Liapunov exponents and absolute continuity for maps of the interval, *Ergodic Theory Dynam. Systems* 3 (1) (1983) 13–46.
- [6] W. de Melo, S. van Strien, *One-dimensional Dynamics*, Springer-Verlag, Berlin, 1993.
- [7] J. Graczyk, G. Świątek, Induced expansions for quadratic polynomials, *Ann. Sci. École Norm. Sup. (4)* (1996) 399–482.
- [8] J. Graczyk, G. Świątek, Generic hyperbolicity in the logistic family, *Ann. of Math. (2)* (1997) 1–52.
- [9] J. Graczyk, G. Świątek, The Real Fatou Conjecture, *Ann. of Math. Stud.*, vol. 144, Princeton University Press, Princeton, 1998.
- [10] J. Graczyk, D. Sands, G. Świątek, Metric attractors for smooth unimodal maps, *Ann. of Math. (2)* 159 (2) (2004) 725–740.
- [11] J. Graczyk, D. Sands, G. Świątek, Decay of geometry for unimodal maps: Negative Schwarzian case, *Ann. of Math. (2)* 161 (2) (2005) 613–677.
- [12] J. Guckenheimer, Sensitive dependence to initial conditions for one-dimensional maps, *Comm. Math. Phys.* 70 (2) (1979) 133–160.
- [13] J. Guckenheimer, S. Johnson, Distortion of S-unimodal maps, *Ann. of Math. (2)* 132 (1) (1990) 71–130.
- [14] F. Hofbauer, G. Keller, Some remarks on recent results about S-unimodal maps, *Ann. Inst. H. Poincaré Phys. Théor.* 53 (4) (1990) 413–425.
- [15] M.V. Jakobson, Absolutely continuous invariant measures for one-parameter families of one-dimensional maps, *Comm. Math. Phys.* 81 (1) (1981) 39–88.
- [16] M.V. Jakobson, G. Świątek, Metric properties of non-renormalizable S-unimodal maps I. Induced expansion and invariant measures, *Ergodic Theory Dynam. Systems* 14 (4) (1994) 721–755.
- [17] G. Keller, T. Nowicki, Fibonacci maps re(al)visited, *Ergodic Theory Dynam. Systems* 15 (1) (1995) 99–120.
- [18] K. Krzyżewski, W. Slenk, On invariant measures for expanding differentiable mappings, *Studia Math.* 3 (1969) 83–92.
- [19] A. Lasota, J.A. Yorke, On the existence of invariant measures for piecewise monotonic transformations, *Trans. Amer. Math. Soc.* 186 (1974) 481–488, (1973).
- [20] G. Levin, Bounds for maps of the interval with one reflecting critical point I, *Fund. Math.* 157 (2–3) (1998) 287–298.
- [21] G. Levin, S. van Strien, Bounds for maps of an interval with one critical point of inflection type. II, *Invent. Math.* 141 (2) (2000) 399–465.
- [22] G. Levin, G. Świątek, Universality of critical circle covers, *Comm. Math. Phys.* 228 (2002) 371–399.
- [23] M. Lyubich, Combinatorics, geometry and attractors of quasi-quadratic maps, *Ann. of Math. (2)* 140 (2) (1994) 347–404.
- [24] M. Lyubich, J. Milnor, The Fibonacci unimodal map, *J. Amer. Math. Soc.* 6 (2) (1993) 425–457.
- [25] M. Lyubich, Dynamics of quadratic polynomials. I, II, *Acta Math.* 178 (2) (1997) 185–247, 247–297.
- [26] M. Martens, Distortion results and invariant Cantor sets of unimodal maps, *Ergodic Theory Dynam. Systems* 14 (2) (1994) 331–349.
- [27] M. Martens, T. Nowicki, Invariant measures for typical quadratic maps, in: *Géométrie complexe et systèmes dynamiques*, Orsay, 1995, Astérisque 261 (2000) 239–252, xiii.
- [28] J. Milnor, On the concept of attractor, *Comm. Math. Phys.* 99 (2) (1985) 177–195.
- [29] M. Misiurewicz, Absolutely continuous measures for certain maps of an interval, *Inst. Hautes Études Sci. Publ. Math.* 53 (1981) 17–51.
- [30] M. Misiurewicz, A. Rodrigues, Double standard maps, *Comm. Math. Phys.* 273 (1) (2007) 37–65.
- [31] T. Nowicki, S. van Strien, Invariant measures exist under a summability condition for unimodal maps, *Invent. Math.* 105 (1) (1991) 123–136.
- [32] D. Ruelle, Applications conservant une mesure absolument continue par rapport à dx sur $[0, 1]$, *Comm. Math. Phys.* 55 (1) (1977) 47–51.
- [33] W. Shen, Decay of geometry for unimodal maps: An elementary proof, *Ann. of Math. (2)* 163 (2) (2006) 383–404.

- [34] S. van Strien, Hyperbolicity and invariant measures for general C^2 interval maps satisfying the Misiurewicz condition, *Comm. Math. Phys.* 128 (3) (1990) 437–495.
- [35] S. van Strien, E. Vargas, Real bounds, ergodicity and negative Schwarzian for multimodal maps, *J. Amer. Math. Soc.* 17 (4) (2004) 749–782 (electronic).