

Global existence for degenerate quadratic reaction–diffusion systems

M. Pierre, R. Texier-Picard *

ENS Cachan Bretagne, IRMAR, CNRS, EUB, av. Robert Schuman, F-35170 Bruz, France

Received 26 March 2008; accepted 27 June 2008

Available online 25 July 2008

Abstract

We consider a class of degenerate reaction–diffusion systems with quadratic nonlinearity and diffusion only in the vertical direction. Such systems can appear in the modeling of photochemical generation and atmospheric dispersion of pollutants. The diffusion coefficients are different for all equations. We study global existence of solutions.

© 2008 Elsevier Masson SAS. All rights reserved.

Résumé

Nous considérons une classe de systèmes dégénérés de réaction–diffusion avec une non-linéarité quadratique et avec diffusion uniquement dans la direction verticale. De tels systèmes peuvent apparaître dans la modélisation de la synthèse de polluants par réactions photochimiques et de leur dispersion atmosphérique. Les coefficients de diffusion sont différents pour chaque équation. Nous étudions l'existence globale de solutions.

© 2008 Elsevier Masson SAS. All rights reserved.

MSC: 35K45; 35K50; 35K57; 35K65; 86A10

Keywords: Reaction–diffusion systems; Degenerate parabolic system; Global existence; Atmosphere modelling; Quadratic nonlinearity

1. Introduction

The goal of this paper is to analyze global existence in time of solutions to reaction–diffusion systems of the type considered in [4] and combined from various models in [21,9,22] and which describe, in particular, the atmospheric dispersion of ozone and other photochemically generated pollutants. Three main coupled difficulties appear in these systems, set in a three-dimensional spatial domain:

- First, the diffusion occurs only in the vertical direction: consequently, the system is ‘degenerate’.
- However, transport of species hold in all directions.
- Even if diffusion occurred in all directions (that is, even if the linear part of the system was strictly parabolic), global existence of solutions would not be obvious because of the structure of the nonlinear reaction terms.

* Corresponding author.

E-mail addresses: michel.pierre@bretagne.ens-cachan.fr (M. Pierre), picard@bretagne.ens-cachan.fr (R. Texier-Picard).

It is proved in [4] that, if the diffusion coefficients are all the same, then global existence of classical solutions does hold. Our goal here is to tackle the more difficult situation *where these coefficients are different from each other* and to give some global existence result.

Besides their interest with respect to the mentioned applications, these systems contain several theoretical questions of interest for lots of other reaction–diffusion systems. Let us describe more precisely an explicit family of these systems, and we will come back to more details on the corresponding difficulties.

Although we will consider more general systems, we focus on the atmospheric diffusion model of [4] to describe and comment the equations. If the geographical area to be considered is represented by a bounded regular open subset Ω of \mathbb{R}^2 , the equations for atmospheric dispersion can be written in a cylindrical domain Q of the form

$$Q = \{(x, y, z, t) \in \mathbb{R}^4, (x, y) \in \Omega, z \in (0, 1), t \in (0, T)\},$$

where $z = 0$ represents the surface of the earth and $z = 1$ the limit of the troposphere.

Let the functions $\phi_i, i = 1, \dots, n$, represent the molecular densities of the different species involved in the photochemical reaction. Then the reaction–advection–diffusion equations can be written in the form

$$\frac{\partial \phi_i}{\partial t} = d_i \frac{\partial^2 \phi_i}{\partial z^2} + \nabla \cdot (\omega \phi_i) + f_i(\phi) + g_i, \tag{1.1}$$

where the velocity field $\omega = (\omega_1, \omega_2, \omega_3)$ models the atmospheric current, d_i is the diffusion coefficient for species i , $f_i(\phi)$ are nonlinear reaction terms, representing the chemistry of the process, and the g_i are source terms. Finally, the notation $\nabla \cdot u$ denotes the divergence of u in the spatial variables (x, y, z) .

Throughout the paper we assume the following

$$\left\{ \begin{array}{l} \forall i = 1, \dots, n, \\ \omega_i : \bar{Q} \rightarrow \mathbb{R} \text{ is continuously differentiable,} \\ f_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ is continuously differentiable,} \\ d_i \in (0, +\infty), \\ g_i \in L^\infty(Q). \end{array} \right. \tag{1.2}$$

In most references mentioned above, an incompressibility condition $\nabla \cdot \omega = 0$ is assumed. In this case, if the functions ϕ_i are regular enough, (1.1) writes

$$\frac{\partial \phi_i}{\partial t} = d_i \frac{\partial^2 \phi_i}{\partial z^2} + \omega_1 \frac{\partial \phi_i}{\partial x} + \omega_2 \frac{\partial \phi_i}{\partial y} + \omega_3 \frac{\partial \phi_i}{\partial z} + f_i(\phi) + g_i. \tag{1.3}$$

We will not need this incompressibility assumption in our analysis. We only need to note that even if it is not satisfied, (1.1) can formally be written in the form (1.3) where the nonlinear term $f_i(\phi)$ is replaced by $f_i(\phi) + (\nabla \cdot \omega)\phi_i$. It can be checked easily that this operation does not affect the assumptions on the nonlinearity presented below. Hence in the following we will always consider Eq. (1.3) instead of (1.1).

Boundary and initial conditions have to be specified. Since the diffusion takes place in the vertical direction, boundary conditions are needed for $z = 0$ and $z = 1$. There is no horizontal diffusion, but conditions on the boundary of Ω are needed where the advection field $-\omega$ is inward (and only there). We denote by $\partial^- Q$ the corresponding part of the boundary of Q . It is defined as follows: let $\nu = (\nu_1, \nu_2) \in \mathbb{R}^2$ be the normal outward unitary vector at a point of the boundary $\partial\Omega$; then

$$\partial^- Q = \{(x, y, z, t) \in \partial\Omega \times (0, 1) \times (0, T); \omega_1 \nu_1 + \omega_2 \nu_2 > 0\}. \tag{1.4}$$

We will choose the same boundary and initial conditions as in the references already mentioned, namely

$$\left. \begin{array}{ll} \phi_i(x, y, z, 0) = \phi_{0i}(x, y, z) & (t = 0), \\ \frac{\partial \phi_i}{\partial z}(x, y, 1, t) = 0 & (z = 1), \\ -\frac{\partial}{\partial z}[d_i \phi_i] + \mu_i \phi_i(x, y, 0, t) = e_i(x, y, 0, t) & (z = 0), \\ \phi_i(x, y, z, t) = \theta_i(x, y, z, t), & (x, y, z, t) \in \partial^- Q. \end{array} \right\} \tag{1.5}$$

The condition at the top of the cylinder is a simple noflux condition, while the condition at the surface of the earth takes into account a smooth nonnegative prescribed flux e_i due to anthropogenic and biogenic emission, and a deposition flux $\mu_i \phi_i$. The coefficient μ_i represents a positive deposition velocity.

Note that our approach could handle as well any kind of “reasonable” boundary conditions. In particular, the following could be chosen instead of (1.5) and may actually be more significant since it takes into account the transport in the flux conditions:

$$\begin{aligned} \left(\frac{\partial}{\partial z} [d_i \phi_i] + \omega_3 \phi_i \right) (x, y, 1, t) &= 0 \quad (\text{at } z = 1), \\ \left(-\frac{\partial}{\partial z} [d_i \phi_i] + \omega_3 \phi_i + \mu_i \phi_i \right) (x, y, 0, t) &= e_i(x, y, 0, t) \quad (\text{at } z = 0). \end{aligned}$$

The following conditions will be throughout assumed on the data:

$$\begin{cases} \forall i = 1, \dots, n, \\ \phi_{0i} \in L^\infty(\Omega \times (0, 1)), \quad e_i \in L^\infty(\Omega \times (0, T)), \quad \theta_i \in L^\infty(\partial^- Q), \\ \phi_{0i} \geq 0, \quad e_i \geq 0, \quad \mu_i \in [0, +\infty], \quad \theta_i \geq 0, \\ \exists \zeta_i \in [H^2 \cap L^\infty](Q) \text{ satisfying the conditions of (1.5).} \end{cases} \tag{1.6}$$

The last regularity condition of (1.6) above could be weakened. It is only needed to reduce more easily the situation to homogeneous boundary conditions, keeping in particular the L^2 -regularity of the various derivatives involved.

Now, we come to the assumptions on the **structure of the nonlinearity** $f = (f_i)_{1 \leq i \leq n}$. We recall in Section 4 the specific 20×20 nonlinearity mentioned in [4] and introduced in [9,22]. We will actually consider more general nonlinearities with the three following properties.

First, we assume that f preserves positivity of the solutions, namely *quasi-positivity*

$$\forall i = 1, \dots, n, \quad \forall \phi \in [0, +\infty)^n \text{ with } \phi_i = 0, \quad f_i(\phi) \geq 0. \tag{1.7}$$

This assumption is quite natural in this context.

Next, as one knows, one cannot expect global existence in time without any structure assumption on the nonlinearities. In [4], a so-called “intermediately quasi-conservative” triangular structure of the system of Section 4 is exploited to prove global existence (assuming also that all the d_i ’s are equal). Namely, five ordered relations are satisfied by the f_i ’s which allow to progressively make L^∞ estimates on the ϕ_i ’s when the d_i ’s are all equal; these five relations are recalled in (4.1).

Here, we will only assume ONE global dissipative property on the nonlinearity f , namely

$$\begin{cases} \exists a_1, \dots, a_n \in \mathbb{R}, \quad \exists b_1, \dots, b_n \in \mathbb{R}, \quad \forall i = 1, \dots, n, \quad a_i > 0, \\ \forall \phi \in \mathbb{R}^+{}^n, \quad \sum_{i=1}^n a_i f_i(\phi) \leq \sum_{i=1}^n b_i \phi_i. \end{cases} \tag{1.8}$$

Obviously, this assumption is satisfied by the nonlinearity in Section 4: add up for instance the five relations of (4.1). We may notice that, when all the coefficients d_i ’s are equal, say $d_i = d$, and when, for instance $\forall i, b_i = 0$ (to simplify), then the function $W = \sum_i a_i \phi_i$ satisfies

$$\frac{\partial W}{\partial t} - d \frac{\partial^2 W}{\partial z^2} + \omega \cdot \nabla W \leq \sum_i a_i g_i.$$

From this and the corresponding boundary conditions on W , we directly deduce an L^∞ -estimate on W and, by nonnegativity of all the ϕ_i ’s, an L^∞ -estimate on all of them. Global existence of solutions then easily follows.

The situation is quite more delicate when the diffusion coefficients are different from each other. It is known that, even for 2×2 systems with good nondegenerate diffusions and one (or even two!) conservative properties of type (1.8), blow up in finite time may occur (see e.g. [19]). Then, two kinds of results may be obtained for nondegenerate diffusions:

- either, we assume more structure on the nonlinearity: for a 2×2 system for instance, we may moreover assume that an L^∞ -estimate is available for the first component; then, an L^∞ -estimate may also be deduced for the second

component (see [15]) and global existence follows. This idea may be generalized to $n \times n$ systems with a so-called “triangular structure” which allow to successively obtain L^∞ estimates on the components, see [16,5]. In these situations, global *classical* solutions are obtained;

- another point of view is to look for global “weak solutions” which may not be in L^∞ for all time. This is the choice made in [18,3] and it leads to global existence results for a quite larger class of nonlinearities. It may be proved in particular that global existence of weak solutions holds for systems with the (only) structure (1.8) and for which the nonlinear growth is at most *quadratic*. A recent work [7] analyzes the size of the possible set of singularities of these weak solutions in a specific quadratic system.

Here, we will restrict our analysis to the (at most) quadratic situation and assume (like for the system of Section 4 and like in many more reaction–diffusion models arising in chemistry, biochemistry, biology, etc.) that

$$\exists k \geq 0, \forall \phi \in R^n, \quad |f(\phi)| \leq k[1 + |\phi|^2]. \quad (1.9)$$

This ‘quadratic assumption’ is consistent with our approach here which is mainly based on an a priori L^2 -estimate. Indeed, as noticed in [19,3], assumption (1.8) together with nonnegativity ensure that the solutions are bounded in $L^2(Q)$ if the diffusions are nondegenerate. Here, the situation is more delicate since diffusion occurs only in one direction. We are not able to prove this L^2 -estimate in general. The considerations made in Section 5 indicate that there may be none in general. Strangely enough, if we assume that ω_1, ω_2 do not depend on the vertical component z (see (2.8)), then this L^2 -estimate does hold. It follows that the nonlinear terms are bounded in L^1 . Since we are able to reduce the problem to a one-dimensional situation, it is then possible to bootstrap the estimates and reach L^∞ . Thus, global existence of classical solutions based on an a priori L^2 -estimate is the main contribution of this paper. Global existence in time is obtained no matter the various values of the d_i ’s in $(0, +\infty)$. Note that assumption (2.8) is discussed in some situations (see e.g. [21, p. 1108] and [6]), when one-dimensional Lagrangian models are considered. We do not know how much it is necessary in our context, but it appears quite naturally in the analysis (see the discussion in Section 5).

Next section is devoted to a (classical) change of variable which reduces the first order part of the differential operator to a single $\partial\Phi/\partial t$ operator. The main point is that, when (2.8) holds, then the diffusion operator in the vertical variable z remains invariant in the new set of variables and the new system reduces to a family of one-dimensional parameter-dependent nondegenerate reaction–diffusion systems to which we can apply more classical approaches. Global existence of solutions is then proved in Section 3. We recall in Section 4 the specific nonlinearity mentioned in [4]. Then the last Section 5 is devoted to some remarks and to some open problems that we find interesting, together with a self-contained presentation of preliminary lemmas.

2. Change of variables

The system (1.3) can be viewed as a parabolic evolution problem, with degeneracies in the horizontal variables x, y , where only advection takes place. Following [4], we can also gather the horizontal advection terms and the time derivative. Our strategy here is not to view (1.3) as a degenerate elliptic system, as in [4], but to use a change of variables to get rid of the degeneracies. In this section we introduce new coordinates and characterize the domain in the new variables, then we write the system of equations in these new variables.

2.1. The domain in the new coordinates

The change of variables introduced in this section is equivalent to a “method of characteristics”, where only the horizontal wind field is taken into account.

Let us define new coordinates (ξ, η) by

$$\begin{cases} x = \beta(\xi, \eta, z, t), \\ y = \gamma(\xi, \eta, z, t) \end{cases}$$

where β, γ are the solutions of the Cauchy problem

$$\frac{\partial \beta}{\partial t} = -\omega_1(\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t), z, t), \quad (2.1)$$

$$\frac{\partial \gamma}{\partial t} = -\omega_2(\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t), z, t), \tag{2.2}$$

$$\beta(\xi, \eta, z, 0) = \xi, \tag{2.3}$$

$$\gamma(\xi, \eta, z, 0) = \eta. \tag{2.4}$$

These solutions are defined for all $t \in (0, T)$ and of class C^1 in all variables, according to the assumptions on ω_1, ω_2 (see (1.2)). In the new variables, the relevant domain is no longer cylindrical. But, if we define

$$D_{z,t} = \{(\xi, \eta) \in \mathbb{R}^2, (\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t)) \in \Omega\},$$

we can assert that, for fixed $(z, t) \in (0, 1) \times (0, T)$, the transformation

$$\Lambda_{z,t} : (\xi, \eta) \mapsto (x = \beta(\xi, \eta, z, t), y = \gamma(\xi, \eta, z, t))$$

defines a diffeomorphism from $D_{z,t}$ to the section Ω of the cylinder Q .

Indeed, for fixed (z, t) , and for $(x, y) \in \Omega$ given, then $(\xi, \eta) = [\Lambda_{z,t}]^{-1}(x, y)$ is uniquely determined by $\xi = \alpha(t), \eta = \delta(t)$, where (α, δ) is the unique solution to the backward Cauchy problem

$$\alpha_s = \omega_1(\alpha(t-s), \delta(t-s), z, t-s),$$

$$\delta_s = \omega_2(\alpha(t-s), \delta(t-s), z, t-s),$$

$$\alpha(0) = x,$$

$$\delta(0) = y.$$

We consider now the whole 4-dimensional domain

$$D = \{(\xi, \eta, z, t) \in \mathbb{R}^2 \times (0, 1) \times (0, T), \text{ s.t. } (\xi, \eta) \in D_{z,t}\}$$

and characterize its boundary. Obviously the faces $z = 0, z = 1, t = 0$ and $t = T$ of Q are simply transported by the diffeomorphism to similar faces of D . To identify the part $\partial^- Q$ of the boundary of Q , we introduce the distance function $d(\cdot)$ to ∂Q . It is regular near $\partial\Omega \times (0, 1) \times (0, T)$ and satisfies $\nabla d = (\nu_1, \nu_2, 0, 0)$ on this part of the boundary. If we denote

$$\tilde{d}(\xi, \eta, z, t) = d(\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t), z, t),$$

then, we have the characterization:

$$[(\xi, \eta, z, t) \in \partial D] \Leftrightarrow [\tilde{d}(\xi, \eta, z, t) = 0].$$

Moreover, using the chain rule and Eqs. (2.1)–(2.2), we obtain

$$\frac{\partial \tilde{d}}{\partial t} = \frac{\partial d}{\partial t} - \omega_1 \frac{\partial d}{\partial x} - \omega_2 \frac{\partial d}{\partial y},$$

so that, on $\partial\Omega \times (0, 1) \times (0, T)$, we have

$$\frac{\partial \tilde{d}}{\partial t} = -\omega_1 \nu_1 - \omega_2 \nu_2. \tag{2.5}$$

According to the definition (1.4) of $\partial^- Q$, we deduce:

Lemma 1. Define the function τ by

$$\forall (\xi, \eta, z, t) \in D, \quad \tau(\xi, \eta, z, t) = \inf\{\sigma \geq 0, \forall s \in [\sigma, t], (\xi, \eta, z, s) \in D\}.$$

Then the following equivalence holds:

$$[(\Lambda_{z,t}(\xi, \eta), z, t) \in \partial^- Q] \Leftrightarrow [(\xi, \eta, z, t) \in \partial^- D],$$

where

$$\partial^- D = \{(\xi, \eta, z, \tau(\xi, \eta, z, t)), (z, t) \in (0, 1) \times (0, T), (\xi, \eta) \in \partial D_{z,t}\}. \tag{2.6}$$

Indeed, we know by (2.5) that

$$\partial^- D = \left\{ (\xi, \eta, z, t) \in \partial D; (z, t) \in (0, 1) \times (0, T), (\xi, \eta) \in \partial D_{z,t}, \frac{\partial \tilde{d}}{\partial t} < 0 \right\}.$$

Since \tilde{d} is constant on ∂D , $\partial \tilde{d} / \partial t$ is the fourth component of the normal outward vector to ∂D . It is negative at a point (ξ, η, z, t) of ∂D if and only if D is “above” this point, that is if $t = \tau(\xi, \eta, z, t)$. Whence the lemma.

2.2. The new system of equations

Now we write the system of Eqs. (1.3)–(1.5) in the new variables ξ, η, z, t . Denoting

$$\phi_i(x, y, z, t) = \phi_i(\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t), z, t) = \Phi_i(\xi, \eta, z, t), \tag{2.7}$$

we differentiate Φ_i with respect to t . Using the chain rule and Eqs. (2.1)–(2.2), we obtain

$$\frac{\partial \Phi_i}{\partial t}(\xi, \eta, z, t) = \left(\frac{\partial \phi_i}{\partial t} - \omega_1 \frac{\partial \phi_i}{\partial x} - \omega_2 \frac{\partial \phi_i}{\partial y} \right) (\beta(\xi, \eta, z, t), \gamma(\xi, \eta, z, t), z, t).$$

We now make the following crucial assumption.

$$\text{The functions } \omega_1, \omega_2 \text{ are independent of the vertical variable } z. \tag{2.8}$$

Then the functions β and γ do not depend either on z and we have the equalities:

$$\begin{aligned} \frac{\partial \Phi_i}{\partial z}(\xi, \eta, z, t) &= \frac{\partial \phi_i}{\partial z}(\beta(\xi, \eta, t), \gamma(\xi, \eta, t), z, t), \\ \frac{\partial^2 \Phi_i}{\partial z^2}(\xi, \eta, z, t) &= \frac{\partial^2 \phi_i}{\partial z^2}(\beta(\xi, \eta, t), \gamma(\xi, \eta, t), z, t). \end{aligned}$$

With these notations we can rewrite Eq. (1.3) in the form

$$\frac{\partial \Phi_i}{\partial t} = d_i \frac{\partial^2 \Phi_i}{\partial z^2} + \omega_3 \frac{\partial \Phi_i}{\partial z} + F_i(\Phi) + G_i, \tag{2.9}$$

where the new nonlinear term is defined by $F_i(\Phi) = f_i(\phi)$ and the function $G_i(\xi, \eta, z, t)$ is defined from g_i as in (2.7).

For simplicity we denote by Λ_t instead of $\Lambda_{z,t}$ the diffeomorphism introduced in Section 2.1 (which is now independent of z). Then the boundary conditions (1.5) can be written in the new coordinates in the form

$$\Phi_i(\xi, \eta, z, 0) = \phi_{0i}(\Lambda_0(\xi, \eta), z), \tag{2.10}$$

$$\frac{\partial \Phi_i}{\partial z}(\xi, \eta, 1, t) = 0, \tag{2.11}$$

$$\left(-\frac{\partial}{\partial z} [d_i \Phi_i] + \mu_i \Phi_i \right) (\xi, \eta, 0, t) = e_i(\Lambda_t(\xi, \eta), 0, t), \tag{2.12}$$

$$\forall (\xi, \eta, z, t) \in \partial^- D, \quad \Phi_i(\xi, \eta, z, t) = \theta_i(\Lambda_t(\xi, \eta), z, t). \tag{2.13}$$

Note that for the problem (2.9)–(2.13), the variables ξ and η play the role of parameters. We have to deal with a usual *nondegenerate* parabolic problem in the variables (t, z) , where the boundary conditions (2.11) and (2.12) have a usual form, namely

$$\begin{cases} \frac{\partial \Phi_i}{\partial t} - d_i \frac{\partial^2 \Phi_i}{\partial z^2} - \omega_3 \frac{\partial \Phi_i}{\partial z} = F_i(\Phi) + G_i, \\ \frac{\partial \Phi_i}{\partial z} = 0 & \text{for } z = 1, \\ -\frac{\partial}{\partial z} [d_i \Phi_i] + \mu_i \Phi_i = e_i & \text{for } z = 0. \end{cases} \tag{2.14}$$

A particularity is that the domain D is not a cylinder in the variable t . For each (ξ, η) , the evolution is to be solved in the open (possibly empty) subset

$$I_{\xi,\eta} := \{t \in (0, T); (\xi, \eta, z, t) \in D\}, \tag{2.15}$$

which is a countable union of intervals, independent of z , each of them being of the form (τ_-, τ^+) where $\tau_- = \tau(\xi, \eta, z, t)$ for some t (again τ_-, τ^+ are independent of z). By virtue of (2.6), we can assert that the “initial” value of Φ_i at the point (ξ, η, z, τ_-) is prescribed as follows

$$\begin{cases} \Phi_i(\xi, \eta, z, \tau_-) = \phi_{0_i}(\Lambda_0(\xi, \eta), z) & \text{if } \tau_- = 0, \\ \Phi_i(\xi, \eta, z, \tau_-) = \theta_i(\Lambda_{\tau_-}(\xi, \eta), z, \tau_-) & \text{if } \tau_- > 0. \end{cases} \tag{2.16}$$

A technical remark. We will often have to make estimates of $\int_Q a(\phi_i)$ for regular functions $a : \mathbb{R} \rightarrow [0, +\infty)$. Then, using the above change of variables, for some constant C depending only on the data (essentially on the sup norm of the Jacobian of $\Lambda_{z,t}$ and of its inverse), we may bound $\mathcal{I} = \int_Q a(\phi_i) dx dy dz dt$ by

$$\mathcal{I} \leq C \int_D a(\Phi_i) d\xi d\eta dz dt = C \int_{\tilde{\Omega}} d\xi d\eta \int_{(0,1) \times I_{\xi,\eta}} a(\Phi_i) dz dt,$$

where

$$\tilde{\Omega} = \bigcup \{ \Lambda_{z,t}(\Omega); (z, t) \in (0, 1) \times (0, T) \} = \{ (\xi, \eta) \in \mathbb{R}^2; I_{\xi,\eta} \neq \emptyset \}.$$

If one has an estimate of the form

$$\int_{\tau_- 0}^{\tau^+ 1} \int a(\Phi_i) dt dz \leq \int_{\tau_- 0}^{\tau^+ 1} \int h(\xi, \eta, z, t) dt dz, \tag{2.17}$$

for some integrable function h and for all (ξ, η) and all intervals $(\tau_-, \tau^+) \in I_{\xi,\eta}$, then, summing up over all intervals (τ_-, τ^+) and integrating in (ξ, η) leads to

$$\mathcal{I} \leq C \int_D h(\xi, \eta, z, t) d\xi d\eta dz dt \leq C \int_Q h(\Lambda_{z,t}^{-1}(x, y), z, t) dx dy dz dt. \tag{2.18}$$

3. Existence of global classical solutions

In this section, we state and prove our main existence result of global classical solutions. For $\psi \in L^2(Q)$, we denote

$$\Lambda\psi = \frac{\partial\psi}{\partial t} - \nabla \cdot (\psi\omega) + \psi\nabla \cdot \omega,$$

computed in the sense of distributions. Note that, we *formally* have

$$\Lambda\psi = \frac{\partial\psi}{\partial t} - \omega \cdot \nabla\psi,$$

this being true only when ψ is regular enough.

We also denote, for $i = 1, \dots, n$,

$$\forall \psi \in L^2(Q), \quad L_i(\psi) = \Lambda\psi - d_i \frac{\partial^2\psi}{\partial z^2}$$

and $L = (L_i)_{1 \leq i \leq n}$.

Theorem 2. Assume that (1.2), (1.6), (1.7), (1.8), (1.9) as well as (2.8) hold. Then, for all $g \in L^\infty(Q)^n$ with $g \geq 0$, there exists a unique nonnegative solution to the following system

$$\begin{cases} \forall i = 1, \dots, n, & \phi_i, \partial\phi_i/\partial z, \partial^2\phi_i/\partial z^2, \Lambda\phi_i \in L^2(Q), \phi_i \in L^\infty(Q), \\ L\phi = f(\phi) + g & \text{on } Q, \\ \forall i = 1, \dots, n, & \text{boundary and initial conditions (1.5) hold.} \end{cases} \tag{3.1}$$

A starting point of the analysis is the following lemma on the linear part of the system.

Lemma 3. Assume that (1.2), (1.6) and (2.8) hold. Then, for all $g_i \in L^2(Q)$, there exists a unique solution to the following system

$$\begin{cases} \phi_i, \partial\phi_i/\partial z, \partial^2\phi_i/\partial z^2, \Lambda\phi_i \in L^2(Q), \\ L_i\phi_i = g_i \quad \text{on } Q, \\ \text{boundary and initial conditions (1.5) hold.} \end{cases} \tag{3.2}$$

Moreover, $[g_i \geq 0] \Rightarrow [\phi_i \geq 0]$ and $[g_i \in L^\infty(Q)] \Rightarrow [\phi_i \in L^\infty(Q)]$.

This lemma follows from classical results that may be found in [17,1,12,13]. However, we give in Section 5 the main steps in its proof, and we point out how assumption (2.8) has to be used even in this lemma.

In order to prepare existence results for the complete nonlinear system, we first *truncate the nonlinearities*. Let $M > 0$ and $\varphi^M \in C^\infty(\mathbb{R})$ such that

$$0 \leq \varphi^M \leq 1, \quad \varphi^M = 1 \quad \text{on } \left[-\frac{M}{2}, \frac{M}{2}\right], \quad \varphi^M = 0 \quad \text{outside } [-M, M].$$

We define

$$\forall \phi \in \mathbb{R}^n, \quad f^M(\phi) = f(\phi)\varphi^M(|\phi|^2).$$

We denote $f^M = (f_i^M)_i$, etc.

Lemma 4. Assume that (1.2), (1.6), (1.7) and (2.8) hold. Then, for all $g \in L^\infty(Q)^n$ with $g \geq 0$, there exists a unique solution to the following truncated system

$$\begin{cases} \forall i = 1, \dots, n, \quad \phi_i, \partial\phi_i/\partial z, \partial^2\phi_i/\partial z^2, \Lambda\phi_i \in L^2(Q), \quad \phi_i \in L^\infty(Q), \\ L\phi = f^M(\phi) + g \quad \text{on } Q, \\ \forall i = 1, \dots, n, \text{ boundary and initial conditions (1.5) hold.} \end{cases} \tag{3.3}$$

Moreover, for all $i = 1, \dots, n$, $\phi_i \geq 0$.

This lemma is obtained in a standard way by a fixed point argument: we indeed have a Lipschitz perturbation of a “good” linear operator. Again, we indicate the main arguments in the last section. Preservation of positivity is due to the quasi-positivity of f_M . Uniqueness is standard for reaction–diffusion systems in the family of classical *uniformly bounded solutions*.

Remark. To prove existence of solutions on Q with the nonlinearity f itself, the game consists in proving L^∞ -estimates on the solution of the approximate system (3.3) which do not depend on M . Then choosing M large enough, a solution of (3.3) will also be a solution of (3.1).

We will now denote by C any constant depending only on the data (but not on M).

For simplicity, from now on, we will often write $\psi_t, \psi_z, \psi_{zz}, \dots$ for the derivatives of a function ψ .

Thanks to the assumption (2.8), we may use the change of coordinates of Section 2. The system (3.3) is equivalent to the system (2.14)–(2.16) with $F(\Phi)$ replaced by $F^M(\Phi) = f^M(\phi)$. For each (ξ, η) , we will make estimates on the set

$$G = \{(z, t) \in (0, 1) \times (\tau_-, \tau^+)\}$$

for all time intervals (τ_-, τ^+) .

We first start by the key L^2 -estimate which is valid essentially only under assumption (1.8).

Proposition 5. Let ϕ be the solution of the truncated system (3.3) and Φ the solution of the corresponding system in the new variables. Then

$$\forall (\xi, \eta), \text{ for all interval } (\tau_-, \tau^+), \quad \|\Phi\|_{L^2(G)} \leq C(\tau^+ - \tau_-)^{1/2}. \tag{3.4}$$

Remark. As we know, Φ is defined for t in the domain $I_{\xi,\eta}$ (see (2.15)) which is the union of the intervals (τ_-, τ^+) . Summing up the above estimate on all of the sub-intervals (τ_-, τ^+) of $I_{\xi,\eta}$ leads to

$$\|\Phi\|_{L^2[(0,1)\times I_{\xi,\eta}]}^2 \leq CT. \tag{3.5}$$

We can then go back to the initial function ϕ and, through the computations (2.17), (2.18), obtain the bound $\|\phi\|_{L^2[(0,1)\times I_{\xi,\eta}]}^2 \leq CT$.

Proof. We set $W = \sum_i a_i \Phi_i$, $Z = \sum_i d_i a_i \Phi_i$, where the coefficients a_i are defined in (1.8). Summing all the equations in the Φ_i 's (see (2.14) with F^M instead of F), we have, using also assumption (1.8):

$$W_t - Z_{zz} - \omega_3 W_z = \sum_i a_i F_i^M(\Phi) + \sum_i a_i G_i \leq \sum_i b_i \Phi_i + C.$$

Since $a_i > 0$ for all i , one has $\sum_i b_i \Phi_i \leq AW$ for some $A > 0$ depending only on the a_i, b_i . Then, we deduce the following inequality from which the L^2 estimate will follow:

$$W_t - Z_{zz} - \omega_3 W_z \leq AW + C. \tag{3.6}$$

We set $\sigma := Z/W$ so that we may write

$$W_t - (\sigma W)_{zz} - \omega_3 W_z \leq AW + C. \tag{3.7}$$

A main point is that, thanks to the positivity of the Φ_i, a_i ,

$$0 < \min_i d_i \leq \sigma \leq \max_i d_i. \tag{3.8}$$

As in [19,3], we will exploit this property to get an $L^2(G)$ -estimate on W which obviously implies the expected L^2 -estimate on Φ . We do it by duality. For this, let $\Theta \in C_0^\infty(G)$, $\Theta \geq 0$. We introduce the nonnegative solution θ of the dual problem

$$\begin{cases} -\theta_t - \sigma \theta_{zz} + (\omega_3 \theta)_z - A\theta = \Theta & \text{on } G, \\ \theta_z = r(2z - 1)\theta & \text{for } z = 0, 1, \\ \theta = 0 & \text{at } t = \tau^+, \end{cases} \tag{3.9}$$

where $r \in (0, +\infty)$ is such that

$$r\sigma - \omega_3 \geq 0 \quad \text{for } z = 1, \quad -r\sigma - \omega_3 \leq 0 \quad \text{for } z = 0. \tag{3.10}$$

This choice of r is possible since σ is bounded from below and $|\omega_3|$ is bounded from above. We denote $R(z) = r(z^2 - z)$ so that $R'(z) = r(2z - 1)$. In particular,

$$[\theta_z = r(2z - 1)\theta] \Leftrightarrow [e^{-R}\theta]_z = 0. \tag{3.11}$$

Multiplying (3.6) by $\theta \geq 0$ and using $Z = \sigma W$ lead to

$$0 \geq \int_G \theta [W_t - Z_{zz} - \omega_3 W_z - AW - C].$$

This writes

$$0 \geq \int_G (W\Theta - C\theta) - \int_0^1 [\theta W](z, \tau_-) dz + a, \tag{3.12}$$

where a denotes the integrated terms in z , namely

$$a = \int_{\tau_-}^{\tau^+} \{ [-\theta Z_z - \omega_3 \theta W + \theta_z Z](1, t) - [-\theta Z_z - \omega_3 \theta W + \theta_z Z](0, t) \} dt,$$

which, according to the boundary conditions on the Φ_i 's and θ , is equal to

$$a = \int_{\tau_-}^{\tau^+} W\theta[r\sigma - \omega_3](1, t) + \theta \left[W(\omega_3 + r\sigma) + \sum_i a_i(\mu_i \Phi_i - e_i) \right](0, t) dt.$$

By the choice of r as in (3.10) and thanks to $\mu_i \geq 0, e_i \in L^\infty$, we deduce that $a \geq -C \int_{\tau_-}^{\tau^+} \theta(0, t) dt$. Together with (3.12) and the fact that $W(\cdot, \tau_-)$ is a given L^∞ -function, this leads to

$$\int_G W\Theta \leq C \left\{ \int_0^1 \theta(z, \tau_-) dz + \int_{\tau_-}^{\tau^+} \theta(0, t) dt + \int_G \theta \right\}. \tag{3.13}$$

The next task is now to prove

$$\int_0^1 \theta(z, \tau_-) dz + \int_{\tau_-}^{\tau^+} \theta(0, t) dt + \int_G \theta \leq C(\tau^+ - \tau_-)^{1/2} \|\Theta\|_{L^2(G)}. \tag{3.14}$$

Then, the expected estimate of Proposition 5 will follow by duality. To obtain (3.14), we actually prove maximal L^2 -regularity for the parabolic operator involved in (3.9), namely

$$\int_G \theta^2 + (\theta_t)^2 + (\theta_z)^2 + (\theta_{zz})^2 \leq C \int_G \Theta^2. \tag{3.15}$$

Indeed, assuming (3.15), we then obtain, by setting $\rho := (\tau^+ - \tau_-)^{1/2}$:

$$\begin{aligned} \int_G \theta &\leq |G|^{1/2} \left\{ \int_G \theta^2 \right\}^{1/2} \leq \rho \left\{ C \int_G \Theta^2 \right\}^{1/2}, \\ \int_0^1 \theta(z, \tau_-) dz &= - \int_{\tau_-}^{\tau^+} \int_0^1 \theta_t \leq \rho \left\{ \int_G (\theta_t)^2 \right\}^{1/2} \leq \rho \left\{ C \int_G \Theta^2 \right\}^{1/2}. \end{aligned}$$

Then, starting from the embedding estimate

$$\forall t \in (\tau_-, \tau^+), \quad \theta(0, t) \leq \left\{ \int_0^1 [\theta^2 + (\theta_z)^2](z, t) dz \right\}^{1/2},$$

we have similarly, using again (3.15)

$$\int_{\tau_-}^{\tau^+} \theta(0, t) dt \leq \int_{\tau_-}^{\tau^+} dt \left\{ \int_0^1 [\theta^2 + (\theta_z)^2] dz \right\}^{1/2} \leq \rho \left\{ C \int_G \Theta^2 \right\}^{1/2}.$$

Whence (3.14). It remains to prove (3.15). We set $U := e^{-R}\theta$ where R is defined in (3.11). Obviously, it is sufficient to prove (3.15) with θ replaced by U . The function U satisfies an equation of the form

$$\left. \begin{aligned} -U_t - \sigma U_{zz} + a(z, t)U + b(z, t)U_z &= \hat{\Theta}, \\ U(\cdot, \tau^+) &= 0, \\ U_z &= 0 \quad \text{at } z = 0 \text{ and } z = 1, \end{aligned} \right\} \tag{3.16}$$

where a, b are uniformly bounded by C and $\hat{\Theta} = e^{-R}\Theta$.

Multiply the equation in U by $-U_{zz}$ and integrate by parts to obtain:

$$\int_0^1 -\partial_t \frac{1}{2} [U_z]^2 + \sigma [U_{zz}]^2 = \int_0^1 [-\hat{\theta} + aU + bU_z] U_{zz}. \tag{3.17}$$

Since σ is bounded from below, we can use Young’s inequality to control the right-hand side as follows:

$$\int_0^1 [\hat{\theta} + aU + bU_z] U_{zz} \leq \int_0^1 \frac{\sigma}{2} [U_{zz}]^2 + C[\hat{\theta}^2 + U^2 + (U_z)^2]. \tag{3.18}$$

Absorbing the term in $(U_{zz})^2$ and exploiting the linear Gronwall’s inequality in the term $(U_z)^2$, we deduce from (3.17), (3.18) and $U(\tau^+) = 0$, the following estimate on $G_t = (t, \tau^+) \times (0, 1)$:

$$\int_{G_t} [U_{zz}]^2, \sup_{s \in (t, \tau^+)} \int_0^1 [U_z]^2(s) \leq C \int_{G_t} \hat{\theta}^2 + U^2. \tag{3.19}$$

Going back to (3.16), we derive $\int_{G_t} (U_t)^2 \leq C \int_{G_t} \hat{\theta}^2 + U^2$ which, together with $U(\cdot, t) = -\int_t^{\tau^+} U_t(\cdot, s) ds$, leads to

$$\int_0^1 U^2(z, t) dz \leq C \int_t^{\tau^+} \int_0^1 (U_t)^2 \leq C \int_t^{\tau^+} \int_0^1 \hat{\theta}^2 + U^2.$$

Integration of this Gronwall’s inequality in $t \rightarrow \int_0^1 U^2(t)$ gives

$$\int_G U^2 + (U_t)^2 \leq C \int_G \hat{\theta}^2.$$

This inequality coupled with (3.19) leads to (3.15) with θ replaced by U , and this ends the proof of Proposition 5. \square

Now, we recall classical estimates on the heat operator in one space dimension (see e.g. [14]):

Lemma 6. *Let ψ be a solution on G of*

$$\begin{cases} \psi_t - d\psi_{zz} - \omega_3\psi_z = F, \\ \psi_z = 0 & \text{for } z = 1, \\ -d\psi_z + \mu\psi = e \in L^\infty(0, T) & \text{for } z = 0, \\ \psi(z, 0) = \psi_0 \in L^\infty(0, 1). \end{cases} \tag{3.20}$$

Then, for $C = C(T, d, \omega_3, \mu, e, \|\psi_0\|_\infty)$,

$$\begin{aligned} \forall r \in (1, 3), \quad & \|\psi\|_{L^r(G)} \leq C[1 + \|F\|_{L^1(G)}], \\ \forall p \in (1, 3/2), q = 3p/(3 - 2p) \quad & \|\psi\|_{L^q(G)} \leq C[1 + \|F\|_{L^p(G)}], \\ \forall p > 3/2, \quad & \|\psi\|_{L^\infty(G)} \leq C[1 + \|F\|_{L^p(G)}]. \end{aligned}$$

Remark. These estimates may be found for instance in [14]. The only remark to be done is that the constants above do not indeed depend on G and in particular do not depend on $\tau^+ - \tau_-$ even if this difference becomes small. A simple way to see it is to notice that a solution on the interval $I = (\tau_-, \tau^+)$ is the restriction to I of a solution on the fixed length interval $I_1 = (\tau_-, \tau_- + T)$ with a left-hand side \tilde{F} equal to F on I and to zero on $I_1 \setminus I$. Then, for instance, the first inequality is obtained as follows from the same inequality on the ‘fixed’ domain $G_1 = (0, 1) \times I_1$:

$$\|\psi\|_{L^r(G)} \leq \|\psi\|_{L^r(G_1)} \leq C[1 + \|\tilde{F}\|_{L^1(G_1)}] = C[1 + \|F\|_{L^1(G)}].$$

Lemma 7. *Let ϕ be the solution of the truncated system (3.3) and Φ the solution of the corresponding system in the new variables. Then*

$$\forall(\xi, \eta), \text{ for all interval } (\tau_-, \tau^+), \quad \|\Phi\|_{L^\infty(G)} \leq C. \tag{3.21}$$

Proof. We start with the L^2 -estimate of Proposition 5 where we even forget the precise dependence in $(\tau^+ - \tau_-)$, that is to say, we start with: $\|\Phi\|_{L^2(G)} \leq C$.

By the ‘quadratic hypothesis’ (1.9): $\|F^M(\Phi)\|_{L^1(G)} \leq C$.

By the first statement of Lemma 6: $\forall r \in (2, 3), \|\Phi\|_{L^r(G)} \leq C$.

By (1.9) again: $\forall p \in (1, 3/2), \|F^M(\Phi)\|_{L^p(G)} \leq C$.

By the second statement of Lemma 6: $\forall q < +\infty, \|\Phi\|_{L^q(G)} \leq C$.

By (1.9): $\forall m < +\infty, \|F(\Phi)\|_{L^m(G)} \leq C$.

Finally, by the last statement of Lemma 6: $\|\Phi\|_{L^\infty(G)} \leq C$.

Lemma 7 follows. \square

Proof of Theorem 2. As explained in the remark following Lemma 4, we apply the uniform estimate of Lemma 7 to the solution of the truncated system (3.3). We choose M larger than the constant C of Lemma 7. Then, the solution of the truncated system (3.3) is also a solution of the true system (3.1). \square

Uniqueness is classical in the class of uniformly bounded solutions.

Remark. In the context of air pollution models, the coefficients d_i are the so-called ‘‘eddy diffusivities’’ introduced to model turbulent mixing, see for example [21] and the references therein. In realistic models, these coefficients may depend on the vertical variable z . In this case, the diffusion part must be written in the divergence form,

$$L_i \psi = \Lambda \psi - \frac{\partial}{\partial z} \left(d_i \frac{\partial \psi}{\partial z} \right).$$

Adapting the proof of Proposition 5, it is easy to show that the results of Theorem 2 remain valid if the coefficients d_i are differentiable functions of z , as long as the following conditions are satisfied:

$$\exists \alpha > 0, \exists C > 0, \forall i, \forall z \in (0, 1), \quad d_i(z) > \alpha, \quad |d'_i(z)| \leq C. \tag{3.22}$$

However, this approach fails as such when the diffusion coefficients vanish at $z = 0$, which is the case in the models stemming from the Monin–Obukhov similarity theory, see [21, p. 869]. Extensions of this method may be made to handle coefficients vanishing inside the domain (see [3]). Degeneracy at $z = 0$ would require further improvements.

4. A specific nonlinearity f

Here, as a main example, we reproduce the 20×20 system mentioned in [4] and introduced in the context of pollutants models in [9] and [22].

The nonlinear part is given as follows:

$$\begin{aligned} f_1(\phi) = & -k_1\phi_1 + k_{22}\phi_{19} + k_{25}\phi_{20} + k_{11}\phi_{13} + k_9\phi_{11}\phi_2 + k_3\phi_5\phi_2 \\ & + k_2\phi_2\phi_4 - k_{23}\phi_1\phi_4 - k_{14}\phi_1\phi_6 + k_{12}\phi_{10}\phi_2 - k_{10}\phi_{11}\phi_1 - k_{24}\phi_{19}\phi_1, \end{aligned}$$

and

$$\begin{aligned} f_2(\phi) = & k_1\phi_1 + k_{21}\phi_{19} - k_9\phi_{11}\phi_2 - k_3\phi_5\phi_2 - k_2\phi_2\phi_4 - k_{12}\phi_{10}\phi_2, \\ f_3(\phi) = & k_1\phi_1 + k_{17}\phi_4 + k_{19}\phi_{16} + k_{22}\phi_{19} - k_{15}\phi_3, \\ f_4(\phi) = & -k_{17}\phi_4 + k_{15}\phi_3 - k_{16}\phi_4 - k_2\phi_2\phi_4 - k_{23}\phi_1\phi_4, \\ f_5(\phi) = & 2k_4\phi_7 + k_7\phi_9 + k_{13}\phi_{14} + k_6\phi_7\phi_6 - k_3\phi_5\phi_2 + k_{20}\phi_{17}\phi_6, \\ f_6(\phi) = & 2k_{18}\phi_{16} - k_8\phi_9\phi_6 - k_6\phi_7\phi_6 + k_3\phi_5\phi_2 - k_{20}\phi_{17}\phi_6 - k_{14}\phi_1\phi_6, \\ f_7(\phi) = & -k_4\phi_7 - k_5\phi_7 + k_{13}\phi_{14} - k_6\phi_7\phi_6, \end{aligned}$$

$$\begin{aligned}
 f_8(\phi) &= k_4\phi_7 + k_5\phi_7 + k_7\phi_9 + k_6\phi_7\phi_6, \\
 f_9(\phi) &= -k_7\phi_9 - k_8\phi_9\phi_6, \\
 f_{10}(\phi) &= k_7\phi_9 + k_9\phi_{11}\phi_2 - k_{12}\phi_{10}\phi_2, \\
 f_{11}(\phi) &= k_{11}\phi_{13} - k_9\phi_{11}\phi_2 + k_8\phi_9\phi_6 - k_{10}\phi_{11}\phi_1, \\
 f_{12}(\phi) &= k_9\phi_{11}\phi_2, \\
 f_{13}(\phi) &= -k_{11}\phi_{13} + k_{10}\phi_{11}\phi_1, \\
 f_{14}(\phi) &= -k_{13}\phi_{14} + k_{12}\phi_{10}\phi_2, \\
 f_{15}(\phi) &= k_{14}\phi_1\phi_6, \\
 f_{16}(\phi) &= -k_{19}\phi_{16} - k_{18}\phi_{16} + k_{16}\phi_4, \\
 f_{17}(\phi) &= -k_{20}\phi_{17}\phi_6, \\
 f_{18}(\phi) &= k_{20}\phi_{17}\phi_6, \\
 f_{19}(\phi) &= -k_{21}\phi_{19} - k_{22}\phi_{19} + k_{25}\phi_{20} + k_{23}\phi_1\phi_4 - k_{24}\phi_{19}\phi_1, \\
 f_{20}(\phi) &= -k_{25}\phi_{20} + k_{24}\phi_{19}\phi_1.
 \end{aligned}$$

As noticed and strongly used in [4], this nonlinearity satisfies the five following relations:

$$\left\{ \begin{aligned}
 f_1 + f_2 + f_{13} + f_{15} + f_{19} + f_{20} &= 0, \\
 f_7 + f_8 + f_9 + f_{10} + f_{11} + f_{12} + f_{13} + f_{14} &= 0, \\
 f_{17} + f_{18} &= 0, \\
 f_3 + f_4 + f_{16} &\leq k_1\phi_1 + k_{22}\phi_{19}, \\
 f_5 + f_6 &\leq 2k_4\phi_7 + k_7\phi_9 + k_{13}\phi_{14} + 2k_{18}\phi_{16}.
 \end{aligned} \right. \tag{4.1}$$

Summing up these five relations yields the assumption (1.8). Let us emphasize that we only use this last global dissipative relation, and not the five ones separately. On the other hand, we exploit the quadratic structure (see next section for more comments on other kinds of systems).

5. Comments and open problems

In this section, we first give direct and elementary proofs of the preliminary Lemmas 3 and 4. Then we make some comments and indicate open problems.

Proof of Lemma 3.

Preliminary remark. One way to prove this linear lemma is to use the results in [1,12,13]. However, in these references, solutions are a priori less regular than stated in the lemma. Roughly speaking, $\Lambda\phi_i, \partial^2\phi_i/\partial z^2$ are found in an H^{-1} -type space rather than in L^2 . We do not know how to prove the L^2 regularity in general (see open problems below). On the other hand, under the assumption (2.8), we are able to do it. For simplicity, we take advantage of the change of variable introduced in Section 2 and we provide a more selfcontained proof.

Let $\zeta_i \in [H^2 \cap L^\infty](Q)$ as in assumption (1.6). Then, setting by translation $\psi_i := \phi_i - \zeta_i$, in order to prove existence of ϕ_i as in Lemma 3, we may assume all the data equal to zero except for the right-hand side $g_i \in L^2(Q)$.

Now by assumption (2.8) and thanks to the change of variables described in Section 2, we are reduced to solving the new following system on each interval $[\tau_-, \tau^+)$:

$$\left. \begin{aligned}
 \frac{\partial \Phi_i}{\partial t} - d_i \frac{\partial^2 \Phi_i}{\partial z^2} - \omega_3 \frac{\partial \Phi_i}{\partial z} &= G_i, \\
 \frac{\partial \Phi_i}{\partial z} &= 0 && \text{for } z = 1, \\
 -\frac{\partial}{\partial z}[d_i \Phi_i] + \mu_i \Phi_i &= 0 && \text{for } z = 0, \\
 \Phi_i(\xi, \eta, z, \tau_-) &= 0.
 \end{aligned} \right\} \tag{5.1}$$

This problem is nondegenerate and it is well known (see e.g. [14]) that for G_i given in $L^2(G)$, it has a (unique) solution with derivatives in L^2 and the corresponding estimates, namely

$$\int_{\tau_- 0}^{\tau^+ 1} \int |(\Phi_i)_t|^2 + |(\Phi_i)_{zz}|^2 + |(\Phi_i)_z|^2 dt dz \leq C \int_{\tau_- 0}^{\tau^+ 1} |G_i|^2 dt dz, \tag{5.2}$$

where C depends only on the data. The function G_i is measurable with respect to (ξ, η) . The above estimate preserves this measurability for the solution Φ : indeed, if G_i depends continuously on (ξ, η) , then so does the solution by (5.2). Now, if G is in L^2 in the four variables, it may be approximated in $L^2(G)$ and a.e. in (ξ, η) by regular functions. Again, (5.2) ensures the convergence and the measurability of the limit. Moreover, according to the computations (2.17), (2.18) and to those of Section 2, we also have

$$\int_Q |\Delta \phi_i|^2 + |(\phi_i)_z|^2 + |(\phi_i)_{zz}|^2 \leq C.$$

For the last statement of Lemma 3, we may again use the change of variable of Section 2. We are back to a system of type (5.1) with

$$0 \leq G_i \leq \|G_i\|_\infty < +\infty$$

and nonnegative bounded boundary data instead of zero data. It is well known that they provide nonnegative bounded solutions. \square

Proof of Lemma 4. As in the previous proof, we use the change of variable of Section 2 and we are lead to the system

$$\left. \begin{aligned} \frac{\partial \Phi_i}{\partial t} - d_i \frac{\partial^2 \Phi_i}{\partial z^2} - \omega_3 \frac{\partial \Phi_i}{\partial z} &= F_i^M(\Phi) + G_i, \\ \frac{\partial \Phi_i}{\partial z} &= 0 && \text{for } z = 1, \\ -\frac{\partial}{\partial z} [d_i \Phi_i] + \mu_i \Phi_i &= e_i && \text{for } z = 0, \\ \Phi_i(\xi, \eta, z, \tau_-) &= \Phi_{0i}(\xi, \eta, z). \end{aligned} \right\} \tag{5.3}$$

We have here a globally Lipschitz perturbation of the previous linear system. Since all data are in L^∞ , by a classical fixed point theorem, we obtain unique uniformly bounded solutions on $(0, T)$. Measurability with respect to (ξ, η) is preserved in this approach.

To prove that the positivity is preserved when the data are nonnegative, instead of solving directly with the right-hand side F^M , we first do it with F^M replaced by $F^M \circ \Pi$ where $\Pi : \mathbb{R}^n \rightarrow (\mathbb{R}^+)^n$ is the orthogonal projection onto the positive cone $(\mathbb{R}^+)^n$ of \mathbb{R}^n . The assumption (1.7) implies that, for all i and all $\Phi \in \mathbb{R}^n$, $[F_i^M(\Pi(\Phi))] \Phi_i^- \geq 0$. Then, multiplying the i th equation of the modified system by Φ_i^- and using boundary conditions lead to

$$-\frac{\partial}{\partial t} \frac{1}{2} \int_0^1 (\Phi_i^-)^2 - d_i \int_0^1 \{(\Phi_i^-)_z\}^2 - \frac{\omega_3}{2} \{(\Phi_i^-)^2\}_z \geq 0.$$

Integrating this inequality in t proves that $\Phi_i^- = 0$, whence the positivity of the solutions of the modified system. But, since $F^M = F^M \circ \Pi$ on the positive cone, Φ is also solution of the initial system. \square

About the L^2 -estimate: The main ingredient in the proof of Theorem 2 is the L^2 -estimate given in Proposition 5. We could ask whether such an estimate would be true without the assumption (2.8). Let us make the question more precise. If, as in the proof of Proposition 5, we set $W = \sum_i a_i \phi_i$, $Z = \sum_i d_i a_i \phi_i$, then summing the n equations in (1.3), we obtain similarly to (3.12):

$$W_t - [\sigma W]_{zz} + \omega \cdot \nabla W - AW \leq C, \tag{5.4}$$

where

$$0 < \min_i d_i \leq \sigma = \frac{Z}{W} \leq \max_i d_i < +\infty.$$

We do not know whether the inequality (5.4) implies an L^2 -estimate on W as in Proposition 5. The dual problem is

$$-\theta_t - \sigma \theta_{zz} - \nabla \cdot (\omega \theta) - A \theta = \Theta. \tag{5.5}$$

The goal would be to prove L^2 -estimates for this equation (see (3.15)). Technically, when we multiply it by $-\theta_{zz}$ as in the proof of Proposition 5, then it is not clear what to do with the new terms of the form $\omega_1 \theta_x \theta_{zz}$. After integration by parts, they become $\theta_z [\omega_{1z} \theta_x + \omega_1 \theta_{xz}]$. The last term can be treated easily, but it is not so clear for the first one... except if $\omega_{1z} = 0$, whence the hypothesis (2.8).

We would like to emphasize that it is not even obvious to obtain L^2 -estimates for the main linear operator involved in the system (1.3). More precisely, let

$$\theta_t - \theta_{zz} - \omega \cdot \nabla \theta = \Theta, \tag{5.6}$$

with ‘good boundary conditions’ and $\Theta \in L^2(Q)$. Does this imply “maximal regularity”, namely

$$\theta_t - \omega \cdot \nabla \theta, \theta_{zz} \in L^2(Q)? \tag{5.7}$$

The above approach (multiplying by θ_{zz}) fails in general in exactly the same way as for the more general operator with the σ -term (and as just explained).

Therefore, here is a first open problem: for which ω does (5.6) imply (5.7) when $\Theta \in L^2(Q)$?

One can find partial results in the literature from the theory of so-called ultra-parabolic operators (see e.g. [2]). Strangely enough, it follows from this theory that, for instance, for $\omega(x, y, z, t) = (z, 0, 0)$, the answer would be positive, although $\omega_{1z} \neq 0$... Indeed, for each fixed y , the operator is then *hypoelliptic* in the variables (x, z, t) as shown by Kolmogorov in [10] and generalized by Hörmander in [8]. As proved in [20] (see also [11]), this implies the expected L^2 (and even L^p) regularity in (x, z, t) and, therefore also in (x, y, z, t) after integration in y (see the technical remark (2.17), (2.18)). This approach may be generalized to operators of the form (5.6) when they are hypoelliptic (see [20]) or, equivalently, when they satisfy Hörmander’s dimension condition for the space generated by the commutators of $\partial_t - \omega \cdot \nabla$ and ∂_z up to an arbitrary order. However, this approach would not apply for instance to $\omega = (z, x - tz, 0)$ in which case the operator is not hypoelliptic (see [8,11]). Note that it is interesting to understand the optimal assumptions needed on ω in order to better handle *nonlinear models* where $\omega = \omega(\theta)$. Many such nonlinear models with degenerate diffusion are mentioned in Section 3 of [11].

Once the previous problem is understood, a *second open problem* would be to understand for which ω the solutions of (5.5) satisfy

$$\Theta \in L^2(Q) \Rightarrow \theta, \theta_{zz}, \theta_t + \nabla \cdot (\omega \theta) \in L^2(Q).$$

This would be a first step in solving our main problem without assumption (2.8).

A *third open problem* is to understand what happens in dimensions greater than 3. A main point in our approach is that we reduced the problem to a one-dimensional one. Assume more generally that we have a problem in dimension $d > 3$ with no diffusion in a number d' of the directions. Then, with an assumption like (2.8), we may reduce the problem to a $(d - d')$ -dimensional problem. We still keep the L^2 -estimate of Proposition 5, but, if $d - d' > 1$, we cannot any more reach L^∞ -estimates as for $d - d' = 1$. It is very likely that we can then obtain global “weak”-solution in the spirit of [18,3], but this needs to be done.

Another *interesting fourth problem* is when the nonlinearities are superquadratic. Several models are mentioned in [21] (see pages 143, 167, 209) which come with this structure. But the question is interesting for itself. Even when the diffusions are not degenerate, the problem is not yet completely understood. Some partial results are however known. For instance, if the right-hand side presents some “triangular structure”, L^∞ -estimates may successively be obtained on all the components (see [15,16]). This approach is based on L^p a priori estimates and it would be necessary to first extend them to a degenerate situation. It is probably necessary to first understand it in the case $p = 2$ (as in the previous problems just mentioned).

Another approach would be to look for global weak solutions. If the nonlinearity allows an a priori L^1 -estimate on the right-hand side, we may expect global weak-solutions as in [18]. But this needs to be done.

One more open problem would be interesting to look at, with respect to applications. The models considered here are approximate ‘simplified’ versions of more elaborate ones: for instance, saying that no diffusion occurs in the horizontal directions is just an approximation of the fact that diffusion does occur, but is small. Therefore, a natural question would be to study what happens exactly for the system with positive diffusion coefficients tending to zero in the horizontal directions. Since very different diffusion coefficients generate sharp behaviours in these systems, this question is not easy, but relevant.

References

- [1] C. Bardos, Problèmes aux limites pour les équations aux dérivées partielles, *Ann. Sci. École Norm. Sup. (4)* 4 (1970) 185–233.
- [2] M. Bramanti, M.C. Cerutti, M. Manfredini, L^p estimates for some ultraparabolic operators with discontinuous coefficients, *J. Math. Anal. Appl.* 200 (1996) 332–354.
- [3] L. Desvillettes, K. Fellner, M. Pierre, J. Vovelle, About global existence for quadratic systems of reaction–diffusion, *Adv. Nonlinear Stud.* 7 (3) (2007) 491–511.
- [4] W.E. Fitzgibbon, M. Langlais, J.J. Morgan, A degenerate reaction–diffusion system modeling atmospheric dispersion of pollutants, *J. Math. Anal. Appl.* (2005).
- [5] W.E. Fitzgibbon, S.L. Hollis, J.J. Morgan, Stability and Lyapunov functions for reaction–diffusion systems, *SIAM J. Math. Anal.* 28 (3) (1997) 595–610.
- [6] N. Ghouali, T.M. Touaoula, A linear model for the dynamics of fish larvae, *Electron. J. Differential Equations* (140) (2004), 10 p. (electronic).
- [7] Th. Goudon, A. Vasseur, Regularity analysis for systems of reaction–diffusion equations, in press.
- [8] L. Hörmander, Hypoelliptic second order differential equations, *Acta Math.* 119 (1967) 147–171.
- [9] J. Kim, S. Cho, Computation accuracy and efficiency of the time splitting method in solving atmospheric transport equations, *Atmos. Environ.* 31 (1997) 2215–2224.
- [10] A. Kolmogorov, Zufällige Bewegungen (zur Theorie der Brownschen Bewegung), *Ann. of Math. (2)* 35 (1) (1934) 116–117 (in German).
- [11] E. Lanconelli, A. Pascucci, S. Polidoro, Linear and nonlinear ultraparabolic equations of Kolmogorov type arising in diffusion theory and in finance, in: *Nonlinear Problems in Mathematical Physics and Related Topics, II*, in: *Int. Math. Ser. (N.Y.)*, vol. 2, Kluwer/Plenum, New York, 2002, pp. 243–265.
- [12] M. Langlais, Solutions fortes pour une classe de problèmes aux limites du second ordre dégénérés, *Comm. Partial Differential Equations* 4 (1979) 869–897.
- [13] M. Langlais, A degenerating elliptic problem with unilateral constraints, *Nonlinear Anal.* 4 (1980) 329–342.
- [14] O.A. Ladyzenskaya, V.A. Solonnikov, N.N. Uralceva, *Linear and Quasi-linear Equations of Parabolic Type*, *Transl. Math. Monographs*, vol. 23, Amer. Math. Soc., Providence, RI, 1968.
- [15] R.H. Martin, M. Pierre, Influence of mixed boundary conditions in some reaction–diffusion systems, *Proc. Roy. Soc. Edinburgh Sect. A* 127 (5) (1997) 1053–1066.
- [16] J. Morgan, Global existence for semilinear parabolic systems, *SIAM J. Math. Anal.* 20 (5) (1989) 1128–1144.
- [17] O.A. Oleinik, E.V. Radlevic, *Second Order Equations with Nonnegative Characteristic Form*, Plenum, 1973.
- [18] M. Pierre, Weak solutions and supersolutions in L^1 for reaction–diffusion systems, *J. Evol. Equ.* 3 (1) (2003) 153–168.
- [19] M. Pierre, D. Schmitt, Blowup in reaction–diffusion systems with dissipation of mass, *SIAM Rev.* 42 (2000) 93–106 (electronic).
- [20] L.P. Rothschild, E.M. Stein, Hypoelliptic differential operators on nilpotent groups, *Acta Math.* 137 (1977) 247–320.
- [21] J. Seinfeld, S. Pandis, *Atmospheric Chemistry and Physics*, Wiley, New York, 1995.
- [22] J.G. Verwer, W. Hundsdorfer, J.G. Blom, Numerical time integration for air pollution models, *Surveys Math. Indust.* 10 (2002) 107–174.