

Entropy conditions for scalar conservation laws with discontinuous flux revisited

Boris Andreianov^{a,b}, Darko Mitrović^{c,*}

^a *Laboratoire de Mathématiques CNRS UMR 6623, Université de Franche-Comté, 25030 Besançon Cedex, France*

^b *Institut für Mathematik, Technische Universität Berlin, Straße des 17. Juni 136, 10623 Berlin, Germany*

^c *University of Montenegro, Faculty of Natural Sciences and Mathematics, Cetinjski put bb, 81000 Podgorica, Montenegro*

Received 14 April 2014; received in revised form 15 August 2014; accepted 31 August 2014

Available online 16 October 2014

Abstract

We propose new entropy admissibility conditions for multidimensional hyperbolic scalar conservation laws with discontinuous flux which generalize one-dimensional Karlsen–Risebro–Towers entropy conditions. These new conditions are designed, in particular, in order to characterize the limit of vanishing viscosity approximations. On the one hand, they comply quite naturally with a certain class of physical and numerical modeling assumptions; on the other hand, their mathematical assessment turns out to be intricate.

The generalization we propose is not only with respect to the space dimension, but mainly in the sense that the “crossing condition” of Karlsen, Risebro, and Towers (2003) [31] is not mandatory for proving uniqueness with the new definition. We prove uniqueness of solutions and give tools to justify their existence via the vanishing viscosity method, for the multi-dimensional spatially inhomogeneous case with a finite number of Lipschitz regular hypersurfaces of discontinuity for the flux function.

© 2014 Elsevier Masson SAS. All rights reserved.

MSC: 35L65; 35L67

Keywords: Inhomogeneous scalar conservation law; Discontinuous flux; Entropy solution; Vanishing viscosity approximation; Well-posedness; Crossing condition

1. Introduction

Conservation laws of the form

$$\partial_t u + \operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, u) = S(t, \mathbf{x}, u) \quad (1)$$

serve as mathematical models for one-dimensional gas dynamics, road traffic, for flows in porous media with neglected capillarity effects, blood flow, radar shape-from-shading problems, and in several other applications. The

* Corresponding author.

E-mail addresses: boris.andreianov@univ-fcomte.fr (B. Andreianov), matematika@t-com.me (D. Mitrović).

multi-dimensional conservation law also appears in coupled models, although in this case the regularity of the flux function f in (t, \mathbf{x}) is often not sufficient to develop a full well-posedness theory. The mathematical theory of (1) is very delicate because, in general, even for regular data classical solutions need not to exist; on the other hand, weak (distributional) solutions are, in general, not unique. The classical theory is best established for Cauchy and boundary-value problems in the case where f is Lipschitz continuous in (t, \mathbf{x}) and uniformly locally Lipschitz continuous in u . The source S can be assumed, e.g., Lipschitz continuous in u uniformly in (t, \mathbf{x}) . In this case the S.N. Kruzhkov definition of entropy solution in [33] and the associated analysis techniques (vanishing viscosity approximation for the existence proof, doubling of variables for the uniqueness proof) provide a well-posedness framework for (1).

1.1. Discontinuous-flux models and rough entropy inequalities

Local Lipschitz assumption on $k \mapsto f(t, \mathbf{x}, k)$ is natural in many applications, but the assumption of regular dependence on the spatial variable \mathbf{x} is very restrictive. Indeed, road traffic with variable number of lanes [19], Buckley–Leverett equation in a layered porous medium (see [29,8]), sedimentation applications (see [23,24,17]) make appear models with piecewise regular, jump discontinuous in \mathbf{x} flux functions. The theory of such problems, called *discontinuous-flux conservation laws*, has been an intense subject of research in the last twenty years. The main goal of this research was to design a suitable approach to definition and numerical approximation of entropy solutions, in relation with the physical context of different discontinuous-flux models. Speaking of “notion of solution” in this context, one usually means weak solutions subject to some additional *admissibility conditions*, cf. [33].

Almost all admissibility conditions designed in the literature were confined to the one-dimensional case. We mention here the minimal jump condition [26], minimal variation condition and Γ -condition [23,24], entropy conditions [31,2], vanishing capillarity limit [29,8], admissibility conditions via adapted entropies [12,18] or via admissible jumps description at the interface [3,4,25]. An extensive overview on the subject as well as a kind of unification of the mentioned approaches in the one-dimensional case was given in [11], where further references can be found.

Adimurthi et al. [3] observed that infinitely many different, though equally mathematically consistent, notions of solution may co-exist in the discontinuous-flux problems; therefore the choice of solution notion is a part of modeling procedure (see, e.g., [8] for an exhaustive study of the vanishing capillarity limits of the one-dimensional Buckley–Leverett equation, where different sets of admissibility conditions are put forward for different choices of physically relevant vanishing capillarity). In the present contribution we limit our attention to characterization of vanishing viscosity limit solutions for problems of kind (1); these approximations were studied in a huge number of works (see, e.g., [26,23,24,42,43,41,31,32,13,18,40,25]) including several works in multiple space dimensions (see [28,27,10,39,16]) and they remain relevant in several models based on discontinuous-flux conservation laws.

The basis of the different definitions of admissibility of solutions is provided by Kruzhkov entropy inequalities [33] in the regions of smoothness of the flux; the main difficulty consists in taking into account the jump discontinuities of the flux. To do so, for a quite general setting one may only assume that

$$\text{for all } k \in \mathbb{R}, \quad f(\cdot, \cdot, k) \in BV_{loc}(\mathbb{R}^+ \times \mathbb{R}^d). \quad (2)$$

This rather weak regularity appears naturally e.g. in the study of triangular systems of conservation laws (see [30] and references therein). In the framework (2), under a non-degeneracy assumption of the fluxes $k \mapsto f(t, \mathbf{x}, k)$, existence of solutions satisfying the family of entropy inequalities

$$\begin{aligned} \forall k \in \mathbb{R} \quad & |u - k|_t + \operatorname{div}_{\mathbf{x}}(\operatorname{sgn}(u - k)(f(t, \mathbf{x}, u) - f(t, \mathbf{x}, k))) - \operatorname{sgn}(u - k)S(t, \mathbf{x}, u) \\ & \leq -\operatorname{sgn}(u - k)(\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k))^{ac} + |(\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k))^s| \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^d) \end{aligned} \quad (3)$$

has been proved by Panov in [40] using vanishing viscosity method; here,

$$\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k) = (\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k))^{ac} + (\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k))^s$$

is the Jordan decomposition of the Radon measure $\operatorname{div}_{\mathbf{x}} f(t, \mathbf{x}, k)$ into its absolutely continuous part and its singular part (cf. (17) below, for a particular but representative case). Let us stress that inequalities (3) use a roughly estimated contribution of the jump singularities in the flux f , which turns out to be a serious obstacle for proving uniqueness of solutions in the sense (3).

In order to explain this more accurately, let us consider the one-dimensional variant of (1) augmented with the initial conditions $u|_{t=0} = u_0 \in L^\infty(\mathbb{R}^d)$. Condition (3) is the general form of the Karlsen–Risebro–Towers admissibility condition (see [42,43,31]; see also [41,13]):

$$\forall k \in [a, b] \quad \partial_t |u - k| + \partial_x \left\{ \operatorname{sgn}(u - k) [H(x)(f(u) - f(k)) + H(-x)(g(u) - g(k))] \right\} - |f(k) - g(k)| \delta_0(x) \leq 0 \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}). \tag{4}$$

This is precisely (3) written for the case with source $S \equiv 0$ and flux $\mathfrak{f}(t, x; u) = f(u)H(x) + g(u)H(-x)$; here and throughout the paper,

$H(\cdot) = \operatorname{sgn}^+(\cdot)$ is the Heavyside function;

δ_0 is the Dirac measure concentrated at 0, i.e., $\delta_0 = H'$ in $\mathcal{D}'(\mathbb{R})$.

The advantage of definition of solution by the family of inequalities (4) lies in a simple and natural existence and stability proof. Using the vanishing viscosity approximations of (1) for which existence is easy even in the discontinuous-flux setting, one readily obtains for the corresponding solutions u_ε the inequalities

$$\partial_t |u_\varepsilon - k| + \partial_x \left\{ \operatorname{sgn}(u_\varepsilon - k) [H(x)(f(u_\varepsilon) - f(k)) + H(-x)(g(u_\varepsilon) - g(k))] \right\} - \operatorname{sgn}(u_\varepsilon(t, 0) - k) (f(k) - g(k)) \delta_0(x) \leq \varepsilon \partial_{xx} |u_\varepsilon - k| \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}), \tag{5}$$

a.e. on $(0, T)$. Roughly estimating the contribution of the term concentrated on $\{x = 0\}$, we can pass to the limit and get (4). In a similar way, stability of solutions in the sense (4) under a.e. convergence can be justified.

Concerning the question of uniqueness of solutions admissible in the Karlsen–Risebro–Towers sense, the following facts are known:

- Given a datum u_0 , uniqueness of an admissible solution in the sense (4) to the above problem holds true, provided f, g satisfy the *crossing condition*

$$f(u) - g(u) < 0 < f(v) - g(v) \quad \Rightarrow \quad u < v \tag{6}$$

on the shape of the fluxes f and g . For the proof, see Towers [42,43] (see also [41]) for the case where the fluxes f, g have no crossing points in (a, b) ; see Karlsen, Risebro and Towers [31] for general uniqueness argument under assumption (6).

- If the crossing condition (6) fails, then at least for some initial data there exist more than one admissible solution in the sense (4), see [11, Sect. 4.7].

Several attempts were made already to improve the Karlsen–Risebro–Towers conditions. Let us mention here the following directions.

1. In the work [34], the idea was to use a transformation of the original equation (cf. [39] and [16]) in order to enforce the crossing condition for the fluxes depending on the transformed unknown; yet such transformation is often artificial with respect to the underlying model.

2. In [25,11,10] the crossing condition was dropped.

(2a) In [25], a new version of the Γ -condition on the solution jumps was proposed, under which the uniqueness proof was achieved without the crossing condition. Admissible solutions were characterized in terms of generalized Oleinik jump inequalities.

(2b) A different though equivalent to [25] admissibility condition was proposed in [11,10] in terms of the *vanishing viscosity germ* \mathcal{G}_{VV} recalled in Section 2.3 below. It was shown that \mathcal{G}_{VV} -entropy solutions are always unique, regardless the shape of the fluxes. The way to express this admissibility condition is rather tricky. For a straightforward approach put forward in [11], one uses intricate interface coupling conditions. Alternatively, a carefully selected family of *adapted entropies* (see [14,12,18]) can be used to characterize admissible solutions, in the place of the classical Kruzhkov entropies exploited in (4). Neither the coupling conditions encoded in the germ \mathcal{G}_{VV} , nor the associated choice of adapted entropies are self-evident; their relation to the vanishing viscosity approximation follows from a lengthy analysis of possible viscosity profiles.

To sum up, none of the aforementioned approaches of admissibility is as intuitive and appealing as (3) or (4). In the present paper, our goal is to give a definition of solution to (1) which, similarly to the definitions (3) or (4), could be seen as a natural one and that would lead to well-posedness without assuming the crossing condition. Notice that in passing, we give an equivalent and somewhat more natural characterization of \mathcal{G}_{VV} -entropy solutions of [11,10].

1.2. Singular values of u and refined entropy inequalities

The idea of this paper is to strengthen the definitions of [31,40] by suggesting a finer way to take into account the contribution of the flux jumps into the entropy inequality.¹ To explain the idea, we go back to the general framework (2); in this case, we suggest to replace (3) by the following less restrictive inequalities:

$$\begin{aligned} \forall k \in \mathbb{R} \quad & |u - k|_t + \operatorname{div}_{\mathbf{x}}(\operatorname{sgn}(u - k)(\mathfrak{f}(t, \mathbf{x}, u) - \mathfrak{f}(t, \mathbf{x}, k))) \\ & \leq -\operatorname{sgn}(p_u - k) \operatorname{div}_{\mathbf{x}} \mathfrak{f}(t, \mathbf{x}, k) \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^d) \end{aligned} \quad (7)$$

where p_u is some globally defined Borel function satisfying the property

$$p_u = u \quad \text{a.e. with respect to the Lebesgue measure.} \quad (8)$$

Introducing everywhere defined p_u means that we tacitly assign a value to u also on the singular set² of (t, \mathbf{x}) where the flux $(t, \mathbf{x}) \mapsto \mathfrak{f}(t, \mathbf{x}, k)$ experiences jumps or Cantor-type singularities.³ In the sequel, we will say that p_u encodes the *singular values* of u .

Remark 1. Let us present several observations concerning p_u .

(i) Introduction of unknown singular values p_u on the jump manifolds Σ may appear as a naive attempt to resolve the difficulty of definition of solution: indeed, in general these values cannot be observable.

(ii) Yet from the modeling viewpoint, existence of singular values p_u on Σ can be put in relation with the fact that the vanishing viscosity approximation ensures global continuity of the approximate solutions u_ε . This approximation is suitable for models where the solution is expected to be continuous across Σ at a finer spatial scale, undergoing rapid transition of interface layer kind. Such fine-scale continuity assumptions are also natural in numerical approximation and modeling: we refer to [21] for an early work based on this idea, and to [9] for a deeper discussion of a wide class of related modeling hypotheses.

(iii) In practice, whenever the values of approximate solutions u_ε on a jump manifold Σ of the flux $(t, \mathbf{x}) \mapsto \mathfrak{f}(t, \mathbf{x}, k)$ happen to exhibit a non-oscillatory behavior, one obtains p_u on Σ as the pointwise limit of $u_\varepsilon|_\Sigma$, for some vanishing sequence of ε . Mathematically, *a priori* justification of strong convergence of $u_\varepsilon|_\Sigma$ to some limit p_u seems out of reach even when strong compactness of $(u_\varepsilon)_\varepsilon$ in L^1_{loc} topology is easy to justify.

(iv) It is also clear that singular values p_u need not be uniquely defined with respect to the Hausdorff measure on Σ in its natural dimension d : to observe this, it is enough to consider the simplest case of converging one-dimensional viscosity profiles that may provide a continuum of different values for p_u . Proposition 11 suggests some canonical choice of p_u .

1.3. A brief description of technical ideas and obtained results

Throughout the paper we assume that

$$\begin{aligned} & \text{at a.e. point of } \Sigma \text{ (with respect to the } d\text{-dimensional Hausdorff measure),} \\ & \Sigma \text{ is not orthogonal to the direction of the time axis.} \end{aligned} \quad (9)$$

As a matter of fact, we are not able to suggest technical tools that would permit to exploit definition (7) in the general setting (2), (9) neither in view of existence nor in view of uniqueness of solutions. We refer to the recent work of Crasta, De Cicco and De Philippis [22] for uniqueness results in the framework of discontinuous SBV flux \mathfrak{f} (this

¹ This approach goes back to the preprint [35] of the second author. Shortly after the completion of the present paper, the work [22] of Crasta, De Cicco and De Philippis has been completed. Results and techniques of [22] appear as complementary to the ours: the flux \mathfrak{f} in [22] is less regular, the solution u is more regular than in our work, moreover, the uniqueness proof of [22] relies on the kinetic formulation, cf. [13].

² The literature on discontinuous-flux problems concentrates on the case where the union in $k \in \mathbb{R}$ of the sets of singularities of $(t, \mathbf{x}) \mapsto \operatorname{div}_{\mathbf{x}} \mathfrak{f}(t, \mathbf{x}, k)$ is of Lebesgue measure zero, typically it is a locally finite union Σ of hypersurfaces in $\mathbb{R}^+ \times \mathbb{R}^d$. In this case one can say that u is defined on the set of full measure excluding singularities, and p_u extends u to the singular set Σ .

³ To deal with the general case, rigorous meaning should be given to the last term of (7). The idea of formulation (7), (8) has been deeply developed in [22], using fine tools of analysis of BV and SBV (see, e.g., [7]) functions.

means that in the Jordan decomposition of $\operatorname{div}_{\mathbf{x}} \mathfrak{f}(t, \mathbf{x}, k)$, the singular part $(\operatorname{div}_{\mathbf{x}} \mathfrak{f}(t, \mathbf{x}, k))^s$ has zero Cantor part: see, e.g. [7]). In [22], the authors rely on BV regularity of solutions, which may fail in general, cf. [1]; further analysis is needed to provide sufficient conditions for existence of BV solutions or to relax assumptions of the uniqueness theorem of [22]. In contrast to [22], we deal with merely L^∞ solutions which existence is proved but the results of our paper only concern the practical case of piecewise Lipschitz, jump discontinuous across a union Σ of Lipschitz hypersurfaces, and genuinely nonlinear fluxes.

In this case, firstly, we are able to prove uniqueness of admissible solutions in the sense (7), (8) without any artificial condition on the flux crossing. Uniqueness result for solutions in the sense of (7), (8) is based upon consideration of constraints that are imposed by inequalities (7) on the couple $(u^-, u^+)(t_0, \mathbf{x}_0)$ of one-sided traces of a solution u at a point $(t_0, \mathbf{x}_0) \in \Sigma$ where Σ (called interface in the sequel) is a jump discontinuity hypersurface of a piecewise smooth flux function $(t, \mathbf{x}) \mapsto \mathfrak{f}(t, \mathbf{x}, k)$ and u^-, u^+ are strong one-sided traces of the admissible solution u in the sense of Definition 1. The traces can be seen as limits, in an appropriate sense, of $u((t_0, \mathbf{x}_0) \pm h\nu|_\Sigma(t_0, \mathbf{x}_0))$, as $h \downarrow 0$. Here, $\nu|_\Sigma(t_0, \mathbf{x}_0)$ is a normal vector to Σ with some orientation fixed by the choice of local coordinates in a neighborhood of (t_0, \mathbf{x}_0) , and one can also define the traces of flux functions

$$\mathfrak{f}^\pm(\cdot, k) := \lim_{h \downarrow 0} \mathfrak{f}(\cdot \pm h\nu(\cdot), k). \tag{10}$$

The essence of the uniqueness argument is to justify the fact that in the global Kato inequality for two solutions u and v , which formally reads

$$\begin{aligned} \partial_t |u - v| + \operatorname{div}_{\mathbf{x}}(\operatorname{sgn}(u - v)(\mathfrak{f}(t, \mathbf{x}, u) - \mathfrak{f}(t, \mathbf{x}, v))) &\leq L|u - v| + I\mathcal{H}^d|_\Sigma \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^d), \\ I &:= \{\operatorname{sgn}(u^+ - v^+)(\mathfrak{f}^+(\cdot, u^+) - \mathfrak{f}^+(\cdot, v^+)) - \operatorname{sgn}(u^- - v^-)(\mathfrak{f}^-(\cdot, u^-) - \mathfrak{f}^-(\cdot, v^-))\} \cdot \nu(\cdot), \end{aligned}$$

the contribution of the interface term I on the right-hand side is non-positive. Here $\mathcal{H}^d|_\Sigma$ is the d -dimensional Hausdorff measure supported on the interface Σ , and L is the uniform in (t, \mathbf{x}) Lipschitz constant of the source term $S(t, \mathbf{x}, \cdot)$. The non-positivity of I follows indeed from the restrictions on u^\pm imposed by (7), see Section 2.2 (cf. [11]).

The genuine nonlinearity assumption on the flux, i.e., the assumption that for every (ξ_0, ξ) in the d -dimensional unit sphere S^d , there holds

for almost every $(t, \mathbf{x}) \in \mathbb{R}^+ \times \mathbb{R}^d$, the mapping

$$k \mapsto \xi_0 k + \xi \cdot \mathfrak{f}(t, \mathbf{x}, k) \text{ is not constant on non-degenerate intervals,} \tag{11}$$

is a technical hypothesis. It ensures existence of strong interface traces u^\pm on Σ , on which our uniqueness proof heavily relies. While this property can be circumvented in the space–time homogeneous setting of [11] and of many related references, in our case it is essential. Indeed, our approach to existence of admissible solutions also relies on existence of u^\pm , but also on invariance of the considered class of equations under variables’ changes. Therefore, we have to work with (t, \mathbf{x}) -dependent fluxes (and with source terms); in this general setting, assumption (11) is the only known condition that guarantees existence of traces (see [6] for more information on traces of entropy solutions to an inhomogeneous conservation law).

Assumption (11) is essential for our existence proof also because it ensures compactness of suitable vanishing viscosity approximations. It is standard, in this context, to assume in addition some conditions that ensure existence of invariant regions for (1); they are needed to obtain uniform L^∞ bounds on sequences of approximate solutions (see, e.g., [11, Sect. 6]). Here, we take the simplest of such assumptions that is:

$$\begin{aligned} \exists [a, b] \subset \mathbb{R} \quad \text{such that} \quad &\text{for a.e. } t, \quad \mathfrak{f}(t, \cdot, a) \equiv \operatorname{const}, \quad \mathfrak{f}(t, \cdot, b) \equiv \operatorname{const}, \\ &\text{for a.e. } (t, \mathbf{x}), \quad \text{one has} \quad S(t, x, a) \geq 0 \quad \text{and} \quad S(t, x, b) \leq 0, \\ &\text{and the initial datum fulfills} \quad a \leq u_0(\cdot) \leq b. \end{aligned} \tag{12}$$

Assumption (12) ensures that $u(t, \mathbf{x}) \equiv a$ (respectively, $u(t, x) \equiv b$) is a sub-solution (resp., a super-solution) to the Cauchy problem for (1) with initial condition u_0 , thus (12) confines to the interval $[a, b]$ the values of solutions (and also the values of suitably constructed approximate solutions) we will consider.

We are able to prove existence of solutions defined by (7), (8) not by surpassing the technical difficulty of passage to the limit in approximate entropy inequalities (5) but using an indirect and rather lengthy argument. This argument

exploits the idea of the existence proof in [10], developed in the simple case of a flat interface; see Remark 3 for an explanation of this choice. The study of the flat case is combined with appropriately chosen changes of variables and a principle of invariance of (7), (8) under such variables' changes (see in particular Proposition 14). We are able to include changes of variables that are singular in a neighborhood of $(d - 1)$ -dimensional singularities such as intersections of different d -dimensional interfaces. As a typical and illuminating example, we consider the case of interface $\Sigma \subset \mathbb{R}^+ \times \mathbb{R}$ consisting in two Lipschitz curves merging into one.

1.4. Outline of the paper

The paper is organized as follows. In Section 2 we treat the case that is fundamental for our techniques. More precisely, in this section we deal with multi-dimensional scalar conservation laws in heterogeneous setting, with a source term but with a single flat discontinuity interface Σ . Definition of solution admissibility is stated in Section 2.1 and further discussed in Section 2.5. Uniqueness of an admissible solution to the Cauchy problem is proved in Section 2.2, then a different proof exploiting the machinery of [11,10] is sketched in Section 2.3. Existence is justified in Section 2.4, with an explicit construction of p_u in terms of interface traces u^\pm of the vanishing viscosity limit u .

Section 3 deals with the general geometry of jump discontinuity interfaces of the flux f in (1). In Section 3.1, we sketch the uniqueness argument, explain and motivate the existence strategy. In Section 3.2, we give main details on the idea of a global singular change of variables that permits to reduce the case of flux f with Lipschitz jump discontinuity manifolds of rather general form (possibly curved, intersecting, and closed) to the locally flat case with singularities that can be ignored. With minor modifications with respect to Section 2.4, existence in this locally flat case is justified. In Section 3.3, examples of changes of variables satisfying the general assumptions of Section 3.2 are presented. In addition, in Section 3.4 we briefly describe an alternative construction of solutions where regular local changes of variables are pieced together using a kind of partition of unity. In Remark 4, we underline the fact that in both approaches, a viscosity approximation adapted to interface geometry is essential for the existence proof.

For readers' convenience, let us point out that Definition 1 with well-posedness results of Theorems 4, 10 summarize our work for the model case with flat interface. For the general case of piecewise Lipschitz, jump discontinuous in (t, \mathbf{x}) flux f , Theorem 15 (see also the conclusion of Section 3.4) provides a set of sufficient conditions under which the new interpretation (7), (8) of Eq. (1) leads to well-posedness of the Cauchy problem.

2. New admissibility conditions for multidimensional heterogeneous scalar conservation laws with a flat discontinuity

This is the fundamental section, indeed, all the other situations that we can resolve are reduced, at least locally, to the Cauchy problem investigated here.

2.1. Definition of solution and interface traces

Consider the problem

$$\partial_t u + \operatorname{div}(F(t, \mathbf{x}, u)H(x_1) + F(t, \mathbf{x}, u)H(-x_1)) = S(t, \mathbf{x}, u), \quad (13)$$

$$u|_{t=0} = u_0, \quad (14)$$

where $F = (F_1, \dots, F_d) : \mathbb{R}^+ \times \mathbb{R}^d \times [a, b] \rightarrow \mathbb{R}^d$ and $G = (G_1, \dots, G_d) : \mathbb{R}^+ \times \mathbb{R}^d \times [a, b] \rightarrow \mathbb{R}^d$. This is (1) with the flux given by

$$f(t, \mathbf{x}, \cdot) = F(t, \mathbf{x}, \cdot)H(x_1) + G(t, \mathbf{x}, \cdot)H(-x_1). \quad (15)$$

We will assume that the flux and the source S satisfy the compatibility conditions at $k = a$ and $k = b$, as required in assumption (12). We also assume that

$$\begin{aligned} &\text{the fluxes } F, G \text{ are globally Lipschitz continuous in all variables,} \\ &\text{the source } S \text{ is globally Lipschitz continuous in } u \in [a, b]. \end{aligned} \quad (16)$$

Here and in the sequel of this paper, $\hat{x}_1 = (x_2, \dots, x_d)$. The unique interface (jump discontinuity hypersurface) for the flux (15) is given by

$$\Sigma = \{(t_0, 0, \hat{x}_{10}) \mid t_0 \in \mathbb{R}^+, \hat{x}_{10} \in \mathbb{R}^{d-1}\}.$$

Throughout this section, somewhat abusively we denote by (t_0, \mathbf{x}_0) both the points of Σ with coordinates (t_0, \hat{x}_{10}) and the points of $\mathbb{R} \times \mathbb{R}^d$ with coordinates $(t_0, 0, \hat{x}_{10})$.

As explained in the introduction, the main goal of this paper is to propose and justify, by proving well-posedness results, the following definition.

Definition 1. We say that a function $u \in C(\mathbb{R}^+; L^1_{loc}(\mathbb{R}^d))$ taking values in $[a, b]$ is an entropy admissible solution to (13), (14) if $u(0, \cdot) = u_0$ and there exists a function $p_u : \Sigma \rightarrow [a, b]$ such that $\forall k \in [a, b]$

$$\begin{aligned} & \partial_t |u - k| + \operatorname{div}_{\mathbf{x}} \left\{ \operatorname{sgn}(u - k) \left(H(x_1) (F(t, \mathbf{x}, u) - F(t, \mathbf{x}, k)) + H(-x_1) (G(t, \mathbf{x}, u) - G(t, \mathbf{x}, k)) \right) \right\} \\ & - \operatorname{sgn}(u - k) S(t, \mathbf{x}, u) + \operatorname{sgn}(u - k) \left(H(x_1) \operatorname{div}_{\mathbf{x}} F(t, \mathbf{x}, k) + H(-x_1) \operatorname{div}_{\mathbf{x}} G(t, \mathbf{x}, k) \right) \\ & - \operatorname{sgn}(p_u(t, \hat{x}_1) - k) (F_1(t, \mathbf{x}, k) - G_1(t, \mathbf{x}, k)) \delta_0(x_1) \leq 0 \quad \text{in } \mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^d). \end{aligned} \tag{17}$$

To shorten the calculations, in the sequel we will assume $S \equiv 0$; the general case is obtained with the help of the Gronwall inequality. The technique of the proofs of this section readily extends to a locally finite number of flat, possibly crossing discontinuity surfaces, but we will stick to the case of one interface $\Sigma = \{x_1 = 0\}$. Let us stress that presence of a source term S and of multiple flat discontinuity hypersurfaces is required in order to reduce the case of curved, crossing or closed interfaces considered in Section 3 to the case of flat interfaces investigated in the present section.

Clearly, inequalities (17) (with $S \equiv 0$) in the one-dimensional situation imply the Karlsen–Risebro–Towers inequalities (4), therefore it is natural to expect that the uniqueness becomes easier to justify, while existence proof will present new and considerable difficulties. Indeed, in the present section we will show that

- an admissible solution in the sense of (17) is unique, with a rather standard proof involving a tedious case-by-case study;
- the standard vanishing viscosity method converges towards this solution, with a quite indirect proof based upon construction of viscosity profiles.

Both results are achieved by looking at the values of one-sided traces of the solutions on the interface, so we start by defining these traces and giving sufficient conditions for their existence.

Definition 2. We say that an integrable function W admits left trace W^- and right trace W^+ on $\Sigma = \{x_1 = 0\}$ if there exist functions $W^\pm : \mathbb{R}^+ \times \mathbb{R}^{d-1} \mapsto \mathbb{R}$ such that for every $\varphi \in C_c(\mathbb{R}^{d-1})$ there holds

$$\lim_{h \rightarrow 0^+} \int_{\mathbb{R}^+ \times \mathbb{R}^{d-1}} |W(t, \pm h, \hat{x}_1) - W^\pm(t, \hat{x}_1)| \varphi(t, \hat{x}_1) dt d\hat{x}_1 = 0. \tag{18}$$

While general L^∞ functions do not admit such traces, local entropy solutions of conservation laws admit traces in the sense (18) under additional, not very restrictive technical assumptions. Under assumption (11) on the flux (15), the arguments from [38] provide existence of traces to local Kruzhkov entropy solutions to (13) in the non-homogeneous situation as well (see also [6]). Indeed, while the proof from [38] is given for case where the flux does not depend on time or space variables (i.e. it is homogeneous), its arguments extend to the general, piecewise Lipschitz (t, \mathbf{x}) -dependent setting with condition (11) (see in particular [6]). Let us mention in passing that in one dimensional homogeneous situation, the linear degeneracy of fluxes can also be treated via introduction of “singular mappings” that have traces in all cases (see [11]) or by considering traces of the flux and of entropy fluxes instead of the traces of the solution itself (see [34]).

Throughout this section, we will denote by $u^-(t_0, \mathbf{x}_0)$, respectively $u^+(t_0, \mathbf{x}_0)$, the left trace, respectively the right trace at $(t_0, \mathbf{x}_0) \in \Sigma$ of an admissible solution u to (13). We readily derive the Rankine–Hugoniot conditions for admissible solutions.

Lemma 3. *Let u be an admissible solution to (13), (14) in the which admits left and right strong traces at the interface u^+ and u^- , respectively. Then, for \mathcal{H}^d -a.e. point $(t_0, \mathbf{x}_0) \in \Sigma$, there holds*

$$F_1(t_0, \mathbf{x}_0, u^+(t_0, \mathbf{x}_0)) = G_1(t_0, \mathbf{x}_0, u^-(t_0, \mathbf{x}_0)). \tag{19}$$

Proof. Taking $k = a$ then $k = b$ in the entropy inequalities (17), one deduces the weak formulation of (13). Then it is enough to test this weak formulation of (13) on the function of the form $\eta = \varphi\mu_h$, $\varphi \in C_c^1(\mathbb{R}^+ \times \mathbb{R}^d)$, where μ_h is given by

$$\mu_h(\mathbf{x}) = \begin{cases} \frac{1}{h}(x_1 + 2h), & x_1 \in [-2h, -h] \\ 1, & x \in [-h, h] \\ \frac{1}{h}(2h - x_1), & x_1 \in [h, 2h] \\ 0, & |x_1| > 2h, \end{cases} \tag{20}$$

which can be taken for test function by approximation. After letting $h \rightarrow 0$, we reach to identity (19) in $\mathcal{D}'(\Sigma)$; since both sides of (19) are bounded functions, the identity also holds pointwise, \mathcal{H}^d -a.e. on Σ . \square

2.2. A uniqueness and L^1 contraction proof

We prove the following result. Let $B(0, R) = \{x \in \mathbb{R}^d \mid |x| < R\}$.

Theorem 4. *Assume that the flux (15) satisfies (11), (12), (16). Let u, v be two entropy admissible solutions to (13) in the sense of Definition 1 with source $S \equiv 0$ and initial data u_0 and v_0 , respectively. Then, for every $R > 0$, $T > 0$ there exists $C > 0$ such that*

$$\int_0^T \int_{B(0,R)} |u(t, x) - v(t, x)| dx dt \leq T \int_{B(0,R+CT)} |u_0(x) - v_0(x)| dx. \tag{21}$$

Remark 2. Notice that the order-preservation result ($u_0 \geq v_0$ implies $u \geq v$) can be proved in the same way. The first proof given below is a self-contained case-by-case study. The second proof that will be sketched in Section 2.3 establishes the equivalence between our new definition and the definition of \mathcal{G}_{VV} -entropy solution (see [11,10]) for which uniqueness follows readily from the general theory developed in [11] (see also [25]).

Proof. We start as in the proof of Lemma 3, but using the entropy inequalities with general k . Namely, insert into (17) the function $\psi = \mu_h\varphi$, where $\varphi \in C_c^1(\mathbb{R}_+^d)$ while μ_h is given by (20). Letting $h \rightarrow 0$, due to arbitrariness of φ , as in Lemma 3 we reach to the following relation for almost every $(t_0, \mathbf{x}_0) \in \Sigma$

$$\begin{aligned} & \operatorname{sgn}(u^+ - k)(F_1(t_0, \mathbf{x}_0, u^+) - F_1(t_0, \mathbf{x}_0, k)) - \operatorname{sgn}(u^- - k)(G_1(t_0, \mathbf{x}_0, u^-) - G_1(t_0, \mathbf{x}_0, k)) \\ & + \operatorname{sgn}(p_u - k)(F_1(t_0, \mathbf{x}_0, k) - G_1(t_0, \mathbf{x}_0, k)) \leq 0, \end{aligned} \tag{22}$$

where for the sake of brevity, we write

$$p_u(t_0, \mathbf{x}_0) = p_u, \quad u^+(t_0, \mathbf{x}_0) = u^+, \quad u^-(t_0, \mathbf{x}_0) = u^-.$$

Notice that in the passage to the limit, we used continuity of the maps

$$\begin{aligned} (t, \mathbf{x}, z) & \mapsto \operatorname{sgn}(z - k)(F(t, \mathbf{x}, z) - F(t, \mathbf{x}, k)) \\ (t, \mathbf{x}, z) & \mapsto \operatorname{sgn}(z - k)(G(t, \mathbf{x}, z) - G(t, \mathbf{x}, k)). \end{aligned} \tag{23}$$

Remark that if

$$k \geq \max\{u^+, u^-, p\} \quad \text{or} \quad k \leq \min\{u^+, u^-, p\}$$

then the left-hand side in (22) is equal to zero according to the Rankine–Hugoniot relation (19). Now, we make precise the information contained in the other possible cases where (22) holds true. To shorten the statements like (19), (22) in the sequel we will use the notation

$$f(\cdot) := F_1(t_0, \mathbf{x}_0, \cdot) \quad \text{and} \quad g(\cdot) := G_1(t_0, \mathbf{x}_0, \cdot). \tag{24}$$

Case I: $u^+ \leq u^-$.

- $u^+ \leq u^- \leq p_u$, where two different cases occur:

$$u^+ \leq u^- \leq k \leq p_u, \quad \text{which implies } f(k) \leq g(k); \tag{25}$$

$$u^+ \leq k \leq u^- \leq p_u, \quad \text{which implies } f(k) \leq f(u^+). \tag{26}$$

- $u^+ \leq p_u \leq u^-$, where we also have two possible cases:

$$u^+ \leq p_u \leq k \leq u^-, \quad \text{which implies } g(k) \leq g(u^-); \tag{27}$$

$$u^+ \leq k \leq p_u \leq u^-, \quad \text{which implies } f(k) \leq f(u^+). \tag{28}$$

- $p \leq u^+ \leq u^-$, here we have the following alternative:

$$p \leq u^+ \leq k \leq u^-, \quad \text{which implies } f(k) \leq f(u^+); \tag{29}$$

$$p \leq k \leq u^+ \leq u^-, \quad \text{which implies } g(k) \leq f(k). \tag{30}$$

Case II: $u^- \leq u^+$.

- $u^- \leq u^+ \leq p_u$, where we have one of the two situations:

$$u^- \leq u^+ \leq k \leq p_u, \quad \text{which implies } f(k) \leq g(k); \tag{31}$$

$$u^- \leq k \leq u^+ \leq p_u, \quad \text{which implies } g(u^-) \leq g(k). \tag{32}$$

- $u^- \leq p_u \leq u^+$, where the alternative is:

$$u^- \leq p_u \leq k \leq u^+, \quad \text{which implies } f(u^+) \leq f(k); \tag{33}$$

$$u^- \leq k \leq p_u \leq u^+, \quad \text{which implies } g(u^-) \leq g(k). \tag{34}$$

- $p_u \leq u^- \leq u^+$, where we have the last two possibilities:

$$p_u \leq u^- \leq k \leq u^+, \quad \text{which implies } f(u^+) \leq f(k); \tag{35}$$

$$p_u \leq k \leq u^- \leq u^+, \quad \text{which implies } g(k) \leq f(k). \tag{36}$$

Now, we are ready to pass to the main part of the uniqueness proof, which consists in derivation of the Kato inequality: for every $\varphi \in C_c^1(\mathbb{R}^+ \times \mathbb{R}^d)$,

$$\int_{\mathbb{R}^+ \times \mathbb{R}^d} \{ |u - v| \varphi_t + \text{sgn}(u - v) ((F(t, \mathbf{x}, u) - F(t, \mathbf{x}, v)) H(x_1) + (G(t, \mathbf{x}, u) - G(t, \mathbf{x}, v)) H(-x_1)) \cdot \nabla \varphi \} d\mathbf{x} dt \geq 0. \tag{37}$$

The classical doubling of variables technique of [33] ensures that (37) holds for $\varphi \in C_c^1((\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Sigma)$. Therefore, given an arbitrary $\psi \in C_c^1(\mathbb{R}^+ \times \mathbb{R}^d)$, inequality (37) is satisfied with the test function $\varphi = (1 - \mu_h)\psi$, where μ_h is defined by (20). Letting $h \rightarrow 0$, we get

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^d} \{ |u - v| \psi_t + \text{sgn}(u - v) ((F(t, \mathbf{x}, u) - F(t, \mathbf{x}, v)) H(x_1) + (G(t, \mathbf{x}, u) - G(t, \mathbf{x}, v)) H(-x_1)) \nabla \psi \} d\mathbf{x} dt \\ & \geq \int_{\Sigma} (-\text{sgn}(u^+ - v^+)) (F(t_0, \mathbf{x}_0, u^+) - F(t_0, \mathbf{x}_0, v^+)) \\ & \quad + \text{sgn}(u^- - v^-) (G(t_0, \mathbf{x}_0, u^-) - G(t_0, \mathbf{x}_0, v^-)) \psi(t_0, \mathbf{x}_0) d\hat{x}_{10} dt_0 \\ & =: \int_{\Sigma} \Delta(u^\pm, v^\pm)(t_0, \mathbf{x}_0) \psi(t_0, \mathbf{x}_0) d\hat{x}_{10} dt_0, \quad \mathbf{x}_0 = (0, \hat{x}_{10}). \end{aligned} \tag{38}$$

Now, we prove that the integrand $\Delta(u^\pm, v^\pm)$ (that is a short-cut notation for $\Delta(t_0, \mathbf{x}_0, u^\pm(t_0, \mathbf{x}_0), v^\pm(t_0, \mathbf{x}_0))$) in the right-hand side of the latter expression is greater than or equal to zero, for almost every $(t_0, \mathbf{x}_0) = (t, 0, \hat{x}_{10}) \in \Sigma$.

The proof is tedious and it is accomplished by considering numerous different possibilities depending on pointwise relations between the values u^\pm , v^\pm and p_u , p_v .

Concerning the relation between u^\pm and v^\pm , we see that, according to the Rankine–Hugoniot conditions, the quantity $\Delta(u^\pm, v^\pm)$ can be non-zero only when

$$u^- > v^- \text{ and } u^+ < v^+ \quad \text{or} \quad u^- < v^- \text{ and } u^+ > v^+ \quad (39)$$

(the proof is the same as in Cases 1–5 in the proof of [31, Theorem 2.1]). On the other hand, the two cases from (39) are symmetric since $\Delta(u^\pm, v^\pm) = \Delta(v^\pm, u^\pm)$, and therefore their analysis is the same. Thus, it is enough to prove the inequality $\Delta(u^\pm, v^\pm) \geq 0$ whenever the first relation from (39) is satisfied. We proceed by considering the following possible sub-cases:

Case 1: $u^+ < v^+ < v^- < u^-$

Case 2: $u^+ < v^- < v^+ < u^-$

Case 3: $v^- < u^+ < v^+ < u^-$

Case 4: $v^- < u^+ < u^- < v^+$

Case 5: $v^- < u^- < u^+ < v^+$.

Notice that, according to the disposition of u^\pm and v^\pm and the Rankine–Hugoniot condition (19), one has

$$\begin{aligned} \Delta(u^\pm, v^\pm) &= -\operatorname{sgn}(u^+ - v^+)(f(u^+) - f(v^+)) + \operatorname{sgn}(u^- - v^-)(g(u^-) - g(v^-)) \\ &= f(u^+) - f(v^+) + g(u^-) - g(v^-) = 2(f(u^+) - f(v^+)) = 2(g(u^-) - g(v^-)). \end{aligned}$$

Thus, we aim to prove that in each case of the above list, there holds

$$f(u^+) - f(v^+) \geq 0 \quad \text{or} \quad g(u^-) - g(v^-) \geq 0. \quad (40)$$

Since $u, v \in L^\infty(\mathbb{R}^+ \times \mathbb{R}^d)$ are two admissible solutions to (13), we consider two functions p_u and p_v from Definition 1 representing the singular values on Σ corresponding to u and v , respectively.

Case 1 For almost every fixed $(t_0, \mathbf{x}_0) \in \Sigma$, we have the following possibilities.

- $u^+ < v^+ < v^- < u^- < p_u$:

The conclusion follows by taking $k = v^+$ in (26).

- $u^+ < v^+ \leq p_u \leq u^-$:

The conclusion follows by taking $k = v^+$ in (28).

- $u^+ \leq p_u \leq v^- < u^-$:

The conclusion follows by taking $k = v^-$ in (27).

- $p_u < u^+ < v^+ < v^- < u^-$:

The conclusion follows by taking $k = v^-$ in (29).

Case 5 This case is symmetric with the previous one. We simply need to consider the position of p_v instead of p_u and to apply (31)–(36) instead of (25)–(30).

Case 2 We have the following possibilities.

- $u^+ < v^- < v^+ < u^- < p_u$:

The conclusion follows by taking $k = v^+$ in (26).

- $u^+ < v^- < v^+ \leq p_u \leq u^-$:

The conclusion follows by taking $k = v^+$ in (28).

- $u^+ < v^- \leq p_u \leq v^+ < u^-$:

Here, we must involve the position of p_v . First, recall that from (27) and (28)

$$\begin{aligned} g(k) &\leq g(u^-), \quad \forall k \in [p_u, u^-] \\ f(k) &\leq f(u^+), \quad \forall k \in [u^+, p_u]. \end{aligned} \quad (41)$$

Now, we have the following possibilities.

- $u^+ < v^- \leq p_u \leq v^+ \leq p_v$.
From (32) (applied on v) and (41), we have $g(v^-) \leq g(p_u) \leq g(u^-)$ which is (40).
- $u^+ < v^- \leq p_u \leq p_v \leq v^+ < u^-$.
From (34) and (41), we have $g(v^-) \leq g(p_u) \leq g(u^-)$.
- $u^+ < v^- \leq p_v \leq p_u \leq v^+ < u^-$.
From (33) and (41), we have $f(v^+) \leq f(p_u) \leq f(u^+)$.
- $p_v \leq v^- \leq p_u \leq v^+ < u^-$.
From (35) and (41), we have $f(v^+) \leq f(p_u) \leq f(u^+)$.

- $u^+ \leq p_u \leq v^- < v^+ < u^-$:
The conclusion follows by taking $k = v^-$ in (27).
- $p_u < u^+ < v^+ < v^- < u^-$:
The conclusion follows by taking $k = v^-$ in (29).

Case 4 This case is symmetric with the previous one. We simply need to consider the position of p_v instead of p_u and to apply (31)–(36) instead of (25)–(30) or vice versa, when needed.

Case 3 We have the following possibilities.

- $v^- < u^+ < v^+ < u^- \leq p_u$:
In this case, the first relation in (40) follows by taking $k = v^+$ in (26).
- $v^- < u^+ < v^+ \leq p_u \leq u^-$:
In this case, (40) follows from (28) by taking $k = v^+$ there.
- $v^- < u^+ \leq p_u \leq v^+ < u^-$:
We must involve the position of p_v again. We have the following possibilities.

- $v^- < u^+ \leq p_u \leq v^+ < u^- \leq p_v$.
From (32), on v , it follows $g(v^-) \leq g(p_u)$ while from (41), $g(p_u) \leq g(u^-)$. Thus, (40) follows.
- $v^- < u^+ \leq p_u \leq v^+ \leq p_v \leq u^-$.
The situation is the same as the previous one.
- $v^- < u^+ \leq p_u \leq p_v \leq v^+ \leq u^-$.
From (34), on v , and (41), it follows $g(v^-) \leq g(p_u) \leq g(u^-)$ which is (40).
- $v^- < u^+ \leq p_v \leq p_u \leq v^+ \leq u^-$.
From (33) and (41), it follows $f(v^+) \leq f(p_u) \leq f(u^+)$ which is (40).
- $v^- \leq p_v \leq u^+ \leq p_u \leq v^+ \leq u^-$.
Relation (40) follows as in the previous case.
- $p_v \leq v^- < u^+ \leq p_u \leq v^+ \leq u^-$.
From (35) and (41), it follows $f(v^+) \leq f(p_u) \leq f(u^+)$.

- $v^- \leq p_u \leq u^+ < v^+ < u^-$:
Relation (40) follows from (29).
- $p_u \leq v^- < u^+ < v^+ < u^-$:
The conclusion is the same as in the previous item.

From the above considerations, we conclude that in all possible cases, there holds $\Delta(u^\pm, v^\pm) \geq 0$. This means that the Kato inequality (37) holds. From here, the proof of the theorem follows in the standard way, see [33,11]. \square

2.3. Uniqueness via reduction to \mathcal{G}_{VV} -entropy solutions

It is possible to reduce uniqueness proof to the setting of [11,10]. Indeed, in these papers uniqueness of so-called \mathcal{G}_{VV} -entropy solutions has been proved. Here, given an ordered couple of continuous functions (f, g) on $[a, b]$, the

vanishing viscosity germ \mathcal{G}_{VV} is the subset of $[a, b]^2$ given by: $(u^-, u^+) \in \mathcal{G}_{VV}$ if $g(u^-) = f(u^+) =: s$ and

either $u^- = u^+$,

or $u^- < u^+$ and there exists

$$u^o \in [u^-, u^+] \quad \text{such that} \quad \begin{cases} g(z) \geq s & \text{for all } z \in [u^-, u^o], \\ f(z) \geq s & \text{for all } z \in [u^o, u^+], \end{cases}$$

or $u^- > u^+$ and there exists

$$u^o \in [u^-, u^+] \quad \text{such that} \quad \begin{cases} g(z) \leq s & \text{for all } z \in [u^o, u^-], \\ f(z) \leq s & \text{for all } z \in [u^+, u^o]. \end{cases} \tag{42}$$

This set is called the *vanishing viscosity germ* associated with the couple (f, g) . We refer to Diehl in [25] for an equivalent description of the set \mathcal{G}_{VV} in terms of Oleinik-kind inequalities. Then, \mathcal{G}_{VV} -entropy solutions are defined as follows.

Definition 5. We say that the function $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R}^d)$ represent \mathcal{G}_{VV} -entropy solution to (13), (14) if

- (17) holds when tested on functions $\varphi \in C_c^1((\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Sigma)$;
- for a.e. $(t_0, \mathbf{x}_0) \in \Sigma$, left $u^-(t_0, \mathbf{x}_0)$ and right $u^+(t_0, \mathbf{x}_0)$ traces of u on Σ are such that $(u^-(t_0, \mathbf{x}_0), u^+(t_0, \mathbf{x}_0))$ belongs to the vanishing viscosity germ associated with the couple (f, g) given by $f : u \mapsto F_1(t_0, \mathbf{x}_0, u)$, $g : u \mapsto G_1(t_0, \mathbf{x}_0, u)$.

Then it is not difficult to check that a solution is admissible in the sense of Definition 1 if and only if it is a \mathcal{G}_{VV} -entropy solution. Indeed, one can take for $p_u(t_0, \mathbf{x}_0)$ in (17) the value $u^o(t_0, \mathbf{x}_0)$ associated with the couple $(u^-, u^+)(t_0, \mathbf{x}_0)$ in definition (42) of \mathcal{G}_{VV} , and vice versa. The complete verification involves essentially the same case studies as in the direct uniqueness proof developed in the previous section, therefore we omit these details.

2.4. The existence proof

Recall that we assumed that the confinement property (12) holds, and that Eq. (13) is genuinely nonlinear in the sense (11). These properties imply that the family of vanishing viscosity approximations for (13), (14) (i.e., solutions of (44) below with initial condition (14)) is $[a, b]$ -valued and that it is strongly precompact [40]; moreover, the corresponding limit admits strong traces at the interface Σ (cf. [6]). Recall that we assume that F, G are given Lipschitz functions on $\mathbb{R}^+ \times \mathbb{R}^d \times [a, b]$. In addition, for the proof of existence we assume that

$$\partial_u F, \partial_u G \text{ are Lipschitz continuous in } (t, x), \mathcal{H}^d\text{-a.e. on } \Sigma. \tag{43}$$

This rather strong, but not restrictive in practice assumption simplifies the proof below; mere continuity and even a uniform in k Lebesgue-point property of $\partial_u F(\cdot, \cdot, k), \partial_u G(\cdot, \cdot, k)$ at \mathcal{H}^d -a.a. point of Σ would be enough for the technique of the proof to work.

Let u be constructed as an accumulation point, as $\varepsilon \rightarrow 0$, of vanishing viscosity approximations $(u_\varepsilon)_\varepsilon$. The essence of the existence proof consists in construction of a function p_u on Σ in order to justify that u is an admissible solution of (13), (14) in the sense of Definition 1. We will need several auxiliary statements before the conclusion can be given. First, we have two lemmas that permit to extract information on $u = \lim_{\varepsilon \rightarrow 0} u_\varepsilon$ from existence of a suitable family of vanishing viscosity profiles $(R_\varepsilon)_\varepsilon$ with prescribed traces of $R := \lim_{\varepsilon \rightarrow 0} R_\varepsilon$ on Σ . We start by justifying a Kato inequality for u_ε and R_ε , where R_ε solves an auxiliary equation with frozen coefficients. Recall that we assumed $S = 0$.

Lemma 6. Assume that in a neighborhood $V(t_0, \mathbf{x}_0)$ of a point $(t_0, \mathbf{x}_0) \in \Sigma$, the family of functions $(u_\varepsilon)_\varepsilon$ in $L^2(0, T; H_{loc}^1(\mathbb{R}^d))$ satisfies equations

$$\partial_t u_\varepsilon + \operatorname{div}_{\mathbf{x}}(H(x_1)F(t, \mathbf{x}, u_\varepsilon) + H(x_1)G(t, \mathbf{x}, u_\varepsilon)) = \varepsilon \Delta u_\varepsilon \tag{44}$$

in the sense of distributions. Assume that the family of $L^2(0, T; H_{loc}^1(\mathbb{R}^d))$ functions $(R_\varepsilon)_\varepsilon$ takes values in $[a, b]$, has its variation is uniformly bounded by a constant M , and R_ε satisfies

$$\partial_t R_\varepsilon + \operatorname{div}_{\mathbf{x}}(F(t_0, \mathbf{x}_0, R_\varepsilon)H(x_1) + G(t_0, \mathbf{x}_0, R_\varepsilon)H(-x_1)) = \varepsilon \Delta R_\varepsilon \tag{45}$$

in the same neighborhood $V(t_0, \mathbf{x}_0)$ of (t_0, \mathbf{x}_0) in the sense of distributions.

Then, the following Kato-type inequality is satisfied:

$$\begin{aligned} \partial_t |u_\varepsilon - R_\varepsilon| + \operatorname{div}_x (\operatorname{sgn}(u_\varepsilon - R_\varepsilon) ((F(t, x, u_\varepsilon) - F(t, x, R_\varepsilon))H(x_1) \\ + (G(t, x, u_\varepsilon) - G(t, x, R_\varepsilon))H(-x_1))) \leq \gamma_\varepsilon(t, \mathbf{x}) + \varepsilon \Delta |u_\varepsilon - R_\varepsilon| \end{aligned} \tag{46}$$

in the sense of distributions in $V(t_0, \mathbf{x}_0)$, where $(\gamma_\varepsilon)_\varepsilon$ is a family of Radon measures in $V(t_0, \mathbf{x}_0)$ verifying

$$\gamma_\varepsilon(t, \mathbf{x}) \leq C(|t - t_0| + |\mathbf{x} - \mathbf{x}_0|)(\delta_0(x_1) + r_\varepsilon(t, \mathbf{x})) + C. \tag{47}$$

Here the uniform in ε constant C depends on the Lipschitz constant of F , G , $\partial_u F$ and $\partial_u G$; while $(r_\varepsilon)_\varepsilon$ is a family of Radon measures with total variation in $V(t_0, x_0)$ bounded by M , uniformly in ε .

Remark 3. The analogous result can be stated for a discontinuous flux \mathfrak{f} having a general Lipschitz jump manifold Σ . In this case, one has to assume that R_ε verify in $V(t_0, \mathbf{x}_0)$ the viscous conservation law with flux coefficients \mathfrak{f}^\pm from (10) frozen at the point (t_0, \mathbf{x}_0) on each side from the jump manifold Σ . Sometimes such profiles R_ε can be constructed, which may involve a source term S_ε in the right-hand side that accounts for the curvature of Σ , see [10] and Section 3.1 for more information. But in general, the construction of R_ε with some prescribed piecewise constant behavior, as $\varepsilon \rightarrow 0$ (this is needed in Lemma 7 below) is very delicate when Σ is not flat; while one-dimensional profiles R_ε with $S_\varepsilon \equiv 0$ are readily constructed for a flat interface. Therefore, in this paper we will stick to the basic choice $R_\varepsilon = R(\frac{x_1}{\varepsilon})$ for a flat boundary $\Sigma = \{(t, \mathbf{x}) \mid x_1 = 0\}$, at a price of rectification arguments developed in Section 3 for reduction of curved manifolds Σ to the flat case.

Proof. First, we subtract (45) from (44), in the resulting expression we add and subtract the measure

$$\operatorname{div}_x (F(t, x, R_\varepsilon)H(x_1) + G(t, x, R_\varepsilon)H(-x_1)).$$

We get

$$\begin{aligned} \partial_t (u_\varepsilon - R_\varepsilon) + \operatorname{div}_x [(F(t, \mathbf{x}, u_\varepsilon) - F(t, \mathbf{x}, R_\varepsilon))H(x_1) - (G(t, \mathbf{x}, u_\varepsilon) - G(t, \mathbf{x}, R_\varepsilon))H(-x_1)] \\ + \operatorname{div}_x (F(t, \mathbf{x}, R_\varepsilon)H(x_1) + G(t, \mathbf{x}, R_\varepsilon)H(-x_1)) - \operatorname{div}_x (F(t_0, \mathbf{x}_0, R_\varepsilon)H(x_1) + G(t_0, \mathbf{x}_0, R_\varepsilon)H(-x_1)) \\ = \varepsilon \Delta (u_\varepsilon - R_\varepsilon). \end{aligned}$$

This equality is understood in the distributional sense in $V(t_0, \mathbf{x}_0)$. Due to the $L^2(0, T; H^1_{loc}(\Omega))$ regularity of $u_\varepsilon, R_\varepsilon$, proceeding by approximation we can multiply this expression by $\varphi \operatorname{sgn}_\alpha(u_\varepsilon - R_\varepsilon)$ where $\operatorname{sgn}_\alpha$ is a Lipschitz regularization of sgn and φ is a localizing test function. Classical chain-rule and passage-to-the-limit in the regularization parameter α arguments apply (see, e.g., [36,20]). In this way, we obtain in $\mathcal{D}'(V(t_0, \mathbf{x}_0))$ the following Kato-like inequality:

$$\begin{aligned} \partial_t |u_\varepsilon - R_\varepsilon| + \operatorname{div}_x [\operatorname{sgn}(u_\varepsilon - R_\varepsilon) ((F(t, \mathbf{x}, u_\varepsilon) - F(t, \mathbf{x}, R_\varepsilon))H(x_1) + (G(t, \mathbf{x}, u_\varepsilon) - G(t, \mathbf{x}, R_\varepsilon))H(-x_1))] \\ \leq |\omega_\varepsilon| + \varepsilon \Delta |u_\varepsilon - R_\varepsilon|, \\ \omega_\varepsilon := \operatorname{div}_x [(F(t, \mathbf{x}, R_\varepsilon) - F(t_0, \mathbf{x}_0, R_\varepsilon))H(x_1)] + \operatorname{div}_x [(G(t, \mathbf{x}, R_\varepsilon) - G(t_0, \mathbf{x}_0, R_\varepsilon))H(-x_1)]. \end{aligned}$$

If we denote $\gamma_\varepsilon = |\omega_\varepsilon|$, we obtain (46); it remains to estimate γ_ε .

We get the bound (47) by computing ω_ε explicitly. First, we estimate the jump term coming from the differentiation of $H(\pm x_1)$, keeping in mind that R_ε is continuous across Σ due to its H^1_{loc} regularity in space. Because $F(t, \mathbf{x}, k) - F(t_0, \mathbf{x}_0, k)$, $G(t, \mathbf{x}, k) - G(t_0, \mathbf{x}_0, k)$ are Lipschitz continuous functions of (t, \mathbf{x}, k) taking value zero at $(t, \mathbf{x}) = (t_0, \mathbf{x}_0)$, the contribution of the jump term is upper bounded by $C \operatorname{dist}((t, \mathbf{x}), (t_0, \mathbf{x}_0))\delta_0(x_1)$. Next, we focus on the terms coming from the differentiation of $F(t, \mathbf{x}, R_\varepsilon) - F(t_0, \mathbf{x}_0, R_\varepsilon)$ (the contributions of the analogous term with G are estimated in the same way). We get the term $\operatorname{div}_x F(t, \mathbf{x}, r)|_{r=R_\varepsilon(t, \mathbf{x})}$ bounded by the Lipschitz constant of F and the term

$$\nabla R_\varepsilon(t, \mathbf{x}) \cdot (\partial_u F(t, \mathbf{x}, r) - \partial_u F(t_0, \mathbf{x}_0, r))|_{r=R_\varepsilon(t, \mathbf{x})}$$

estimated by $r_\varepsilon = |\nabla R_\varepsilon|$ times the modulus of continuity of $\partial_u F, \partial_u G$ (that is how the distance between (t, \mathbf{x}) and (t_0, \mathbf{x}_0) enters the bound (47)). By assumption, $\partial_u F, \partial_u G$ are Lipschitz continuous and the integral of r_ε is bounded by the constant M . This leads to (47) and concludes the proof of the lemma. \square

Next, we infer at the limit $\varepsilon \rightarrow 0$ some information on traces of $u = \lim_{\varepsilon \rightarrow 0} u_\varepsilon$ that can be extracted from information available for the traces of $R = \lim_{\varepsilon \rightarrow 0} R_\varepsilon$:

Lemma 7. *Let $V(t_0, \mathbf{x}_0)$ be a neighborhood of $(t_0, \mathbf{x}_0) \in \Sigma$. In addition to the assumptions and notations of the previous lemma, assume*

- $u_\varepsilon \rightarrow u$ while $R_\varepsilon \rightarrow R$ a.e. in $V(t_0, \mathbf{x}_0)$;
- u and R admit strong left and right traces u^- and R^- , and u^+ and R^+ , respectively, defined on $\Sigma \cap V(t_0, \mathbf{x}_0)$;
- and (t_0, \mathbf{x}_0) is the Lebesgue point of u_\pm and of R_\pm on Σ .

With the notation f, g introduced in (24), writing u^\pm for $u^\pm(t_0, \mathbf{x}_0)$ and R^\pm for $R^\pm(t_0, \mathbf{x}_0)$, there holds

$$\operatorname{sgn}(u^+ - R^+)(f(u^+) - f(R^+)) - \operatorname{sgn}(u^- - R^-)(g(u^-) - g(R^-)) \leq 0. \tag{48}$$

Proof. We first let $\varepsilon \rightarrow 0$ in (46), using the a.e. convergence and uniform boundedness of $u_\varepsilon, R_\varepsilon$ as well as (47). We find that in $\mathcal{D}'(V(t_0, \mathbf{x}_0))$, there holds

$$\begin{aligned} \partial_t |u - R| + \operatorname{div}_x (\operatorname{sgn}(u - R) ((f(t, \mathbf{x}, u) - f(t, \mathbf{x}, R))H(x_1) + (g(t, \mathbf{x}, u) - g(t, \mathbf{x}, R))H(-x_1))) \\ \leq C(|t - t_0| + |\mathbf{x} - \mathbf{x}_0|)(\delta_0(x_1) + r(t, \mathbf{x})) + C. \end{aligned} \tag{49}$$

Here, r is a measure defined as a weak-* limit, along a subsequence, of the bounded sequence $(r_\varepsilon)_\varepsilon$ of Radon measures defined in Lemma 6. Now, proceeding by approximation with C_c^∞ functions we test (49) with $\eta_h(t, \mathbf{x}) = \varphi(t, \hat{x}_1)\mu_h(x_1)$, where the function μ_h is defined by (20) and φ is regular, such that for h small enough, η_h is a Lipschitz function supported in $V(t_0, \mathbf{x}_0)$. By letting $h \rightarrow 0$, using the existence of one-sided strong traces u_\pm, R_\pm of u, R on Σ we get (with $\mathbf{x} = (0, \hat{x}_1)$)

$$\begin{aligned} \int_\Sigma \varphi(t, \mathbf{x}) \left(\operatorname{sgn}(u^+(t, \mathbf{x}) - R^+(t, \mathbf{x})) (F(t, \mathbf{x}, u^+(t, \mathbf{x})) - F(t, \mathbf{x}, R^+(t, \mathbf{x}))) \right. \\ \left. - \int_\Sigma \varphi(t, \mathbf{x}) \operatorname{sgn}(u^-(t, \mathbf{x}) - R^-(t, \mathbf{x})) (G(t, \mathbf{x}, u^-(t, \mathbf{x})) - G(t, \mathbf{x}, R^-(t, \mathbf{x}))) \right) dt d\hat{x}_1 \\ \leq \int_\Sigma C(|t - t_0| + |\mathbf{x} - \mathbf{x}_0|)(1 + M)\varphi(t, \mathbf{x}) dt d\hat{x}_1. \end{aligned}$$

Indeed, the contribution of the term C to the right-hand side vanishes as $h \rightarrow 0$ and the contribution of the measure $r(t, \mathbf{x})$ is estimated via the bound M on its total variation.

By taking for φ an approximation of the Dirac measure concentrated at (t_0, \mathbf{x}_0) taking into account the fact that (t_0, \mathbf{x}_0) is the Lebesgue point of the functions u_\pm and R_\pm , we get the desired result. \square

To continue, we need two more lemmas. The first one provides a rough information on interface traces u^\pm of u , in the spirit of [31]. The second one uses (48) of Lemma 7 to describe more precisely the couples of possible interface traces, ensuring existence of a suitable value p_u to be used in (17).

Lemma 8. *Assume that u is an L^1_{loc} -limit (along a subsequence) of the family $(u_\varepsilon)_\varepsilon$ of solutions to (44), and it admits strong one-sided traces u^\pm on Σ , as defined above. With the notation of Lemma 7, the Rankine–Hugoniot condition $g(u^-) = f(u^+)$ holds, moreover, for all $k \in [a, b]$ there holds*

$$\operatorname{sgn}(u^+ - k)(f(u^+) - f(k)) - \operatorname{sgn}(u^- - k)(g(u^-) - g(k)) \leq |f(k) - g(k)|. \tag{50}$$

Proof. The passage to the limit in the weak formulation (44) ensures that u is a weak solution of (13), therefore the Rankine–Hugoniot condition (19) of Lemma 3 holds. Moreover, the following (rough) entropy inequality (which is precisely (3) written for the case of Eq. (13), see also (4)) holds for every $k \in [a, b]$:

$$\begin{aligned} & \partial_t |u - k| + \operatorname{div}_{\mathbf{x}}(\operatorname{sgn}(u - k)((F(t, \mathbf{x}, u) - F(t, \mathbf{x}, k))H(x_1) + (G(t, \mathbf{x}, u) - G(t, \mathbf{x}, k))H(-x_1))) \\ & + \operatorname{sgn}(u - k)(H(x_1) \operatorname{div}_{\mathbf{x}} F(t, \mathbf{x}, k) + H(-x_1) \operatorname{div}_{\mathbf{x}} G(t, \mathbf{x}, k)) \leq |F_1(t, \mathbf{x}, k) - G_1(t, \mathbf{x}, k)| \delta_0(x_1), \end{aligned} \tag{51}$$

in the sense of distributions. To prove (51), following [33] and [40] it is enough to multiply (44) by a regularization of $\operatorname{sgn}(u_\varepsilon - k)$ (cf. the proof of Lemma 6) with the rough estimation of the measure term charging Σ ; then to let $\varepsilon \rightarrow 0$ along the subsequence such that u_ε strongly converges toward the function u . Then, taking in (51) test functions $\mu_h(x)\xi(t, \mathbf{x})$ with $\xi \in \mathcal{D}(\Sigma)$ and μ_h given by (20), letting $h \rightarrow 0$ one finds that (50) holds in $\mathcal{D}'(\Sigma)$. The left-hand side of (50) being an $L^\infty(\Sigma)$ function, (50) also holds pointwise, a.e. on Σ . \square

Lemma 9. *Assume that u is an L^1_{loc} -limit (along a subsequence) of the family $(u_\varepsilon)_\varepsilon$ of solutions to (44), and it admits strong one-sided traces u^\pm on Σ , as defined above. Given \mathcal{H}^d -a.e. point $(t_0, \mathbf{x}_0) \in \Sigma$, let f, g and u^\pm be defined as in Lemma 7.*

Then there exists a measurable function $p_u : \Sigma \rightarrow [a, b]$ such that for \mathcal{H}^d -a.e. $(t_0, \mathbf{x}_0) \in \Sigma$, the value $p_u = p_u(t_0, \mathbf{x}_0)$ satisfies

$$\operatorname{sgn}(u^+ - k)(f(u^+) - f(k)) - \operatorname{sgn}(u^- - k)(g(u^-) - g(k)) + \operatorname{sgn}(p_u - k)(f(k) - g(k)) \leq 0, \tag{52}$$

and p_u lies in the (closed) interval with the endpoints u^+ and u^- .

Proof. Fix $(t_0, \mathbf{x}_0) \in \Sigma$, a Lebesgue point of traces u^\pm defined on Σ . We will define pointwise $p_u(t_0, \mathbf{x}_0)$; observe that the construction given below is based on a case study of inequalities for functions $f = F(t_0, \mathbf{x}_0, \cdot)$, $g = G(t_0, \mathbf{x}_0, \cdot)$ that depend continuously on (t_0, \mathbf{x}_0) ; therefore it provides a measurable function p_u defined on Σ . To simplify the analysis, assume that $u^+ \geq u^-$; the other case is symmetric. Remark that it is enough to prove (52) for $k \in [u^-, u^+]$ since for other values of k the result is evident. Indeed, if $k \notin (u^-, u^+)$ then with any choice of $p_u \in [u^-, u^+]$, inequality (52) follows from the Rankine–Hugoniot relation $f(u^+) = g(u^-)$ of Lemma 8.

We have the following cases:

- If the crossing condition (6) is satisfied for the couple of functions (f, g) on the interval $[u^-, u^+]$, then we define $p_u(t_0, \mathbf{x}_0)$ (denoted p_u in the sequel) by

$$p_u := \begin{cases} u^+, & f(k) \geq g(k) \ \forall k \in [u^-, u^+] \\ u^-, & f(k) \leq g(k) \ \forall k \in [u^-, u^+] \\ u^o, & \text{if } u^o \in (u^-, u^+) \text{ is a crossing point of } f \text{ and } g. \end{cases}$$

Indeed, due to the crossing condition (6) the above choice yields

$$\operatorname{sgn}(p_u - k)(f(k) - g(k)) = -|f(k) - g(k)|,$$

which reduces (52) to the already proved property (50).

- Assume next that the couple of functions (f, g) does not satisfy the crossing condition (6), but there exists some intersection point (denoted again u^o) such that

$$\begin{aligned} g(u^-) & \leq g(k), & k & \in [u^-, u^o], \\ f(u^+) & \leq f(k), & k & \in [u^o, u^+]. \end{aligned}$$

Then we can still proceed as above, taking $p_u = u^o$.

- Now, if none of the above possibilities holds, then there exists a crossing point $u^o \in (u^-, u^+)$ of f and g such that there exist $k_- \in (u^-, u^o)$, $k_+ \in (u^o, u^+)$ satisfying

$$g(u^-) > g(k_-) \quad \text{and} \quad f(u^+) > f(k_+). \tag{53}$$

We will prove that this case is impossible, arguing by contradiction. First, notice that it holds:

$$g(u^-) = f(u^+) \leq f(u^o) = g(u^o). \tag{54}$$

This relation is obtained by putting $k = u^o$ in (50) and using the Rankine–Hugoniot relation of Lemma 8. From (54) and the assumptions (53), we see that we can choose values $k^- \in [k_-, u^o)$ and $k^+ \in (u^o, k_+]$ such that

$$g(k^-) = f(k^+) < f(u^+) = g(u^-) \leq g(u^o) = f(u^o); \quad (55)$$

moreover, we will take

$$\begin{aligned} k^- &= \max\{k \in (u^-, u^o) : g(k) \leq \max\{g(k_-), f(k_+)\}\}, \\ k^+ &= \min\{k \in (u^o, u^+) : f(k) \leq \max\{g(k_-), f(k_+)\}\}. \end{aligned}$$

Now, using the procedure from the proof of [11, Proposition 5.1], we argue that there exists a stationary solution to (45) R_ε under the form of one-dimensional profile $R(\frac{\cdot}{\varepsilon})$ satisfying

$$\lim_{z \rightarrow -\infty} R(z) = k^-, \quad \lim_{z \rightarrow +\infty} R(z) = k^+, \quad R(0) = u^o. \quad (56)$$

Indeed, R is given by

$$R(z) = \begin{cases} R^l(z), & z \leq 0 \\ R^r(z), & z \geq 0, \end{cases}$$

where R^l and R^r are solutions to the ODEs

$$(R^l)' = g(k^-) - g(R^l), \quad (R^r)' = f(R^r) - f(k^+)$$

respectively, with the initial data $R^l(0) = u^o = R^r(0)$. Remark that this does ensure $R^l(-\infty) = k^-$ and $R^r(+\infty) = k^+$. Indeed, according to the choice of k^- and k^+ , there holds $g(R^l) - g(k^-) > 0$, $R^l \in (k^-, u^o]$, and $f(k^+) - f(R^r) < 0$, $R^r \in [u^o, k^+)$. This actually means that R^l will decrease from 0 to $-\infty$ until it asymptotically reaches the stationary point k^- of the corresponding ODE; while R^r will decrease from 0 to ∞ tending to k^+ .

This construction provides $R \in C(\mathbb{R}) \cap C^1(\mathbb{R} \setminus \{0\})$ and thus the corresponding rescaled profile R_ε belongs to $L^2_{loc}(\mathbb{R}^+; H^1_{loc}(\mathbb{R}^d))$ and it is readily checked that it represents a solution to (45) in the sense of distributions. Remark that such $(R_\varepsilon)_\varepsilon$ is a family of functions of uniformly bounded variation; in view of the explicit formula (56), R_ε converges to $R(x_1) = k^-$ for $x_1 < 0$, and to $R(x_1) = k^+$ for $x_1 > 0$. Therefore the family of functions $(R_\varepsilon)_\varepsilon$ satisfies conditions of Lemmas 6, 7 with $S_\varepsilon = 0$, the corresponding traces R^\pm of $R = \lim_{\varepsilon \rightarrow 0} R_\varepsilon$ being equal to k^\pm . The family $(u_\varepsilon)_\varepsilon$ satisfies as well the assumptions of these lemmas, therefore we conclude from (48) (recalling the meaning of notation for f, g, u^\pm, R^\pm) that the following relation holds

$$\operatorname{sgn}(u^+ - k^+)(f(u^+) - f(k^+)) - \operatorname{sgn}(u^- - k^-)(g(u^-) - g(k^-)) \leq 0.$$

Since $u^- \leq k^- \leq k^+ \leq u^+$, this reduces to

$$f(u^+) + g(u^-) \leq f(k^+) + g(k^-),$$

which contradicts (55) implying that case (53) is not possible. This concludes the proof. \square

Now, it is easy to prove existence of the entropy admissible solution to (13), (14).

Theorem 10. *Let (11), (12), (16) and (43) be fulfilled. Then, there exists a solution u to (13), (14) in the sense of Definition 1.*

Proof. Existence of weak solutions to (44) in $L^2_{loc}(\mathbb{R}^+; H^1_{loc}(\mathbb{R}^d))$ (as required in Lemma 6) with a given initial datum u_0 can be justified with rather classical arguments. Let us give a very brief sketch of stages of one among many possible construction arguments.

- One can start with $L^2(\mathbb{R}^d)$ data u_0 and construct solutions u_ε in the energy space $L^2(0, T; H^1(\mathbb{R}^d))$ with Galerkin approximations, see, e.g., [36] and references therein, see also [11, Sect. 6.2].

- Further, assumption (12) along with the comparison principle for u_ε (that can be proved as in [36,20,44,15]) ensures the confinement property $a \leq u_\varepsilon \leq b$, which yields the uniform L^∞ bound on $(u_\varepsilon)_\varepsilon$.

- Using, by approximation, the test function $\exp(-|x|)u_\varepsilon$, one finds estimates on u_ε in the space $L^2(0, T; H^1(\mathbb{R}^d, w))$ for all $T > 0$, where $H^1(\mathbb{R}^d, w)$ is the weighted H^1 space with the norm defined by $\|u\|_{1,w}^2 := \int_{\mathbb{R}^d} \exp(-|x|)(|u|^2 + |\nabla u|^2)$; as soon as u_0 remains $[a, b]$ -valued, these estimates do not depend on $\|u_0\|_{L^2(\mathbb{R}^d)}$.

• With a.e. approximation by $[a, b]$ -valued $L^2(\mathbb{R}^d)$ functions, using space and time compactness that follows from the above uniform estimates and the variational formulation of (44) in $(L^2(0, T; H^1(\mathbb{R}^d, w)))^*$ (cf., e.g., [36]), one can extend the solution construction to any $[a, b]$ -valued u_0 and get solutions $u_\varepsilon \in L^2(0, T; H^1(\mathbb{R}^d, w))$. Clearly, these solutions lie in $L^2_{loc}(\mathbb{R}^+; H^1_{loc}(\mathbb{R}^d))$.

Then, (11) guarantees existence of an accumulation point u of $(u_\varepsilon)_\varepsilon$, as $\varepsilon \rightarrow 0$, thanks to strong precompactness theorems ([40], see also [5]).

By construction, the result of the previous lemma can be applied to u . It is not difficult to check that the function p_u defined in the previous lemma and the strong limit u of $(u_\varepsilon)_\varepsilon$ the vanishing viscosity approximations of (44) with fixed initial datum u_0 satisfy conditions of Definition 1.

Indeed, given $k \in [a, b]$, denote by L_k the element of the dual of $C^1_c(\mathbb{R}^+ \times \mathbb{R}^d)$ defined by the left-hand side of (17). For any nonnegative compactly supported function $\varphi \in C^1(\mathbb{R}^+ \times \mathbb{R}^d)$, using μ_h defined by (20) we have:

$$L_k(t, x)(\varphi) = L_k((1 - \mu_h)\varphi) + L_k(\mu_h\varphi). \tag{57}$$

The first summand in the right-hand side of (57) is less than or equal to zero due to (51), since the truncation function $(1 - \mu_h)$ is supported on a subset of $\mathbb{R}^+ \times \mathbb{R}^d$ where the left-hand sides of (51) and (17) coincide. Further, due to existence of traces u^\pm and to the continuity of the maps (23) the limit as $h \rightarrow 0$ of the second summand in the right-hand side of (57) can be written as

$$\int_\Sigma \Delta_u(t_0, \mathbf{x}_0)\varphi(t_0, \mathbf{x}_0) dt_0 d\hat{x}_{10}$$

which is non-positive, since the quantity

$$\begin{aligned} \Delta_u(t_0, \mathbf{x}_0) &:= \operatorname{sgn}(u^+(t_0, \mathbf{x}_0) - k)(f(u^+(t_0, \mathbf{x}_0)) - f(k)) - \operatorname{sgn}(u^-(t_0, \mathbf{x}_0) - k)(g(u^-(t_0, \mathbf{x}_0)) - g(k)) \\ &\quad + \operatorname{sgn}(p_u(t_0, \mathbf{x}_0) - k)(f(k) - g(k)) \end{aligned}$$

is non-positive due to inequality (48) proved in Lemma 9. Thus, as $h \rightarrow 0$ we find that L_k is a non-positive distribution, which proves (17).

It remains to prove continuity of the map $t_0 \mapsto u(t_0, \cdot)$ with values in $L^1_{loc}(\mathbb{R}^d)$. First, as a consequence of entropy inequalities away from Σ and of assumption (9), there exist strong L^1_{loc} one-sided traces $u(t_0+, \cdot)$ and $u(t_0-, \cdot)$ at hypersurfaces $\{(t, \mathbf{x}) \mid t = t_0\}$. Indeed, existence of a right trace $u(t_0+, \cdot)$ for every $t_0 \geq 0$ is justified in [37] in the homogeneous case and in [6] in the (t, x) -dependent case. Existence of the left trace $u(t_0-, \cdot)$ for every $t_0 > 0$ follows from the rescaling method used in [37,38,6] and the isentropicity (which implies reversibility) of the solution obtained as the limit of the rescaled sequence. Second, the weak formulation of the evolution equation (13) satisfied by u readily implies the equality $u(t_0-, \cdot) = u(t_0+, \cdot)$ for all t_0 . Indeed, we simply multiply (13) by $\varphi \in C^1_c(\mathbb{R})$, pass to the weak formulation with respect to $\mathbf{x} \in \mathbb{R}^d$, integrate over $(t_0 - \varepsilon, t_0 + \varepsilon)$ and let $\varepsilon \rightarrow 0$. Due to arbitrariness of φ , the conclusion follows. Moreover, the weak formulation of (13), (14) entails that $u(0+, \cdot) = u_0$, i.e., the initial condition is satisfied. \square

2.5. Remark to the definition of admissibility

Let us underline that a shortcoming of the admissibility concept of Definition 1 is that the singular values given by a function $p_u : \mathbb{R}^+ \rightarrow [a, b]$ are not explicitly determined on the set Σ , see Remark 1(iv). To weaken this indetermination, let us show that the function p_u on Σ can be chosen so that it takes values only from the set of the crossing points of the fluxes f, g defined in (24) (observe that a, b always belong to the set of crossing points, due to assumption (12)). Indeed, we have

Proposition 11. *Assume that the function $u : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow [a, b]$ represents an entropy admissible solution to (13), i.e., (17) holds with some \mathcal{H}^d -measurable function p_u on Σ . Then there exists another \mathcal{H}^d -measurable function $p_u : \Sigma \rightarrow [a, b]$ such that for \mathcal{H}^d -a.e. point $(t_0, \mathbf{x}_0) \in \Sigma$, one has*

$$p_u(t_0, \mathbf{x}_0) \in \mathcal{C}(t_0, \mathbf{x}_0) := \{k \in [a, b] : F_1(t_0, \mathbf{x}_0, k) = G_1(t_0, \mathbf{x}_0, k)\},$$

such that (17) is still satisfied with this new function p_u .

Proof. Notice that it is enough to prove that there exists required function p_u such that (52) holds for the right and left traces u^+ and u^- , respectively, of the function u . Since u^- and u^+ must satisfy (19), we have the following possible situations (\mathcal{H}^d -a.e. point of $(t_0, \mathbf{x}_0) \in \Sigma$ being fixed in the arguments that follow), classified as in the case study (25)–(36):

Case 1: $u^+ \leq u^-$.

- $u^+ \leq u^- \leq p_u$.

From (25) and (26) we see that instead of p_u we can take $u^o \in \mathcal{C}$ such that it is the minimal element from \mathcal{C} satisfying $u^o \geq p_u$.

- $u^+ \leq p_u \leq u^-$.

From (27) and (28) and the Rankine–Hugoniot conditions, we see that there exists $u^o \in \mathcal{C}$ such that $u^+ \leq u^o \leq u^-$. Such u^o can be taken instead of the given p_u .

- $p_u \leq u^+ \leq u^-$.

From (29) and (30) we see that instead of p_u we can take $u^o \in \mathcal{C}$ such that it is the maximal element from \mathcal{C} satisfying $u^o \leq p_u$.

Case 2: $u^- \leq u^+$.

- $u^- \leq u^+ \leq p_u$.

From (31) and (32) we see that instead of p_u we can take $u^o \in \mathcal{C}$ such that it is the minimal element from \mathcal{C} satisfying $u^o \geq p_u$.

- $u^- \leq p_u \leq u^+$.

From (33) and (34) and the Rankine–Hugoniot conditions, we see that there exists $u^o \in \mathcal{C}$ such that $u^+ \leq u^o \leq u^-$. Such u^o can be taken instead of the given p_u .

- $p_u \leq u^- \leq u^+$.

From (35) and (36) we see that instead of p_u we can take $u^o \in \mathcal{C}$ such that it is the maximal element from \mathcal{C} satisfying $u^o \leq p_u$.

This concludes the proof, the value p_u being re-defined in each case by $p_u := u^o \in \mathcal{C}$. Observe that the measurability of the so re-defined p_u follows from the continuity of F_1, G_1 as functions of $(t_0, \mathbf{x}_0) \in \Sigma$ and from measurability of the traces u^\pm and of the original function p_u defined on Σ . \square

3. Piecewise regular discontinuous-flux Cauchy problems

Clearly, the flat case of the previous section is a model case. More realistic problems of the kind (1) can present flux discontinuities along multiple curved, possibly intersecting hypersurfaces: one example is the case of flows in a homogeneous porous medium with inclusions of another homogeneous porous medium. In this section, we justify well-posedness of (1), (14) in the case where the discontinuity interfaces Σ are not necessarily flat, with two different approaches to construction of solutions under *ad hoc* technical restrictions on the singularities of the flux \mathfrak{f} .

3.1. Guidelines for extension of well-posedness results

First, let us explain why the uniqueness proof is generalized in a straightforward way, and what are the difficulties of the existence proof and the ideas to overcome these difficulties.

We start with state-of-the-art observations. Approaches to uniqueness and existence in the case of multiple, non-flat and possibly crossing jump manifolds Σ of the flux \mathfrak{f} have already been developed (in the setting of the Karlsen–Risebro–Towers definition, [31]; and in the setting of \mathcal{G}_{VV} -entropy solutions, [10]). In a part, they can be exploited with our new definition (7), (8).

- Both the definition of admissible solution and the uniqueness proof of the flat case are easily extended to the setting of piecewise Lipschitz regular, genuinely nonlinear fluxes with a locally finite number of jump discontinuities along Lipschitz hypersurfaces (see [10], see also [31]).

For instance, following the approach of Section 2.3, one readily extends the definitions to the case of a curved interface Σ : it is enough to replace the couple $(f, g) = (F_1, G_1)$ in Definition 5 by the couple of normal flux functions with coefficients frozen at (t_0, \mathbf{x}_0) :

$$\begin{aligned}
 f &: u \mapsto F(t_0, \mathbf{x}_0, u) \cdot \nu(t_0, \mathbf{x}_0) \quad (\text{flux in the direction of } \nu(t_0, \mathbf{x}_0)), \\
 g &: u \mapsto G(t_0, \mathbf{x}_0, u) \cdot \nu(t_0, \mathbf{x}_0) \quad (\text{flux in the opposite direction}),
 \end{aligned}$$

where $\nu(t_0, \mathbf{x}_0)$ is some fixed unit normal vector to Σ at the point \mathbf{x}_0 , and traces $u^\pm(t_0, \mathbf{x}_0)$ are taken according to this fixed orientation of $\nu(t_0, \mathbf{x}_0)$. As in the flat case, being solution in the sense of Definition 5 is equivalent to being admissible solution in the sense of (7), (8). Indeed, the arguments that ensure constraints on the couple $(u^-, u^+)(t_0, \mathbf{x}_0)$ starting from (7), (8) reduce to a pointwise discussion.

For the uniqueness proof, first observe that the analogue of the Kato inequality (38) is easily obtained for a locally finite union Σ of Lipschitz interfaces. Then the constraints on $(u^-, u^+)(t_0, \mathbf{x}_0)$ derived from (7) (constraints that amount to the fact that $(u^-, u^+)(t_0, \mathbf{x}_0)$ belongs to the vanishing viscosity germ associated with the point (t_0, \mathbf{x}_0) of Σ) give the desired sign to the quantity $\Delta(u^\pm, v^\pm)(t_0, \mathbf{x}_0)$ in the analogue of inequality (38). This results in the L^1 -contraction principle (with an exponential growth term, if a non-zero, Lipschitz continuous in u source term S is present in Eq. (1)) and to uniqueness of an admissible solution in the sense (7), (8).

- At the same time, in order to justify the existence of solutions in the sense (7), (8), unlike for the Karlsen–Risebro–Towers definition ([31]) we cannot hope for a simple existence proof based on straightforward passage to the limit from approximate solutions: nothing seems to ensure strong compactness of singular values $(p_{u_\varepsilon}(\cdot))_\varepsilon$ on Σ . An indirect existence proof analogous to that of the flat case in Section 2.4 is known ([10], see also Remark 3) under the additional local regularity of jump manifolds: for instance, piecewise convexity or concavity of these manifolds is enough. The construction was proposed in [10] where it was presented under the simplifying assumption that the flux is piecewise constant in (t, \mathbf{x}) and the jump manifold is given by the graph of a t -independent C^2 function: $x_1 = \phi(\hat{x}_1)$. It is straightforward to extend this method to (t, \mathbf{x}) -dependent genuinely nonlinear fluxes with jump manifolds given as portions Σ_i of graphs of the form $x_1 = \phi_i(t, \hat{x}_1)$ (up to relabeling the axes), where functions ϕ_i are Lipschitz continuous in (t, \mathbf{x}) and such that, additionally, $\Delta_{\hat{x}_1} \phi_i$ is a Radon measure on $\mathbb{R}^+ \times \mathbb{R}^{d-1}$. We refer in particular to [10, estimate (28)] for the details of the computations.

- To sum up, uniqueness of solutions in the sense (7), (8) for (1), (12) can be proved for the case of piecewise Lipschitz f with locally finite number of Lipschitz components in the jump manifolds Σ . However, existence in this sense requires unnatural assumptions, if we follow the method of [10]; while direct existence arguments used, e.g., in [31], are not applicable.

In this paper, we want to drop the unnatural restriction on $\Delta_{\hat{x}_1} \phi_i$ in the existence proof based on the approach of [10] and Section 2.4. The method mainly consists in comparison of the solution with limits of appropriate viscosity profiles. Yet we do not attempt to construct approximate viscosity profiles (cf. Remark 3) satisfying the original equation up to a measure source term not charging the interface (such construction is successful for the flat case or under the above mentioned restrictions on $\Delta_{\hat{x}_1} \phi_i$, but it is very delicate in the general case). Instead, we rely upon the invariance of the notion of solution – in the sense (7), (8) – under Lipschitz changes of coordinates in Eq. (1). This invariance permits us to exploit the possibility to rectify Σ , at least locally, with a change of variables. Actually we propose two methods for constructing solutions: the one in Section 3.2 (given with the essential details and examples) and the one of Section 3.4 (this method is more general but the details of construction are quite tedious, therefore they are only sketched).

As a matter of fact, in both cases the construction of solutions via a change of variables amounts to a specific “adapted” viscosity approximation of the original equation (1). Such approximation is considerably more involved than the isotropic homogeneous viscosity $\varepsilon \Delta u_\varepsilon$ used in most of the previous works on the subject (see, e.g., in [31, 40, 25, 10]). The use of an involved viscosity regularization operator need not be seen as a drawback of the method: indeed, we put forward the following remark.

Remark 4. As observed in Remark 1(ii), an important feature of the vanishing viscosity approximation is to ensure the global continuity of the approximate solutions (this contrasts with the properties of the vanishing capillarity regularizations, see, e.g., [29, 8]). Regularizing the strongly heterogeneous first-order model (1) with introduction of the isotropic homogeneous diffusion $\varepsilon \Delta u$ may look un-realistic. Heuristically, along with this basic vanishing heterogeneous diffusion operator one should consider, for the existence proof, the possibility to approximate solutions via more involved heterogeneous diffusion operators that still ensure global continuity of u_ε .




Fig. 1. Change of variables transforming square into circle.

3.2. Existence for almost rectifiable sets of singularities

The proof developed here relies on three ingredients. First, we have already known how to construct solutions (where the main difficulty is to construct p_u) for the case of a flat portion of the interface; moreover, the arguments put forward in Lemmas 6–9 are local arguments that work in a neighborhood of \mathcal{H}^d -a.e. interface point, provided the corresponding portion of the interface is flat. Second, we point out the invariance of the notion of admissible solution with respect to a family of (possibly singular) changes of variables. Third, it is possible to deal with interface-rectifying changes of variables involving lower-dimensional singularities.

The combination of the two latter ideas leads to the definition of *almost rectifiability* for jump manifolds, illustrated by several examples that show that this notion is in fact rather general.

Definition 12. A union of hypersurfaces Σ in $\mathbb{R}^+ \times \mathbb{R}^d$ is called almost rectifiable if there exists a direction $\vec{e}_1 \in \mathbb{R}^d$ (that, up to a rotation of coordinate axes, can be assumed to be the direction of the first canonical basis vector, i.e., $\partial_{\vec{e}_1} = \partial_{x_1}$) and a closed set $\gamma \subset \Sigma$ consisting of a locally finite number of disjoint Lipschitz manifolds of codimension in \mathbb{R}^{d+1} greater than one, satisfying the following $(\Sigma \setminus \gamma)$ -rectification property:

There exists a continuous surjective map

$$\Phi : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}^+ \times \mathbb{R}^d, \quad \Phi : (\tilde{t}, \tilde{x}_1, \tilde{\mathbf{x}}_1) \mapsto (t, x_1, \hat{x}_1)$$

that only acts in the direction \vec{e}_1 , i.e., it has the form

$$t = \tilde{t}, \quad \hat{x}_1 = \tilde{x}_1, \quad x_1 = \phi(\tilde{t}, \tilde{\mathbf{x}}) \tag{58}$$

with some scalar function ϕ on $\mathbb{R}^+ \times \mathbb{R}^d$, and such that Φ fulfills the following properties:

- (a) There exists a closed subset Γ of $\mathbb{R}^+ \times \mathbb{R}^d$ such that:
 1. One has $\Phi(\Gamma) = \gamma$, moreover, the restriction of Φ to $\tilde{\Omega} := (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Gamma$ is injective;
 2. the map Φ is Locally Lipschitz continuous on $\tilde{\Omega}$ and the inverse of Φ , defined on $\Omega := (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma$, is locally Lipschitz continuous as well;
 3. in addition, the second-order derivatives $\partial_{\tilde{t}\tilde{x}_1}^2 \phi$ and $\partial_{\tilde{x}_j\tilde{x}_1}^2 \phi$, $2 \leq j \leq d$, are locally bounded on $\tilde{\Omega}$;
- (b) There exists a set $\tilde{\Sigma} \subset \tilde{\Omega}$ consisting of flat manifolds orthogonal to the direction \tilde{x}_1 such that $\Phi(\tilde{\Sigma}) = \Sigma \setminus \gamma$.

Heuristically, the almost-rectifiability property of Σ means that, up to a union of lower-dimensional sub-manifolds γ of the union Σ of hypersurfaces of \mathbb{R}^{d+1} , a change of variables rectifies Σ . To do this, one may need Φ^{-1} to be multivalued. For instance, a circle can be transformed into a square by rectifying the two open half-circles, but then the two poles of the circle have to be stretched into segments (see Fig. 1).

Remark 5. To construct such transformation in practice, having chosen coordinates in such a way that \vec{e}_1 is transversal to Σ in a.e. point with respect to the d -dimensional Hausdorff measure on Σ , the map $\phi_{t, \tilde{x}_1}(\cdot) := \phi(t, \cdot, \tilde{x}_1)$ from \mathbb{R} to \mathbb{R} can taken to be a non-strictly increasing, piecewise polynomial function; moreover, whenever Σ is a union of Lipschitz hypersurfaces, one can ask for a Lipschitz dependence of the function $\phi_{t, \tilde{x}_1}(\cdot)$ on the parameters t, \tilde{x}_1 (such construction leads to the desired bounds on the mixed second derivatives $\partial_{\tilde{t}\tilde{x}_1}^2 \phi$ and $\partial_{\tilde{x}_j\tilde{x}_1}^2 \phi$).

The simplest explicit example is the case where $\Sigma := \{(t, x_1, \hat{x}_1) \mid x_1 = \phi_0(t, \hat{x}_1)\}$ in which case it is enough to set $x_1 = \tilde{x}_1 + \phi_0(t, \tilde{x}_1)$. The properties (a)(1)–(3) and (b) are readily checked in this case. Further examples are presented in Section 3.3.

In the sequel, whenever convenient we will denote $(\tilde{x}_1, \tilde{\hat{x}}_1)$ by \mathbf{y} and we will not distinguish \tilde{t} and t since they are equal; to summarize the definition, we can write

$$\begin{aligned} (t, \mathbf{x}) &= \Phi(\tilde{t}, \tilde{x}_1, \tilde{\hat{x}}_1) = \Phi(t, \mathbf{y}), \\ \Phi &\text{ is a locally bi-Lipschitz bijection from } \tilde{\Omega} = (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Gamma \text{ to } \Omega = (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma \\ &\text{ such that } \Phi^{-1} \text{ rectifies } \Sigma \setminus \gamma, \end{aligned}$$

with specific mixed second derivatives of Φ that are locally bounded on $\tilde{\Omega}$.

In the sequel, we assume that the flux \mathbf{f} in (1) has its jump manifolds contained in an almost rectifiable set Σ and it is Lipschitz continuous in all variables on each connected component of the complementary of Σ . Notice that one can drop the assumption $\gamma \subset \Sigma$ whenever this is convenient for construction of the change of variables Φ : indeed, one can always extend Σ by including “ghost” interfaces across which the flux $(t, \mathbf{x}) \mapsto \mathbf{f}(t, \mathbf{x}, k)$ has jump zero, for all k .

Using the change of variables Φ described in Definition 12, we transform Eq. (1) set up in $\mathbb{R}^+ \times \mathbb{R}^d$ into an analogous equation set up in the domain $\tilde{\Omega} = \Phi^{-1}(\Omega)$, where Ω is the set $(\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma$ which complementary γ is of codimension 2 or more. Indeed, observe that considering the first-order equation (1) (in a weak or in an entropy sense) in the whole space $\mathbb{R}^+ \times \mathbb{R}^d$ is equivalent to considering it (in the same sense) restricted to Ω : cf. the proof of Proposition 14 below. Then, by an explicit calculation we see that under the change of variables $(t, \mathbf{x}) = \Phi(t, \mathbf{y})$ Eq. (1) becomes the equation on $\tilde{\Omega}$ under the analogous conservative form

$$\partial_t \tilde{u} + \operatorname{div}_{\mathbf{y}} \tilde{\mathbf{f}}(t, \mathbf{y}, \tilde{u}) = \tilde{S}(t, \mathbf{y}, \tilde{u}). \tag{59}$$

Here

$$\tilde{\mathbf{f}}(t, \mathbf{y}, k) = \mathcal{D}(t, \mathbf{y}) f(\Phi(t, \mathbf{y}), k) + k \partial_t \phi(t, \mathbf{y}) \vec{e}_1$$

($\mathcal{D}(t, \cdot)$ being the $d \times d$ Jacobian matrix of the change of variables \mathbf{x} into \mathbf{y} given by $\mathbf{x} = \Phi(t, \mathbf{y})$) and \tilde{S} is computed using \mathbf{f} , S and second-order derivatives of ϕ that appear in assumption (a)(3) of Definition 12. For the Ansatz (58) and under the assumption (a) of Definition 12, the scalar function $\tilde{S}(t, \mathbf{y}, k)$ defined on $\tilde{\Omega} \times [a, b]$ is measurable with respect to (t, \mathbf{y}) and locally Lipschitz continuous in k . Further, $\tilde{\mathbf{f}} = \tilde{\mathbf{f}}(t, \mathbf{y}, k)$ is a piecewise continuous vector-function defined on $\tilde{\Omega} \times [a, b]$ which is discontinuous with respect to (t, \mathbf{y}) along the union of flat manifolds $\tilde{\Sigma}$ contained in $\tilde{\Omega}$ (see (b) of Definition 12), and which is Lipschitz with respect to (t, \mathbf{y}, k) , locally in $(t, \mathbf{y}) \in \tilde{\Omega}$ and globally in $k \in [a, b]$. Similarly, if one imposes the (t, \mathbf{x}) -Lipschitz property (43) of the derivatives $\partial_u \mathbf{f}$ in each component of $(\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Sigma$, this property persists for the flux $\tilde{\mathbf{f}}$ in variables (t, \mathbf{y}) in each component of $\tilde{\Omega} \setminus \tilde{\Sigma}$, locally in $\tilde{\Omega}$. This means in particular that the Lipschitz constant of $\partial_u \tilde{\mathbf{f}}$ may blow up as (t, \mathbf{y}) approaches Γ ; this, however, will not harm the analysis as seen from the proof of Proposition 14.

Furthermore, the genuine nonlinearity property (11) is inherited by the flux $\tilde{\mathbf{f}}$ on $\tilde{\Omega} \times [a, b]$ provided it is satisfied by \mathbf{f} on $(\mathbb{R}^+ \times \mathbb{R}^d) \times [a, b]$. Finally, property (12) for \mathbf{f} , S ensures that for Eq. (59),

$$\text{the constant } a \text{ (resp., } b) \text{ is a weak sub- (resp., super-) solution.} \tag{60}$$

Because Eq. (59) has to be considered only in $\tilde{\Omega} = (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Gamma$ (recall that $\Omega \cap \{(t, \mathbf{x}) \mid t > 0\}$ is an open subset of $\mathbb{R}^+ \times \mathbb{R}^d$ which can have holes or cracks, see Fig. 1 and examples below), i.e., with test functions that belong to $\mathcal{D}(\tilde{\Omega})$, we propose the following definition of which Definition 1 is a particular case corresponding to $\Gamma = \emptyset$ and $\Sigma = \{(t, \mathbf{x}) \mid x_1 = 0\}$.

Definition 13. We say that a function \tilde{u} taking values in $[a, b]$ is an admissible solution to (59) with initial datum \tilde{u}_0 if \tilde{u} it satisfies the analogue of (7) within $\tilde{\Omega}$:

$$\forall k \in \mathbb{R} \quad |\tilde{u} - k|_t + \operatorname{div}_{\mathbf{y}} (\operatorname{sgn}(\tilde{u} - k) (\tilde{\mathbf{f}}(t, \mathbf{y}, \tilde{u}) - \tilde{\mathbf{f}}(t, \mathbf{y}, k))) \leq -\operatorname{sgn}(p_{\tilde{u}} - k) \operatorname{div}_{\mathbf{y}} \tilde{\mathbf{f}}(t, \mathbf{y}, k) \quad \text{in } \mathcal{D}'(\tilde{\Omega}) \tag{61}$$

with $\tilde{u}(0, \cdot) = \tilde{u}_0$, where

$$p_u \text{ is some Borel function that coincides with } \tilde{u} \text{ a.e. on } \tilde{\Omega}. \tag{62}$$

As in Definition 1, the initial datum entering the entropy inequalities can be included in $C(\mathbb{R}^+; L^1_{loc}(\mathbb{R}^d))$ sense. The following easy observation follows:

Proposition 14. *Assume that the flux \tilde{f} in (1) is piecewise Lipschitz, discontinuous in (t, x) across a set Σ almost rectifiable in the sense of Definition 12.*

Given a measurable function u_0 on \mathbb{R}^d taking values in $[a, b]$, consider initial condition \tilde{u}_0 defined on $\tilde{\Omega}_0 = \partial\tilde{\Omega} \cap \{(t, \mathbf{y}) \mid t = 0\}$ by $\tilde{u}_0(\mathbf{y}) := u_0(\mathbf{x})$ where \mathbf{y} is such that $\mathbf{x} = \Phi(0, \mathbf{y})$. Let \tilde{f}, \tilde{S} be defined so that (1) transforms into (59) under the change of variables Φ .

A function \tilde{u} on $\tilde{\Omega}$ is an admissible solution of (59) with initial datum \tilde{u}_0 in the sense of Definition 13 if and only if the function u on $\mathbb{R}^+ \times \mathbb{R}^d$ such that $u(t, \mathbf{x}) = \tilde{u}(t, \mathbf{y})$ with $\mathbf{x} = \Phi(t, \mathbf{y})$ is an admissible solution of (1) in the sense (7), (8) with the initial datum u_0 .

Proof. Since $u, \tilde{f}(\cdot, u(\cdot))$ and $S(\cdot, u(\cdot))$ belong to $L^\infty(\mathbb{R}^+ \times \mathbb{R}^d)$ and $\tilde{u}, \tilde{f}(\cdot, \tilde{u}(\cdot))$ and $\tilde{S}(\cdot, \tilde{u}(\cdot))$ belong at least to $L^\infty_{loc}(\tilde{\Omega})$, the spaces $\mathcal{D}(\Omega)$ and $\mathcal{D}(\tilde{\Omega})$ for test functions can be replaced by the spaces $W^{1,1}_{loc}$ of locally Lipschitz, compactly supported in Ω and in $\tilde{\Omega}$, respectively, test functions. The latter spaces are transported the one onto the other under the change of variables Φ that maps $\tilde{\Omega}$ onto Ω .

Therefore, first, the claim of the proposition holds true when the inequalities (7), (8) for u are restricted to $\mathcal{D}'(\Omega)$ test functions. Indeed, the change of variables transforms one inequality into the other, with the transformed test function. In particular, under the change Φ the values p_u on $\Sigma \setminus \gamma$ featuring in inequalities (7), (8) become the values of $p_{\tilde{u}}$ on $\tilde{\Sigma}$ in inequalities (61), (62) and vice versa.

Second, it is easily seen that the entropy formulations (7), (8) on $\Omega = (\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma$ and on $\mathbb{R}^+ \times \mathbb{R}^d$ are in fact equivalent. Indeed, sets in \mathbb{R}^{d+1} of codimension at least 2 have zero \mathcal{H}^d Hausdorff measure and zero $W^{1,1}$ capacity. In particular, the measure in the right-hand side of (7) does not charge γ ; the space $W^{1,1}_c((\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma)$ of Lipschitz continuous compactly supported functions, extended by 0 on γ , is dense in $W^{1,1}_c(\mathbb{R}^+ \times \mathbb{R}^d)$; and passage to the limit in (7) with a sequence of $W^{1,1}_c((\mathbb{R}^+ \times \mathbb{R}^d) \setminus \gamma)$ test functions converging in $W^{1,1}(\mathbb{R}^+ \times \mathbb{R}^d)$ leads to the same inequality with the limit test function. Therefore, considering $W^{1,1}_c(\Omega)$ test functions and considering $\mathcal{D}(\mathbb{R}^+ \times \mathbb{R}^d)$ test functions in the context of integral inequalities (7) is equivalent.

Combining the first and the second observation, we justify our claim. \square

Due to Proposition 14, existence of admissible solutions for (1) with almost rectifiable jump manifolds Σ is reduced to existence of solutions, in the sense of Definition 13, to analogous conservation law (59) set up in $\tilde{\Omega}$. The crucial advantage of the latter is that it admits only flat jump manifolds. Because coefficients of (59) may become singular in a neighborhood of $\partial\tilde{\Omega}$, existence of admissible solutions to (59) is ensured by a combination of vanishing viscosity with a suitable truncation of $\tilde{\Omega}$. Indeed, one can approximate $\tilde{\Omega}$ from inside by a sequence of open domains $\tilde{\omega}_h$; their convergence to $\tilde{\Omega}$ is understood in the sense that

$$\bigcup_{\varepsilon > 0} \tilde{\omega}_\varepsilon = \tilde{\Omega}, \quad \bigcup_{\varepsilon > 0} \partial\tilde{\omega}_\varepsilon \supset \tilde{\Omega}_0, \quad \text{and} \quad (\tilde{\omega}_\varepsilon)_\varepsilon \text{ are embedded in the sense that } \tilde{\omega}_{\varepsilon_0} \subset \tilde{\omega}_\varepsilon \text{ for } 0 < \varepsilon < \varepsilon_0. \tag{63}$$

Then \tilde{f} is globally Lipschitz continuous on $\tilde{\omega}_\varepsilon \times [a, b]$, $(t, x, k) \mapsto \tilde{S}(t, x, k)$ is uniformly Lipschitz continuous in k on $\tilde{\omega}_\varepsilon \times [a, b]$. We can choose $\tilde{\omega}_\varepsilon$ with the property that the boundary of $\partial\tilde{\omega}_\varepsilon$ is piecewise parallel, piecewise orthogonal to the time axis. This can be achieved by covering compact subsets of $\tilde{\Omega} \cup \tilde{\Omega}_0$ by a finite number of cylinders contained in $\tilde{\Omega} \cup \tilde{\Omega}_0$. We will denote by $\partial^{par}\tilde{\omega}_\varepsilon$ the parabolic boundary of $\tilde{\omega}_\varepsilon$, i.e., the part of $\partial\tilde{\omega}_\varepsilon$ where the exterior unit normal vector is either orthogonal to the time direction (the lateral boundary), or it points in the direction of increasing time (this is the union of lower boundaries of $\tilde{\omega}_\varepsilon$, that contains $\tilde{\Omega}_0 \cap \partial\tilde{\omega}_\varepsilon$; but in general, it can also contain pieces of hyperplanes $\{(t, \mathbf{y}) \mid t = const\}$). Then, in the place of the initial-value viscosity regularized problem for the equation

$$\partial_t \tilde{u} + \operatorname{div}_{\mathbf{y}} \tilde{\mathbf{f}}(t, \mathbf{y}, \tilde{u}) = \tilde{S}(t, \mathbf{y}, \tilde{u}) + \varepsilon \Delta_{\mathbf{y}} \tilde{u}, \tag{64}$$

we consider the more general problem with the condition $u = a$ prescribed on the part $\partial^{par} \tilde{\omega}_\varepsilon \setminus \tilde{\Omega}_0$ of the parabolic boundary:

$$\tilde{u}|_{\partial^{par} \tilde{\omega}_\varepsilon \setminus \tilde{\Omega}_0} = a \tag{65}$$

and initial condition on the lower boundary:

$$\tilde{u}(0, \cdot)|_{\partial^{par} \tilde{\omega}_\varepsilon \cap \tilde{\Omega}_0} = \tilde{u}_0 \in [a, b]. \tag{66}$$

While this is not a classical initial–boundary value problem because $\tilde{\omega}_\varepsilon$ is not cylindrical, solutions can be constructed in the same way as for the classical setting of cylindrical domain (cf. the proof of [Theorem 10](#)). To account for the specificity of the domain geometry, one can combine a Galerkin semi-discretization scheme with a restarting procedure that accounts for a finite number of changes of geometry of the sections $\tilde{\omega}_\varepsilon^t := \{\mathbf{y} \in \mathbb{R}^d \mid (t, \mathbf{y}) \in \tilde{\omega}_\varepsilon\}$. We denote by \tilde{u}_ε the so constructed solution of (64), (65), (66).

The so constructed solutions obey the comparison property, which is also proved in the same way as for the classical initial–boundary value problems, cf. the proof of [Lemma 6](#). Recall that due to the choice of $[a, b]$ -valued boundary and initial conditions, property (60) is still valid for the generalized initial–boundary value problem (59), (65), (66) set in $\tilde{\omega}_\varepsilon$. This guarantees that \tilde{u}_ε is $[a, b]$ -valued. Due to the genuine nonlinearity property (11) of $\tilde{\mathbf{f}}$ and the embedding property of $(\tilde{\omega}_\varepsilon)_\varepsilon$, for all fixed $\varepsilon_0 > 0$ $(\tilde{u}_\varepsilon)_\varepsilon$ has an accumulation point u defined in $\tilde{\omega}_{\varepsilon_0}$. Then, the diagonal extraction procedure and the convergence of $\tilde{\omega}_\varepsilon$ to $\tilde{\Omega}$ in the sense explained above permit us to define u globally on $\tilde{\Omega}$.

Therefore, arguments developed in the proof of [Theorem 10](#) and the related lemmas prove that $\tilde{u} = \lim_{\varepsilon \rightarrow 0} \tilde{u}_\varepsilon$ verifies (61) with test functions in $\mathcal{D}'(\tilde{\omega}_{\varepsilon_0})$ and some values $p_{\tilde{u}}$ as in (62). It is easily seen that the initial data pass to the limit, i.e., $\tilde{u}(0, \cdot) = \tilde{u}_0$. Due to convergence of $\tilde{\omega}_{\varepsilon_0}$ to $\tilde{\Omega}$ as $\varepsilon \rightarrow 0$, this ensures that u is a solution to (59) in the sense of [Definition 13](#) with initial datum \tilde{u}_0 .

Observe that the above approach gives not only an existence proof but also a uniqueness proof, due to the equivalence stated in [Proposition 14](#) and to an elementary extension of the calculations of Section 2.2 to the case of several flat interfaces (to be precise, for the uniqueness of solutions in the sense of [Definition 13](#), a uniform Lipschitz condition of dependence on u of $\tilde{\mathbf{f}}, \tilde{S}$ should be imposed; it is verified in the examples we provide below). Yet the direct uniqueness proof for solutions of (1) can be conducted without making appeal to a global change of variables, see Section 3.1 and [31,10]. Therefore, we can summarize our result as follows.

Theorem 15. *Consider Eq. (1) with initial datum u_0 . We impose:*

- the confinement assumption (12) on the flux \mathbf{f} , source S and initial datum u_0 ;
- the genuine nonlinearity assumption (11) of the flux \mathbf{f} ;
- the almost-rectifiability assumption ([Definition 12](#)) on the set Σ of jump discontinuities in (t, \mathbf{x}) of the flux \mathbf{f} ;
- on every connected component of $(\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Sigma$, the assumptions of Lipschitz continuity of the flux \mathbf{f} with respect to all variables (t, \mathbf{x}, k) and of $\partial_u \mathbf{f}$ with respect to (t, \mathbf{x}) ;
- the assumption of global Lipschitz continuity of $k \mapsto S(t, \mathbf{x}, k)$ for the source term.

Then there exists a unique weak solution to Eq. (1) that is admissible in the sense of (7), (8) and takes the initial value u_0 .

Now, we will provide two examples explaining how to check the almost-rectifiability assumptions introduced above. Along with the example of [Fig. 1](#), this gives an idea of how large is the area of applicability of [Theorem 15](#).

3.3. Two examples of almost rectifiable jump manifolds

Now, we will explicitly construct the almost-rectifying maps Φ for two basic examples that permit to overview the ideas helpful in construction of almost-rectifying maps of [Definition 12](#).

First, we deal with a family of curved non-crossing surfaces (for the case $d = 2$, with jump surfaces parallel to the time axis) assuming that they have a common, globally defined transversal direction. The crossing is permitted




Fig. 2. Case of several non-intersecting hypersurfaces in Σ .

in the second example, where we focus on the case of two merging curves (for the case $d = 1$, with t -dependent jump curves). Clearly, the techniques for these examples can be combined. For example, the case of jump across the hyper-cylinder $\Sigma = \{(t, \mathbf{x}) \mid t \geq 0, |\mathbf{x}| = 1\}$ (which is a prototype of the situation not covered by the first example) can be treated by considering Σ as the union of two hypersurfaces that merge along the lines $\{(t, 0, \hat{x}_1) \mid t \geq 0, |\hat{x}_1| = 1\}$, in the spirit of the second example.

3.3.1. The case of several non-intersecting simple discontinuity manifolds

For simplicity, we will work in two-dimensional space, where we select the preferred direction x_1 ; we have $\hat{x}_1 = x_2$. Assume that the flux discontinuity corresponding to the t -independent flux \mathbf{f} from (1) consists of several curves (that become hypersurfaces, if the time variable is also considered) prescribed via Lipschitz continuous functions $\phi_i : \mathbb{R} \rightarrow \mathbb{R}, i = 1, 2, \dots, n$:

$$\Sigma = \bigcup_{i=1}^n \Sigma_i \equiv \mathbb{R}^+ \times \left(\bigcup_{i=1}^n \sigma_i \right), \quad \sigma_i = \{x \in \mathbb{R} : x_1 = \phi_i(x_2)\},$$

with a strict separation (non-intersection) condition $\phi_i - \phi_{i-1} \geq \delta > 0$ for $i = 1, \dots, n - 1$.

The equation that we are considering is thus (with the notation χ_Ω for the characteristic function of the set Ω)

$$\partial_t u + \sum_{i=1}^{n+1} \operatorname{div}_{\mathbf{x}}(F_i(t, \mathbf{x}, u)\chi_{\omega_i}(\mathbf{x})) = 0, \tag{67}$$

$$u|_{t=0} = u_0, \quad a \leq u_0 \leq b, \tag{68}$$

where $F_i(t, \mathbf{x}, u) = (F_{1,i}(t, \mathbf{x}, u), F_{2,i}(t, \mathbf{x}, u))$ are Lipschitz continuous and ω_i is the open region between the curves σ_{i-1} and $\sigma_i, i = 1, \dots, n + 1$, with the convention $\sigma_0 = \{(x_1, x_2) \mid x_1 = -\infty\}$ and $\sigma_{n+1} = \{(x_1, x_2) \mid x_1 = +\infty\}$. We can also add a source term here, but we will omit it because it plays a passive role in the construction of the almost-rectifying map for Σ .

Now, the transformation Φ from Definition 12 is constructed as follows (see Fig. 2). In the domain $\omega_1 = \{(x_1, x_2) \mid x_1 \leq \phi_1(x_2)\}$ we set

$$y_1 = x_1 - \phi_1(x_2), \quad y_2 = x_2.$$

In the domains $\omega_i = \{(x_1, x_2) \mid \phi_{i-1}(x_2) \leq x_1 \leq \phi_i(x_2)\}, i \in \{2, \dots, n - 1\}$, we set

$$y_1 = \frac{x_1 - \phi_i(x_2)}{\phi_{i-1}(x_2) - \phi_i(x_2)} + i - 2, \quad y_2 = x_2.$$

In the domain $\omega_{n+1} = \{(x_1, x_2) \mid \phi_n(x_2) \leq x_1\}$, we take

$$y_1 = x_1 - \phi_n(x_2) + (n - 1), \quad y_2 = x_2.$$

Since the curves are non-intersecting, the above defined transform is bi-Lipschitz globally. Its inverse, given by is the formula




Fig. 3. Case of two merging curves in Σ .

$$\begin{aligned}
 x_1 &= (\phi_1(y_2) + y_1)H(-y_1) \\
 &+ \sum_{i=1}^{n-1} (\phi_{i+1}(y_2) - (y_1 - i)(\phi_i(y_2) - \phi_{i+1}(y_2)))H(y_1 - (i - 1))H(i - y_1) \\
 &+ (\phi_n(y_2) + y_1 - (n - 1))H(y_1 - (n - 1))
 \end{aligned}$$

with H denoting the Heaviside function, satisfies the properties required in Definition 12 with $\gamma = \emptyset$, $\Gamma = \emptyset$.

3.3.2. The case of two merging lines

Consider a scalar conservation law whose flux has discontinuities disposed along two curves $\{(t, \phi_{-1}(t)) \mid t \in [0, t_*]\}$ and $\{(t, \phi_1(t)) \mid t \in [0, t_*]\}$ merging into one curve $\{(t, \phi_0(t)) \mid t \in [t_*, +\infty)\}$ at the point (t_*, x_*) (see Fig. 3).

Let $\phi_{-1}, \phi_1, \phi_0$ be three functions of t that coincide at $t = t_*$, with the common value x_* . We assume that for all $t < t_*$, $\phi_{-1}(t) < \phi_1(t)$. The functions $\phi_{-1}, \phi_1, \phi_0$ are assumed to be Lipschitz continuous.

Then we define $\gamma := \{(t_*, x_*)\}$, whose codimension with respect to \mathbb{R}^{d+1} , $d = 1$, is equal to two. We construct the map from $(t, y) \in \mathbb{R}^+ \times \mathbb{R}$ defined as follows. The straight rays or segments composing the set $\Gamma := \{(t_*, y) \mid y \in [-1, 1]\}$ and the set

$$\tilde{\Sigma} := ([0, t_*] \times \{-1\}) \cup ([0, t_*] \times \{+1\}) \cup ((t_*, +\infty) \times \{0\})$$

split $\mathbb{R}^+ \times \mathbb{R}$ into five regions A, B, B', C and C' . The map $\Phi : (t, y) \mapsto (t, x) = (t, x(t, y))$ is defined per region (we consider all the regions as closed, because the values of Φ match on the boundaries between the regions).

A. In the region $t \leq t_*, -1 \leq y \leq 1$,

$$x(t, y) = \phi_{-1}(t) + \frac{y+1}{2}(\phi_1(t) - \phi_{-1}(t)).$$

The image of this region is the space between the curves $\{(t, \phi_{-1}(t)) \mid t \leq t_*\}$ and $\{(t, \phi_1(t)) \mid t \leq t_*\}$.

B. In the region $t \leq t_*, y \leq -1$,

$$x(t, y) = \phi_{-1}(t) + (1 + |t - t_*|)(y + 1).$$

The image is the region $\{(t, x) \mid t \leq t_*, x \leq \phi_{-1}(t)\}$.

B'. In the region $t \geq t_*, y \leq 0$,

$$x(t, y) = \phi_0(t) + (y + 1)H(-(y + 1)) + |t - t_*|y.$$

The image is the region $\{(t, x) \mid t \geq t_*, x \leq \phi_0(t)\}$.

C. In the region $t \leq t_*, y \geq 1$,

$$x(t, y) = \phi_1(t) + (1 + |t - t_*|)(y - 1).$$

The image is the region $\{(t, x) \mid t \leq t_*, x \geq \phi_1(t)\}$.

C' . In the region $t \geq t_*, y \geq 0$,

$$x(t, y) = \phi_0(t) + (y - 1)H(y - 1) + |t - t_*|y.$$

The image is the region $\{(t, x) \mid t \geq t_*, x \geq \phi_0(t)\}$.

This is an example of map constructed according to the recipe of Remark 5. It is Lipschitz continuous (observe that each of the regions B', C' is split into two subregions with different polynomials defining the map) matching on the boundaries between (sub)regions. Both the map Φ and its inverse are Lipschitz away from any neighborhood of γ or Γ , respectively. The second derivatives whose local boundedness is required in Definition 12(a)(3) are easily computed and controlled.

3.4. An alternative method for construction of solutions

Let us briefly indicate another way of constructing solutions.

Definition 16. A union of hypersurfaces Σ in $\mathbb{R}^+ \times \mathbb{R}^d$ is called locally almost rectifiable if up to a set of codimension greater than 1, Σ can be included into the locally finite union of disjoint open sets U_i such that for all i , $\Sigma \cap U_i$ is a portion of the graph of a Lipschitz function $\psi_i : \mathbb{R}^d \mapsto \mathbb{R}$.

This property is easily checked in practice. For example, the cylinder $\Sigma = \{(t, x_1, x_2) \mid t \geq 0, x_1^2 + x_2^2 = 1\}$ can be covered, up to the union of the four lines $\{(t, 0, \pm 1)\} \cup \{(t, \pm 1, 0)\}$ and the curve $\{(0, x_1, x_2) \mid x_1^2 + x_2^2 = 1\}$, by the union of the four regions

$$U_i = \{(t, x_1, x_2) \mid t > 0, \theta_i x_1 > 0, \theta^i x_2 > 0\}, \quad \theta_i, \theta^i \in \{-1, 1\}.$$

In the sequel, we consider fluxes which singularities are included in a locally almost rectifiable set Σ . According to assumption (9), we exclude the possibility that the discontinuity hypersurface is orthogonal to the time direction. Therefore up to an i -dependent rotation of space coordinates in U_i we can assume that

$$\Sigma \cap U_i = \{(t, x_1, \hat{x}_1) \mid x_1 = \psi_i(t, \hat{x}_1)\},$$

with Lipschitz continuous ψ_i . The fact that U_i are disjoint permits to consider a family of nonnegative $C_c^\infty(U_i)$ functions $(\lambda_i^\varepsilon)_\varepsilon$ with supports of $\lambda_i^\varepsilon, \lambda_j^\varepsilon$ that are disjoint, for all $i \neq j$, and such that for all i , λ_i^ε converges to 1 a.e. on U_i as $\varepsilon \rightarrow 0$; moreover, we can assume that $\lambda_i^\varepsilon \equiv 1$ inside U_i except in the ε -neighborhood of ∂U_i .

Now, we sketch the following construction that can be seen as another “adapted viscosity” approximation of (1).

3.4.1. An adapted diffusion operator

For every U_i , one can make the change of coordinates $x_1 = \tilde{x}_1 + \psi_i(t, \hat{x}_1)$ that rectifies $\Sigma \cap U_i$; by analogy with the construction of Definition 12, let us write $(t, \mathbf{x}) = \Phi_i(t, \mathbf{y})$, with $U_i = \Phi_i(\tilde{U}_i)$, but keeping in mind that Φ_i has a much simpler structure than in Section 3.2 (it is bi-Lipschitz, of jacobian 1, and the mixed derivatives featuring in (a)(2) of Definition 12 are equal to zero). In the sequel, whenever u is considered at $(t, \mathbf{x}) \in U_i$, we mean that $(t, \mathbf{x}) = \Phi_i(t, \mathbf{y})$ with $(t, \mathbf{y}) \in \tilde{U}_i$, and we write \tilde{u} for the transformed function $\tilde{u}(t, \mathbf{y}) = u(t, \mathbf{x})$.

Denote by $\tilde{\lambda}_i^\varepsilon$ the function in the transformed variables such that $\tilde{\lambda}_i^\varepsilon(t, \mathbf{y}) = \lambda_i^\varepsilon(t, \mathbf{x})$. In the new variables \mathbf{y} in \tilde{U}_i , consider the following degenerate elliptic operator in conservative form:

$$\tilde{A}_i^\varepsilon : \tilde{u} \mapsto \varepsilon \sum_{k=1}^d \partial_{y_k} (\tilde{\lambda}_i^\varepsilon \partial_{y_k} \tilde{u});$$

observe that, by the definition of $\lambda_i^\varepsilon, \tilde{A}_i^\varepsilon$ acts as ε times the Laplacian in new variables \mathbf{y} in a large portion of \tilde{U}_i . Let A_i^ε be the corresponding operator acting on functions $u = \tilde{u} \circ \Phi_i^{-1}$ defined on U_i , this operator is implicitly defined by the change of variables Φ_i . For a global definition of A_i^ε , observe that values of $(A_i^\varepsilon u)(\cdot)$ outside U_i can be set to zero, because λ_i^ε are supported in U_i . Therefore A_i^ε is a degenerate anisotropic heterogeneous diffusion operator (in general, in non-divergence form) with coefficients that are Lipschitz continuous.

Due to the fact that the supports of $(\lambda_i^\varepsilon)_i$ are disjoint, the operators A_i^ε , implicitly defined on U_i in the original coordinates, can be just pieced together to yield a second-order degenerate diffusion operator $A^\varepsilon = \sum_i A_i^\varepsilon$. By construction, near Σ this operator represents the homogeneous isotropic Laplacian diffusion in coordinates adapted to the geometry of Σ , except in an ε -neighborhood \mathcal{N}^ε of $\Sigma \cap (\bigcup_i \partial U_i)$ in which the diffusion can degenerate.

Then we claim that the original equation (1) regularized with the following sum of diffusion operators:

$$u \mapsto \delta \Delta u + A^\varepsilon u$$

permits to construct a solution of (1) in the sense of (7), by letting first $\delta \downarrow 0$ with $\varepsilon > 0$ fixed, then letting $\varepsilon \downarrow 0$.

Indeed, to start with, given $\delta, \varepsilon > 0$, existence of $L^2(\mathbb{R}^+; H_{loc}^1(\mathbb{R}^d))$ solutions $u^{\delta, \varepsilon}$ with prescribed initial datum u_0 is ensured by classical theories of uniformly parabolic equations in non-divergence form. It remains to pass to the limit and to characterize $\lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} u^{\delta, \varepsilon}$ by inequalities (7), with singular values (8).

3.4.2. Auxiliary problem with degenerate adapted diffusion

In the first step, the compactness argument based on (11), as used in Section 2.4, permits to obtain an accumulation point u^ε from $(u^{\delta, \varepsilon})_\delta$, as $\delta \rightarrow 0$. Away from an ε -neighborhood \mathcal{N}^ε of $\Sigma \cap (\bigcup_i \partial U_i)$, the function u^ε is an entropy solution to Eq. (1) regularized with degenerate diffusion operator A^ε :

$$\partial_t u + \operatorname{div}_{\mathbf{x}} \mathbf{f}(t, \mathbf{x}, u) = A^\varepsilon u + S(t, \mathbf{x}, u). \tag{69}$$

Observe that in this limit problem, non-degenerate diffusion adapted to the geometry of Σ persists in a vicinity of $\Sigma \setminus \mathcal{N}^\varepsilon$.

To be precise, the limit u^ε verifies in $(\mathbb{R}^+ \times \mathbb{R}^d) \setminus \mathcal{N}^\varepsilon$ the entropy formulation of (69) analogous to the one put forward in the work [15]; in particular, u^ε is a local Kruzhkov entropy solution of (1) in the regions where the flux \mathbf{f} is Lipschitz continuous and A^ε degenerates. The uniqueness theory of [15] only covers the case of degenerate anisotropic diffusion but at a price of lengthy technicalities, it can be generalized in a straightforward way to the heterogeneous case (cf. [44] for analogous extension of the isotropic homogeneous theory of [20] to the isotropic heterogeneous case). Therefore, the (possibly local) formulation of [15] for Eq. (69) leads to the (local) Kato inequality for every couple of (local) entropy solutions $u^\varepsilon, \hat{u}^\varepsilon$ of (69).

3.4.3. Convergence of adapted diffusion approximations

Starting from this point, our analysis mimics the one of Section 2.4, using a family of explicit local viscosity profiles for (69); these are easily obtained in transformed coordinates (t, \mathbf{y}) , considering separately each of the sets U_i .

Namely, in the second step we apply the analogous (11)-based compactness argument to $(u^\varepsilon)_\varepsilon$. An accumulation point u , as $\varepsilon \rightarrow 0$, is a local Kruzhkov entropy solution of (1) away from Σ . Indeed, the contribution of A^ε to the entropy dissipation is nonnegative, and its contribution to the entropy flux vanishes, as $\varepsilon \rightarrow 0$, because there holds a uniform estimate on $\sum_i \|\sqrt{\varepsilon} \lambda_i^\varepsilon \nabla_{\mathbf{y}} \tilde{u}^\varepsilon\|_{L^2}$. Further, since U_i cover Σ up to a lower-dimensional set, in order to see that u is a solution of (1) in the sense (7), (8) it is enough to justify the entropy inequalities (7) with test functions supported in U_i , for every i , and some p_u satisfying (8). The simplicity of the change of coordinates defined on U_i implies the invariance principle analogous to the one shown in Proposition 14: u verifies (7), (8) with test functions supported in U_i if and only if \tilde{u} verifies the analogous inequalities with test functions supported in \tilde{U}_i and appropriate singular values $p_{\tilde{u}}$. In variables (t, \mathbf{y}) of \tilde{U}_i , the interface $\tilde{\Sigma}_i := \phi_i^{-1}(\Sigma \cap U_i)$ is flat, orthogonal to the direction \tilde{x}_1 ; and the diffusion that appears in the equation on \tilde{u}^ε is the isotropic Laplacian $\varepsilon \Delta_{\mathbf{y}}$ except in the set $\Phi^{-1}(U_i \cap \mathcal{N}^\varepsilon)$ that vanishes, as $\varepsilon \rightarrow 0$. Therefore in a neighborhood of every point of $\tilde{\Sigma}_i$ one can construct profiles \tilde{R}^ε that only depend on \tilde{x}_1/ε , as in the construction of Section 2.4. The analogue of Lemma 6 follows from the Kato inequality involving \tilde{u}^ε and \tilde{R}^ε . Then or all i , the arguments of Section 2.4 permit to characterize the one-sided traces of \tilde{u} on $\tilde{\Sigma}_i$ and to define singular values $p_{\tilde{u}}$, \mathcal{H}^d -a.e. on $\tilde{\Sigma}_i$. This provides entropy inequalities (7), (8) with support of the test functions restricted to U_i , where the corresponding singular values p_u are obtained from $p_{\tilde{u}}$ with the help of the transformation ϕ_i . Combining those with classical Kruzhkov entropy inequalities in $(\mathbb{R}^+ \times \mathbb{R}^d) \setminus \Sigma$, we justify (7), (8) globally.

This concludes the sketch of construction of a solution u in the sense (7), (8) under the assumption of local rectifiability of Σ up to a lower-dimensional subset (Definition 16). In this way, the result fully analogous to the one of Theorem 15 can be obtained for a different (at least as large as the class of almost rectifiable Σ 's, see Definition 12) class of jump singularities Σ in the flux \mathbf{f} .

4. Conclusion

We have presented a new definition of solution to some discontinuous-flux problems, along with detailed uniqueness and existence proofs for the case of a flat interface (a flux discontinuity hypersurface being seen as an interface). These results can be put in close correspondence with the results of [10], where a simple particular case has been considered and quite different (rather artificial) approach to solutions characterization has been pursued. We think that, although our proofs in this paper do not appear any simpler than the proofs of [10], the new definition of solution may be particularly useful in engineering applications because it complies with the physical and numerical intuition. Further, we demonstrate that our well-posedness arguments extend to a general configuration in spatially inhomogeneous media with a locally finite number of Lipschitz-regular interfaces, by providing two original constructions of adapted viscosity approximate solutions.

Conflict of interest statement

There is no conflict of interest.

Acknowledgements

Boris Andreianov is supported by the French ANR project CoToCoLa (ANR-11-JS01-0006). During the preparation of the article, Darko Mitrović was engaged as a part time researcher at the University of Bergen funded by the Research Council of Norway whose support we gratefully acknowledge. His work is also supported in part by the Croatian Science Foundation's funding of the project "Weak convergence methods and applications" number 9780, and by the project "Advection–diffusion equations in highly heterogeneous media" of the Montenegrin Ministry of Science (number 01-471).

The authors are grateful to the Berlin Mathematical School and to Prof. Etienne Emmrich from the Technical University of Berlin for the invitation for a research stay during which the paper has been written.

References

- [1] Adimurthi, R. Dutta, S.S. Ghoshal, G.D. Veerappa Gowda, Existence and nonexistence of TV bounds for scalar conservation laws with discontinuous flux, *Commun. Pure Appl. Math.* 64 (1) (2011) 84–115.
- [2] Adimurthi, G.D. Veerappa Gowda, Conservation laws with discontinuous flux, *J. Math. Kyoto Univ.* 43 (2003) 27–70.
- [3] Adimurthi, S. Mishra, G.D. Veerappa Gowda, Optimal entropy solutions for conservation laws with discontinuous flux functions, *J. Hyperbolic Differ. Equ.* 2 (2005) 783–837.
- [4] Adimurthi, S. Mishra, G.D. Veerappa Gowda, Existence and stability of entropy solutions for a conservation law with discontinuous non-convex fluxes, *Netw. Heterog. Media* 2 (2007) 127–157.
- [5] J. Aleksić, D. Mitrović, On the compactness for two dimensional scalar conservation law with discontinuous flux, *Commun. Math. Sci.* 4 (2009) 963–971.
- [6] J. Aleksić, D. Mitrović, Strong traces for averaged solutions of heterogeneous ultra-parabolic transport equations, *J. Hyperbolic Differ. Equ.* 10 (4) (2013) 659–676.
- [7] L. Ambrosio, N. Fusco, D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Univ. Press, New York, 2000.
- [8] B. Andreianov, C. Cancès, Vanishing capillarity solutions of Buckley–Leverett equation with gravity in two-rocks medium, *Comput. Geosci.* 17 (3) (2013) 551–572.
- [9] B. Andreianov, C. Cancès, On interface transmission conditions for conservation laws with discontinuous flux of general shape, preprint, HAL hal.archives-ouvertes.fr/hal-00940756, 2014.
- [10] B. Andreianov, K.H. Karlsen, N.H. Risebro, On vanishing viscosity approximation of conservation laws with discontinuous flux, *Netw. Heterog. Media* 5 (3) (2010) 617–633.
- [11] B. Andreianov, K.H. Karlsen, N.H. Risebro, A theory of L^1 -dissipative solvers for scalar conservation laws with discontinuous flux, *Arch. Ration. Mech. Anal.* 201 (2011) 27–86.
- [12] E. Audusse, B. Perthame, Uniqueness for scalar conservation law via adapted entropies, *Proc. R. Soc. Edinb. A* 135 (2005) 253–265.
- [13] F. Bachmann, J. Vovelle, Existence and uniqueness of entropy solution of scalar conservation laws with a flux function involving discontinuous coefficients, *Commun. Partial Differ. Equ.* 31 (2006) 371–395.
- [14] P. Baiti, H.K. Jenssen, Well-posedness for a class of 2×2 conservation laws with L^∞ data, *J. Differ. Equ.* 140 (1) (1997) 161–185.
- [15] M. Bendahmane, K.H. Karlsen, Renormalized entropy solutions for quasi-linear anisotropic degenerate parabolic equations, *SIAM J. Math. Anal.* 36 (2) (2004) 405–422.
- [16] M. Bulicek, P. Gwiazda, A. Świerczewska-Gwiazda, Multi-dimensional scalar conservation laws with fluxes discontinuous in the unknown and the spatial variable, *Math. Models Methods Appl. Sci.* 3 (2013) 407–439.

- [17] R. Burger, K.H. Karlsen, J. Towers, A model of continuous sedimentation of flocculated suspensions in clarifier-thickener units, *SIAM J. Appl. Math.* 65 (3) (2005) 882–940.
- [18] R. Burger, K.H. Karlsen, J. Towers, On Enquist–Osher-type scheme for conservation laws with discontinuous flux adapted to flux connections, *SIAM J. Numer. Anal.* 3 (2009) 1684–1712.
- [19] R. Burger, K.H. Karlsen, J. Towers, A conservation law with discontinuous flux modelling traffic flow with abruptly changing road surface conditions, in: *Hyperbolic Problems: Theory, Numerics and Applications, Part 2*, in: Proc. Symp. Appl. Math., vol. 67, Amer. Math. Soc., Providence, 2009, pp. 455–464.
- [20] J. Carrillo, Entropy solutions for nonlinear degenerate problems, *Arch. Ration. Mech. Anal.* 147 (4) (1999) 269–361.
- [21] G. Chavent, G. Cohen, J. Jaffré, A finite-element simulator for incompressible two-phase flow, *Transp. Porous Media* 2 (1987) 465–478.
- [22] G. Crasta, V. De Cicco, G. De Philippis, Kinetic formulation and uniqueness for scalar conservation laws with discontinuous flux, preprint, arXiv:1404.5837, 2014.
- [23] S. Diehl, On scalar conservation law with point source and discontinuous flux function modelling continuous sedimentation, *SIAM J. Math. Anal.* 6 (1995) 1425–1451.
- [24] S. Diehl, A conservation law with point source and discontinuous flux function modelling continuous sedimentation, *SIAM J. Appl. Math.* 56 (2) (1996) 388–419.
- [25] S. Diehl, A uniqueness condition for non-linear convection–diffusion equations with discontinuous coefficients, *J. Hyperbolic Differ. Equ.* 6 (2009) 127–159.
- [26] T. Gimse, N.H. Risebro, Riemann problems with discontinuous flux function, in: Proc. 3rd Int. Conf. Hyperbolic Problems Studentlitteratur, Uppsala, 1991, pp. 488–502.
- [27] J. Jimenez, Mathematical analysis of a scalar multidimensional conservation law with discontinuous flux, *J. Evol. Equ.* 11 (3) (2011) 553–576.
- [28] J. Jimenez, L. Lévi, Entropy formulations for a class of scalar conservations laws with space-discontinuous flux functions in a bounded domain, *J. Eng. Math.* 60 (3–4) (2008) 319–335.
- [29] E. Kaasschieter, Solving the Buckley–Leverett equation with gravity in a heterogeneous porous media, *Comput. Geosci.* 3 (1999) 23–48.
- [30] K.H. Karlsen, S. Mishra, N.H. Risebro, Convergence of finite volume schemes for triangular systems of conservation laws, *Numer. Math.* 111 (4) (2009) 559–589.
- [31] K.H. Karlsen, N.H. Risebro, J. Towers, L^1 -stability for entropy solutions of nonlinear degenerate parabolic connection–diffusion equations with discontinuous coefficients, *Skr. - K. Nor. Vidensk. Selsk.* 3 (2003) 1–49.
- [32] K. Karlsen, J. Towers, Convergence of the Lax–Friedrichs scheme and stability for conservation laws with a discontinuous space–time dependent flux, *Chin. Ann. Math., Ser. B* 3 (2004) 287–318.
- [33] S.N. Kruzhkov, First order quasilinear equations in several independent variables, *Mat. Sb.* 81 (1970) 217–243.
- [34] D. Mitrovic, New entropy conditions for scalar conservation laws with discontinuous flux, *Discrete Contin. Dyn. Syst.* 30 (2011) 1191–1210.
- [35] D. Mitrovic, Proper entropy conditions for scalar conservation laws with discontinuous flux, technical report, 2012.
- [36] F. Otto, L^1 -contraction and uniqueness for quasilinear elliptic–parabolic equations, *J. Differ. Equ.* 131 (1) (1996) 20–38.
- [37] E.Yu. Panov, Existence of strong traces for generalized solutions of multidimensional scalar conservation laws, *J. Hyperbolic Differ. Equ.* 2 (2005) 885–908.
- [38] E.Yu. Panov, Existence of strong traces for quasi-solutions of multidimensional conservation laws, *J. Hyperbolic Differ. Equ.* 4 (2007) 729–770.
- [39] E.Yu. Panov, On existence and uniqueness of entropy solutions to the Cauchy problem for a conservation law with discontinuous flux, *J. Hyperbolic Differ. Equ.* 3 (2009) 525–548.
- [40] E.Yu. Panov, Existence and strong pre-compactness properties for entropy solutions of a first-order quasilinear equation with discontinuous flux, *Arch. Ration. Mech. Anal.* 195 (2010) 643–673.
- [41] N. Seguin, J. Vovelle, Analysis and approximation of a scalar conservation law with a flux function with discontinuous coefficients, *Math. Models Methods Appl. Sci.* 13 (2) (2003) 221–257.
- [42] J.D. Towers, Convergence of a difference scheme for conservation laws with a discontinuous flux, *SIAM J. Numer. Anal.* 38 (2) (2000) 681–698.
- [43] J.D. Towers, A difference scheme for conservation laws with a discontinuous flux: the nonconvex case, *SIAM J. Numer. Anal.* 39 (4) (2001) 1197–1218.
- [44] G. Vallet, Dirichlet problem for a degenerated hyperbolic–parabolic equation, *Adv. Math. Sci. Appl.* 15 (2) (2005) 423–450.