ANNALES SCIENTIFIQUES DE L'É.N.S.

SPENCER BLOCH

On the Chow groups of certain rational surfaces

Annales scientifiques de l'É.N.S. 4^e série, tome 14, n° 1 (1981), p. 41-59

http://www.numdam.org/item?id=ASENS_1981_4_14_1_41_0

© Gauthier-Villars (Éditions scientifiques et médicales Elsevier), 1981, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'É.N.S. » (<http://www.elsevier.com/locate/ansens>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE CHOW GROUPS OF CERTAIN RATIONAL SURFACES

BY SPENCER BLOCH ⁽¹⁾

Let X be a smooth projective surface defined over a field k . X is said to be *rational* if X_k is birational with $\mathbb{P}_k^2$ for some k'/k . I will say X is *split* over k' if X_k can be obtained by blowing up a finite number of k' -points starting from $\mathbb{P}_k^2$ and then blowing down a finite number of exceptional curves of the first kind defined and absolutely irreducible over k' . Any rational surface over k admits a splitting field k' finite over k . The following notation will be used throughout: X =rational surface/ k , k' =splitting field for X , and:

$X' = X_{k'}$; N =Néron-Severi group of X' .

F =quotient field of X ; $F' = F k'$.

$CH_0(X)$ =Chow group of zero cycles on X modulo rational equivalence.

$A_0(X) = \text{Ker}(CH_0(X) \xrightarrow{\deg} \mathbb{Z}) = \text{Ker}(CH_0(X) \rightarrow CH_0(X'))$.

X^i =points of codim i on X ; $K_2(\cdot) = K_2$ group Milnor [11].

$\text{Br}(\cdot)$ =Brauer group; $k_*(\cdot) = \text{Mod } 2$ Milnor ring [10] [e. g. $k_1(k) = k^*/k^{*2}$].

Finally, to avoid technical problems associated with characteristic p , we will work only with rational surfaces X which are *separably split*, i. e. such that the splitting field k' can be taken to be separable over k .

The objective of this paper is to study a certain map Φ associating to a cycle of degree 0 in the Chow group $A_0(X)$ of a rational surface X a *torseur* for the Néron-Severi torus:

$$\Phi : A_0(X) \rightarrow H^1(k'/k, N \otimes k'^*). \quad (2)$$

We show Image Φ is finite when k is local or global, and that Image $\Phi=0$ if k is non-archimedean local and X has good reduction over the closed fibre. When X is a *conic bundle surface*, i. e., when there exists $\pi : X \rightarrow \mathbb{P}_k^1$ a rational map with generic fibre a conic curve we

⁽¹⁾ Partially supported by National Science Foundation under NSF MCS77-01931.

⁽²⁾ Colliot-Thélène has pointed out to me that the existence of such a map Φ is immediate from [5] and [15], where results like (0.2) and an even stronger version of the finiteness of $X(k)/B$ are also announced. The reader of the present paper should perhaps focus on the role of K-theory in evaluating $\text{Ker } \Phi$ and $\text{coker } \Phi$ as in (0.1.1), (0.3) and (0.4).

also show $\text{Ker } \Phi$ is finite when k is local or global and indeed $\text{Ker } \Phi = 0$ if $k \notin \mathbb{R}$. Interesting examples of such surfaces are the *Châtelet surfaces* $X : T_0^2 - a T_1^2 = x(x - a_1)(x - a_2)$. It is possible to give examples of such surfaces with $A_0(X) \neq (0)$ [4]. In an appendix, we verify a compatibility between Φ and the notion of *Brauer equivalence* defined in [9]. Using this and the above results, we show that the set $X(k)/B$ of k -points of X modulo Brauer equivalence is finite for k global and X any rational surface.

Here, in more precise form are the main results.

THEOREM (0.1). — *Let X be a smooth projective surface over a field k . Assume X is rational, split by a galois extension k'/k . Then there is an exact sequence:*

$$(0.1.1) \quad \Gamma(k'/k, N \otimes k'^*) \rightarrow H^1(k'/k, K_2(F')/K_2(k')) \xrightarrow{\mu} A_0(X) \\ \xrightarrow{\Phi} H^1(k'/k, N \otimes k'^*) \xrightarrow{\theta} H^2(k'/k, K_2(F')/K_2(k')).$$

If the g.c.d. of the degrees of all closed points on X is 1, then there is a map:

$$H^2(k'/k, K_2(F')/K_2(k')) \rightarrow \coprod_{x \in X^1} \text{Br}(k(x)),$$

extending the above exact sequence one place to the right.

THEOREM (0.2). — *With notation as above, if k a local field and k'/k is an arbitrary (not necessarily finite) galois extension, then $H^1(k'/k, N \otimes k'^*)$ is finite. If, moreover, k is non-archimedean and there exists a lifting of X to $\tilde{X}$ smooth and projective over the ring of integers of k , then $\text{Im } \Phi = 0$. Finally, if k is global and k'/k is finite galois, then $\text{Im } \Phi$ is finite.*

THEOREM (0.3). — *Let k be a field of characteristic $\neq 2$, which is either local, global, or C_i for $i \leq 3$, and let $\bar{k}$ be the separable closure of k . Let X be a separably split conic bundle surface over k . Then $H^1(\bar{k}/k, K_2(F\bar{k})/K_2(\bar{k}))$ is subquotient of $k_3(k)$. In particular, for k non-archimedean local, C_1, C_2 , or totally imaginary global this group is zero. For k arbitrary local or global the group is finite.*

THEOREM (0.4). — *Let X be a conic bundle surface defined over a field k of characteristic $\neq 2$:*

- (i) *If k is a C_1 field then $A_0(X) = (0)$.*
- (ii) *If k is a local or global field, then $A_0(X)$ is finite.*
- (iii) *If k is non-archimedean local and X has good reduction in the sense that there exists $\tilde{X}$ as in (0.2), then $A_0(X) = (0)$.*

The conic bundle surfaces given birationally by:

$$X : T_1^2 - a T_2^2 = \prod_{i=1}^3 (x - a_i) \cdot T_0^2 \quad \text{in } \mathbb{P}_k^2 \times \mathbb{A}_k^i, \quad a, a_i \in k^*$$

with homogeneous coordinates T_0, T_1, T_2 and affine coordinate x were studied by Châtelet [3]. Finiteness of $A_0(X)$ for such surfaces with k a number field was proven by Colliot-Thélène and Coray ([4], Remarques 6.7) using results of Châtelet [3] which can be found in Manin [9].

Manin ([8], [9]) constructed a pairing $A_0(X) \times H^1(k'/k, N) \rightarrow \text{Br}(k)$ which he used to show (among other things) non-triviality of $A_0(X)$. From our point of view, this pairing arises *via*:

$$A_0(X) \otimes H^1(k'/k, N) \rightarrow H^1(k'/k, N \otimes k'^*) \otimes H^1(k'/k, N) \xrightarrow[\text{intersection product}]{\quad} H^2(k'/k, k'^*).$$

In considering cycles on rational varieties of dimension > 2 one wants to define the Néron-Severi torus to be the torus with character group N , i.e.:

$$\text{Néron-Severi torus}(k) = \text{Hom}_{\mathbb{Z}}(N, k'^*)^{\text{Gal}(k'/k)}.$$

This definition coincides with the one used in this paper because for X rational of dimension 2, N is self-dual. A very beautiful paper of Colliot-Thélène and Sansuc [5] contains among other results a proof that for X a smooth compactification of a torus T of arbitrary dimension, the group of torseurs for the Néron-Severi torus is isomorphic to the group $T(k)/R$. Here points on T are said to be R -equivalent if they can be connected by a chain of rational curves.

Among the open questions remaining in this area let me single out the question of finiteness of $A_0(X)$ for any smooth rational surface over a local or global field, as well as the question of injectivity of $A_0(X) \rightarrow H^1(k'/k, N \otimes k'^*)$. I know of no counter-example to these assertions over *any* ground field. The relation between the proof of (0.3) and the Eichler norm Theorem suggests one should study the injectivity question when $k = \mathbb{R}$.

I should like to acknowledge many helpful conversations with J.-L. Colliot-Thélène, D. Coray and J.-J. Sansuc ⁽³⁾.

1. The fundamental exact sequence

The following complex is defined for any variety X [12]:

$$(1.1) \quad K_2(F) \xrightarrow{\text{tame}} \coprod_{x \in X^1} k(x)^* \rightarrow \coprod_{x \in X^2} \mathbb{Z}.$$

(Here “tame” denotes the tame symbol [11].) Suppose for simplicity X is a smooth surface over k and let $\pi : Y \rightarrow X$ be obtained by blowing up a k -point on X . Let $E \cong \mathbb{P}_k^1 \subset Y$ be the

⁽³⁾ *Added in proof:* in a recent paper, On the Chow groups of certain rational surfaces: complements to a paper of S. Bloch, Colliot-Thélène and Sansuc have strengthened a number of results found here. In particular they calculate the H^1 group in (3.4) for conic bundle surfaces and give examples where it is $\neq 0$.

exceptional fibre, and let $e \in E$ be the generic point. The top row and middle and right-hand columns of the following diagram are exact:

$$(1.2) \quad \begin{array}{ccccccc} & & & 0 & & 0 & \\ & & & \downarrow & & \downarrow & \\ 0 & \rightarrow & k^* & \rightarrow & k(e)^* & \rightarrow & \prod_{E^1}^0 \mathbb{Z} \rightarrow 0 \\ & & & \downarrow & & \downarrow & \\ & & & \prod_{y \in Y^1} k(y)^* & \rightarrow & \prod_{Y^2} \mathbb{Z} & \\ || & & & \downarrow & & \downarrow & \\ K_2(F) & \rightarrow & \prod_{x \in X^1} k(x)^* & \rightarrow & \prod_{X^2} \mathbb{Z} & \\ & & & \downarrow & & \downarrow & \\ & & & 0 & & 0 & \end{array}$$

[Here $\prod^0 \mathbb{Z} = \text{Ker} (\prod \mathbb{Z} \xrightarrow{\text{sum}} \mathbb{Z})$.]

PROPOSITION (1.3). — (i) The complex (1.1) computes the cohomology of the Zariski sheaf $\mathcal{K}_2$ on X , i.e., it has cohomology groups reading from left to right $\Gamma(X, \mathcal{K}_2)$, $H^1(X, \mathcal{K}_2)$, and $H^2(X, \mathcal{K}_2) \simeq \text{CH}_0(X)$.

(ii) With Y as above, $\Gamma(Y, \mathcal{K}_2) \simeq \Gamma(X, \mathcal{K}_2)$ and $H^1(Y, \mathcal{K}_2) \simeq H^1(X, \mathcal{K}_2) \oplus k^*$.

Proof. — (i) is a standard result in algebraic K-theory [12]. For (ii), using the contravariant functoriality of $\Gamma(\dots, \mathcal{K}_2)$ and the covariant functoriality of (1.1) for proper maps ⁽⁴⁾, one gets:

$$\begin{array}{ccc} \Gamma(Y, \mathcal{K}_2) & & \\ \pi_* \updownarrow \pi^* & \searrow & K_2(F) \\ \Gamma(X, \mathcal{K}_2) & \nearrow & \end{array}$$

where $\Gamma(X, \mathcal{K}_2) \simeq \Gamma(Y, \mathcal{K}_2)$. The isomorphism $H^1(X, \mathcal{K}_2) \oplus k^* \simeq H^1(Y, \mathcal{K}_2)$ now follows from a diagram chase using (1.2).

Q.E.D.

COROLLARY (1.4). — Assume X is a smooth projective rational surface split by k . Then $\Gamma(X, \mathcal{K}_2) \simeq K_2(k)$ and $H^1(X, \mathcal{K}_2) \simeq N \otimes k^*$, where $N = \text{Néron-Severi group of } X$.

Proof. — Follows from (1.3) and the corresponding result for $X = \mathbb{P}_k^2$ [14].

Remark (1.5). — With X as above, we have $N \cong H^1(X, \mathcal{O}_X^*)$ and the isomorphism $H^1(X, \mathcal{O}_X^*) \otimes k^* \rightarrow H^1(X, K_2)$ is given by the symbol map.

⁽⁴⁾ This functionality results from covariant functoriality of the localization sequence. The reader who wishes a detailed discussion in an analogous (cohomological) context is referred to S. BLOCH and A. OGUS, *Gersten's Conjecture and the Homology of Schemes* (Ann. scient. Éc. Norm. Sup., T. 7, fasc. 2, 1974).

THEOREM (1.6). — *Let X be a smooth projective rational surface over a field k , and let k' be galois over k splitting X . Then there is an exact sequence:*

$$\Gamma(k'/k, \underset{\mathbb{Z}}{N \otimes k'^*}) \rightarrow H^1(k'/k, K_2(F')/K_2(k')) \xrightarrow{\mu} A_0(X) \xrightarrow{\Phi} H^1(k'/k, N \otimes k'^*) \xrightarrow{\theta} H^2(k'/k, K_2(F')/K_2(k')).$$

Proof. — One knows that $CH_0(X') \cong \mathbb{Z}$, so there are exact sequences (defining $\mathbb{Z}$):

$$(1.7) \quad \begin{cases} 0 \rightarrow \mathbb{Z} \rightarrow \coprod_{x' \in X^1} k'(x')^* \rightarrow \coprod_{X^2}^0 \mathbb{Z} \rightarrow 0, \\ 0 \rightarrow K_2(F')/K_2(k') \rightarrow \mathbb{Z} \rightarrow N \otimes k'^* \rightarrow 0. \end{cases}$$

One has:

$$H^*(k'/k, \coprod_{X^1} k'(x')^*) \cong \coprod_{x \in X^1} H^*(k'/k_x, k'(x')^*),$$

where $k_x = k' \cap k(x)$, and on the right $x' \in X^1$ is chosen lying above $x \in X$. Also:

$$\Gamma\left(k'/k, \coprod_{X^2}^0 \mathbb{Z}\right) \cong \coprod_{X^2}^0 \mathbb{Z}.$$

It follows from the first sequence in (1.7) that $A_0(X) \cong H^1(k'/k, \mathbb{Z})$, and substituting this value in the cohomology sequence from the second exact sequence yields the Theorem.

Remark (1.7). — If the g. c. d. of the degrees of all closed points of X is 1 [e. g., if $X(k) \neq \emptyset$], then $H^1\left(k'/k, \coprod_{X^2}^0 \mathbb{Z}\right) = 0$ and the above sequence can be extended one more term to the right via a map:

$$H^2(k'/k, K_2(F')/K_2(k')) \rightarrow \coprod_{x \in X^1} \text{Br}(k(x)).$$

2. Finiteness of $\text{Im } \Phi$

THEOREM (2.1). — *Let k be a local field, k'/k a (not necessarily finite) galois extension. Let N be a free f.g. $\mathbb{Z}$ -module with a continuous $\text{Gal}(k'/k)$ -action. Let $N^v = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$. Then $H^1(k'/k, \underset{\mathbb{Z}}{N \otimes k'^*})$ is a finite group, dual to $H^1(k'/k, N^v)$.*

Proof. — This is a standard result in local duality [13].

THEOREM (2.2). — *Let k be a global field, k'/k a finite galois extension. Let X be a smooth projective rational surface defined over k split over k' . Then the kernel of the map:*

$$\theta : H^1(k'/k, N \otimes k'^*) \rightarrow H^2(k'/k, K_2(F')/K_2(k')),$$

defined in (1.6) is finite.

Proof. — The geometric situation, viz. X together with an isomorphism $X_k \cong (\mathbb{P}_k^2$ with a finite number of k' points blown up and down), can be spread out over the ring of S -integers $\mathcal{O}_{k,S}$ for some finite set S of primes of k . Thus for each $p \notin S$ we may assume the reclusion $X(p)$ is a smooth projective rational surface over $k(p)$, and that we are given $X(p) \times_{k(p)} k'(p') \cong [\mathbb{P}_{k'(p')}^2$ with finite number of $k'(p')$ -points blown up and down]. Enlarging S , we may assume it contains all primes which ramify in k' , and that the class group of $\mathcal{O}_{k',S'}$ is trivial, where $S' = \{\text{primes lying over } S\}$.

Tensoring the exact sequence:

$$0 \rightarrow \mathcal{O}_{k',S'}^* \rightarrow k'^* \rightarrow \coprod_{p' \notin S'} \mathbb{Z} \rightarrow 0,$$

with N , we obtain a map $\psi : H^1(k'/k, N \otimes k'^*) \rightarrow \coprod_{p \notin S} H^1(D_p, N)$, where $D_p \subset \text{Gal}(k'/k)$ is a decomposition group at p . Note $\text{Ker } \psi$ is a quotient of $H^1(k'/k, N \otimes \mathcal{O}_{k',S'}^*)$ which is finite since $\mathcal{O}_{k',S'}^*$ is f.g. Thus it will suffice to show $\text{Ker } \theta \subset \text{Ker } \psi$.

We may think now of X' as smooth and projective over $\text{Sp } \mathcal{O}_{k',S'}$, so $p' \notin S'$ defines a discrete valuation on F' , and hence a tame symbol $K_2(F') \rightarrow F'(p')^*/k'(p')^*$, where $F'(p')$ = function field of $X'(p')$. Our objective is to define injective maps:

$$\tau_p : H^1(D_p, N) \hookrightarrow H^2(D_p, F'(p')^*/k'(p')^*)$$

(where p' denotes a fixed prime above p) fitting into a commutative square:

$$(2.2.1) \quad \begin{array}{ccc} H^1(k'/k, N \otimes k'^*) & \xrightarrow{\theta} & H^2(k'/k, K_2(F')/K_2(k')) \\ \psi \downarrow & & \downarrow \text{tame} \\ \coprod_{p \notin S} H^1(D_p, N) & \xrightarrow{\coprod \tau_p} & \coprod_{p \notin S} H^2(D_p, F'(p')^*/k'(p')^*) \end{array}$$

The existence of such a commutative square (2.2.1) will prove $\text{Ker } \theta \subset \text{Ker } \psi$ as claimed.

LEMMA (2.2.2). — Define τ_p to be the boundary map associated to the exact sequence:

$$0 \rightarrow F'(p)^*/k'(p')^* \rightarrow \text{Div}(X'(p')) \rightarrow N \rightarrow 0.$$

Then τ_p is injective and the diagram (2.2.1) commutes.

Proof. — Notice first that

$$N = \text{Néron-Severi}(X') \cong \text{Néron-Severi}(X'(p'))$$

because both surfaces are split over their fields of definition, so the above exact sequence is defined. Moreover, if $C' \subset X'$ is an irreducible curve it can be “closed-up” and normalized to a normal family of curves $\tilde{C}$ flat over $\mathcal{O}_{k',S'}$, so p' defines a divisor $C'(p')$ on $\tilde{C}$, hence a map:

$$k'(C')^* \xrightarrow{\alpha_p} \coprod \mathbb{Z},$$

where the sum runs over irreducible components of $C'(p')$.

Now let C' run through all irreducible curves on X' and identify components of $C'(p')$ with divisors on $X'(p')$ via cycle-theoretic direct image to get a square:

$$\begin{array}{ccc} K_2(F') & \xrightarrow{\text{tame}} & \coprod_{x' \in X'^{-1}} k'(x')^* \\ \text{tame} \downarrow & & \downarrow \alpha \\ F'(p')^* & \longrightarrow & \text{Div}(X'(p')) \end{array}$$

which commutes up to sign ⁽⁵⁾. We can now consider a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_2(F')/K_2(k') & \longrightarrow & Z & \longrightarrow & N \otimes k'^* \longrightarrow 0 \\ & & \text{tame} \downarrow & & \downarrow & & \downarrow V_{p'} & (6) \\ 0 & \longrightarrow & F'(p')^*/k'(p')^* & \longrightarrow & \text{Div}(X'(p')) & \longrightarrow & N \longrightarrow 0 \end{array}$$

where the middle vertical arrow is defined via α in the square above. This suffices to show that (2.2.1) commutes. Finally, note $\text{Div}(X'(p))$ is a direct limit of permutation modules so $H^1(D_p, \text{Div}(X'(p)))=0$ and τ_p is injective.

Q.E.D.

On reflection, it seems that the same argument should yield something in the local case as well:

THEOREM (2.3). — *Let X be a smooth, projective rational surface defined over a non-archimedean local field k with residue field $\mathbb{F}_q$ and ring of integers $\mathcal{O}$. Assume X has good reduction in the sense that there exists an $\tilde{X}$ smooth and projective over $\text{Sp } \mathcal{O}$ with generic fibre X . Then there exists k'/k finite unramified splitting X , and:*

$$\text{Im } \Phi = \text{Ker } (\theta : H^1(k'/k, N \otimes k'^*) \rightarrow H^2(k'/k, K_2(F')/K_2(k'))).$$

is zero.

Proof. — Let $X_0 \rightarrow \text{Sp } \mathbb{F}_q$ be the closed fibre of $\tilde{X}$. Then X_0 is rational [the canonical bundle has negative degree and $q=0$, since both these properties specialize from $X^{(7)}$]. The fact that X has an unramified splitting field follows from:

LEMMA (2.3.1). — *Let R be a complete discrete valuation ring with residue field $\mathbb{F}$ and quotient field k . Let $f: \tilde{X} \rightarrow \text{Sp } R$ be smooth, projective, geometrically irreducible, and*

⁽⁵⁾ The referee is dubious. Quite generally, for a noetherian scheme Y , there are complexes

$$\coprod_{y^0} K_n(k(y)) \rightarrow \coprod_{y^1} K_{n-1}(k(y)) \rightarrow \coprod_{y^2} K_{n-2}(k(y)) \rightarrow \dots \quad [12].$$

Apply this with $n=2$ and Y = "spreading out" of X' over the local ring at p' on $\mathcal{O}_k$.

⁽⁶⁾ If the curve $C' \subset X'$ discussed above has good reduction at p' , and if $a \in k'^*$, then clearly $\alpha_0(a) = V_{p'}(a)$. Since N is generated by such curves, we get $V_{p'}$ here.

⁽⁷⁾ Let ω = canonical bundle of $\tilde{X}/\text{Sp } \mathcal{O}$, $\omega_k = \omega|_X$, $\omega_0 = \omega|_{X_0}$. Then $0 > (\omega_k, L_k) = (\omega_0, L_0)$. The assertion about q follows because q is one half the first Betti number in etale cohomology, and the etale Betti numbers are constant in smooth families.

assume the closed fibre X_0 is rational and split over $\mathbb{F}$. Then the generic fibre X is rational and split over k .

Proof. — An $\mathbb{F}$ -valued point of X_0 lifts by smoothness to an R -valued point of $\tilde{X}$, so a morphism $X_0(1) \rightarrow X_0$ obtained by blowing up an $\mathbb{F}$ -point lifts to a diagram:

$$\begin{array}{ccc} X_0(1) & \subset & \tilde{X}(1) \\ \downarrow & & \downarrow \text{blow up an } R\text{-point.} \\ X_0 & \subset & \tilde{X} \end{array}$$

We may thus build a figure:

$$\begin{array}{ccc} & \tilde{Y} \subset Y_0 & \\ \text{blowings up} \swarrow & & \searrow \text{blowings down} \\ \tilde{X} \subset X_0 & & \mathbb{P}_{\mathbb{F}}^2 \subset \mathbb{P}_{\mathbb{R}}^2 \end{array}$$

Let $E_0 \subset Y_0$ be an exceptional curve of the first kind. There exists a unique family of exceptional curves of the first kind $\tilde{E} \subset \tilde{Y}$ lifting E_0 . Indeed, the normal bundle of $E_0 \cong \mathbb{P}^1$ in Y_0 is $\mathcal{O}(-1)$, so both the zeroth and first cohomology of the normal bundle vanish, implying existence and uniqueness of lifting⁽⁸⁾. We may then blow down $\tilde{E}$ on $\tilde{Y}$ in much the same way as E_0 on Y_0 ⁽⁹⁾. Iterating we get a diagram:

$$\begin{array}{ccc} Y_0 \subset \tilde{Y} & & \\ \downarrow & \downarrow & \\ \mathbb{P}_{\mathbb{F}}^2 \subset \tilde{Z} & & \end{array}$$

with $\tilde{Z}$ smooth and projective over $\text{Spec } R$ with closed fibre $\mathbb{P}_{\mathbb{F}}^2$. Since $\mathbb{P}_{\mathbb{F}}^2$ is rigid also, it follows that $\tilde{Z} \cong \mathbb{P}_{\mathbb{R}}^2$. Passing now to the generic fibres we get $X_k \leftarrow Y_k \rightarrow Z_k \cong \mathbb{P}_k^2$, both morphisms being obtained by successive blowings up of k -points.

Q.E.D.

Returning to the proof of (2.3), let k'/k be unramified with residue field splitting X_0 . The units $\mathcal{O}'^*$ in the ring of integers $\mathcal{O}' \subset k'$ are cohomologically trivial so $N \otimes \mathcal{O}'^*$ is cohomologically trivial ([13], Thm. 9, p. 152) and:

$$H^1(k'/k, N \otimes k'^*) \cong H^1(k'/k, N).$$

⁽⁸⁾ For this sort of argument, cf. A. GROTHENDIECK, *SGA*, fasc. I, exp. III, Paris, 1960-1961.

⁽⁹⁾ Let $\eta \rightarrow \text{Sp } R$ be the generic point. Let A_η be the ring of germs of regular functions on Zariski neighborhoods of $E_\eta \subset Y_\eta$. Since E_η is rational with self-intersection -1 it blows down so A_η is smooth of $\dim .2$ over $k(\eta)$. Let $A \subset A_\eta$ be the R -algebra of functions along $\tilde{E} \subset \tilde{Y}$, $I \subset A$ the ideal of functions vanishing along $\tilde{E}$. Then the germ of $\tilde{E} \in \tilde{Y}$ comes by blowing up $I \subset A$.

Writing $\mathbb{F}'$ and F'_0 for the residue field of k' and the function field of $X_0 \times_{\mathbb{F}} \mathbb{F}'$ respectively we obtain as in (2.2.1):

$$\begin{array}{ccc} H^1(k'/k, N \otimes k'^*) & \xrightarrow{\sim} & H^2(k'/k, K_2(F')/K_2(k')) \\ \downarrow \parallel & & \downarrow \text{tame} \\ 0 \rightarrow H^1(k'/k, N) & \xrightarrow{\tau} & H^2(k'/k, F'_0^*/\mathbb{F}'^*) \end{array}$$

It follows that $\text{Ker } \theta = 0$.

Q.E.D.

3. Applications to conic bundle surfaces

PROPOSITION (3.1). — *Let X be as in paragraph 2 and let $k' \subset k''$ be galois extensions of k splitting X . Then the triangle:*

$$\begin{array}{ccc} H^1(k'/k, K_2(F')/K_2(k')) & \longrightarrow & H^1(k''/k, K_2(F'')/K_2(k'')) \\ \mu' \searrow & & \swarrow \mu'' \\ & A_0(X) & \end{array}$$

commutes. In particular, $\text{Image } \mu'' \supset \text{Image } \mu'$.

The proof is immediate.

We assume henceforth when discussing conic bundle surfaces that $\text{char } k \neq 2$.

THEOREM (3.2). — *Let X be a conic bundle surface, i. e., suppose given a rational map $\pi : X \rightarrow \mathbb{P}_k^1$ with generic fibre a conic curve (Severi-Brauer variety of dimension 1). Assume k is a local field, a global field or a C_i field for $i \leq 3$. Then $H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k}))$ is a subquotient of $k_3(k)$, the third graded piece of the Milnor ring of k . (Here $\bar{k}$ is the separable closure of k , $\bar{F} = F\bar{k}$.)*

COROLLARY (3.3). — *With hypotheses as above, if k is a C_2 -field, a local field $\neq \mathbb{R}$, or a totally imaginary global field, then $H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k})) = 0$. If k is any number field, or $k = \mathbb{R}$, then $H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k})) \cong \bigoplus_s \mathbb{Z}/2\mathbb{Z}$, where $s \leq$ the number of real places of k .*

Proof. — With the exception of the C_2 case, all assertions follow from (3.2) together with results about $k_3(k)$ proved in [10]. For k C_2 (or non-archimedean local) any quadratic form in five variables represents zero. Let

$$l : k^* \rightarrow k_1(k) = k^*/k^{*2}$$

and consider an element

$$l(a) l(b) l(c) \in k_3(k).$$

Let A be the quaternion algebra $T^2 = a, U^2 = b, TU = -UT$. The reduced norm $N : A \rightarrow k$ is a quadratic form in four variables, so it is surjective and there exists $\alpha \in A, N(\alpha) = c$. If $\alpha \in k$, then $c \in k^{*2}$ so $l(a) l(b) l(c) = 0$. If $\alpha \notin k$, then $k(\alpha) \subset A$ is a commutative subfield splitting A . If k [and hence also $k(\alpha)$] is C_2 or local, the map $k_2(k) \rightarrow {}_2\text{Br}(k), l(a) l(b) \rightarrow [A]$

is known to be injective [6], so the symbol $l(a)l(b) \rightarrow 0$ in $k_2(k(\alpha))$. Since the reduced norm on A coincides with the field norm from $k(\alpha)$ to k , we may use the projection formula [2]:

$$l(a)l(b)l(c) = l(a)l(b)N(l(\alpha)) = N_{k_3(k(\alpha))/k_3(k)}(l(a)l(b)l(\alpha)) = 0.$$

Q.E.D.

Remark (3.4). — I don't know whether $H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k}))$ is ever non-zero.

COROLLARY (3.5). — *Let X be a conic bundle surface over a local or global field k . Then $A_0(X)$ is finite.*

Proof. — Combine (3.1), (3.3), the results in paragraph 2 on finiteness of $\text{Im } \Phi$, and (1.6).

COROLLARY (3.6). — *Let X be a conic bundle surface over a non-archimedean local field, and assume X has good reduction over the residue field in the sense of (2.3). Then $A_0(X) = (0)$.*

Proof. — Combine (2.3), (3.1), (3.3), and (1.6).

For the proof of (3.2), fix a conic bundle structure $\pi : X^* \rightarrow \mathbb{P}_k^1$ on X . Let $K = k(\mathbb{P}^1)$, $\bar{K} = \bar{k}(\mathbb{P}^1)$. Note that $X_{\bar{K}} \cong \mathbb{P}_{\bar{K}}^1$ so there exists an exact sequence of $\text{Gal}(\bar{k}/k)$ -modules:

$$(3.7) \quad 0 \rightarrow K_2(\bar{K}) \rightarrow K_2(\bar{F}) \rightarrow \coprod_{\bar{x} \in X_{\bar{K}}} \bar{K}(\bar{x})^* \xrightarrow{N} \bar{K}^* \rightarrow 0.$$

Note also:

$$(3.8) \quad K_2(\bar{K}) \cong K_2(\bar{k}) \oplus \coprod_{\bar{y} \in \mathbb{P}_k^1} \bar{k}(\bar{y})^*.$$

Let $\mathcal{N} \subset K^*$ be the image under the norm map of $\coprod_{\bar{x} \in X_K} K(\bar{x})^*$. It follows from (3.7) and (3.8) that there is an exact sequence:

$$(3.9) \quad 0 \rightarrow H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k})) \rightarrow K^*/\mathcal{N} \xrightarrow{\Psi} \coprod_{y \in \mathbb{P}_k^1} \text{Br}(k(y)).$$

Fix $a, b \in K^*$ such that X_K is isomorphic to the conic curve defined by $T_0^2 - aT_1^2 - bT_2^2 = 0$. We will work with $l(a)l(b) \in k_2(K)$ as well as with the quaternion algebra A defined by $T^2 = a, U^2 = b, TU = -UT$.

LEMMA (3.10). — *Let $N : A^* \rightarrow K^*$ be the reduced norm. Then $NA^* \subset \mathcal{N}$.*

Proof. — If $\alpha \in A, \alpha \notin K$ then $K(\alpha)$ splits A so X_K has a $K(\alpha)$ -rational point. Since the norm from A coincides on $K(\alpha)$ with the field norm we get $N(\alpha) \in \mathcal{N}$. If $\alpha \in K$ then $N(\alpha) = \alpha^2 \in \mathcal{N}$.

Q.E.D.

Returning to the proof of (3.2) we now have:

$$(3.11) \quad 0 \rightarrow H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k})) \rightarrow K^*/\mathcal{N} \xrightarrow{\psi} \coprod_{y \in \mathbb{P}_k^1} \text{Br}(k(y))$$

$$\begin{array}{ccc} & \uparrow & \nearrow \psi' \\ & K^*/\mathcal{N}^* & \end{array}$$

and it will suffice for the proof to show ψ' injective.

LEMMA (3.12). — *Let k be as in (3.2). The diagram below is commutative:*

$$\begin{array}{ccc} K^*/\mathcal{N}^* & \xrightarrow{\psi'} & \coprod \text{Br}(k(y)) \\ \downarrow l(a).l(b) & & \uparrow \text{galois symbol} \\ k_3(K) & \xrightarrow{\text{tame symbol}} & \coprod k_2(k(y)). \end{array}$$

Note that $\mathcal{N}^*.l(a).l(b)=(0)$ by the sort of projection formula argument already used in the proof of (3.3). I will postpone the proof of (3.12) to go directly to the main point.

LEMMA (3.13). — *The maps in the above square have the following properties:*

- (i) “galois symbol” is injective.
- (ii) “tame symbol” has kernel $k_3(k) \subset k_3(K)$.
- (iii) $l(a).l(b)$ is injective.

Proof. — Injectivity of the galois symbol follows from results of [6] together with our assumptions on k . The fact that $\text{Ker}(\text{tame})=k_3(k)$ is proved in [10]. To show injectivity of the left-hand arrow, we consider the map defined in *op. cit.* :

$$k_3(K) \rightarrow I^3/I^4,$$

$$l(a).l(b).l(c) \rightarrow (\langle c \rangle + \langle -1 \rangle)(\langle b \rangle + \langle -1 \rangle)(\langle a \rangle + \langle -1 \rangle),$$

where l is the augmentation ideal in the Witt ring of K . The above quadratic form is a *Pfister form* in the terminology of [7]. In particular, this form lies in I^4 if and only if it is *hyperbolic* (Theorem of Arason-Pfister, *op. cit.*, Cor. 3.4, p. 290). In particular, $l(a).l(b).l(c)=0$ in $k_3(K)$ implies there exists $x_1, \dots, x_8 \in K$ not all zero such that:

$$abcx_1^2 - abx_2^2 - acx_3^2 - bcx_4^2 + ax_5^2 + bx_6^2 + cx_7^2 - x_8^2 = 0.$$

Formally, then we may factor and write:

$$c = \frac{abx_2^2 - ax_5^2 - bx_6^2 + x_8^2}{abx_1^2 - ax_3^2 - bx_4^2 + x_7^2}.$$

Notice that both numerator and denominator are norms from A^* . We may assume $A \not\cong M_2(K)$ else $X_K \cong \mathbb{P}_K^1$ and the whole discussion is silly. Thus the denominator in the above equation vanishes if and only if $x_1=x_3=x_4=x_7=0$. But vanishing of the

denominator implies vanishing of the numerator, and hence of all the x_i , a contradiction. Hence, neither numerator nor denominator vanishes and we have written c as a norm from A^* .

Q.E.D.

The proof of (3.2) is now immediate. Indeed, $H^1(\bar{k}/k, K_2(\bar{F})/K_2(\bar{k}))$ is a quotient of $\text{Ker } \psi'$ by (3.11), and $\text{Ker } \psi' \subset k_3(k)$ by (3.12).

Q.E.D.

It remains to prove (3.12).

LEMMA (3.14). — Let X_K be a conic curve over $\text{Sp } K$ with $\bar{K}/K$ galois splitting X_K . Let $[X_K] \in H^2(\bar{K}/K, \bar{K}^*)$ be the class of X_K as a Severi-Brauer variety, and assume $[X_K] \neq 0$. Let $\bar{F}$ = quotient field of $X_{\bar{K}}$ and consider the exact sequence of $\text{Gal}(\bar{K}/K)$ -modules:

$$0 \rightarrow \bar{K}^* \rightarrow \bar{F}^* \rightarrow \coprod_{\bar{x} \in X_{\bar{K}}} \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow 0.$$

Then $[X_K] = \partial_2 \partial_1(1)$ where ∂_i are the boundary maps associated to this exact sequence.

Proof. — $(\coprod_{\bar{x} \in X_{\bar{K}}} \mathbb{Z})^{\text{Gal}(\bar{K}/K)} = \coprod_{x \in X_K} \mathbb{Z}$. Since any $x \in X_K$ has even degree over $\text{Sp } K$ [this follows from a norm argument using the fact that X_K splits over $K(x)$], we get an exact sequence:

$$0 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow H^2(\bar{K}/K, \bar{K}^*) \rightarrow H^2(\bar{F}/F, \bar{F}^*),$$

$$1 \mapsto \partial_2 \partial_1(1).$$

Since $[X_K] \rightarrow 0$ in $H^2(\bar{F}/F, \bar{F}^*)$, we are done.

Q.E.D.

Tensoring the sequence in (3.14) with K^* and using the symbol map, we obtain a commutative diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & K^* \otimes K^* & \rightarrow & \bar{F}^* \otimes K^* & \rightarrow & \coprod_{\bar{x} \in X_{\bar{K}}} K^* \rightarrow K^* \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow & \downarrow \\ 0 & \rightarrow & K_2(\bar{K}) & \rightarrow & K_2(\bar{F}) & \rightarrow & \coprod \bar{K}(\bar{x})^* \rightarrow \bar{K}^* \rightarrow 0. \end{array}$$

Writing ∂'_i for the boundary maps on cohomology associated to the bottom row, we find a commutative triangle:

$$\begin{array}{ccc} K^* & \xrightarrow{[X_K] \otimes} & H^2(\bar{K}/K, \bar{K}^*) \otimes K^* \\ \partial'_0 \circ \partial'_1 \searrow & & \swarrow \text{symbol} \\ & H^2(\bar{K}/K, K_2(\bar{K})) & \end{array}$$

Now take $K = k(\mathbb{P}^1)$, $\bar{K} = \bar{k}(\mathbb{P}^1)$ as in (3.12) and compose with the tame symbol to get a commutative square:

$$\begin{array}{ccc} K^* & \xrightarrow{\psi} & \prod_{y \in \mathbb{P}^1_k} \text{Br}(k(y)) \\ |x_K| \otimes \downarrow & & \uparrow \text{tame} \\ H^2(\bar{k}/k, \bar{K}^*) \otimes K^* & \longrightarrow & H^2(\bar{k}/k, K_2(\bar{K})) \end{array}$$

The proof of (3.12) now follows from:

LEMMA (3.15). — Let k be a field of characteristic $\neq 2$, $\bar{k}$ = separable closure of k . Let K be an extension field of transcendence degree 1 over k , and write $\bar{K} = K\bar{k}$. Let $a, b \in K^*$ and write $l(a)l(b) \in k_2(K)$, $(a, b) \in {}_2\text{Br}(K)$. Then for torsion prime to the characteristic:

$$\text{Br}(K) \cong H^2(\bar{k}/k, \bar{K}^*).$$

Moreover, if y is a place of K over k with residue field $k(y)$, the diagram:

$$(3.15.1) \quad \begin{array}{ccccc} & & H^2(\bar{k}/k, \bar{K}^*) \otimes K^* & & \\ & \nearrow (a, b) \otimes & & \searrow & \\ K^* & & & & H^2(\bar{k}/k, K_2(\bar{K})) \\ & \searrow l(a):l(b) & & \nearrow T_1 & \\ & k_3(K) & \xrightarrow{\text{Gal} \circ T_2} & \text{Br}(k(y)) & \end{array}$$

(where T_1 and T_2 are tame symbols) commutes.

Proof. — Replacing k by its perfect closure, we may assume $k(y)$ separable over k . Next replacing K by its completion at y we may suppose $k(y) \subset K$ and $\bar{K} = \prod \bar{K}_i$, one copy for each place lying over y . Now replacing the galois group by the decomposition group for one of the $\bar{K}_i$ we may assume $k = k(y)$, $K = k((\pi))$.

We now have split exact sequences:

$$0 \rightarrow \mathcal{O}^* \rightarrow K^* \rightarrow \mathbb{Z} \rightarrow 0, \quad 0 \rightarrow \bar{\mathcal{O}}^* \rightarrow \bar{K}^* \rightarrow \mathbb{Z} \rightarrow 0.$$

$\pi \swarrow \quad \searrow 1$

Note that $1 + \pi\bar{\mathcal{O}}$ is cohomologically trivial, so:

$$H^2(\bar{k}/k, \bar{K}^*) \cong H^2(\bar{k}/k, \bar{k}^*) \oplus H^2(\bar{k}/k, \mathbb{Z}).$$

Case 1. — $a, b \in \mathcal{O}^*$. Let $a_0, b_0 \in k^*$ denote the mod π reductions of a, b . In this case $(a, b) = (a_0, b_0) \in H^2(\bar{k}/k, \bar{k}^*)$. It is easy enough to see that going either way around (3.15.1), $f \in K^*$ gets taken to $\text{ord}(f) \cdot (a_0, b_0) \in \text{Br}(k)$.

Using linearity, it remains only to consider.

Case 2. — $b = \pi, a \in \mathcal{O}^*$. In this case let $G = \text{Gal}(\bar{k}/k)$ and let $\rho: G \times G \rightarrow \mathbb{Z}$ be a 2-cocycle representing the image of a under the composition:

$$(3.15.2) \quad \left\{ \begin{array}{l} \mathcal{O}^* \rightarrow H^1(\bar{K}/K, \mu_2) \xrightarrow{\delta} H^2(\bar{K}/K, \mathbb{Z}), \\ a \mapsto \chi_a. \end{array} \right.$$

Here δ is the coboundary from the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0$. Then (a, b) is represented by the cocycle:

$$G \times G \rightarrow \bar{K}^*, \quad (g_1, g_2) \mapsto \pi^{p(g_1, g_2)}.$$

Indeed, one knows (cf. [13], p. 214) that:

$$(a, \pi) = \pi \cdot \delta \chi_a. \quad (\text{cup product}).$$

With reference to (3.15.1) we must show:

$$T_1 \{ \pi^{p(g_1, g_2)}, f \} = T_2 (l(a) l(\pi) l(f)).$$

If $f = -\pi$ this is clear as both sides are trivial. If, on the other hand, f is a unit with residue class f_0 , we reduce to showing that $l(a_0) l(f_0) \in k_2(k)$ maps to the element in $\text{Br}(k)$ represented by the cocycle $f_0^{p(g_1, g_2)}$. This follows as above with K in (3.15.2) replaced by k and $\bar{K}$ replaced by $\bar{k}$. This completes the proof of (3.15) and (3.12).

Brauer equivalence. Appendix

Let X be a rational surface over k , and let $\coprod_X^0 \mathbb{Z}$ be the group of 0-cycles of degree 0 on X . Following Manin ([8], [9]), we define a pairing:

$$(\quad): \coprod_X^0 \mathbb{Z} \times (\text{Br}(X)/\text{Br}(k)) \rightarrow \text{Br}(k),$$

as follows. Given a cycle:

$$\sum n_i(x_i) \quad \text{and} \quad a \in \text{Br}(X), \quad \left(\sum n_i(x_i), a \right) = \prod_i \text{cor}_{k(x_i)/k}(a(x_i))^{n_i}.$$

Two k -points x_1, x_2 are said to be Brauer equivalent (written $x_1 \sim_B x_2$) if $(x_1 - x_2, a) = 1$ for all $a \in \text{Br}(X)$. Manin shows (*op. cit.*, Thm. 44.2) that $X(k)/B$ is finite for X a smooth cubic surface over a global field. In this appendix, I want to establish a compatibility between the Manin pairing and the map $\Phi: A_0(X) \rightarrow H^1(\bar{k}/k, N \otimes \bar{k}^*)$. It will follow that $X(k)/B$ is finite for *any* smooth rational surface over a global field, and that all points are Brauer equivalent at good reduction places. This explains and generalizes a number of Manin's calculations (*op. cit.*, 45.5, 45.6, 45.11, 45.12).

The key fact is:

PROPOSITION (A.1). — *The following diagram commutes:*

$$\begin{array}{ccc}
 \prod_{p \in X}^0 \mathbb{Z} \times (\text{Br}(X)/\text{Br}(k)) & \xrightarrow{(\cdot, \cdot)} & \text{Br}(k) = H^2(\bar{k}/k, \bar{k}^*) \\
 \downarrow & & \uparrow \text{Intersection pairing} \\
 A_0(X) \times H^1(\bar{k}/k, N) & \xrightarrow{\Phi \times 1} & H^1(\bar{k}/k, N \otimes \bar{k}^*) \times H^1(\bar{k}/k, N)
 \end{array}$$

[Here the map $\text{Br}(X)/\text{Br}(k) \rightarrow H^1(\bar{k}/k, N)$ arises from the spectral sequence $H^p(\bar{k}/k, H^q(\bar{X}, \mathbb{G}_m)) \Rightarrow H^{p+q}(X, \mathbb{G}_m)$.]

COROLLARY (A.2). — *Let k be a local or global field. Then $X(k)/B$ is finite.*

Proof. — Fix $x_0 \in X(k)$ [if $X(k) = \emptyset$, of course, there is not much to prove] and define $\Psi: X(k) \rightarrow H^1(\bar{k}/k, N \otimes \bar{k}^*)$ by $\Psi(x) = \Phi((x) - (x_0))$. Of course $\text{Image } \Psi \subset \text{Image } \Phi$ is finite. On the other hand, if $x, y \in X(k)$ and $x \not\sim_B y$ then $\Psi(x) \neq \Psi(y)$. Hence $\#(X(k)/B) \leq \# \text{Image } \Phi$. Q.E.D.

COROLLARY (A.3). — *If k is non-archimedean local and X has good reduction over the residue field of k , then $\#(X(k)/B) \leq 1$.*

Proof. — As above, using $\#(\text{Image } \Phi) = 1$. Q.E.D.

We turn now to the proof of (A.1). Let $S \subset X$ be a union of divisors, and let $\bar{S} = \bar{S}_k \subset \bar{X}$. Let $\text{Div}_{\bar{S}}(\bar{X}) \subset \text{Div}(\bar{X}) = \prod_{\bar{D} \subset \bar{X}} \mathbb{Z}$ be the group generated by divisors supported on $\bar{S}$, and assume S sufficiently large so $\text{Div}_{\bar{S}}(\bar{X}) \twoheadrightarrow N$. There is then an exact sequence:

$$(A.4) \quad 0 \rightarrow \Gamma(\bar{X} - \bar{S}, \mathcal{O}_{\bar{X}}^*/\bar{k}^*) \rightarrow \text{Div}_{\bar{S}}(\bar{X}) \rightarrow N \rightarrow 0.$$

Given $\bar{D} \subset \bar{X}$ an irreducible divisor with $\bar{D} \not\subset \bar{S}$, let $\pi: \tilde{D} \rightarrow \bar{X}$ be the normalization, and let $\bar{k}(\tilde{D})_S^*$ denote those meromorphic functions on $\tilde{D}$ which are invertible at points of $\pi^{-1}(\bar{S})$. Define $Z_S \subset Z$ and $(\prod_X^0 \mathbb{Z})_S \subset \prod_X^0 \mathbb{Z}$ so the sequence:

$$(A.5) \quad 0 \rightarrow Z_S \rightarrow \prod_{\bar{D} \not\subset \bar{S}} \bar{k}(\tilde{D})_S^* \rightarrow \left(\prod_X^0 \mathbb{Z} \right)_S \rightarrow 0,$$

is exact. We define a pairing:

$$(A.6) \quad \langle \cdot, \cdot \rangle: \prod_{\bar{D} \not\subset \bar{S}} \bar{k}(\tilde{D})_S^* \times \text{Div}_{\bar{S}}(\bar{X}) \rightarrow \bar{k}^*,$$

as follows. Let f be a function on $\tilde{D}$ with divisor (f) supported off $\pi^{-1}(\bar{S})$. Let $\Delta \subset \bar{S}$ be a divisor and write:

$$\pi^{-1} \Delta = \sum n_i (d_i) \quad (\text{as a cycle}).$$

Define:

$$(A.7) \quad \langle f|_{\bar{D}}, \Delta \rangle = \prod f(d_i)^{n_i}.$$

LEMMA (A.8). — *The following diagram is a commutative diagram of pairings:*

$$\begin{array}{ccc} \coprod_{\bar{D} \neq \bar{S}} \bar{k}(\bar{D})_{\bar{S} \times \text{Div}_{\bar{S}}(\bar{X})}^* & \longrightarrow & \bar{k}^* \\ \downarrow & \uparrow \text{div} & \parallel \\ (\coprod_{\bar{X}}^0 \mathbb{Z})_{\bar{S}} \times \Gamma(\bar{X} - \bar{S}, \mathcal{O}_{\bar{X}}^*) / \bar{k}^* & \longrightarrow & \bar{k}^*. \end{array}$$

Here the top pairing is defined in (A.6) and the bottom is evaluation. The left hand vertical arrow is from (A.5).

Proof. — Given $g \in \Gamma(\bar{X} - \bar{S}, \mathcal{O}_{\bar{X}}^*)$ and given

$$\beta = \sum n_j(b_j) \in (\coprod_{\bar{S}}^0 \mathbb{Z})_{\bar{S}},$$

there exists $\sum f_i|_{\bar{D}_i} \mapsto \beta$ with $f_i \in \bar{k}(\bar{D}_i)_{\bar{S}}^*$. We have:

$$\langle g, \beta \rangle = h(0)h(\infty)^{-1}, \quad h = \prod N_{g, \bar{D}_i}(f_i).$$

Note the equality of divisors $(h) = g_*(\beta) = \sum n_j g(b_j)$. Thus $h = C \cdot \prod (t - g(b_j))^{n_j}$, where t is the standard parameter on $\mathbb{P}^1$ and C is some constant. It follows that:

$$h(0)h(\infty)^{-1} = \prod g(b_j)^{n_j}.$$

Q.E.D.

The following abstract result in group cohomology comes from setting $C = C'$, $C'' = 0$ in ([16], Thm. 6, p. 112).

LEMMA (A.9). — *Let G be a group, A, A', A'', B, B', B'' , and C all G -modules. Suppose given exact sequences:*

$$\begin{aligned} 0 &\rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0, \\ 0 &\rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0 \end{aligned}$$

and a G -pairing $A \times B \xrightarrow{\langle \rangle} C$. Assume $\langle A', B' \rangle = 0$ so there are induced pairings:

$$A' \times B'' \rightarrow C, \quad A'' \times B' \rightarrow C.$$

Then given $a \in H^p(G, A'')$, $b \in H^q(G, B'')$ we have:

$$\langle \partial_A a, b \rangle = (-1)^{p+1} \langle a, \partial_B b \rangle \in H^{p+q+1}(G, C).$$

(Here ∂_A and ∂_B are boundary maps for the corresponding long exact sequence of cohomology.)

Applying this result to the sequences (A.4), (A.5) and the pairing into $\bar{k}^*$ described above, we get a commutative diagram of pairings:

$$(A.10) \quad \begin{array}{ccc} H^0(\bar{k}/k, (\prod \mathbb{Z})_{\bar{S}}) \times H^2(\bar{k}/k, \Gamma(\bar{X} - \bar{S}, \mathcal{O}^*)/\bar{k}^*) & & \\ \downarrow \partial & \uparrow \partial & \searrow \text{evaluate} \\ H^1(\bar{k}/k, Z_{\bar{S}}) \times H^1(\bar{k}/k, N) & \xrightarrow{\langle \rangle} & Br(k) = H^2(\bar{k}/k, \bar{k}^*) \end{array}$$

[Commutativity means $\langle \partial z, n \rangle = \text{evaluation}(z, \partial n)$.]

The next step is to show that the pairing:

$$Z_{\bar{S}} \times N \rightarrow \bar{k}^*,$$

factors through the intersection pairing $N \otimes \bar{k}^* \otimes N \rightarrow \bar{k}^*$ by means of $Z_{\bar{S}} \rightarrow Z \rightarrow N \otimes \bar{k}^*$. This is essentially a consequence of Weil reciprocity as follows. Think of $N \otimes \bar{k}^* \simeq H^1(\bar{X}, \mathcal{K}_2)$. Given $\Delta \subset \bar{S}$ an irreducible divisor, let $\pi: \tilde{\Delta} \rightarrow \bar{X}$ be the normalization. There is a diagram:

$$\begin{array}{ccccccc} K_2(\bar{k}(\tilde{\Delta})) & \rightarrow & \prod_{x \in \tilde{\Delta}} \bar{k}(x)^* & \rightarrow & H^1(\tilde{\Delta}, \mathcal{K}_2) & \rightarrow & 0 \\ & \searrow 0 & \downarrow \text{norm} & \swarrow & & & \\ & & \bar{k}^* & & & & \end{array}$$

with exact top row. The diagonal arrow is zero by the reciprocity [cf. Lemma (A.8)], whence an induced map $H^1(\tilde{\Delta}, \mathcal{K}_2) \rightarrow \bar{k}^*$.

If we associate to Δ the composed map:

$$\varphi_{\Delta}: H^1(\bar{X}, \mathcal{K}_2) \rightarrow H^1(\tilde{\Delta}, \mathcal{K}_2) \rightarrow \bar{k}^*,$$

we obtain a pairing:

$$H^1(\bar{X}, \mathcal{K}_2) \times \text{Div}_{\bar{S}}(\bar{X}) \rightarrow \bar{k}^*.$$

One verifies quite easily ⁽¹⁰⁾ that the diagram of pairings:

$$(A.11) \quad \begin{array}{ccc} & N \otimes \bar{k}^* \times N & \\ \uparrow \parallel & \uparrow & \searrow \text{intersection pairing} \\ H^1(\bar{X}, \mathcal{K}_2) \times \text{Div}_{\bar{S}}(\bar{X}) & \longrightarrow & \bar{k}^* \\ \uparrow & \parallel & \nearrow \langle \rangle \\ \mathbb{Z}_{\bar{S}} \times \text{Div}_{\bar{S}}(\bar{X}) & & \end{array}$$

commutes.

Finally, noting that $\text{Pic}(\bar{X} - \bar{S}) = (0)$, one obtains a commutative diagram up to sign ([5], appendice):

$$(A.12) \quad \begin{array}{ccc} \text{Br}(X)/\text{Br}(k) & \longrightarrow & \text{Br}(X - S)/\text{Br}(k) \\ \downarrow & & \uparrow \\ H^1(\bar{k}/k, N) & \xrightarrow{\partial} & H^2(\bar{k}/k, \Gamma(\bar{X} - \bar{S}, \mathcal{O}^*)/\bar{k}^*). \end{array}$$

Combining (A.10), (A.11), (A.12) one obtains the assertion of (A.1).

Q.E.D.

REFERENCES

- [1] H. BASS, *Algebraic K-Theory*, New York, W. A. Benjamin Inc., 1968.
- [2] H. BASS, and J. TATE, *The Milnor Ring of a Global Field*, in *Algebraic K-Theory II* (Springer Lecture Notes in Math., No. 342, Springer-Verlag, 1973).
- [3] F. CHÂTELET, *Points rationnels sur certaines courbes et surfaces cubiques* (*Enseignement math.*, Vol. 5, 1959, pp. 153-170).
- [4] J.-L. COLLIOT-THÉLÈNE and D. CORAY, *L'équivalence rationnelle sur les points fermés des surfaces rationnelles fibrées en coniques* (*Compositio Mathematica*, Vol. 39, 1979, pp. 301-332).
- [5] J.-L. COLLIOT-THÉLÈNE and J.-J. SANSUC, *La R-équivalence sur les tores* (*Ann. scient. Éc. Norm. Sup.*, T. 10, 1977, pp. 175-230).
- [6] R. ELMAN and T. Y. LAM, *On the Quaternion Symbol Homomorphism $g_F: k_2 F \rightarrow B(F)$* , in *Algebraic K-Theory II* (Springer Lecture Notes in Math., No. 342, Springer-Verlag, 1973).
- [7] T. Y. LAM, *The Algebraic Theory of Quadratic Forms*, W. A. Benjamin, Inc., New York, 1973.
- [8] Yu. MANIN, *Le groupe de Brauer-Grothendieck en géométrie diophantienne* (*Actes Congrès Int. Math.*, Nice, 1970, pp. 401-411).
- [9] Yu. MANIN, *Cubic Forms*, North Holland, Amsterdam, 1974.
- [10] J. MILNOR, *Algebraic K-Theory and Quadratic Forms* (*Inventiones Math.*, Vol. 9, 1970, pp. 318-344).
- [11] J. MILNOR, *Introduction to Algebraic K-Theory* (*Ann. Math.*, Studies, No. 72, Princeton University Press, 1971).

⁽¹⁰⁾ Use functoriality of the map $\text{pic}(\bar{X}) \otimes \bar{k}^* \rightarrow H^1(\bar{X}, \mathcal{K}_2)$ for the morphism $\tilde{\Delta} \rightarrow X$, the fact that $\text{Pic } \bar{X} \rightarrow \text{Pic } \tilde{\Delta} \xrightarrow{\deg} \mathbb{Z}$ is the intersection number with $\tilde{\Delta}$, and the above diagram.

- [12] D. QUILLEN, *Higher Algebraic K-Theory*, in *Algebraic K-Theory I* (*Lecture Notes in Math.*, No. 341, Springer-Verlag, 1973).
- [13] J.-P. SERRE, *Corps Locaux*, Hermann, Paris, 1962.
- [14] C. SHERMAN, *K-Cohomology of Regular Schemes*, (to appear in *Communications in Algebra*).
- [15] J.-L. COLLIOT-THÉLÈNE and J.-J. SANSUC, A Series of Notes in *C. R. Acad. Sc.*, T. 282, série A, 1976; T. 284, série A, 1977; T. 284, 1977; T. 287, série A, 1978.
- [16] S. LANG, *Rapport sur la cohomologie des groupes*, Benjamin, New York, 1966.

Spencer BLOCH,
Princeton University,
Fine Hall, Department of Math.,
Box 37,
Princeton, N.J. 08540,
U.S.A.

(Manuscrit reçu le 15 mars 1980,
révisé le 15 juillet 1980.)