

# ANNALES SCIENTIFIQUES DE L'É.N.S.

HISAYOSI MATUMOTO

**Erratum : “ $C^{-\infty}$ -Whittaker vectors for complex semisimple Lie groups, wave front sets, and Goldie rank polynomial representations”**

*Annales scientifiques de l'É.N.S. 4<sup>e</sup> série*, tome 23, n° 4 (1990), p. 668 (supplément)

[http://www.numdam.org/item?id=ASENS\\_1990\\_4\\_23\\_4\\_S668\\_0](http://www.numdam.org/item?id=ASENS_1990_4_23_4_S668_0)

© Gauthier-Villars (Éditions scientifiques et médicales Elsevier), 1990, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'É.N.S. » (<http://www.elsevier.com/locate/ansens>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## ERRATUM

### $C^{-\infty}$ -Whittaker vectors for complex semisimple Lie groups, wave front sets, and Goldie rank polynomial representations

Hisayosi MATUMOTO <sup>(1)</sup>

---

*Ann. scient. Éc. Norm. Sup.*, 4<sup>e</sup> série, t. 23, 1990, p. 311 à 367.

---

In p355 line 5, I claimed the existence of a filtration  $1 \otimes S_w = L_q \subseteq L_{q-1} \subseteq \cdots \subseteq L_0 = E_p$  such that  $\bar{n}L_j \subseteq L_{j+1}$ . However, this statement is incorrect. A correct statement is : there exists a filtration  $1 \otimes S_w = L_q \subseteq L_{q-1} \subseteq \cdots \subseteq L_0 = E_p$  such that  $L_j/L_{j+1}$  ( $0 \leq j < q$ ) is equivalent to a quotient of  $S_w$ . Let  $B$  be the smashed product  $C[y, x] \# U(\bar{n})$  defined in 4.4. Then we can easily see the above filtration can be that of  $B$ -modules and  $L_j/L_{j+1}$  is equivalent to a quotient  $B$ -module of  $S_w$ . So, Theorem 4.1.1 is reduced to Lemma 4.3.16 and the following refinement of Lemma 4.3.17.

#### Lemma

Let  $\psi$  be a permissible character on  $\bar{n}$ . Let  $V$  be a  $B$ -submodule of  $S(\bar{U}_m/\bar{U}_m^w)'$ . Then we have

$$H^p(\bar{n}, (S(\bar{U}_m/\bar{U}_m^w)'/V) \otimes C_{-\psi}) = 0$$

for all  $p \in \mathbb{N}$  and  $w \in W_m^S - \{e\}$ .

The proof of the above lemma is just the same as that of Lemma 4.3.17.

At p361 line 24, in order to apply Nullstellensatz, we should regard  $y_0$  as a complex vector. It makes sense, since  $\psi' = \text{Ad}(e(y_0, 0)^{-1})\psi$  is permissible.

---

<sup>(1)</sup> Mathematical Sciences Research Institute, 1000 Centennial Drive, Berkeley, CA 94720 U.S.A.