

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

WILLIAM F., JR. DONOGHUE

A coerciveness inequality

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 20,
n° 3 (1966), p. 589-593

http://www.numdam.org/item?id=ASNSP_1966_3_20_3_589_0

© Scuola Normale Superiore, Pisa, 1966, tous droits réservés.

L'accès aux archives de la revue « *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A COERCIVENESS INEQUALITY

WILLIAM F. DONOGHUE, Jr.

Let Ω be an open bounded set in R^n with smooth boundary; by $H^1(\Omega)$ we denote the Hilbert space which is the completion of the smooth functions in Ω under the norm $\|u\|_1$ where

$$\|u\|_1^2 = \|u\|_0^2 + d_1(u).$$

Here $\|u\|_0$ is the usual L^2 norm over Ω and $d_1(u)$ is the Dirichlet integral given by

$$d_1(u) = \int_{\Omega} |\text{grad } u|^2 dx.$$

It is well known that the elements of $H^1(\Omega)$ are equivalence classes of functions and that the study of these functions requires some elementary potential theory. We recall that the capacity associated with the space $H^1(\Omega)$ is the set function $\text{cap}(A) = \inf \|u\|_1^2$ the infimum being taken over all smooth $u(x)$ which are ≥ 1 on A and that this function is an outer measure. The elements of $H^1(\Omega)$ are then determined as functions up to a set of capacity zero. If $u_n(x)$ is a minimizing sequence for the capacity of A , the u_n converge to a well defined element v_A of $H^1(\Omega)$ called the capacity potential of A , and which may be taken equal to 1 on A . By a simple variational argument one finds that there corresponds to v_A a positive measure μ_A supported by the closure of A called the capacity distribution such that $(u, v_A)_1 = \int u(x) d\mu_A$ for all u in $H^1(\Omega)$. Clearly $\|v_A\|_1^2 = \text{cap}(A) = (v_A, v_A)_1 = \int v_A(x) d\mu_A(x) = \int 1 d\mu_A = |\mu_A|$. If \mathcal{M}_A is the closure in $H^1(\Omega)$ of the smooth functions vanishing on a set A then this subspace is proper if and only if $\text{cap}(A) > 0$.

G. Stampacchia has conjectured that the following coerciveness assertion holds: when $\text{cap}(A)$ is positive, the quadratic norms $\|u\|_1$ and $\sqrt{d_1(u)}$ are

equivalent norms on \mathcal{M}_A . It is our purpose here to establish this conjecture. Since it is obvious that $d_1(u) \leq \|u\|_1^2$, what must be shown is the existence of a constant C (depending on A) such that for u in \mathcal{M}_A

$$(1) \quad \|u\|_0^2 \leq Cd_1(u).$$

Since we have supposed the boundary of Ω smooth, the Rellich theorem holds, i. e. the quadratic form $\|u\|_0^2$ is completely continuous relative to $\|u\|_1^2$. We may therefore write $\|u\|_0^2 = (Hu, u)_1$ where H is a positive operator which is completely continuous and of bound at most 1. It is easy to see that H has no null space, while H does have the eigenvalue 1 associated with the eigenfunction $u(x) = \text{constant}$. That eigenvalue is simple, since $Hv = v$ implies $\|v\|_0 = \|v\|_1$ and therefore $d_1(v) = 0$, from which we infer that $v = \text{constant}$, since its derivatives vanish almost everywhere.

Let P be the projection on the subspace \mathcal{M}_A ; then the operator PHP is positive, completely continuous and has bound $\lambda = \|PHP\|$ at most 1. Since $PHPu = u$ implies $Hu = u$ and therefore $u = \text{constant}$, and since the only constant function in \mathcal{M}_A vanishes identically, we see that $\lambda < 1$. It follows that for u in \mathcal{M}_A

$$\|u\|_0^2 \leq \lambda \|u\|_1^2 = \lambda \|u\|_0^2 + \lambda d_1(u)$$

and we obtain the desired inequality with $C = \frac{\lambda}{1 - \lambda}$.

It is possible to obtain an estimate for C in terms of the Lebesgue measure of Ω , the capacity of A and the number $\omega =$ smallest non-zero eigenvalue of the free membrane problem in Ω . For this purpose we write the eigenvalues of H in monotone decreasing order:

$$1 = \lambda_1 > \lambda_2 \geq \lambda_3 \geq \dots$$

and take $e_n(x)$ as the corresponding normalized eigenfunctions. Thus $e_1(x) = 1/\sqrt{m}$ where m is the Lebesgue measure of Ω . For the second eigenfunction we have $\|e_2\|_0^2 = \lambda_2 \|e_2\|_1^2$, whence $\omega \|e_2\|_0^2 = d_1(e_2)$ where $\omega = \lambda_2^{-1} - 1$, and this, by a classical argument, implies $-\Delta e_2(x) = \omega e_2(x)$ with the normal derivative of $e_2(x)$ vanishing on the boundary. Thus $e_2(x)$ is the eigenfunction of the free membrane problem for Ω and ω is the corresponding eigenvalue.

If v_A and μ_A are the capacity potential and distribution associated with A we have

$$(e_1, v_A)_1 = \int e_1(x) d\mu_A = \frac{|\mu_A|}{\sqrt{m}} = \frac{\text{cap}(A)}{\sqrt{m}}.$$

Our object is to estimate λ and hence C . We have $\lambda = \sup \frac{\|u\|_0^2}{\|u\|_1^2}$ the supremum being taken over all non-trivial u in \mathcal{M}_A .

Let \mathcal{M} be the subspace consisting of all u in H^1 orthogonal to v_A . Since the capacity is positive, v_A is not 0 and \mathcal{M} is proper; moreover \mathcal{M} contains \mathcal{M}_A since, for u in \mathcal{M}_A we have $(u, v_A)_1 = \int u(x) d\mu_A(x) = 0$.

Let $\lambda^* = \sup \frac{\|u\|_0^2}{\|u\|_1^2}$ the supremum being taken over all non zero u in \mathcal{M} . We then have $\lambda^* \geq \lambda$, and since \mathcal{M} contains no constant function other than 0, $\lambda^* < 1$. If Q is the projection on \mathcal{M} , λ^* is the largest eigenvalue of the positive, completely continuous operator QHQ . We estimate λ^* by the standard Aronszajn-Weinstein method. Let $R_\zeta = (H - \zeta I)^{-1}$ be the resolvent of H ; λ^* is then the (unique) zero of the function $(R_\zeta v_A, v_A)_1$ in the interval $\lambda_2 < \zeta < 1$. For the sake of completeness, we give an elementary proof for this special case. If $QHQw = \xi w$, then $Hw = \xi w + cv_A$ where the coefficient c may be 0. If $c = 0$, w is an eigenvector of H orthogonal to v_A , and therefore not e_1 . The number ξ is then one of the $\lambda_n < 1$, hence $\xi \leq \lambda_2$. If c is not 0 we have $(H - \xi I)w = cv_A$ whence $R_\xi v_A = c^{-1}w$, and therefore, since w in \mathcal{M} is orthogonal to v_A , $(R_\xi v_A, v_A)_1 = 0$ and ξ is a zero of the function $(R_\zeta v_A, v_A)_1$. Thus the spectrum of QHQ is a subset of the zeros and poles of this function. Conversely, if $(R_\xi v_A, v_A)_1 = 0$ for some ξ , we write $w = R_\xi v_A$ which is in \mathcal{M} and obtain $(H - \xi I)w = v_A$ or $Hw = \xi w + v_A$, whence $QHQw = \xi w$ and therefore ξ is an eigenvalue of QHQ . We seek the largest eigenvalue of that operator, and note that the function $(R_\zeta v_A, v_A)_1$ is monotone increasing and assumes all real values in the interval $\lambda_2 < \zeta < 1$, and is negative to the right of 1; hence λ^* is the (unique) zero of the function in that interval.

We therefore write out the function explicitly:

$$(R_\zeta v_A, v_A)_1 = \sum_{n=1}^{\infty} \frac{|(v_A, e_n)_1|^2}{\lambda_n - \zeta}$$

and note that the root λ^* is surely to the left of the root of

$$\frac{\|v_A\|_1^2 - |(v_A, e_1)_1|^2}{\lambda_2 - \zeta} + \frac{|(v_A, e_1)_1|^2}{1 - \zeta}.$$

The root is easily computed, and we find

$$\lambda \leq \lambda^* \leq 1 - (1 - \lambda_2) \frac{|(v_A, e_1)_1|^2}{\|v_A\|_1^2}$$

and therefore

$$C \leq (1 - \lambda_2)^{-1} \frac{\|v_A\|_1^2}{|(v_A, e_1)_1|^2} - 1 < (1 + 1/\omega) \frac{\text{meas } (\Omega)}{\text{cap } (A)}.$$

The foregoing estimate for the constant in Stampacchia's inequality has the disadvantage that it involves the capacity of A relative to the space $H^1(\Omega)$ and that this function is not known. However, as we shall presently show, this function is equivalent to the usual capacity for the corresponding Bessel potentials, a set function usually written $\gamma_2(A)$. There exists a constant M depending only on Ω such that

$$M\gamma_2(A) \leq \text{cap } (A) \leq \gamma_2(A)$$

for all subsets A of Ω , and therefore the constant occurring in inequality (1) involves a numerator which depends only on the domain Ω and a denominator $\gamma_2(A)$; it therefore is independent of any other property of A , for example, the distance of that set from the boundary.

The equivalence of the set functions $\gamma_2(A)$ and $\text{cap } (A)$ is a consequence of the smoothness hypothesis made concerning the boundary of Ω ; there exists a continuous linear transformation $u \rightarrow \tilde{u}$ mapping $H^1(\Omega)$ into $P^1(\mathbb{R}^n)$, the space of Bessel potentials on \mathbb{R}^n such that $u(x) = \tilde{u}(x)$ for all x in Ω . The transformation is bounded; thus there exists a positive M such that $\|u\|_1^2 \geq M \|\tilde{u}\|_1^2$. If v_A is the capacity potential for A in the space $P^1(\mathbb{R}^n)$ we have

$$\gamma_2(A) = \|v_A\|_1^2 \geq \|v^*\|_1^2 \geq \text{cap } (A)$$

where v^* is the restriction of v_A to Ω considered as an element of $H^1(\Omega)$. Conversely, if v_A in $H^1(\Omega)$ is the capacity potential of A ,

$$\text{cap } (A) = \|v_A\|_1^2 \geq M \|\tilde{v}_A\|_1^2 \geq M \gamma_2(A).$$

It is natural to enquire to what extent inequality (1) is valid for the spaces $H^\alpha(\Omega)$ where the norm is defined by

$$\|u\|_\alpha^2 = \|u\|_0^2 + d_\alpha(u)$$

$$\text{with } d_\alpha(u) = \iint_{\Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy \quad \text{when } 0 < \alpha < 1$$

and $d_\alpha(u) = \Sigma d_1(D_k u)$ when α is an integer, the summation being taken over all derivatives of order α , and finally when $\alpha > 1$ is not an integer, $d_\alpha(u) = \Sigma d_\beta(D_k u)$ where the summation is taken over all derivatives of order k , k being the largest integer $< \alpha$ and β defined by $\alpha = k + \beta$.

All of the arguments we have given carry over to the case $\alpha < 1$; Stampacchia's inequality is valid with a constant which depends only on the domain Ω and the reciprocal of the capacity $\gamma_{2\alpha}(A)$, this being the capacity for the corresponding space of Bessel potentials $P^\alpha(R^n)$.

The situation is essentially more complex when $\alpha > 1$; if we repeat our analysis we find that the operator H which represents the L^2 norm in the space $H^\alpha(\Omega)$ is positive, completely continuous and with bound 1, however, the eigenvalue 1 is no longer simple. The eigenspace corresponding to that eigenvalue consists of all polynomials of sufficiently low degree, and such a polynomial may vanish on a set of positive capacity. Thus the inequality does not hold, unless a further hypothesis is made, viz. that the set A is not contained in the set of zeros of a polynomial of degree $\leq m =$ the largest integer strictly smaller than α . In this case inequality (1) is valid, but the constant depends in an essential way on the other data than simply the capacity $\gamma_{2\alpha}(A)$.

Let us remark that the surface on the unit sphere in R^n is a set of positive capacity, but is contained in the null set of the polynomial $1 - |x|^2$.

Throughout our discussion we have made use of the hypothesis that the boundary of Ω is smooth in order that the Rellich theorem guaranteeing the complete continuity of $\|u\|_0^2$ should hold. We have also used that hypothesis to have the extension theorem embedding $H^1(\Omega)$ into $P^1(R^n)$. The careful study of these questions given in [1] shows that the regularity hypotheses needed are very mild.

1. R. ADAMS, N. ARONSZAJN and K. T. SMITH, « *Theory of Bessel Potentials II* » Annales Institut Fourier, to appear.

*Michigan State University
East Lansing, Mich.*