

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

S. K. CHATTERJEA

Operational representations for the Laguerre polynomials

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 20,
n° 4 (1966), p. 739-744

<http://www.numdam.org/item?id=ASNSP_1966_3_20_4_739_0>

© Scuola Normale Superiore, Pisa, 1966, tous droits réservés.

L'accès aux archives de la revue « *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

OPERATIONAL REPRESENTATIONS FOR THE LAGUERRE POLYNOMIALS

By S. K. CHATTERJEA

1. Carlitz [1] gave the following operational representation for the Laguerre polynomials:

$$(1.1) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} \prod_{j=1}^n (x D - x + \alpha + j) \cdot 1.$$

Recently Al-Salam [2] has given the operational formula

$$(1.2) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha-n} e^x \{x(1+x D)\}^n x^\alpha e^{-x},$$

which is closely related to the formula of Chak [3]:

$$(1.3) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha-n-1} e^x (x^2 D)^n x^{\alpha+1} e^{-x}.$$

In a recent paper [4], M. K. Das has proved the operational formula

$$(1.4) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha-n-k} e^x \{x(x D - k + 1)\}^n x^{\alpha+k} e^{-x}.$$

It may be noted that (1.2) and (1.3) are special cases of (1.4).

Moreover, one may note from (1.4) the following special case

$$(1.5) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha-n-2} e^x \{x(x D - 1)\}^n x^{\alpha+2} e^{-x}.$$

In [2, pp. 129-136] Al-Salam has proved several results involving Laguerre polynomials by the operational formula (1.2). The object of this paper is to derive some formulas of Laguerre polynomials by employing formulas of Carlitz and Chak, and such type of derivation does not seem to appear in the earlier investigation.

2. For our purpose, we write (1.1) in the form :

$$(2.1) \quad L_n^{(\alpha)}(x) = \frac{1}{n!} e^x (\delta + \alpha + 1)_n e^{-x}; \quad \delta \equiv x D$$

For Chak's operator we notice that [5]

$$(2.2) \quad e^{tx\delta} f(x) = f\left(\frac{x}{1-tx}\right)$$

and

$$(2.3) \quad e^{t\delta} f(x) = f(x e^t).$$

Again it is easy to prove that

$$(2.4) \quad (x \delta)^n x^{\alpha+k} = (\alpha + k)_n x^{\alpha+k+n}.$$

First we shall prove the following generating function :

$$(2.5) \quad \sum_{n=0}^{\infty} \frac{t^n}{(\alpha + 1)_n} L_n^{(\alpha)}(x) = e^t {}_0F_1[-; \alpha + 1; -x t].$$

We have

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{t^n}{(\alpha + 1)_n} L_n^{(\alpha)}(x) \\ &= \sum_{n=0}^{\infty} \frac{t^n}{n! (\alpha + 1)_n} e^x (\delta + \alpha + 1)_n e^{-x} \\ &= e^x {}_1F_1[\delta + \alpha + 1; \alpha + 1; t] e^{-x} \\ &= e^{x+t} {}_1F_1[-\delta; \alpha + 1; -t] e^{-x} \\ &= e^{x+t} \sum_{n=0}^{\infty} \frac{(-\delta)_n}{(\alpha + 1)_n n!} (-t)^n e^{-x} \\ &= e^{x+t} \sum_{n=0}^{\infty} \frac{(x t)^n}{(\alpha + 1)_n n!} D^n e^{-x} \end{aligned}$$

$$\begin{aligned} &= e^{x+t} \sum_{n=0}^{\infty} \frac{(-x t)^n}{(\alpha + 1)_n n!} e^{-x} \\ &= e^t {}_0F_1[-; \alpha + 1; -x t]. \end{aligned}$$

Another way of obtaining (2.5) is the following:

From (1.3) we have

$$\begin{aligned} &\sum_{n=0}^{\infty} \frac{(x t)^n}{(\alpha + 1)_n} L_n^{(\alpha)}(x) \\ &= \sum_{n=0}^{\infty} \frac{t^n}{(\alpha + 1)_n n!} x^{-\alpha-1} e^x (x \delta)^n x^{\alpha+1} e^{-x} \\ &= x^{-\alpha-1} e^x \sum_{n=0}^{\infty} \frac{t^n}{(\alpha + 1)_n n!} (x \delta)^n x^{\alpha+1} e^{-x} \\ (2.6) \quad &= x^{-\alpha-1} e^x {}_0F_1[-; \alpha + 1; t x \delta] x^{\alpha+1} e^{-x}. \end{aligned}$$

But

$$\begin{aligned} &{}_0F_1[-; \alpha + 1; t x \delta] x^{\alpha+1} e^{-x} \\ &= {}_0F_1[-; \alpha + 1; t x \delta] \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} x^{\alpha+r+1} \\ &= x^{\alpha+1} \sum_{r=0}^{\infty} \frac{(-x)^r}{r!} {}_1F_1[\alpha + r + 1; \alpha + 1; x t] \\ &= x^{\alpha+1} e^{xt} \sum_{r=0}^{\infty} \frac{(-x)^r}{r!} {}_1F_1[-r; \alpha + 1; -x t] \\ (2.7) \quad &= x^{\alpha+1} e^{xt-x} {}_0F_1[-; \alpha + 1; -x^2 t]. \end{aligned}$$

It follows therefore from (2.6) and (2.7) that

$$(2.8) \quad \sum_{n=0}^{\infty} \frac{(x t)^n}{(\alpha + 1)_n} L_n^{(\alpha)}(x) = e^{xt} {}_0F_1[-; \alpha + 1; -x^2 t],$$

which is (2.5).

We remark therefore that the method adopted at first in proving (2.5) is quite simple and interesting.

Next we shall prove the following generating function:

$$(2.9) \quad \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha + 1)_n} t^n L_n^{(\alpha)}(x) = (1 - t)^{-c} {}_1F_1\left[c; \alpha + 1; \frac{-x t}{1 - t}\right].$$

Here we have

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha+1)_n} t^n L_n^{(\alpha)}(x) \\
 &= \sum_{n=0}^{\infty} \frac{(c)_n t^n}{(\alpha+1)_n n!} e^x (\delta + \alpha + 1)_n e^{-x} \\
 &= e^x {}_2F_1[\delta + \alpha + 1, c; \alpha + 1; t] e^{-x} \\
 &= e^x (1-t)^{\alpha+1-c-\delta-\alpha-1} {}_2F_1[-\delta, \alpha + 1 - c; \alpha + 1; t] e^{-x} \\
 &= e^x (1-t)^{-c} \left(\frac{1}{1-t}\right)^{\delta} \sum_{n=0}^{\infty} \frac{(\alpha+1-c)_n}{(\alpha+1)_n} \frac{(-xt)^n}{n!} D^n e^{-x} \\
 &= e^x (1-t)^{-c} \left(\frac{1}{1-t}\right)^{\delta} {}_1F_1[\alpha+1-c; \alpha+1; xt] e^{-x}.
 \end{aligned}$$

From (2.3) we notice that

$$a^{\delta} f(x) = f(ax).$$

Thus we derive

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha+1)_n} t^n L_n^{(\alpha)}(x) \\
 &= e^x (1-t)^{-c} {}_1F_1\left[\alpha+1-c; \alpha+1; \frac{xt}{1-t}\right] e^{-\frac{x}{1-t}} \\
 &= e^x (1-t)^{-c} e^{\frac{xt}{1-t}} {}_1F_1\left[c; \alpha+1; \frac{-xt}{1-t}\right] e^{-\frac{x}{1-t}} \\
 &= (1-t)^{-c} {}_1F_1\left[c; \alpha+1; \frac{-xt}{1-t}\right].
 \end{aligned}$$

Alternatively we derive (2.9) in the following manner:

From (1.3) we have

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha+1)_n} (xt)^n L_n^{(\alpha)}(x) \\
 &= \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha+1)_n} \frac{t^n}{n!} x^{-\alpha-1} e^x (x\delta)^n x^{\alpha+1} e^{-x} \\
 (2.10) \quad &= x^{-\alpha-1} e^x {}_1F_1[c; \alpha+1; tx\delta] x^{\alpha+1} e^{-x}.
 \end{aligned}$$

Now

$$\begin{aligned}
 & {}_1F_1 [c; \alpha + 1; tx\delta] x^{\alpha+1} e^{-x} \\
 &= x^{\alpha+1} \sum_{r=0}^{\infty} \frac{(-x)^r}{r!} {}_2F_1 [r + \alpha + 1, c; \alpha + 1; xt] \\
 &= x^{\alpha+1} \sum_{r=0}^{\infty} \frac{(-x)^r}{r!} (1 - xt)^{-c-r} {}_2F_1 [-r, \alpha + 1 - c; \alpha + 1; xt] \\
 &= x^{\alpha+1} (1 - xt)^{-c} \sum_{r=0}^{\infty} \frac{\left(\frac{-x}{1 - xt}\right)^r}{r!} {}_2F_1 [-r, \alpha + 1 - c; \alpha + 1; xt] \\
 &= x^{\alpha+1} (1 - xt)^{-c} e^{\frac{-x}{1 - xt}} {}_1F_1 \left[\alpha + 1 - c; \alpha + 1; \frac{x^2 t}{1 - xt} \right] \\
 (2.11) \quad &= x^{\alpha+1} (1 - xt)^{-c} e^{\frac{-x}{1 - xt} + \frac{x^2 t}{1 - xt}} {}_1F_1 \left[c; \alpha + 1; -\frac{x^2 t}{1 - xt} \right].
 \end{aligned}$$

Thus it follows from (2.10) and (2.11) that

$$(2.12) \quad \sum_{n=0}^{\infty} \frac{(c)_n}{(\alpha + 1)_n} (xt)^n L_n^{(\alpha)}(x) = (1 - xt)^{-c} {}_1F_1 \left[c; \alpha + 1; \frac{-x^2 t}{1 - xt} \right],$$

which is (2.9).

Using the operational formula of Carlitz, we shall finally prove the Hardy-Hille formula:

$$\begin{aligned}
 (2.13) \quad & \sum_{n=0}^{\infty} \frac{n!}{(\alpha + 1)_n} L_n^{(\alpha)}(x) L_n^{(\alpha)}(y) t^n \\
 &= (1 - t)^{-\alpha-1} \exp \left[-\frac{(x + y)t}{1 - t} \right] {}_0F_1 \left[-; \alpha + 1; \frac{xyt}{(1 - t)^2} \right].
 \end{aligned}$$

We observe that

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \frac{n!}{(\alpha + 1)_n} t^n L_n^{(\alpha)}(x) L_n^{(\alpha)}(y) \\
 &= \sum_{n=0}^{\infty} \frac{n!}{(\alpha + 1)_n} t^n \frac{1}{n!} e^{x(\delta + \alpha + 1)_n} e^{-x} \frac{1}{n!} e^{y(\theta + \alpha + 1)_n} e^{-y} \\
 & \quad \left(\text{where } \delta = x \frac{d}{dx}, \text{ and } \theta = y \frac{d}{dy} \right)
 \end{aligned}$$

$$\begin{aligned}
&= e^{x+y} \sum_{n=0}^{\infty} \frac{(\delta + \alpha + 1)_n (\theta + \alpha + 1)_n t^n}{(\alpha + 1)_n n!} e^{-(x+y)} \\
&= e^{x+y} {}_2F_1[\delta + \alpha + 1, \theta + \alpha + 1; \alpha + 1; t] e^{-(x+y)} \\
&= e^{x+y} (1-t)^{-\delta-\theta-\alpha-1} {}_2F_1[-\delta, -\theta; \alpha + 1; t] e^{-(x+y)} \\
&= e^{x+y} (1-t)^{-\alpha-1} (1-t)^{-\delta-\theta} {}_0F_1[-; \alpha + 1; xy t] e^{-(x+y)} \\
&= e^{x+y} (1-t)^{-\alpha-1} \left(\frac{1}{1-t}\right)^\delta \left(\frac{1}{1-t}\right)^\theta {}_0F_1[-; \alpha + 1; xy t] e^{-(x+y)} \\
&= e^{x+y} (1-t)^{-\alpha-1} {}_0F_1\left[-; \alpha + 1; \frac{xy t}{(1-t)^2}\right] e^{\frac{-(x+y)}{1-t}} \\
&= (1-t)^{-\alpha-1} e^{\frac{-(x+y)t}{1-t}} {}_0F_1\left[-; \alpha + 1; \frac{xy t}{(1-t)^2}\right].
\end{aligned}$$

REFERENCES

- [1] L. CARLITZ, *A note on the Laguerre polynomials*, Michigan Math. Jour., vol. 7 (1960) pp. 219-223.
- [2] W. A. AL-SALAM, *Operational representations for the Laguerre and other polynomials*, Duke Math. Jour., vol. 31 (1964), pp. 127-142.
- [3] A. M. CHAK, *A class of polynomials and a generalization of Stirling numbers*, Duke Math. Jour., vol. 23 (1956), pp. 45-56.
- [4] M. K. DAS, *Operational representations for the Laguerre polynomials*, - to appear elsewhere.
- [5] E. STEPHENS, *The elementary theory of operational mathematics*, (New York, 1937), pp. 184.