

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

JOSÉ FRANCISCO RODRIGUES

LISA SANTOS

**A parabolic quasi-variational inequality arising in a
superconductivity model**

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 29,
n° 1 (2000), p. 153-169

http://www.numdam.org/item?id=ASNSP_2000_4_29_1_153_0

© Scuola Normale Superiore, Pisa, 2000, tous droits réservés.

L'accès aux archives de la revue « *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A Parabolic Quasi-Variational Inequality Arising in a Superconductivity Model

JOSÉ FRANCISCO RODRIGUES – LISA SANTOS

Abstract. We consider the existence of solutions for a parabolic quasilinear problem with a gradient constraint which threshold depends on the solution itself. The problem may be considered as a quasi-variational inequality and the existence of solution is shown by considering a suitable family of approximating quasilinear equations of p -Laplacian type. *A priori* estimates on the time derivative of the approximating solutions and on the nonlinear diffusion coefficients are used in the passage to the limit, as well as a suitable sequence of convex sets with variable gradient constraint. The asymptotic behaviour as $t \rightarrow \infty$ is also considered, and the solutions of the quasi-variational inequality are shown to converge, at least for subsequences, to a solution of a stationary quasi-variational inequality. These results can be applied to the critical-state model of type-II superconductors in longitudinal geometry.

Mathematics Subject Classification (1991): 35K85 (primary), 35K55, 35R35 (secondary).

1. – Introduction

In a critical-state model of type-II superconductors with a longitudinal geometry, the main unknown is the magnetic field $H = (0, 0, u(x, t))$, where $x = (x_1, x_2) \in \Omega \subseteq \mathbb{R}^2$. By Maxwell's equations,

$$\begin{cases} \mu H_t + \nabla \times E = 0, \\ \nabla \times H = J, \end{cases}$$

where E denotes the electric field, the unknown density of the induced current J is given by

$$J = \left(\frac{\partial u}{\partial x_2}, -\frac{\partial u}{\partial x_1}, 0 \right) = \nabla^\perp u .$$

In some superconductivity models (see [3]), it is assumed that the electric field E inside the superconductor depends on the current density J through

a power law (which generalizes the classical Ohm's law $E = \rho J$) with the scalar resistivity given by $\rho = \rho(|J|) = \rho_0 |J|^{p-2} = \rho_0 |\nabla u|^{p-2}$, where $\rho_0 > 0$ is a constant and $p \geq 2$. Hence, the Maxwell's equation for the longitudinal component of the magnetic field implies that $u = u(x, t)$ satisfies the two-dimensional quasilinear parabolic equation

$$\mu u_t - \rho_0 \nabla \cdot (|\nabla u|^{p-2} \nabla u) = 0.$$

In Bean's critical-state model (see [11]), the current density cannot exceed the critical value $j_c > 0$ and it has been suggested that this threshold may also depend on $|H| = |u|$ (see Kim *et al.* [6]).

The constitutive relation for E is then modified to

$$E = \begin{cases} \rho_0 |\nabla u|^{p-2} \nabla^\perp u & \text{if } |\nabla u| < j_c(|u|), \\ (\rho_0 j_c^{p-2} + \lambda) \nabla^\perp u & \text{if } |\nabla u| = j_c(|u|), \end{cases}$$

being $\lambda \geq 0$ an unknown Lagrange multiplier.

Imposing initial and boundary conditions (in a bounded domain Ω in \mathbb{R}^2), it may be easily shown that this problem is equivalent to the following quasi-variational inequality (for details about this derivation see [11], where only the degenerate case $\rho_0 = 0$ was considered) to a new variable $h = u - h_e$,

$$\left\{ \begin{array}{l} h(t) \in \mathbb{K}_{h(t)} \text{ for a.e. } t \in [0, T], \quad h(0) = h_0 \in \mathbb{K}_{h_0}, \\ \int_{\Omega} h_t(t) (v - h(t)) + \frac{\rho_0}{\mu} \int_{\Omega} |\nabla h(t)|^{p-2} \nabla h(t) \cdot \nabla (v - h(t)) \\ \geq \int_{\Omega} f(t) (v - h(t)), \quad \forall v \in \mathbb{K}_{h(t)} \text{ for a.e. } t \in [0, T], \end{array} \right.$$

being

$$\mathbb{K}_{h(t)} = \left\{ v \in W_0^{1,p}(\Omega) : |\nabla v| \leq j_c(|h(t) + h_e(t)|) \text{ a.e.} \right\}$$

where $h_e = h_e(t)$ is related to the density of external currents, as well as f .

This model is the main motivation for the study of this new type of quasi-variational inequalities. This kind of problems seem not well studied in the literature and are not considered, for instance, in the classical references on quasi-variational inequalities [1] or [2]. Recent works in this area are, for example, the modelling works [10], [11] or a very special case of a one-dimensional quasi-variational inequality considered in [7].

In Section 2 we present the main results of this paper, a result on the existence of solutions for the quasi-variational inequality and a result about the asymptotic behaviour in time of these solutions.

In Section 3 we consider a family of approximating solutions and we establish *a priori* estimates. Section 4 includes the passage to the limit in the family of solutions of the approximated problems, concluding with the proof of the existence of at least a solution for the quasi-variational inequality in the N -dimensional case.

Section 5 studies the asymptotic behaviour in time of the solutions, which, in particular, also yields the existence of a solution to the corresponding elliptic quasi-variational inequality, extending a result of [8].

2. – Main results

Let Ω be a bounded, open subset of \mathbb{R}^N , with smooth boundary $\partial\Omega$, and let $T \in \mathbb{R}^+$, $I =]0, T[$, $Q_T = \Omega \times I$, $\Sigma = \partial\Omega \times I$, $\Omega_0 = \overline{\Omega} \times \{0\}$. Let $\nabla = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_N})$ denote the spatial gradient and suppose that F , f and h are given functions such that

- (1) $F \in C^0(\mathbb{R})$, $\exists m > 0$: $F \geq m$,
- (2) $f \in L^\infty(Q_T)$; $f_t \in M(Q_T)$,
- (3) $h \in W_0^{1,\infty}(\Omega)$, $|\nabla h| \leq F(h)$ a.e. in Ω , $\Delta_p h \in M(\Omega)$,

where $\Delta_p h = \nabla \cdot (|\nabla h|^{p-2} \nabla h)$ denotes the p -Laplacian, $1 < p < \infty$ and $M(\Omega)$, $M(Q_T)$ denote the spaces of bounded measures in Ω and Q_T , respectively, $M(\Omega) = [C^0(\overline{\Omega})]'$ and $M(Q_T) = [C^0(\overline{Q_T})]'$. We also denote by $W_0^{1,\infty}(\Omega)$ the space of Lipschitz functions that vanish on $\partial\Omega$.

Given $u \in L^\infty(\Omega)$, we can define a convex set

$$(4) \quad \mathbb{K}_u = \left\{ v \in W_0^{1,p}(\Omega) : |\nabla v| \leq F(u) \text{ a.e. in } \Omega \right\}$$

and the quasi-variational problem as follows:

To find u in an appropriate class of functions, such that:

$$(5) \quad \begin{cases} u(t) \in \mathbb{K}_{u(t)} \cap L^\infty(\Omega) \text{ for a.e. } t \in I , & u(0) = h , \\ \int_{\Omega} u_t(t) (v - u(t)) + \int_{\Omega} |\nabla u(t)|^{p-2} \nabla u(t) \cdot \nabla (v - u(t)) \\ \geq \int_{\Omega} f(t) (v - u(t)) , \forall v \in \mathbb{K}_{u(t)} , & \text{for a.e. } t \in I . \end{cases}$$

Our first aim is to prove existence of a solution to the problem (5). For that, we consider, for $\varepsilon > 0$, f_ε , h_ε and F_ε smooth functions approximating, respectively, f , h and F in the norms of $L^q(Q_T)$, $W_0^{1,q}(\Omega)$, for any $q < \infty$ and in $C^0(\mathbb{R})$, with $h^\varepsilon|_{\partial\Omega} = 0$, $\|\Delta_p h^\varepsilon + \varepsilon \Delta h^\varepsilon\|_{L^1(\Omega)} \leq C$ (C independent of ε) and a C^2 function $k_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$, non decreasing and such that

$$(6) \quad k_\varepsilon = 1 \text{ if } s \leq 0, \quad k_\varepsilon = e^{\frac{s}{\varepsilon}} \text{ if } s \geq \varepsilon.$$

We consider a sequence of the following approximated problems

$$(7) \quad \begin{cases} u_t^\varepsilon - \nabla \cdot \left[k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-2}{2}} \nabla u^\varepsilon + \varepsilon \nabla u^\varepsilon \right] = f_\varepsilon \text{ in } Q_T , \\ u^\varepsilon(0) = h^\varepsilon , \\ u^\varepsilon|_{\Sigma} = 0 . \end{cases}$$

We noticed that a similar approximation was introduced by Gerhardt (see [5]) in the treatment of the elastic-plastic torsion problem with multiply connected cross-section, which is an elliptic variational inequality and it was also used for studying a parabolic variational inequality with gradient constraint in [12].

THEOREM 1. *With the assumptions (1), (2) and (3), for any $1 < p < \infty$, the problem (5) has at least a solution u belonging to $L^\infty(0, T; W_0^{1,\infty}(\Omega)) \cap C^0(\overline{Q}_T)$, such that $u_t \in L^\infty(0, T; M(\Omega))$. In addition, u is the weak limit in $L^q(0, T; W_0^{1,q}(\Omega))$ (for any $q < \infty$) of a subsequence $\{u^{\varepsilon_n}\}_n$ of solutions of the family of approximated problems (7), $u^{\varepsilon_n} \rightharpoonup_n u$ in $C^0(\overline{Q}_T)$ and also $u_t^{\varepsilon_n} \rightharpoonup u_t$ in $L^\infty(0, T; M(\Omega))$ weak-**.

REMARK 1. Since u is continuous and bounded, for each $t \in [0, T]$, $\mathbb{K}_{u(t)} \subset W_0^{1,\infty}(\Omega) \subset C^0(\overline{\Omega})$ and the first integral of the left hand side of (5) should be understood in the duality sense between $M(\Omega)$ and $C^0(\overline{\Omega})$, taking into account that $u_t(t)$ is a.e. a bounded measure.

Consider now the corresponding stationary quasi-variational inequality:

To find

$$(8) \quad \begin{cases} u^\infty \in \mathbb{K}_{u^\infty} \cap L^\infty(\Omega) , \\ \int_\Omega |\nabla u^\infty|^{p-2} \nabla u^\infty \cdot \nabla (v - u^\infty) \geq \int_\Omega f^\infty (v - u^\infty) , \quad \forall v \in \mathbb{K}_{u^\infty} , \end{cases}$$

where we assume $f^\infty \in L^1(\Omega)$, F satisfies (1) and

$$\mathbb{K}_{u^\infty} = \left\{ v \in W_0^{1,p}(\Omega) : |\nabla v| \leq F(u^\infty) \text{ a.e. in } \Omega \right\} .$$

Existence of a solution of the problem (8) may be proved as in [3].

We may also consider the problem (5) with $T = +\infty$. Then we have the following

THEOREM 2. *Suppose that the assumption (2) is satisfied for $T = +\infty$, i.e., there exists $C > 0$ such that*

$$(9) \quad \sup_{0 < t < \infty} \|f(t)\|_{L^\infty(\Omega)} + \int_0^{+\infty} \int_\Omega |f_t| \leq C ,$$

and, in addition,

$$(10) \quad \frac{1}{t} \int_0^t \int_\Omega |f_t(\tau)| \tau d\tau \longrightarrow 0 \quad \text{when } t \rightarrow \infty ,$$

$$(11) \quad f(t) \rightharpoonup f_\infty \text{ in } L^1(\Omega)\text{-weak, when } t \rightarrow +\infty .$$

Let u denote a solution of the problem (5). Then there exists a sequence $\{t_n\}_n$, $t_n \xrightarrow[n]{+}$ $+\infty$, and a function $u^\infty \in W_0^{1,\infty}(\Omega)$ such that

$$\begin{cases} u(t_n) \xrightarrow[n]{+} u^\infty \text{ in } W_0^{1,\infty}(\Omega) \text{ weak-* and uniformly in } \overline{\Omega}, \\ u^\infty \text{ is a solution of the problem (8).} \end{cases}$$

3. – The approximating problems

Let us consider the family of approximated problems defined by (7).

PROPOSITION 1. *With the assumptions (1), (2) and (3), the problem (7) has a unique solution $u^\varepsilon \in C_{\alpha, \alpha/2}^{2,1}(Q_T) \cap C^0(\overline{Q}_T)$, $0 < \alpha < 1$.*

PROOF. This is a direct consequence of well known results for quasilinear parabolic equations (see [9]). \square

We shall establish some a priori estimates for the solutions of the approximated problems.

LEMMA 1. *Suppose that (1), (2) and (3) are satisfied and that u^ε is the solution of the problem (7). Then there exists a constant $M > 0$ independent of $\varepsilon \in]0, 1[$ such that*

$$(12) \quad |u^\varepsilon(x, t)| \leq M \quad \text{for all } (x, t) \in \overline{Q}_T .$$

PROOF. This estimate is a consequence of the assumptions and global boundedness for weak solutions of the Dirichlet problem for quasilinear degenerate parabolic equations (see [4] and its references). We need only to remark that in (7) $k_\varepsilon \geq 1$ and this is the crucial fact for deriving the weak maximum principle estimate. \square

REMARK 2. We remark that the constant M that appears in (12) depends on T only through $\|f(t)\|_{L^\infty(\Omega)}$, and therefore it will be independent of T if $f \in L^\infty(0, \infty; L^\infty(\Omega)) = L^\infty(Q_\infty)$, i.e., under the assumption (9).

LEMMA 2. *Suppose that (1), (2) and (3) are satisfied and u^ε is the solution of the problem (7). Then there exists a constant $C > 0$ independent of $\varepsilon \in]0, 1[$ such that*

$$\|u_t^\varepsilon\|_{L^\infty(0,T;L^1(\Omega))} \leq C .$$

PROOF. We differentiate the equation of the approximated problem (7) in order to t and set $v = u_t^\varepsilon$. Then

$$(13) \quad \begin{aligned} & v_t - \nabla \cdot \left[k_\varepsilon' (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (2 \nabla u^\varepsilon \cdot \nabla v - 2 F_\varepsilon F_\varepsilon' v) (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla u^\varepsilon \right. \\ & + k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) \frac{p-2}{2} (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-4}{2}} 2 \nabla u^\varepsilon \cdot \nabla v \nabla u^\varepsilon \\ & \left. + k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla v + \varepsilon \nabla v \right] = f_t^\varepsilon . \end{aligned}$$

Consider the sign function, denoted by sgn^0 and defined as follows:

$$\text{sgn}^0(\tau) = \begin{cases} -1 & \text{if } \tau < 0, \\ 0 & \text{if } \tau = 0, \\ 1 & \text{if } \tau > 0. \end{cases} \quad (\tau \in \mathbb{R}) ,$$

Let $s_\delta : \mathbb{R} \rightarrow \mathbb{R}$, $\delta > 0$, be a sequence of C^2 functions, approximating pointwise sgn^0 when $\delta \rightarrow 0$, such that

$$s_\delta|_{\{0\} \cup \mathbb{R} \setminus]-\delta, \delta[} = \text{sgn}^0|_{\{0\} \cup \mathbb{R} \setminus]-\delta, \delta[}, \quad s'_\delta \geq 0.$$

Notice that, if $\zeta_\delta(\tau) = \int_0^\tau s_\delta(\sigma) d\sigma$, then

$$\forall \tau \in \mathbb{R} \quad \lim_{\delta \rightarrow 0} \tau s'_\delta(\tau) = 0, \quad \lim_{\delta \rightarrow 0} \zeta_\delta(\tau) = |\tau|.$$

Multiply the equation (13) by $s_\delta(v)$ and integrate over Q_t . Then, since $v|_{\partial\Omega \times]0, t[} = 0$ and $s_\delta(v)|_{\partial\Omega \times]0, t[} = 0$, we have

$$\begin{aligned} & \int_{\Omega} \zeta_\delta(v(t)) - \int_{\Omega} \zeta_\delta(v(0)) + \int_{Q_t} 2k'_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (\nabla u^\varepsilon \cdot \nabla v)^2 (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} s'_\delta(v) \\ & + \int_{Q_t} -2k'_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) F_\varepsilon F'_\varepsilon (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla u^\varepsilon \cdot \nabla v \, v s'_\delta(v) \\ (14) \quad & + \int_{Q_t} k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (p-2) (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-4}{2}} (\nabla u^\varepsilon \cdot \nabla v)^2 s'_\delta(v) \\ & + \int_{Q_t} k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} |\nabla v|^2 s'_\delta(v) \\ & + \int_{Q_t} k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) \varepsilon |\nabla v|^2 s'_\delta(v) = \int_{Q_t} f_t^\varepsilon s_\delta(v). \end{aligned}$$

Notice that, since k_ε is monotone nondecreasing,

$$\begin{aligned} & \int_{Q_t} 2k'_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) (\nabla u^\varepsilon \cdot \nabla v)^2 (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} s'_\delta(v) \geq 0 \\ & \int_{Q_t} k_\varepsilon (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) \varepsilon |\nabla v|^2 s'_\delta(v) \geq 0, \end{aligned}$$

and

$$\int_{Q_t} -2k'_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) F_\varepsilon F'_\varepsilon (|\nabla u^\varepsilon|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla u^\varepsilon \cdot \nabla v \, v s'_\delta(v) \xrightarrow{\delta \rightarrow 0} 0.$$

On the other hand,

$$\begin{aligned}
& \int_{Q_t} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) (p-2) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-4}{2}} (\nabla u^\varepsilon \cdot \nabla v)^2 s'_\delta(v) \\
& \quad + \int_{Q_t} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-2}{2}} |\nabla v|^2 s'_\delta(v) \\
& = \int_{Q_t} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-4}{2}} \\
& \quad \times s'_\delta(v) \left[(p-2) (\nabla u^\varepsilon \cdot \nabla v)^2 + \left(|\nabla u^\varepsilon|^2 + \varepsilon \right) |\nabla v|^2 \right].
\end{aligned}$$

For $p \geq 2$, obviously, the second member of this equation is nonnegative. For $1 < p < 2$, using Cauchy-Schwartz inequality, we have

$$\begin{aligned}
& \int_{Q_t} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-4}{2}} s'_\delta(v) \left[(p-2) (\nabla u^\varepsilon \cdot \nabla v)^2 + \left(|\nabla u^\varepsilon|^2 + \varepsilon \right) |\nabla v|^2 \right] \\
& \geq \int_{Q_t} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{\frac{p-4}{2}} s'_\delta(v) |\nabla u^\varepsilon|^2 |\nabla v|^2 (1 - (2-p)) \geq 0.
\end{aligned}$$

So, since $\int_{Q_t} f_t^\varepsilon s_\delta(v) \leq \int_{Q_t} |f_t^\varepsilon|$, neglecting the nonnegative terms of (14) and letting $\delta \rightarrow 0$ in (14), we obtain

$$(15) \quad \int_{\Omega} |v(t)| - \int_{\Omega} |v(0)| \leq \int_{Q_t} |f_t^\varepsilon| ,$$

and so

$$\|u_t^\varepsilon\|_{L^1(\Omega)} \leq \|\Delta_p h^\varepsilon + \varepsilon \Delta h^\varepsilon\|_{L^1(\Omega)} + \|f_t^\varepsilon\|_{L^1(Q_T)} \leq C ,$$

where $C > 0$ may be chosen independent of ε . \square

LEMMA 3. *Suppose that assumptions (1), (2) and (3) are satisfied and u^ε is the solution to the problem (7). Then there is a constant $C > 0$ independent of $\varepsilon \in]0, 1[$ such that*

$$(16) \quad \left\| k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \right\|_{L^1(Q_T)} \leq C .$$

PROOF. We consider the first equation of the problem (7), multiply it by u^ε and integrate over Q_T . Then

$$\int_{Q_T} u_t^\varepsilon u^\varepsilon + \int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \left[\left(|\nabla u^\varepsilon|^2 + \varepsilon \right)^{p-2} |\nabla u^\varepsilon|^2 + \varepsilon |\nabla u^\varepsilon|^2 \right] = \int_{Q_T} f^\varepsilon u^\varepsilon .$$

Since f^ε and u^ε are uniformly bounded, independently of ε , in $L^\infty(Q_T)$,

$$\left| \int_{Q_T} f^\varepsilon u^\varepsilon \right| \leq C, \quad C \text{ independent of } \varepsilon$$

and, since u_t^ε is uniformly bounded in $L^\infty(0, T; L^1(\Omega))$, by Lemma 2,

$$\left| \int_{Q_T} u_t^\varepsilon u^\varepsilon \right| \leq C, \quad C \text{ independent of } \varepsilon$$

and so, it follows that

$$\int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) |\nabla u^\varepsilon|^p \leq C', \quad C' \text{ constant independent of } \varepsilon.$$

But, on the other hand,

$$\begin{aligned} \int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) &= \int_{\{|\nabla u^\varepsilon|^2 \leq F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \\ &\quad + \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \\ &\leq |Q_T| e + \int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \frac{|\nabla u^\varepsilon|^p}{m^p} \end{aligned}$$

because in $\{|\nabla u^\varepsilon|^2 \leq F_\varepsilon^2(u^\varepsilon) + \varepsilon\}$ we have $k_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)) \leq e$ and in $\{|\nabla u^\varepsilon|^2 \geq F_\varepsilon^2(u^\varepsilon) + \varepsilon\}$ we have $\frac{|\nabla u^\varepsilon|}{m} \geq 1$.

So, we conclude

$$\begin{aligned} \int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) &\leq |Q_T| e + \frac{1}{m^p} \int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) |\nabla u^\varepsilon|^p \\ &\leq |Q_T| e + \frac{C'}{m^p}, \quad \text{independently of } \varepsilon. \quad \square \end{aligned}$$

LEMMA 4. *Suppose that the assumptions (1), (2) and (3) are satisfied and u^ε is the solution of the problem (7). Then, for any $q \in [1, +\infty[$ there exists a constant $C_q > 0$ independent of $\varepsilon \in]0, 1[$ such that*

$$\|\nabla u^\varepsilon\|_{L^q(Q_T)} \leq C_q.$$

PROOF. We know, from (16), that there exists a constant C , independent of ε , such that, for any $\varepsilon \in]0, 1[$,

$$\int_{Q_T} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) \leq C.$$

So, we have

$$C \geq \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} k_\varepsilon \left(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right) = \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} e^{\frac{|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)}{\varepsilon}} .$$

Recalling that

$$\forall s \in \mathbb{R}^+ \quad \forall j \in \mathbb{N} \quad e^s \geq \frac{s^j}{j!}$$

we obtain, for each $j \in \mathbb{N}$,

$$(17) \quad \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} \left[|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right]^j \leq j! \varepsilon^j \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} e^{\frac{|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon)}{\varepsilon}} \\ \leq j! \varepsilon^j C .$$

Given $q \in \mathbb{N}$, we have

$$(18) \quad \int_{Q_T} |\nabla u^\varepsilon|^{2q} = \int_{\{|\nabla u^\varepsilon|^2 \leq F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} |\nabla u^\varepsilon|^{2q} + \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} |\nabla u^\varepsilon|^{2q} ,$$

Since there exists a constant $M > 0$, not depending on ε , such that $\|u^\varepsilon\|_{L^\infty(Q_T)} \leq M$, we can estimate the first integral in the second member of (18) as follows:

$$\int_{\{|\nabla u^\varepsilon|^2 \leq F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} |\nabla u^\varepsilon|^{2q} \leq \int_{Q_T} \left[F_\varepsilon^2(u^\varepsilon) + \varepsilon \right]^q \leq \max_{s \in [-M, M]} \left\{ F^2(s) + 1 \right\}^q |Q_T| .$$

On the other hand the second integral of the second member of (18) satisfies

$$\int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} |\nabla u^\varepsilon|^{2q} \\ = \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} \left[|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) + F_\varepsilon^2(u^\varepsilon) \right]^q \\ \leq \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} \sum_{j=0}^q \binom{q}{j} \max_{x \in [-M, M]} \left[F_\varepsilon^2(x) \right]^{q-j} \left[|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right]^j \\ \leq \sum_{j=0}^q D_j \int_{\{|\nabla u^\varepsilon|^2 > F_\varepsilon^2(u^\varepsilon) + \varepsilon\}} \left[|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) \right]^j ,$$

where D_j is a constant independent of ε .

So, using (17), we easily conclude that

$$\|\nabla u^\varepsilon\|_{L^q(Q_T)} \leq C_q , \quad C_q \text{ depending on } q, \text{ but not on } \varepsilon . \quad \square$$

4. – The limiting process and further remarks

In this section we will let $\varepsilon \rightarrow 0$ and prove that if we define

$$(19) \quad \mathbb{K}_{u^\varepsilon(t)} = \left\{ v \in W_0^{1,p}(\Omega) : |\nabla v| \leq F(u^\varepsilon(t)) \quad \text{a.e. in } \Omega \right\}$$

we have, at least for a subsequence $\varepsilon \rightarrow 0$, for any given $v(t) \in \mathbb{K}_{u(t)}$ for every $t \in I$, the convergence of a sequence of functions v^ε , with $v^\varepsilon(t) \in \mathbb{K}_{u^\varepsilon(t)}$, to the function v , when $\varepsilon \rightarrow 0$.

Recalling Lemma 4, let u be the weak limit in $L^q(0, T; W_0^{1,q}(\Omega))$, $\forall q < \infty$, of a subsequence of $\{u^\varepsilon\}_{\varepsilon \in]0,1[}$ as $\varepsilon \rightarrow 0$, i.e.,

$$(20) \quad u^\varepsilon \rightharpoonup_\varepsilon u \quad \text{in } L^q(0, T; W_0^{1,q}(\Omega)) .$$

Then

LEMMA 5. *Suppose that the assumptions (1), (2) and (3) are satisfied and that u^ε is the solution of the problem (7). Let $\mathbb{K}_{u^\varepsilon(t)}$ be the convex set defined in (19) and u the function defined in (20), weak limit in $L^q(0, T; W_0^{1,q}(\Omega))$ of the subsequence $\{u^\varepsilon\}_\varepsilon$.*

Then for any $v \in L^\infty(0, T; W_0^{1,p}(\Omega))$, $1 \leq p \leq \infty$ such that $v(t) \in \mathbb{K}_{u(t)}$ for a.e. $t \in [0, T]$, there exists a $v^\varepsilon \in L^\infty(0, T; W_0^{1,p}(\Omega))$ with $v^\varepsilon(t) \in \mathbb{K}_{u^\varepsilon(t)}$ for a.e. $t \in [0, T]$ such that

$$v^\varepsilon \xrightarrow{\varepsilon} v \quad \text{in } L^\infty(0, T; W_0^{1,p}(\Omega)), \quad 1 \leq p \leq \infty.$$

PROOF. Since there are constants C_1 and C_q such that

$$\|u_t^\varepsilon\|_{L^\infty(0,T;L^1(\Omega))} \leq C_1 \quad \text{and} \quad \|u^\varepsilon\|_{L^q(0,T;W_0^{1,q}(\Omega))} \leq C_q$$

(for $q > N$), by a well known compactness theorem ([13], page 84), $\{u^\varepsilon\}_\varepsilon$ is relatively compact in $C([0, T]; C(\overline{\Omega}))$ and so, at least for another subsequence of the subsequence in (20), still denoted by $\varepsilon \rightarrow 0$, we have

$$(21) \quad u^\varepsilon \xrightarrow{\varepsilon} u \quad \text{uniformly in } C^0(\overline{Q}_T)$$

and therefore also

$$F(u^\varepsilon) \xrightarrow{\varepsilon} F(u) \quad \text{uniformly in } C^0(\overline{Q}_T) .$$

Let $\alpha_\varepsilon(t) = \|F(u^\varepsilon(t)) - F(u(t))\|_{L^\infty(\Omega)}$ and notice that

$$\alpha_\varepsilon(t) \xrightarrow{\varepsilon} 0 \quad \text{uniformly in } t \in [0, T] .$$

Define $\psi_\varepsilon(t) = 1 + \frac{\alpha_\varepsilon(t)}{m}$ and, given $v \in L^\infty(0, T; W_0^{1,p}(\Omega))$ such that $v(t) \in \mathbb{K}_{u(t)}$ for a.e. $t \in [0, T]$, define $v^\varepsilon(t) = \frac{1}{\psi_\varepsilon(t)} v(t) \in L^\infty(0, T; W_0^{1,p}(\Omega))$. Then,

$$|\nabla v^\varepsilon(t)| = \frac{1}{\psi_\varepsilon(t)} |\nabla v(t)| \leq \frac{1}{\psi_\varepsilon(t)} F(u(t)) \leq F(u^\varepsilon(t))$$

because

$$\begin{aligned} \psi_\varepsilon(t) &= 1 + \frac{\alpha_\varepsilon(t)}{m} \geq 1 + \frac{\alpha_\varepsilon(t)}{F(u^\varepsilon(t))} \\ &\geq \frac{F(u^\varepsilon(t)) + F(u(t)) - F(u^\varepsilon(t))}{F(u^\varepsilon(t))} = \frac{F(u(t))}{F(u^\varepsilon(t))}. \end{aligned}$$

So, we have $v^\varepsilon(t) \in \mathbb{K}_{u^\varepsilon(t)}$ for a.e. $t \in [0, T]$ and $v^\varepsilon \xrightarrow{\varepsilon} v$ in $L^\infty(0, T; W_0^{1,p}(\Omega))$ since

$$\|v^\varepsilon(t) - v(t)\|_{W_0^{1,p}(\Omega)} = \left| \frac{1}{\psi_\varepsilon(t)} - 1 \right| \|v\|_{L^\infty(0,T;W_0^{1,p}(\Omega))} \xrightarrow{\varepsilon} 0$$

uniformly in $t \in [0, T]$. □

PROOF OF THEOREM 1. Recalling (20) and (21), for some subsequence $\varepsilon \rightarrow 0$ we have

$$\begin{aligned} u^\varepsilon &\xrightarrow{\varepsilon} u \quad \text{uniformly in } C^0(\overline{Q}_T), \\ u^\varepsilon &\rightharpoonup_\varepsilon u \quad \text{weakly in } L^q(0, T; W^{1,q}(\Omega)), \quad \forall q < \infty, \end{aligned}$$

and, besides that, this subsequence may be chosen in such a way that

$$u_t^\varepsilon \rightharpoonup_\varepsilon u_t \quad \text{in } L^\infty(0, T; M(\Omega)) \text{ weak-}^* .$$

Now, given $w \in \mathbb{K}_{u^\varepsilon(t)}$ for a.e. $t \in [0, T]$, we have

$$\begin{aligned} &\int_\Omega (|\nabla w|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla w \cdot \nabla(w - u^\varepsilon(t)) \\ &= \int_\Omega k_\varepsilon (|\nabla w|^2 - F^2(u^\varepsilon(t))) (|\nabla w|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla w \cdot \nabla(w - u^\varepsilon(t)) \\ &\geq \int_\Omega k_\varepsilon (|\nabla u^\varepsilon(t)|^2 - F^2(u^\varepsilon(t))) (|\nabla u^\varepsilon(t)|^2 + \varepsilon)^{\frac{p-2}{2}} \nabla u^\varepsilon(t) \cdot \nabla(w - u^\varepsilon(t)) \end{aligned}$$

since, by monotonicity,

$$\begin{aligned} & \int_{\Omega} k_{\varepsilon} \left(|\nabla w|^2 - F^2(u^{\varepsilon}(t)) \right) \left(|\nabla w|^2 + \varepsilon \right)^{\frac{p-2}{2}} \nabla w \cdot \nabla (w - u^{\varepsilon}(t)) \\ & - \int_{\Omega} k_{\varepsilon} \left(|\nabla u^{\varepsilon}(t)|^2 - F^2(u^{\varepsilon}(t)) \right) \left(|\nabla u^{\varepsilon}(t)|^2 + \varepsilon \right)^{\frac{p-2}{2}} \nabla u^{\varepsilon}(t) \cdot \nabla (w - u^{\varepsilon}(t)) \geq 0. \end{aligned}$$

Given $v \in L^{\infty}(0, T; W_0^{1,p}(\Omega))$ such that $v(t) \in \mathbb{K}_{u(t)}$, for a.e. $t \in [0, T]$, we consider $v^{\varepsilon} \in L^{\infty}(0, T; W_0^{1,p}(\Omega))$ as in Lemma 5. Since $v^{\varepsilon}(t) \in \mathbb{K}_{u^{\varepsilon}(t)}$, multiplying the equation (7) by $v^{\varepsilon}(t) - u^{\varepsilon}(t)$ and integrating over $\Omega \times]s, t[$, $0 < s < t < T$, we have

$$\int_s^t \int_{\Omega} u_t^{\varepsilon} (v^{\varepsilon} - u^{\varepsilon}) + \int_s^t \int_{\Omega} \left(|\nabla v^{\varepsilon}|^2 + \varepsilon \right)^{\frac{p-2}{2}} \nabla v^{\varepsilon} \cdot \nabla (v^{\varepsilon} - u^{\varepsilon}) \geq \int_s^t \int_{\Omega} f^{\varepsilon} (v^{\varepsilon} - u^{\varepsilon})$$

and letting $\varepsilon \rightarrow 0$, since $v^{\varepsilon} \rightarrow v$ in $L^{\infty}(0, T; W_0^{1,p}(\Omega))$ strongly, we obtain

$$\int_s^t \int_{\Omega} u_t (v - u) + \int_s^t \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla (v - u) \geq \int_s^t \int_{\Omega} f (v - u).$$

Since s and t are arbitrary, we may conclude

$$\begin{aligned} (22) \quad & \int_{\Omega} u_t(t) (v - u(t)) + \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla (v - u(t)) \\ & \geq \int_{\Omega} f(t) (v - u(t)), \quad \forall v \in \mathbb{K}_{u(t)} \text{ for a.e. } t \in I. \end{aligned}$$

It still remains to prove that $|\nabla u(t)| \leq F(u(t))$ a.e. in Ω , for a.e. $t \in I$. If we prove this, a variant of Minty's lemma will show that (22) implies (5).

To prove that $|\nabla u(t)| \leq F(u(t))$ a.e. in Ω , for a.e. $t \in I$ is equivalent to prove that the set $\{(x, t) \in Q_T : |\nabla u(x, t)| > F(u(x, t))\}$ has measure zero.

We recall from Lemma 3 that

$$\exists C > 0 \quad \forall \varepsilon \in]0, 1[\quad \int_{Q_T} k_{\varepsilon} \left(|\nabla u^{\varepsilon}|^2 - F_{\varepsilon}^2(u^{\varepsilon}) \right) \leq C.$$

Given $\varepsilon \in]0, 1[$ and $M \in \mathbb{R}^+$, $M \geq \varepsilon$, we define

$$A_{M,\varepsilon} = \left\{ (x, t) \in Q_T : |\nabla u^{\varepsilon}(x, t)|^2 - F_{\varepsilon}^2(u^{\varepsilon}(x, t)) \geq M \right\}$$

and then, since $k_{\varepsilon}(|\nabla u^{\varepsilon}|^2 - F^2(u^{\varepsilon})) \geq e^{\frac{M}{\varepsilon}}$ in $A_{M,\varepsilon}$, we have

$$\forall \varepsilon \in]0, 1[\quad \forall M \in [\varepsilon, +\infty[\quad |A_{M,\varepsilon}| e^{\frac{M}{\varepsilon}} \leq \int_{A_{M,\varepsilon}} k_{\varepsilon} \left(|\nabla u^{\varepsilon}|^2 - F_{\varepsilon}^2(u^{\varepsilon}) \right) \leq C,$$

where $|A|$ denotes the Lebesgue measure of the set A in \mathbb{R}^{N+1} .

So we have

$$|A_{M,\varepsilon}| \leq C e^{-\frac{M}{\varepsilon}}$$

and, choosing $M = \sqrt{\varepsilon}$, we get

$$|A_{\sqrt{\varepsilon},\varepsilon}| \leq C e^{-\frac{1}{\sqrt{\varepsilon}}}$$

and

$$\begin{aligned} \int_{Q_T} (|\nabla u|^2 - F_\varepsilon^2(u))^+ &\leq \liminf_{\varepsilon \rightarrow 0} \int_{Q_T} (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) - \sqrt{\varepsilon})^+ \\ &= \liminf_{\varepsilon \rightarrow 0} \int_{A_{\sqrt{\varepsilon},\varepsilon}} (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) - \sqrt{\varepsilon}) \\ &\leq \lim_{\varepsilon \rightarrow 0} D |A_{\sqrt{\varepsilon},\varepsilon}|^{\frac{1}{2}} = 0, \end{aligned}$$

being D a constant that bounds from above $\left[\int_{Q_T} (|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon) - \sqrt{\varepsilon})^2 \right]^{1/2}$, independently of ε .

So, $(|\nabla u|^2 - F^2(u))^+ = 0$ a.e. in Q_T or, equivalently,

$$|\nabla u| \leq F(u) \quad \text{a.e. in } Q_T.$$

Substitute now in (22), for a.e. $t \in I$, v by $u(t) + \theta(w - u(t))$, where w belongs to $\mathbb{K}_{u(t)}$, $\theta \in]0, 1]$ and divide both members by θ . We obtain

$$\begin{aligned} \int_{\Omega} u_t(t)(w - u(t)) + \int_{\Omega} \left| \nabla (u(t) + \theta(w - u(t))) \right|^{p-2} \nabla (u(t) + \theta(w - u(t))) \cdot \nabla (w - u(t)) \\ \geq \int_{\Omega} f(t)(w - u(t)), \quad \forall w \in \mathbb{K}_{u(t)} \quad \text{for a.e. } t \in I. \end{aligned}$$

Letting $\theta \rightarrow 0$ we conclude that u is a solution to the quasivariational inequality (5). \square

5. – Asymptotic behaviour in time

Consider $T = +\infty$. We first show that there exists a global solution of the quasivariational inequality (5) when $T = +\infty$ and then, in some sense, there exists a limit function, when $t \rightarrow +\infty$, solution of a stationary quasivariational inequality.

LEMMA 6. *Suppose that (9) is satisfied. Then the problem (5) has a solution u such that*

$$u \in L^\infty(0, +\infty; W_0^{1,\infty}(\Omega)) \cap C^0(\mathbb{R}_0^+ \times \overline{\Omega}), \quad u_t \in L^\infty(0, +\infty; M(\Omega)),$$

being u the weak limit in $L^q(0, +\infty; W^{1,q}(\Omega))$ (for any $q < \infty$) of a subsequence $\{u^\varepsilon\}_\varepsilon$ of solutions of the family of approximated problems (7), $u_t^\varepsilon \rightharpoonup u_t$ in $L^\infty(0, +\infty; M(\Omega))$ weak- and, for all $T > 0$, there exists a subsequence of $\varepsilon \rightarrow 0$, called $\varepsilon_T \rightarrow 0$, such that $u^{\varepsilon_T} \xrightarrow{\varepsilon_T} u$ in $C^0(\overline{Q}_T)$. In addition, we have the estimate*

$$(23) \quad t \int_\Omega |u_t(t)| \leq \int_0^t \left(\int_\Omega |f_t(\tau)| \right) (1 + \tau) d\tau, \quad \text{for } 0 < t < \infty.$$

PROOF. The assumption (9) implies that the estimates of Lemmas 1 and 2 may be obtained when $T = +\infty$. The L^∞ boundedness of $\{u^\varepsilon\}_\varepsilon$, independently of ε and T and its boundedness in $L^\infty(0, +\infty; W^{1,q}(\Omega))$ can also be obtained easily from the boundedness of $k_\varepsilon(|\nabla u^\varepsilon|^2 - F_\varepsilon^2(u^\varepsilon))$ in $L^\infty(0, +\infty; L^1(\Omega))$, which can be obtained as in Lemma 3, integrating only on Ω , for a.e. $t \in [0, T]$. So, we can pass to the limit in a subsequence $\varepsilon \rightarrow 0$ and get a function $u \in L^\infty(0, +\infty; W^{1,\infty}(\Omega))$, $u_t \in L^\infty(0, +\infty; M(\Omega))$ which is the weak-limit of u^ε in these spaces.

The uniform convergence of a subsequence of u^ε is obtained only in $C^0(\overline{Q}_T)$, for each T , which is enough to pass to the limit in the approximated problem and to obtain that u is a solution of the quasivariational inequality for almost every $t \in \mathbb{R}^+$

In order to show (23) we multiply the approximated equation (7) by $ts_\delta(u_t^\varepsilon)$ and, arguing as in the proof of Lemma 2, we obtain, after letting $\delta \rightarrow 0$,

$$\begin{aligned} \int_\Omega (t|v(t)| - |v(0)| + |v(t)|) &= \int_0^t \left(\int_\Omega \frac{d|v|}{d\tau} \right) \tau d\tau \\ &\leq \int_0^t \left(\int_\Omega |f_t^\varepsilon| \right) \tau d\tau, \quad \text{for any } 0 < t < \infty. \end{aligned}$$

Using (15) we obtain

$$t \int_\Omega |v(t)| \leq \int_0^t \left(\int_\Omega |f_t^\varepsilon| \right) d\tau + \int_0^t \left(\int_\Omega |f_t^\varepsilon| \right) \tau d\tau,$$

and letting $\varepsilon \rightarrow 0$ we conclude (23). □

PROOF OF THEOREM 2. Let u be a solution of the problem (5), with $T = +\infty$.

Since

$$\exists M > 0 \quad \|u\|_{L^\infty(\mathbb{R}^+ \times \Omega)} \leq M$$

and

$$|\nabla u(x, t)| \leq F(u(x, t)) \leq \max_{|s| \leq M} \{F(s)\} \quad \text{for a.e. } (x, t) \in \mathbb{R}^+ \times \Omega$$

we have

$$\exists C > 0 \quad \|u\|_{L^\infty(\mathbb{R}^+; W^{1,\infty}(\Omega))} \leq C$$

and so, for a subsequence $(t_n)_n$, $t_n \xrightarrow[n]{\nearrow} +\infty$ and some function $u^\infty \in W_0^{1,\infty}(\Omega)$,

we have

$$\begin{aligned} u(t_n) &\xrightarrow[n]{\rightharpoonup} u^\infty \quad \text{in } W^{1,\infty}(\Omega) \text{ weak-}^* , \\ u(t_n) &\xrightarrow[n]{\rightarrow} u^\infty \quad \text{in } C^0(\overline{\Omega}) . \end{aligned}$$

Since for every measurable subset $\omega \subset \Omega$ we have $(1 < q < \infty)$

$$\int_\omega |\nabla u^\infty|^q \leq \liminf_n \int_\omega |\nabla u(t_n)|^q \leq \lim_n \int_\omega F^q(u(t_n)) = \int_\omega F^q(u^\infty),$$

we find that

$$|\nabla u^\infty| \leq F(u^\infty) \quad \text{a.e. in } \Omega,$$

and, therefore, we have

$$(24) \quad u^\infty \in \mathbb{K}_{u^\infty} .$$

It remains to show that u^∞ satisfies the inequality in (8).

By monotonicity, we rewrite (5) for a.e. $t \in \mathbb{R}^+$ in the form

$$(25) \quad \int_\Omega u_t(v-u(t)) + \int_\Omega |\nabla v|^{p-2} \nabla v \cdot \nabla(v-u(t)) \geq \int_\Omega f(t)(v-u(t)), \quad \forall v \in \mathbb{K}_{u(t)}.$$

We consider a given function φ satisfying

$$\varphi \in C^1(\mathbb{R}_0^+), \quad 0 \leq \varphi(t) \leq 1, \quad \varphi(0) = 1, \quad \varphi(t) \xrightarrow[t \rightarrow +\infty]{} 0, \quad \varphi_t(t) \xrightarrow[t \rightarrow +\infty]{} 0 ,$$

and we define, for $0 < \delta < 1$,

$$v^\delta(x, t) = \varphi(t) u(x, t) + (1 - \delta)(1 - \varphi(t)) w^\infty$$

with w^∞ arbitrarily given in \mathbb{K}_{u^∞} .

We have

$$|\nabla v^\delta(t)| \leq \varphi(t) F(u(t)) + (1 - \delta)(1 - \varphi(t)) F(u^\infty) ,$$

and, since $F(u(t_n)) \xrightarrow[n]{\rightarrow} F(u^\infty)$ uniformly, there exists $n_0 \in \mathbb{N}$ such that

$$\frac{F(u^\infty)}{F(u(t_n))} \leq \frac{1}{1 - \delta} \quad \text{for } n \geq n_0$$

and so,

$$|\nabla v^\delta(t_n)| \leq \varphi(t_n) F(u(t_n)) + (1 - \varphi(t_n)) F(u(t_n)) = F(u(t_n)) \quad \text{for } n \geq n_0.$$

Substituting in (25), at time t_n , v by $v^\delta(t_n) \in \mathbb{K}_{u(t_n)}$ we obtain

$$\begin{aligned} \int_{\Omega} u_t(t_n) (v^\delta(t_n) - u(t_n)) + \int_{\Omega} |\nabla v^\delta(t_n)|^{p-2} \nabla v^\delta(t_n) \cdot \nabla (v^\delta(t_n) - u(t_n)) \\ \geq \int_{\Omega} f(t_n) (v^\delta(t_n) - u(t_n)) \end{aligned}$$

and, letting $t_n \rightarrow +\infty$, we obtain

$$\int_{\Omega} |(1-\delta) \nabla w^\infty|^{p-2} (1-\delta) \nabla w^\infty \cdot \nabla ((1-\delta) w^\infty - u^\infty) \geq \int_{\Omega} f^\infty ((1-\delta) w^\infty - u^\infty),$$

since $f(t_n) \rightharpoonup_n f^\infty$ in $L^1(\Omega)$ -weak and $\int_{\Omega} |u_t(t_n)| \rightarrow_n 0$ by (23) and the assumptions (9) and (10). Letting now $\delta \rightarrow 0$, we get

$$\int_{\Omega} |\nabla w^\infty|^{p-2} \nabla w^\infty \cdot \nabla (w^\infty - u^\infty) \geq \int_{\Omega} f^\infty (w^\infty - u^\infty), \quad \forall w^\infty \in \mathbb{K}_{u^\infty}$$

and, applying Minty's lemma, we see that u^∞ is a solution to the problem (8), since we have already proved that $u^\infty \in \mathbb{K}_{u^\infty}$. \square

REFERENCES

- [1] C. BAIOCCHI – A. CAPELO, “Variational and Quasivariational Inequalities. Applications to Free Boundary Problems”, J. Wiley & Sons, Chischester-New York, 1984.
- [2] A. BENSOUSSAN – J. L. LIONS, “Contrôle impulsional et inéquations quasivariationelles”, Dunod, Paris, 1982.
- [3] E. H. BRANDT, *Electric field in superconductors with regular cross-sections*, Phys. Rev. B **52** (1995), 15442-15457.
- [4] E. DiBENEDETTO, “Degenerate Parabolic Equations”, Springer-Verlag, New York, 1993.
- [5] C. GERHARDT, *On the existence and uniqueness of a warpening function in the elastic-plastic torsion of a cylindrical bar with multiply connected cross-section*, A. Dold – B. Eckmann – P. Germain – B. Nayroles (eds.), Lecture Notes in Math. 503, Springer, Berlin, 1976.
- [6] Y. B. KIM – C. F. HEMPSTEAD – A. R. STRNAD, *Critical persistent currents in hard superconductors*, Phys. Rev. Lett. **9** (1962), 306-309.

- [7] M. M. KUNZE – M. D. P. MONTEIRO MARQUES, *A note on Lipschitz continuous solutions of a parabolic quasi-variational inequality*, In “Nonlinear Evolutionary Equations and their Applications”, T. T. Li – L. W. Lin – J. F. Rodrigues (eds.), World Scientific, Singapore, 1999, pp. 109-115.
- [8] M. M. KUNZE – J. F. RODRIGUES, *An elliptic quasi-variational inequality with gradient constants and some of its applications*, Math. Methods. Appl. Sci., to appear.
- [9] O. A. LADYZENSKAJA – V. A. SOLONNIKOV – N. N. URAL’CEVA, “Linear and quasilinear equations of parabolic type”, Translations of Mathematical Monographs 23, AMS, 1968.
- [10] L. PRIGHOZIN, *Variational model of sandpile growth*, European J. Appl. Math. **7** (1996), 225-235.
- [11] L. PRIGHOZIN, *On the Bean critical state model in superconductivity*, European J. Appl. Math. **7** (1996), 237-247.
- [12] L. SANTOS, *A diffusion problem with gradient constraint and evolutive Dirichlet condition*, Portugaliae Mathematica **48** (1991), 441-468.
- [13] J. SIMON, *Compact sets in the space $L^p(0, T; B)$* , Ann. Mat. Pura Appl. **146** (1987), 65-96.

CMAF/Univ. de Lisboa
Av. Prof. Gama Pinto, 2
1649-003 Lisboa, Portugal

Centro de Matemática
University of Minho
Campus de Gualtar
4700-030 Braga, Portugal