

Combinatorial realization of the Thom-Smale complex via discrete Morse theory

ÉTIENNE GALLAIS

Abstract. In the case of smooth manifolds, we use Forman's discrete Morse theory to realize combinatorially any Thom-Smale complex coming from a smooth Morse function by a pair triangulation-discrete Morse function. As an application, we prove that any class of homologous vector fields on a smooth oriented closed 3-manifold can be realized by a perfect matching on the Hasse diagram of a triangulation of the manifold.

Mathematics Subject Classification (2010): 57R25 (primary); 57R05 (secondary).

1. Introduction

R. Forman defines a combinatorial analog of smooth Morse theory in [4–6] for simplicial complexes and more generally for CW-complexes. Discrete Morse theory has many applications (computer graphics [10], graph theory [3]). An important problem is the research of optimal discrete Morse functions in the sense that they have the minimal number of critical cells ([8, 9, 18] for the minimality of hyperplane arrangements).

Thanks to a combinatorial Morse vector field V , Forman constructs a combinatorial Thom-Smale complex (C^V, ∂^V) whose homology is the simplicial homology of the simplicial complex. The differential is defined by counting algebraically V -paths between critical cells. Nevertheless, the proof of $\partial^V \circ \partial^V = 0$ is an indirect proof (see [5]). We give two proofs of $\partial^V \circ \partial^V = 0$, one which focusses on the geometry and another one which focusses on the algebraic point of view (compare with [2] and [19]). In fact, the algebraic proof gives also the property that the combinatorial Thom-Smale complex is a chain complex homotopy equivalent to the simplicial chain complex. After that, we investigate one step forward the relation between the smooth Morse theory and the discrete Morse theory. We prove that any Thom-Smale complex has a combinatorial realization. We use this to prove that any class of homologous vector fields on a closed oriented 3-manifold can be realized by a perfect matching on a triangulation of this manifold.

This article is organised as follows. In Section 2, we recall the discrete Morse theory from the viewpoint of combinatorial Morse vector field. Section 2.2.3 is devoted to the proofs of $\partial^V \circ \partial^V = 0$ and that the Thom-Smale complex is a chain complex homotopy equivalent to the simplicial chain complex. In Section 3, we prove that any combinatorial Thom-Smale complex is realizable as a combinatorial Thom-Smale complex. In Section 4, we obtain as a corollary the existence of triangulations with perfect matchings on their Hasse diagram and prove that any class of homologous vector fields on a closed oriented 3-manifold can be realized by such complete matchings.

2. Discrete Morse theory

2.1. Combinatorial Morse vector field

First of all, instead of considering discrete Morse functions on a simplicial complex, we will only consider combinatorial Morse vector fields. In fact, working with discrete Morse functions or combinatorial vector fields is exactly the same [5, Theorem 9.3].

In the following, X is a finite simplicial complex and K is the set of cells of X . A cell $\sigma \in K$ of dimension k is denoted $\sigma^{(k)}$. Let $<$ be the partial order on K given by $\sigma < \tau$ iff $\sigma \subset \bar{\tau}$. Given a simplicial complex, one associates its *Hasse diagram*: the set of vertices is the set of cells K , an edge joins two cells σ and τ if $\sigma < \tau$ and $\dim(\sigma) + 1 = \dim(\tau)$.

Definition 2.1. A *combinatorial vector field* V on X is an oriented matching on the associated Hasse diagram of X that is a set of edges \mathcal{M} such that

- (1) any two distinct edges of \mathcal{M} do not share any common vertex,
- (2) every edge belonging to \mathcal{M} is oriented toward the top dimensional cell.

A cell which does not belong to any edge of the matching is said to be critical.

Remark 2.2. The original definition of a combinatorial vector field is the following one: given a matching on the Hasse diagram define

$$V : K \rightarrow K \cup \{0\}$$

$$\sigma \mapsto V(\sigma) = \begin{cases} \tau & \text{iff } (\sigma, \tau) \text{ is an edge of the matching and } \sigma < \tau, \\ 0 & \text{otherwise.} \end{cases}$$

We will use both these points of view in the following.

A V -path of dimension k is a sequence of cells $\gamma : \sigma_0, \sigma_1, \dots, \sigma_r$ of dimension k such that

- (1) $\sigma_i \neq \sigma_{i+1}$ for all $i \in \{0, \dots, r-1\}$,
- (2) for every $i \in \{0, \dots, r-1\}$, $\sigma_{i+1} < V(\sigma_i)$.

A V -path γ is said to be *closed* if $\sigma_0 = \sigma_r$, and *non-stationary* if $r > 0$.

Definition 2.3. A combinatorial vector field V which has no non-stationary closed path is called a *combinatorial Morse vector field*. In this case, the corresponding matching is called a *Morse matching*.

The terminology *Morse matching* first appeared in [2].

Remark 2.4. Let V be a combinatorial (respectively combinatorial Morse) vector field. If we remove an edge from the underlying matching, it remains a combinatorial (respectively combinatorial Morse) vector field (there are two extra critical cells).

2.2. The combinatorial Thom-Smale complex

2.2.1. Definition of the combinatorial Thom-Smale complex

The following data are necessary to define the combinatorial Thom-Smale complex (see [5]). First, let X be a finite simplicial complex, K its set of cells and V a combinatorial Morse vector field. Suppose that every cell $\sigma \in K$ is oriented.

Let $\gamma : \sigma_0, \sigma_1, \dots, \sigma_r$ be a V -path. Then the *multiplicity* of γ is given by the formula

$$m(\gamma) = \prod_{i=0}^{r-1} - \langle \partial V(\sigma_i), \sigma_i \rangle \langle \partial V(\sigma_i), \sigma_{i+1} \rangle \in \{\pm 1\}$$

where for every cell $\sigma, \tau, \langle \sigma, \tau \rangle \in \{-1, 0, 1\}$ is the incidence number between the cells σ and τ (see [11]) and ∂ is the boundary map when we consider X as a CW-complex. In fact, one can think of the multiplicity as checking if the orientation of the first cell σ_0 moved along γ coincides or not with the orientation of the last cell σ_r .

Let $\Gamma(\sigma, \sigma')$ be the set of V -paths starting at σ and ending at σ' and $\text{Crit}_k(V)$ be the set of critical cells of dimension k .

Definition 2.5. The *combinatorial Thom-Smale complex* associated with (X, V) is (C_*^V, ∂^V) where:

- (1) $C_k^V = \bigoplus_{\sigma \in \text{Crit}_k(V)} \mathbb{Z} \cdot \sigma$,
- (2) if $\tau \in \text{Crit}_{k+1}(V)$ then

$$\partial^V \tau = \sum_{\sigma \in \text{Crit}_k(V)} n(\tau, \sigma) \cdot \sigma$$

where

$$n(\tau, \sigma) = \sum_{\tilde{\sigma} < \tau} \langle \partial \tau, \tilde{\sigma} \rangle \sum_{\gamma \in \Gamma(\tilde{\sigma}, \sigma)} m(\gamma).$$

Thus, this complex is exactly in the same spirit as the Thom-Smale complex for smooth Morse functions (see Section 3): it is generated by critical cells and the differential is given by counting algebraically V -paths.

Theorem 2.6 (Forman [5]). $\partial^V \circ \partial^V = 0$.

Theorem 2.7 (Forman [5]). (C_*^V, ∂^V) is homotopy equivalent to the simplicial chain complex. In particular, its homology is equal to the simplicial homology.

We will give a direct proof of both of these theorems. The proof of Theorem 2.6 is done by looking at V -paths and understanding their contribution to $\partial^V \circ \partial^V$. Then, we prove Theorem 2.7 (which gives another proof of Theorem 2.6) using Gaussian elimination (this idea first appears in [2], see also [19]).

2.2.2. Proof of Theorem 2.6

Let X be a simplicial complex, K be the set of its cells and K_n the set of cells of dimension n . The proof is by induction on the number on edges belonging to the Morse matching.

Initialization: *matching with no edge.* In this case, every cell is critical and the combinatorial Thom-Smale complex coincides with the well-known simplicial chain complex. Therefore, $\partial^V \circ \partial^V = 0$.

Heredity: *suppose the property is true for every matching with at most k edges defining a combinatorial Morse vector field.* Let V be a combinatorial Morse vector field with corresponding matching consisting of $k + 1$ edges. In particular, there is no non-stationary closed V -path. Let (σ, τ) be an edge of this matching with $\sigma < \tau$ and let \bar{V} be the combinatorial Morse vector field corresponding to the original matching with the edge (σ, τ) removed. By induction hypothesis $\partial^{\bar{V}} \circ \partial^{\bar{V}} = 0$. In particular, for every $n \in \mathbb{N}$, every $\tau \in K_{n+1}$ and every $v \in K_{n-1}$ when there is a cell $\sigma_1 \in K_n$ such that there is a \bar{V} -path from an hyperface of τ to σ_1 and another \bar{V} -path from an hyperface of σ_1 to v there exists another cell $\sigma_2 \in K_n$ with the same property so that their contribution to $\partial^{\bar{V}} \circ \partial^{\bar{V}}$ are opposite.

First, we will prove that $\partial^V \circ \partial^V = 0$ when the chain complex is with coefficients in $\mathbb{Z}/2\mathbb{Z}$ and after we will take care of signs.

Suppose that the distinguished edge of the matching (σ, τ) is such that $\dim(\sigma) + 1 = \dim(\tau) = n + 1$. Therefore, $C_i^V = C_i^{\bar{V}}$ for $i \neq n, n + 1$ and $\partial_{|C_i^V}^V = \partial_{|C_i^{\bar{V}}}^{\bar{V}}$ for $i \notin \{n, n + 1, n + 2\}$. So we have $\partial^V \circ \partial^V(\mu) = 0$ for all $\mu \in K - (K_n \cup K_{n+1} \cup K_{n+2})$.

Remark that it is also true for every $\sigma' \in \text{Crit}_n(V)$ that $\partial^V \circ \partial^V(\sigma') = 0$ (since with respect to \bar{V} it is true and $\sigma' \neq \sigma$).

There are two cases left.

Case 1. Let $\tau' \in \text{Crit}_{n+1}(V)$. To see that $\partial^V \circ \partial^V(\tau') = 0$ we must consider two cases. First case is when the two \bar{V} -paths which annihilates don't go through σ . Then, nothing is changed and contributions to $\partial^V \circ \partial^V(\tau')$ cancel by pair. The second case is when at least one the \bar{V} -path which cancel by pair for $\partial^{\bar{V}}$ go through σ . The \bar{V} -paths which go from τ' to v are of two types: those who go via σ and the others. Let $\tau' \rightarrow \sigma_2 \rightarrow v$ be a juxtaposition of two \bar{V} -paths which cancel with the juxtaposition of

\overline{V} -path $\tau' \rightarrow \sigma \rightarrow \nu$. Since $\partial^{\overline{V}} \circ \partial^{\overline{V}}(\tau) = 0$, there must be a critical cell σ_1 such that the juxtaposition of \overline{V} -paths $\tau \rightarrow \sigma \rightarrow \nu$ and $\tau \rightarrow \sigma_1 \rightarrow \nu$ cancels. Therefore, when considering $\partial^{\overline{V}}$, three juxtapositions of \overline{V} -paths disappear and one is created: $\tau' \rightarrow (\sigma \rightarrow \tau) \rightarrow \sigma_1 \rightarrow \nu$. It cancels with $\tau' \rightarrow \sigma_2 \rightarrow \nu$.

It may happen that two juxtapositions of \overline{V} -paths go through σ but this case works exactly in the same way.

Case 2. This case is similar to the previous case. Let ζ be a cell in K_{n+2} . There are two cases to see that $\partial^{\overline{V}} \circ \partial^{\overline{V}}(\zeta) = 0$. The first case is when the two \overline{V} -paths whose contributions are opposite don't go through τ . Then, nothing is changed and contributions to $\partial^{\overline{V}} \circ \partial^{\overline{V}}(\tau')$ cancel by pair. The second case is when the \overline{V} -path which disappears is replaced by exactly a new one which goes through the edge (σ, τ) . The result follows similarly.

Note that to deal with this two cases we used the fact that there is no non-stationary closed V -path (and so \overline{V} -path). Now, let's deal with the signs. We will only consider the case 1. above, other cases work similarly. Denote $n(\alpha \rightarrow \beta)$ (respectively $\overline{n}(\alpha \rightarrow \beta)$) the sign of the contribution in the differential $\partial^{\overline{V}}$ (respectively $\partial^{\overline{V}}$) of a path going from α to β where both cells are critical of consecutive dimension. While considering \overline{V} , we have by induction hypothesis

$$\overline{n}(\tau' \rightarrow \sigma_2) \cdot \overline{n}(\sigma_2 \rightarrow \nu) = -\overline{n}(\tau' \rightarrow \sigma) \cdot \overline{n}(\sigma \rightarrow \nu) \quad (2.1)$$

and

$$\overline{n}(\tau \rightarrow \sigma_1) \cdot \overline{n}(\sigma_1 \rightarrow \nu) = -\overline{n}(\tau \rightarrow \sigma) \cdot \overline{n}(\sigma \rightarrow \nu). \quad (2.2)$$

Since the juxtaposition of the \overline{V} -paths $\tau' \rightarrow \sigma_2 \rightarrow \nu$ don't go through σ we have that

$$\overline{n}(\tau' \rightarrow \sigma_2) \cdot \overline{n}(\sigma_2 \rightarrow \nu) = n(\tau' \rightarrow \sigma_2) \cdot n(\sigma_2 \rightarrow \nu). \quad (2.3)$$

By definition of the multiplicity of paths we have

$$n(\tau' \rightarrow \sigma_1) = \overline{n}(\tau' \rightarrow \sigma) \cdot (-\langle \partial\tau, \sigma \rangle) \cdot \overline{n}(\tau \rightarrow \sigma_1). \quad (2.4)$$

Combining equations (2.1)-(2.4) we obtain the following equalities

$$\begin{aligned} n(\tau' \rightarrow \sigma_1) \cdot n(\sigma_1 \rightarrow \nu) &= \overline{n}(\tau' \rightarrow \sigma) \cdot (-\langle \partial\tau, \sigma \rangle) \cdot \overline{n}(\tau \rightarrow \sigma_1) \cdot n(\sigma_1 \rightarrow \nu) \quad (2.4) \\ &= \overline{n}(\tau' \rightarrow \sigma) \cdot (-\langle \partial\tau, \sigma \rangle) \cdot \overline{n}(\tau \rightarrow \sigma_1) \cdot \overline{n}(\sigma_1 \rightarrow \nu) \\ &= \overline{n}(\tau' \rightarrow \sigma) \cdot \langle \partial\tau, \sigma \rangle \cdot \overline{n}(\tau \rightarrow \sigma) \cdot \overline{n}(\sigma \rightarrow \nu) \quad (2.2) \\ &= \langle \partial\tau, \sigma \rangle \cdot \overline{n}(\tau \rightarrow \sigma) \cdot \overline{n}(\tau' \rightarrow \sigma) \cdot \overline{n}(\sigma \rightarrow \nu) \\ &= \overline{n}(\tau' \rightarrow \sigma) \cdot \overline{n}(\sigma \rightarrow \nu) \quad \text{by definition} \\ &= -\overline{n}(\tau' \rightarrow \sigma_2) \cdot \overline{n}(\sigma_2 \rightarrow \nu) \quad (2.1) \\ &= -n(\tau' \rightarrow \sigma_2) \cdot n(\sigma_2 \rightarrow \nu) \quad (2.3) \end{aligned}$$

which concludes the proof of the theorem.

2.2.3. Proof of Theorem 2.7

The main ingredient of the proof is thinking about combinatorial Morse vector field as an instruction to remove acyclic complexes from the original simplicial chain complex, as done by Chari [2, Proposition 3.3] or Sköldbberg [19]. Given a matching between two cells $\sigma < \tau$, we would like to remove the following short complex (which is acyclic)

$$0 \rightarrow \mathbb{Z}.\tau \xrightarrow{\langle \partial\tau, \sigma \rangle} \mathbb{Z}.\sigma \rightarrow 0$$

where ∂ is the boundary operator of the simplicial chain complex. To do this, we use Gaussian elimination (see e.g. [1]):

Lemma 2.8 (Gaussian elimination). *Let $\mathcal{C} = (C_*, \partial)$ be a chain complex over \mathbb{Z} freely generated. Let $b_1 \in C_i$ (respectively $b_2 \in C_{i-1}$) be such that $C_i = \mathbb{Z}.b_1 \oplus D$ (respectively $C_{i-1} = \mathbb{Z}.b_2 \oplus E$). If $\phi : \mathbb{Z}.b_1 \rightarrow \mathbb{Z}.b_2$ is an isomorphism of \mathbb{Z} -modules, then the four term complex segment of \mathcal{C}*

$$\dots \rightarrow [C_{i+1}] \xrightarrow{\begin{pmatrix} \alpha \\ \beta \end{pmatrix}} \begin{bmatrix} b_1 \\ D \end{bmatrix} \xrightarrow{\begin{pmatrix} \phi & \delta \\ \gamma & \varepsilon \end{pmatrix}} \begin{bmatrix} b_2 \\ E \end{bmatrix} \xrightarrow{\begin{pmatrix} \mu & \nu \end{pmatrix}} [C_{i-2}] \rightarrow \dots \quad (2.5)$$

is isomorphic to the following chain complex segment

$$\dots \rightarrow [C_{i+1}] \xrightarrow{\begin{pmatrix} 0 \\ \beta \end{pmatrix}} \begin{bmatrix} b_1 \\ D \end{bmatrix} \xrightarrow{\begin{pmatrix} \phi & 0 \\ 0 & \varepsilon - \gamma\phi^{-1}\delta \end{pmatrix}} \begin{bmatrix} b_2 \\ E \end{bmatrix} \xrightarrow{\begin{pmatrix} 0 & \nu \end{pmatrix}} [C_{i-2}] \rightarrow \dots \quad (2.6)$$

Both these complexes are homotopy equivalent to the complex segment

$$\dots \rightarrow [C_{i+1}] \xrightarrow{\begin{pmatrix} \beta \end{pmatrix}} [D] \xrightarrow{\begin{pmatrix} \varepsilon - \gamma\phi^{-1}\delta \end{pmatrix}} [E] \xrightarrow{\begin{pmatrix} \nu \end{pmatrix}} [C_{i-2}] \rightarrow \dots \quad (2.7)$$

Here we used matrix notation for the differential ∂ .

Proof. Since $\partial^2 = 0$ in \mathcal{C} , we obtain $\phi\alpha + \delta\beta = 0$ and $\mu\phi + \nu\gamma = 0$. By doing the following change of basis $A = \begin{pmatrix} 1 & \phi^{-1}\delta \\ 0 & 1 \end{pmatrix}$ on $\begin{bmatrix} b_1 \\ D \end{bmatrix}$ and $B = \begin{pmatrix} 1 & 0 \\ -\gamma\phi^{-1} & 1 \end{pmatrix}$ on $\begin{bmatrix} b_2 \\ E \end{bmatrix}$ we see that the complex segments 2.5 and 2.6 are isomorphic. Then, we remove the short complex $0 \rightarrow [b_1] \xrightarrow{\phi} [b_2] \rightarrow 0$ which is acyclic. \square

Now, we are ready to prove Theorem 2.7.

Proof of Theorem 2.7. Like for the proof of Theorem 2.6, we make an induction on the number of edges belonging to the matching defining the combinatorial Morse vector field. Let X be a simplicial complex, K be the set of its cells.

Initialization: *matching with no edge.* In this case, there is nothing to prove since the combinatorial Thom-Smale complex is exactly the simplicial chain complex.

Heredity: *suppose the property is true for every matching with at most k edges defining a combinatorial Morse vector field.* Let V be a combinatorial Morse vector field whose underlying matching consists of $k + 1$ edges. Let $\sigma^{(n)} < \tau^{(n+1)}$ be an element of this matching and \bar{V} be the combinatorial Morse vector field equal to V with the matching $\sigma < \tau$ removed (it is actually a combinatorial Morse vector field). So, $C_i^V = C_i^{\bar{V}}$ for all $i \neq n, n + 1$ and $\partial^V = \partial^{\bar{V}}$ when restricted to C_i^V for all $i \notin \{n, n + 1, n + 2\}$. Moreover, we have the following equalities: $(\partial^{\bar{V}})|_{C_{n+1}^V} = (\partial^V)|_{C_{n+1}^V}$ and $\partial|_{C_n^{\bar{V}}} = \partial|_{C_n^V}$. By induction hypothesis, the combinatorial Thom-Smale complex $(C_*^{\bar{V}}, \partial^{\bar{V}})$ is a chain complex homotopy equivalent to the simplicial chain complex of X . Thus, the combinatorial Thom-Smale complex associated with \bar{V} is equal to the one of V except on the following chain segment (where $\varepsilon = (\partial^{\bar{V}})|_{C_{n+1}^V}$):

$$\dots \rightarrow [C_{n+2}^V] \rightarrow \begin{pmatrix} \alpha \\ \partial^V \end{pmatrix} [C_{n+1}^V] \rightarrow \begin{pmatrix} \langle \partial\tau, \sigma \rangle & \delta \\ \gamma & \varepsilon \end{pmatrix} [C_n^V] \rightarrow (\mu \partial^V)[C_{n-1}^V] \rightarrow \dots \quad (2.8)$$

Since X is a simplicial complex we have $\langle \partial\tau, \sigma \rangle \in \{\pm 1\}$. Applying Lemma 2.8, we obtain the following new combinatorial chain complex which is homotopy to the combinatorial Thom-Smale complex of \bar{V}

$$\dots \rightarrow [C_{n+2}^V] \xrightarrow{(\partial^V)} [C_{n+1}^V] \xrightarrow{(\alpha)} [C_n^V] \xrightarrow{(\partial^V)} [C_{n-1}^V] \rightarrow \dots \quad (2.9)$$

where $\alpha = \varepsilon - \gamma \langle \partial\tau, \sigma \rangle \delta = \partial^V$. Thus, the only thing to prove is that $\alpha = \partial^V$ over C_{n+1}^V . To do this, we investigate ∂^V . There are two types of contributions. First type correspond to V -paths which do not go through σ , and they are counted in ε . Second type are V -paths which go through σ . They begin at an hyperface of a critical cell τ' and go through σ : this is the contribution of δ . Then, they jump to τ : this is the contribution of $\langle \partial\tau, \sigma \rangle$. Finally they begin at an hyperface of τ and go to a critical cell in C_n^V : this is the contribution of γ . It remains to check that the sign is correct, but this is exactly the same as in the first proof of Theorem 2.6. \square

Corollary 2.9. *Let X be a finite simplicial complex, $\mathcal{C} = (C_*, \partial)$ be the corresponding simplicial chain complex and \mathcal{M} be a matching $(\sigma_i < \tau_i)_{i \in I}$ on its Hasse diagram defining a combinatorial vector field V . Then the following properties are equivalent:*

- (1) \mathcal{M} is a Morse matching,
- (2) for any sequence $(\sigma_{i_1} < \tau_{i_1}), (\sigma_{i_2} < \tau_{i_2}), \dots, (\sigma_{i_{|I|}} < \tau_{i_{|I|}})$ such that $i_j \neq i_k$ if $j \neq k$, Gaussian eliminations can be performed in this order.

In particular, any sequence of Gaussian eliminations corresponding to \mathcal{M} lead to the same chain complex which is the combinatorial Thom-Smale complex of V .

Proof.

1 \Rightarrow 2 This is an immediate consequence of the proof of Theorem 2.7 and the fact that it leads to the Thom-Smale complex associated to (X, V) .

2 \Rightarrow 1 It is enough to show that there is no non-stationary closed path under the hypothesis. Suppose there is a closed V -path $\gamma : \sigma_1, \dots, \sigma_r, \sigma_1$ and consider any sequence of Gaussian elimination which coincides with $(\sigma_j < V(\sigma_j))$ until step r . In particular, $r \geq 3$ since X is a simplicial complex. Let V be the corresponding combinatorial vector field. Let $\gamma' : \sigma_1, \dots, \sigma_r$ be the V -path with length decrease by one. Then

$$\langle \partial^{r-1} V(\sigma_r), \sigma_r \rangle = \langle \partial V(\sigma_r), \sigma_r \rangle + m(\gamma').$$

Since $m(\gamma') = \pm 1$, $\langle \partial^{r-1} V(\sigma_r), \sigma_r \rangle$ is not invertible over \mathbb{Z} and the Gaussian elimination cannot be performed (see Lemma 2.8). This is a contradiction. \square

3. Relation between smooth and discrete Morse theories

In this section, we investigate the link between smooth and discrete Morse theories. We first recall briefly the main ingredients of smooth Morse theory. In particular, we describe the Thom-Smale complex and prove the following:

Theorem 3.1 (Combinatorial realization). *Let M be a smooth closed oriented Riemannian manifold and $f : M \rightarrow \mathbb{R}$ be a generic Morse function. Suppose that every stable manifold has been given an orientation so that the smooth Thom-Smale complex is defined. Then, there exists a C^1 -triangulation T of M and a combinatorial Morse vector field V on it which realize the smooth Thom-Smale complex (after a choice of orientation of each cells of T) in the following sense:*

- (1) *there is a bijection between the set of critical cells and the set of critical points,*
- (2) *for each pair of critical cells σ_p and σ_q such that $\dim(\sigma_p) = \dim(\sigma_q) + 1$, V -paths from hyperfaces of σ_p to σ_q are in bijection with integral curves of v up to renormalization connecting q to p ,*
- (3) *this bijection induce an isomorphism between the smooth and the combinatorial Thom-Smale complexes.*

Throughout this section, we follow conventions of Milnor ([12, 13]).

3.1. Smooth Morse theory

Let M be a smooth closed oriented Riemannian manifold of dimension n . Given a smooth function $f : M \rightarrow \mathbb{R}$, a point $p \in M$ is said *critical* if $Df(p) = 0$. Let $\text{Crit}(f)$ be the set of critical points. At a critical point p , we consider the bilinear form $D^2f(p)$. The number of negative eigenvalues of $D^2f(p)$ is called the *index* of p (denoted $\text{ind}(p)$). We denote $\text{Crit}_k(f)$ the set of critical points of index k .

Definition 3.2. A smooth map $f : M \rightarrow \mathbb{R}$ is called a *Morse function* if at each critical point p of f , $D^2f(p)$ is non-degenerate.

More generally, a Morse function on a (smooth) cobordism $(M; M_0, M_1)$ is a smooth map $f : M \rightarrow [a, b]$ such that

- (1) $f^{-1}(a) = M_0, f^{-1}(b) = M_1$,
- (2) all critical points of f are interior (lie in $M - (M_0 \cup M_1)$) and are non-degenerate.

For technical reasons, we must consider the following object:

Definition 3.3. Let f be a Morse function on a cobordism $(M^n; M_0, M_1)$. A vector field v on M^n is a *gradient-like vector field* for f if

- (1) $v(f) > 0$ throughout the complement of the set of critical points of f ,
- (2) given any critical point p of f there is a Morse chart in a neighbourhood U of p so that

$$f(x) = f(p) - \sum_{i=1}^k x_i^2 + \sum_{i=k+1}^n x_i^2$$

and v has coordinates $v(x) = (-x_1, \dots, -x_k, x_{k+1}, \dots, x_n)$.

Given any Morse function, there always exists a gradient-like vector field (see [13]). In the following, we shall abbreviate “gradient-like vector field” by “*gradient*”. Thus, when needed, we will assume that we have chosen one.

Given any $x_0 \in M$, we consider the following Cauchy problem

$$\begin{cases} \gamma'(t) &= v(\gamma(t)) \\ \gamma(0) &= x_0 \end{cases}$$

and call *integral curve* (denoted γ_{x_0}) the solution of this Cauchy problem. The *stable manifold* of a critical point p is by definition the set $W^s(p, v) := \{x \in M \mid \lim_{t \rightarrow +\infty} \gamma_x(t) = p\}$. The *unstable manifold* of a critical point p is by definition the set $\{x \in M \mid \lim_{t \rightarrow -\infty} \gamma_x(t) = p\}$. When stable and unstable manifolds are transverse (this is called *Morse-Smale condition*), we called v a *Morse-Smale gradient*: such gradient always exists in a neighbourhood of a gradient (see e.g. [16]). We shall call a Morse function f *generic* if we have chosen for f a Morse-Smale gradient.

To define the smooth Thom-Smale complex we need the following data:

- a generic Morse function f ,
- an orientation of each stable manifold.

Under these conditions, the number of integral curves of v up to renormalization (that is $\gamma_x \sim \gamma_y$ iff there exists $t \in \mathbb{R}$ such that $\gamma_x(t) = y$) connecting two critical points of consecutive index is finite. Moreover, when we consider an integral curve from q to p where $\text{ind}(p) = \text{ind}(q) + 1$, it carries a coorientation induced by the orientation of the stable manifold and the orientation of the integral curve. One can move this coorientation from p to q along the integral curve and compare it with the orientation of the stable manifold of q . This gives the sign which is carried by the integral curve connecting q to p .

The Thom-Smale complex (C_*^f, ∂^f) is defined as:

- (1) $C_k^f = \bigoplus_{p \in \text{Crit}_k(f)} \mathbb{Z} \cdot p$,
- (2) if $p \in \text{Crit}_k(f)$ then $\partial p = \sum_{q \in \text{Crit}_{k-1}(f)} n(p, q) \cdot q$ where $n(p, q)$ is the algebraic number of integral curves up to renormalization connecting q to p .

Theorem 3.4. *The homology of the Thom-Smale complex is equal to the singular homology of M .*

The proof of this theorem can be extracted from [13].

3.2. Elementary cobordisms

In this subsection, we will prove that we can realize combinatorially the smooth Thom-Smale complex of any elementary cobordisms. Thus, by cutting the manifold M into elementary cobordism we will obtain the first part of Theorem 3.1: there exists a bijection between the set of critical cells and the set of critical points.

We will only consider C^1 -triangulation of manifolds for technical reasons (see [21]). So, whenever we use the word triangulation it means C^1 -triangulation. A triangulation of a $n + 1$ -cobordism $(M^{n+1}; M_0, M_1)$ is a triplet $(T; T_0, T_1)$ such that T is a C^1 -triangulation of M , T_0 (respectively T_1) is a subcomplex of T which is a C^1 -triangulation of M_0 (respectively M_1).

A combinatorial Morse vector field V on a triangulated $n + 1$ -cobordism $(T; T_0, T_1)$ is a combinatorial Morse vector field on T such that no cells of T_1 is critical and every cell of T_0 is critical.

Definition 3.5. Let V be a combinatorial Morse vector field on a triangulated $n + 1$ -cobordism $(T; T_0, T_1)$. V satisfies the *ancestor's property* if given any n -cell $\sigma_0 \in T_0$, there exists an n -cell $\sigma_1 \in T_1$ and a V -path starting at σ_1 and ending at σ_0 .

Remark 3.6. There is a key difference between integral curves up to renormalization of a gradient v and V -paths. Given a point $x \in M$, there is only one solution to the Cauchy problem. Moreover, the past and the future of a point pushed along the flow is uniquely determined. *A contrario* given a cell σ , there are (in general)

many V -paths starting at σ . The ancestor's property characterises $n + 1$ -cobordism equipped with a combinatorial Morse vector field which knows its history in maximal dimension $(n, n + 1)$.

To prove that elementary cobordism can be realized, we need a combinatorial description of being a deformation retract. Let X be a simplicial complex and σ be an hyperface of τ which is free (that is σ is a face of no other cell). In this case, we say that X collapses to $X - (\sigma \cup \tau)$ by an elementary collapsing and write $X \searrow X - (\sigma \cup \tau)$. A collapsing is a finite sequence of such elementary collapsings. In particular, a collapsing defines a matching on the Hasse diagram of the simplicial complex. Moreover, one can prove that $X - (\sigma \cup \tau)$ is a deformation retract of X .

Proposition 3.7. *Let X be a simplicial complex and X_0 be a subcomplex. Suppose $X \searrow X_0$. Then the matching given by this collapsing defines a combinatorial Morse vector field whose set of critical cells is the set of cells of X_0 .*

Proof. The only thing to check is that there is no non-stationary closed path. Since elementary collapsings are performed by choosing a free hyperface of a cell, there is no non-stationary closed path. \square

Let $\Delta^m = (a_0, \dots, a_m)$ be the standard simplex of dimension m . The cartesian product $X = \Delta^m \times \Delta^n$ is the cellular complex whose set of cells is $\{\mu \times \nu\}$ where μ (respectively ν) is a cell of Δ^m (respectively Δ^n) (see [22]).

Proposition 3.8 ([17, Proposition 2.9]). *The cartesian product $\Delta^m \times \Delta^n$ has a simplicial subdivision without any new vertex. More generally, the cartesian product of two simplicial complexes has a simplicial subdivision without any new vertex.*

Lemma 3.9. *Let $X_1 = \Delta^k$ be the standard simplicial complex of dimension k and X_0 be a simplicial subdivision of X_1 . Consider the CW-complex which is equal to the cartesian product $\Delta^k \times \Delta^1$ and where we subdivide $\Delta^k \times \{0\}$ so that it is equal to X_0 . Then, there exists a simplicial subdivision X of this CW-complex such that $X|_{\Delta \times \{i\}} = X_i$ for $i \in \{0, 1\}$.*

Moreover for $i \in \{0, 1\}$ there exists a collapsing $X \searrow X_i$ and the combinatorial Morse vector field associated V_i satisfies the ancestor's property on $(X; X_{\bar{j}}, X_{\bar{j}+1})$ (\bar{j} is the class in $\mathbb{Z}/2\mathbb{Z}$).

Proof. The simplicial subdivision and the collapsing is constructed by induction on k . If $k = 0$, choose a new vertex in the interior of the simplex $\Delta^0 \times \Delta^1$ and the elementary collapsing $\Delta^1 \searrow \{i\}$ gives the two collapsing $\Delta^0 \times \Delta^1 \searrow \Delta^0 \times \{i\}$ for $i \in \{0, 1\}$. In particular, the corresponding combinatorial Morse vector field satisfies the ancestor's property.

Suppose the lemma is true until rank $k - 1$. At rank k , let Y be the corresponding CW-complex and x be a point in the interior of the cell of dimension $k + 1$. By induction hypothesis $Y|_{\partial \Delta^k \times \Delta^1}$ admits a simplicial subdivision. Therefore, $Y|_{\partial(\Delta^k \times \Delta^1)}$ admits a simplicial subdivision denoted Z (just add the simplexes $\Delta^k \times \{i\}$ which are equal to X_i for $i \in \{0, 1\}$). The simplicial subdivision X is

given by making the join of the simplicial subdivision of the boundary over $\{x\}$: $X = Z * \{x\}$.

Now, the collapsing $X \searrow X_0$ is performed in three steps.

Step 1. The cell $\sigma \in X_1$ of dimension k is the free hyperface of the cell $\sigma * \{x\}$. We do the following elementary collapsing:

$$X \searrow X - (\sigma \cup \sigma * \{x\}) \tag{3.1}$$

Step 2. By induction hypothesis, $X|_{\partial\Delta^k \times \Delta^1} \searrow X|_{\partial\Delta^k \times \{0\}}$. Performing the join over x induces the following collapsing:

$$X|_{\partial\Delta^k \times \Delta^1} * \{x\} \searrow X|_{\partial\Delta^k \times \{0\}} * \{x\} \tag{3.2}$$

Step 3. It remains to collapse $X_0 * \{x\}$ on X_0 . Let y be a vertex in X_0 which is a vertex of the original simplex X_1 . Since X_0 is a simplicial subdivision of Δ^k , there exists a collapsing $X_0 \searrow \{y\}$. This collapsing gives the following collapsing:

$$X_0 * \{x\} \searrow X_0 \cup (\{y\} \times \{0\}) * \{x\} \searrow X_0 \tag{3.3}$$

Combining collapsings (3.1), (3.2) and (3.3) gives $X \searrow X_0$. The corresponding combinatorial Morse vector field satisfies the ancestor's property by construction.

The collapsing $X \searrow X_1$ is constructed in the same way and conclusions of lemma follows. □

Remark 3.10. The proof of Lemma 3.9 is by induction. Let $\delta^{(j)}$ be the j -th skeleton of Δ^k . Denote by $X^{(j)}$ (respectively $X_i^{(j)}$) the simplicial complex $X|_{\delta^{(j)} \times \Delta^1}$ (respectively $(X_i)|_{\delta^{(j)} \times \Delta^1}$). For $i \in \{0, 1\}$, the collapsing $X \searrow X_i$ can be restricted to $X^{(j)} \searrow X_i^{(j)}$ for any $0 \leq j \leq k$ and the induced combinatorial Morse vector field satisfies the ancestor's property.

The next two lemmas are technical lemmas. The first one is the basic tool to glue together triangulated cobordisms. The second one will be useful to construct a combinatorial realization of a cobordism with exactly one critical point and is a generalization of Lemma 3.9.

Lemma 3.11. *Let (T_i^M, T_i^N) be two C^1 -triangulations of the pair (M, N) where N^k is a submanifold (possibly with boundary) of M^n ($k \leq n$). Then, there exists a C^1 -triangulation T of $(M \times [0, 1], N \times [0, 1])$ such that*

$$(T|_{M \times \{i\}}, T|_{N \times \{i\}}) = (T_i^M, T_i^N)$$

for $i \in \{0, 1\}$ and 2 collapsings

$$T \searrow T_0^M \cup T|_{N \times [0, 1]} \tag{3.4}$$

$$T|_{N \times [0, 1]} \searrow T_0^N. \tag{3.5}$$

Moreover, the induced combinatorial Morse vector fields V satisfies the ancestor's property on the cobordisms $(T|_{N \times [0, 1]}; T_0^N, T_1^N)$ and $(T; T_0^M, T_1^M)$.

Proof. First, suppose $N = \emptyset$. Both triangulations T_0 and T_1 are C^1 -triangulation of the same manifold therefore they have a common simplicial subdivision $T^{1/2}$ [21] (this is where we use the fact that triangulations are C^1 -triangulations). Subdivide $\Delta^1 = [0, 1]$ in two standard simplexes $[0, 1/2]$ and $[1/2, 1]$. Lemma 3.9 gives a C^1 -triangulation of $M \times [0, 1/2]$ (respectively $M \times [1/2, 1]$) denoted $T^{[0,1/2]}$ (respectively $T^{[1/2,1]}$). The union $T^{[0,1/2]} \cup T^{[1/2,1]}$ is a triangulation of $M \times [0, 1]$ denoted T . By construction, $T_{|M \times \{i\}} = T_i^M$ for $i \in \{0, 1\}$ and we have the two following collapsings:

$$\begin{aligned} T^{[1/2,1]} &\searrow T^{1/2} \\ T^{[0,1/2]} &\searrow T^0. \end{aligned}$$

Composing these two collapsings give the desired collapsing and Lemma 3.9 gives the ancestor's property.

In the case where the submanifold N is non-empty, the construction above gives a triangulation T of the pair $(M \times [0, 1], N \times [0, 1])$ and we have $(T_{|M \times \{i\}}, T_{|N \times \{i\}}) = (T_i^M, T_i^N)$ for $i \in \{0, 1\}$. The collapsing $T \searrow T_0$ can be restricted to $T_{|N \times [0,1]}$. We remove from the matching edges corresponding to $T_{N \times [0,1]} \searrow T_{N \times \{0\}}$ to obtain the desired collapsing. Again Lemma 3.9 give the ancestor's property. \square

Lemma 3.12. *Let (m, n) be a pair of positive integers. Let $\Delta^n = (a_0, \dots, a_n)$ be the standard simplex of dimension n and $\delta^{n-1} = (\widehat{a_0}, \dots, a_n)$ be the hyperface which does not contain a_0 . In particular $\Delta^n = \{a_0\} * \delta^{n-1}$. Then there exists a simplicial subdivision X of the cartesian product $\Delta^m \times \Delta^n$ such that*

- $X_{|\Delta^m \times \delta^{n-1}}$ is a simplicial subdivision without any new vertex given by Lemma 3.8,
- $X_{|\Delta^m \times \{a_0\}} = \Delta^m$,
- $X \searrow X_{|(\partial \Delta^m \times \Delta^n) \cup (\Delta^m \times \{a_0\})}$.

Moreover, for each simplex $\Delta_i^1 = (a_0, a_i)$ ($i \neq 0$),

- $X_{|\Delta^m \times \Delta_i^1}$ coincides with the simplicial complex given by Lemma 3.9,
- the collapsing $X \searrow X_{|(\partial \Delta^m \times \Delta^n) \cup (\Delta^m \times \{a_0\})}$ restricted to $X_{|\Delta^m \times \Delta_i^1}$ coincides with the collapsing of Lemma 3.9,
- the induced combinatorial Morse vector field satisfies the ancestor's property on $(X_{|\Delta^m \times \Delta_i^1}; X_{|\Delta^m \times \{a_0\}}, X_{|\Delta^m \times \{a_i\}})$.

Proof. The proof is by induction on $k = m + n > 0$. At rank $k = 1$ there are two cases. The case $m = 0$ and $n = 1$ is trivial: there is nothing to prove. The case $m = 1$ and $n = 0$ is given by Lemma 3.9.

Suppose the lemma is true until rank $k - 1$. Let $(m, n) \in \mathbb{N}^2$ be such that $m + n = k$. We will first subdivide the boundary of $\Delta^m \times \Delta^n$. Since

$$\begin{aligned} \partial(\Delta^m \times \Delta^n) &= (\partial \Delta^m \times \Delta^n) \cup (\Delta^m \times \partial \Delta^n) \\ &= (\partial \Delta^m \times \Delta^n) \cup (\Delta^m \times (\{a_0\} * \partial \delta^{n-1})) \cup (\Delta^m \times \delta^{n-1}) \end{aligned}$$

we define for each cellular complex above a simplicial subdivision.

- The simplicial subdivision of $\Delta^m \times \delta^{n-1}$ is given by Proposition 3.8: in particular, we do not create any new vertex.
- The induction hypothesis gives a simplicial subdivision of

$$(\partial \Delta^m \times (\{a_0\} * \delta^{n-1})) \cup (\Delta^m \times (\{a_0\} * \partial \delta^{n-1})).$$

Let x be a point which is in the interior of the $(m + n)$ -cell of $\Delta^m \times \Delta^n$. The simplicial subdivision X of $\Delta^m \times \Delta^n$ is given by making the cone over $\{x\}$ of the simplicial subdivision of the boundary of $\Delta^m \times \Delta^n$.

By construction we have the following collapsing

$$X_{|\{x\}*(\Delta^m \times \delta^{n-1})} \searrow X_{|\{x\}*\partial(\Delta^m \times \delta^{n-1})} \tag{3.6}$$

which is realized by a downward induction on the dimension of cells of $\Delta^m \times (\delta^{n-1} - \partial \delta^{n-1})$: every cell $\sigma \in \Delta^m \times (\delta^{n-1} - \partial \delta^{n-1})$ is a free hyperface of $\{x\} * \sigma$.

The induction hypothesis says that there exists a simplicial subdivision Y of $\Delta^u \times \Delta^v$ such that $Y \searrow Y_{|(\partial \Delta^u \times \Delta^v) \cup (\Delta^u \times \{a_0\})}$ whenever $u + v < k$ (a_0 is the first vertex of Δ^v). In fact, we have also the following collapsing since the construction is made by induction:

$$Y \searrow Y_{|\Delta^u \times \{a_0\}}$$

Therefore, we have the following collapsings

$$X_{|\partial \Delta^m \times (\{a_0\} * \delta^{n-1})} \searrow X_{|\partial \Delta^m \times \{a_0\}} \tag{3.7}$$

$$X_{|\Delta^m \times \{a_0\} * \partial \delta^{n-1}} \searrow X_{|\Delta^m \times \{a_0\}}. \tag{3.8}$$

Collapsing (3.6) followed by the cone over x of the collapsing (3.7) and the cone over x of the collapsing (3.8) give the following collapsing:

$$X \searrow X_{|(\partial \Delta^m \times \Delta^n) \cup (\{x\} * (\Delta^m \times \{a_0\}))}.$$

Finally there exists a collapsing $\{x\} * (\Delta^m \times \{a_0\}) \searrow \Delta^m \times \{a_0\}$ (by choosing a vertex $y \in \Delta^m$ and considering the collapsing $\Delta^m \searrow \{y\}$) which gives the result.

In case $n = 1$, this construction is the same as the one of Lemma 3.9. □

Theorem 3.13. *Let f be a generic Morse function on a cobordism $(M; M_0, M_1)$ with exactly one critical point p of index k . Then, there exists a C^1 -triangulation of the cobordism $(T; T_0, T_1)$ such that*

- (1) *the stable manifold of p is a subcomplex of T denoted T_p^s and $T \searrow T_p^s \cup T_0$,*
- (2) *there is a cell σ_p of dimension $\text{ind}(p)$ such that $p \in \sigma_p \subset T_p^s$ and $T_p^s - \sigma_p \searrow (T_p^s \cap T_0)$.*

In particular, the combinatorial Morse vector field given by these two collapsings has exactly one critical cell σ_p outside cells of T_0 .

Proof. Suppose a Morse–Smale gradient v for f is fixed. Let $W^s(p, v)$ be the corresponding stable manifold of p . We follow the proof of Milnor which proves that $M_0 \cup W^s(p, v)$ is a deformation retract of M (see the proof of Theorem 3.14 [12]). Let C be a (small enough) tubular neighbourhood of $W^s(p, v)$. The original proof consists of two steps. First, $M_0 \cup C$ is a deformation retract of M : this is done by pushing along the gradient lines of v . Then, $M_0 \cup W^s(p, v)$ is a deformation retract of $M_0 \cup C$. We prove the theorem in two steps.

First step: construction of a good triangulation of C

The tubular neighbourhood C is diffeomorphic to $D^k \times D^{n-k}$ (for $i \in \mathbb{N}^*$, D^i is the unit disk in \mathbb{R}^i). Thanks to this diffeomorphism, the stable manifold is identified with $D^k \times \{0\}$ and the adherence of the unstable manifold is identified with $\{0\} \times D^{n-k}$. Triangulate the stable manifold by the standard simplex Δ^k and denote σ_p its interior (so $T_p^s = \bar{\sigma}_p$).

We triangulate D^{n-k} by choosing an arbitrary triangulation of $\partial D^{n-k} = S^{n-k-1}$ and considering D^{n-k} as the cone over its center $\{0\}$: this gives a triangulation of D^{n-k} . The triangulation of $\bar{\sigma}_p \times D^{n-k}$ is the following one: choose a simplicial subdivision of $\bar{\sigma}_p \times \partial D^{n-k}$ without any new vertex given by proposition 3.8. Then, triangulate the cartesian product $\bar{\sigma}_p \times D^{n-k}$ with the triangulation of $\bar{\sigma}_p \times \partial D^{n-k}$ already fixed thanks to Lemma 3.12:

- for each simplex $v \in \partial D^{n-k}$, the lemma constructs a triangulation of $\bar{\sigma}_p \times (\{0\} * v)$,
- for each pair of simplexes $(v_0, v_1) \in (\partial D^{n-k})^2$, the simplicial subdivisions of $\bar{\sigma}_p \times (\{0\} * v_i)$ coincides over $\bar{\sigma}_p \times (\{0\} * (v_0 \cap v_1))$.

Let T^C be the triangulation of $\bar{\sigma}_p \times D^{n-k}$ constructed above. By construction, we have the following collapsing

$$T^C \searrow T_p^s \cup T_{|\partial\bar{\sigma}_p \times D^{n-k}}^C \tag{3.9}$$

Second step: combinatorial realization of the first retraction

Let T_0 be a triangulation of M_0 which coincides over $M_0 \cap C$ with the triangulation above. Consider the following submanifolds with boundary: $\partial C_- = M_0 \cap C$, $M_0^{\bar{C}} = M_0 - \text{Int}(\partial C_-)$ and $\partial C_+ = \partial C - \text{Int}(\partial C_-)$. Let V be $\partial C_- \cap \partial C_+$: it is diffeomorphic to $\partial D^k \times \partial D^{n-k} = S^{k-1} \times S^{n-k-1}$ a manifold of dimension $n - 2$.

The manifold $(\partial C_+, V)$ is a manifold with boundary which is triangulated. The gradient lines of v starting at any point of this manifold are transverse to it: we push along the gradient lines of v the triangulation until it meets M_1 . It gives a triangulation of $(M_1^{\partial C_+}, M_1^V)$ which is a submanifold of M_1 with boundary. This triangulation is C^1 since pushing along the flow in this case is a diffeomorphism. Then, we get a product cobordism (with boundary) with triangulation of the top and the bottom already fixed: Lemma 3.11 gives a triangulation of this cobordism with the desired collapsing.

The same construction holds for $(M_0^{\overline{C}}, V)$ (we suppose that the triangulation of $V \times [0, 1]$ is the same as the one given above). Let T be the corresponding triangulation of M . Then, we have the following collapsings

$$T \searrow T_0 \cup T^C. \tag{3.10}$$

Conclusion

The composition of collapsings (3.10) and (3.9) give

$$T \searrow T_0 \cup T_p^s.$$

Since $T_p^s = \overline{\sigma}_p$ we get the following collapsing: $T_p^s - \sigma_p \searrow \partial T_p^s$. Thus a combinatorial Morse vector field which satisfies the conclusion of the theorem has been constructed. Nevertheless, note that the triangulation above is not C^1 : the triangulation of the stable manifold done by Δ^k gives only a topological triangulation. To correct this, push the level M_0 (denote this level M'_0) along the gradient line a little inside the cobordism so that the stable manifold can be C^1 -triangulated by the standard simplex. Then, we endow the cobordism whose boundary is $M_0 \cup M'_0$ with a C^1 -triangulation given by Lemma 3.9. \square

Corollary 3.14. *Let f be a generic Morse function on a Riemannian closed manifold M . Then, there exists T a C^1 -triangulation of M and a combinatorial Morse vector field V defined on T such that for every $k \in \mathbb{N}$ the set of critical points of index k is in bijection with the set of critical cells of dimension k .*

Proof. Since the Morse function f is generic, we have that for any critical points $p \neq q$, $f(p) \neq f(q)$. Let $a_1 < a_2 < \dots < a_l$ be the ordered set of critical values of f . For each $k \in \{1, \dots, l\}$, let $\varepsilon_k > 0$ be small enough so that the cobordism

$$(M^{a_k}; M_-^{a_k}, M_+^{a_k}) = (f^{-1}([a_k - \varepsilon_k, a_k + \varepsilon_k]); f^{-1}(a_k - \varepsilon_k), f^{-1}(a_k + \varepsilon_k))$$

is a cobordism with exactly one critical point. Define for $k \in \{1, \dots, l - 1\}$ the product cobordisms

$$(M^{b_k}; M_+^{a_{k-1}}, M_-^{a_k}) = (f^{-1}([a_{k-1} + \varepsilon_{k-1}, a_k - \varepsilon_k]); f^{-1}(a_{k-1} + \varepsilon_{k-1}), f^{-1}(a_k - \varepsilon_k)).$$

The manifold M is equal to:

$$M^{a_1} \cup M^{b_1} \cup \dots \cup M^{b_{l-1}} \cup M^{a_l}.$$

Theorem 3.13 gives for $k = 1, \dots, l$ a combinatorial realization of the cobordism $(M^{a_k}; M_-^{a_k}, M_+^{a_k})$. Lemma 3.11 gives a combinatorial realization of each cobordism $(M^{b_k}; M_+^{a_{k-1}}, M_-^{a_k})$ for $k = 1, \dots, l - 1$ (with the convention that $M^{a_0} = \emptyset$). Then, we construct a C^1 -triangulation of M and define on it a combinatorial vector field. It is in fact a combinatorial Morse vector field since along V -paths we only can go down and the conclusion of the corollary follows. \square

3.3. Proof of Theorem 3.1

Since f is generic, we use the Rearrangement Theorem [13, Theorem 4.8] to consider g a generic self-indexed Morse function such that

- the set of critical points of index k of g coincides with the one of f for every $k \in \mathbb{N}$,
- for each pair of critical points p and q of successive index, the set of integral curves up to renormalization connecting q to p for g is in bijection with the corresponding set for f (we suppose here that Morse–Smale gradients have been chosen for f and for g),
- this bijection induces an isomorphism between the Thom–Smale complexes of f and g (we suppose that orientations of stable manifolds have been chosen).

Thus, we suppose that $f : M^n \rightarrow \mathbb{R}$ is a generic self-indexed Morse function i.e. for every $k \in \mathbb{N}$, for every $p \in \text{Crit}_k(f)$, $f(p) = k$. In particular $f(M) = [0, n]$. We suppose whenever we need it that a Morse–Smale gradient v for f is given.

One more time, we will cut M in cobordisms (almost) elementary and control combinatorially the behavior of V -paths. For $i \in \{0, \dots, n\}$ choose $0 < \varepsilon_i < 1/2$. For $i \in \{0, \dots, n\}$, let $(M^i; M_-^i, M_+^i)$ be the cobordism

$$(f^{-1}([i - \varepsilon_i, i + \varepsilon_i]); f^{-1}(i - \varepsilon_i), f^{-1}(i + \varepsilon_i)).$$

Similarly, define $(M^{i,i+1}; M_+^i, M_-^{i+1})$ the product cobordism equal to

$$(f^{-1}([i + \varepsilon_i, i + 1 - \varepsilon_{i+1}]); M_+^i, M_-^{i+1}).$$

Then

$$M = M^0 \cup M^{0,1} \cup M^1 \cup \dots \cup M^{n-1,n} \cup M^n.$$

For all $i \in \{0, \dots, n\}$, $(M^i; M_-^i, M_+^i)$ is a cobordism with $|\text{Crit}_i(f)|$ critical points of index i (maybe there is no critical point). The triangulation of M is constructed in the following way:

- (1) triangulation of cobordisms $(M^i; M_-^i, M_+^i)$ for all $i \in \{0, \dots, n\}$ given by Theorem 3.13,
- (2) triangulation of cobordisms $(M^{i,i+1}; M_+^i, M_-^{i+1})$ for all $i \in \{0, \dots, n\}$ given by Lemma 3.11.

Remark 3.15. Theorem 3.13 is proved in the case where there is exactly one critical point. This proof extends directly to the case of k critical points of the same index under the condition that tubular neighbourhoods of stable manifolds are chosen to be disjoint one from each other.

Let p be a critical point of index k and $C(p)$ be a tubular neighbourhood (small enough) of the stable manifold of p in the corresponding cobordism. Denote $\partial C_-(p)$ (respectively $\partial C_+(p)$) the submanifold diffeomorphic to $\partial D^k \times D^{n-k}$ (respectively $D^k \times \partial D^{n-k}$). Denote σ_p the critical cell of dimension k corresponding to p (see Theorem 3.13).

Hypothesis on the triangulation of $\partial C_+(p)$

- (1) stable manifolds of critical points of index $k + 1$ intersect $\partial C_+(p)$ along a subcomplex of dimension k and intersect $\sigma_p \times \partial D^{n-k}$ along cells of dimension k of the type $\sigma \times \{a_i\}$ where a_i is a vertex of ∂D^{n-k} ,
- (2) each integral curve up to renormalization γ from p to $q \in \text{Crit}_{k+1}(f)$ intersects $\partial C_+(p)$ in the interior of a k -cell $\sigma_\gamma \in \sigma_p \times \partial D^{n-k}$,
- (3) given two distinct integral curves up to renormalization γ and γ' from p to critical points of index $k + 1$ then $\sigma_\gamma \neq \sigma_{\gamma'}$.

Remark 3.16. The first hypothesis is satisfied by choosing small enough ε_k and since stable and unstable manifolds are transverse. For such a small enough ε_k the last hypothesis will be satisfied. The second hypothesis is automatically satisfied if the first hypothesis is satisfied.

In each triangulated cobordism $(M^k; M_-^k, M_+^k)$, stable manifolds of critical points of index k are subcomplexes. Following notations of Theorem 3.13, we have the following collapsing

$$T_p^s - \sigma_p \searrow \partial T_p^s.$$

Using Lemma 3.11, we obtain the following collapsing

$$M^{k-1,k} \searrow M_+^{k-1}$$

which can be restricted to the stable manifold of p since it is a submanifold of M_-^k . With respect to the stable manifold, the combinatorial Morse vector field satisfies the ancestor’s property.

Let γ be an integral curve of v up to renormalization from $q \in \text{Crit}_{k-1}(f)$ to p . It intersects $\partial C_+(q)$ in a point which by hypothesis belongs to a cell $\sigma_q \times \{a_\gamma\}$. There is a 1 – 1 correspondance between the set of integral curves up to renormalization from q to p (with $\text{ind}(p) = \text{ind}(q) + 1$) and V -paths from hyperfaces of σ_p to σ_q given by $\gamma \longleftrightarrow \sigma_\gamma$.

From σ_γ , there is a unique V -path ending at σ_q . Since V satisfies the ancestor’s property in the stable manifold and σ_γ is a cell of dimension $k - 1$ there is an ancestor of σ_γ which is an hyperface of σ_p . This gives a V -path between an hyperface of σ_p to σ_q which corresponds to γ .

We endow each critical cell with the orientation of the corresponding stable manifold and every other cell is endowed with an arbitrary orientation.

By construction, the multiplicity of V -path coincides with the sign of the corresponding gradient path and the theorem follows.

4. Perfect matchings and homologous vector fields

In this section, we use Theorem 3.1 to prove the following: given a closed oriented 3-manifold and a class of homologous vector fields, there is a triangulation such that a perfect matching on the Hasse diagram of the triangulation realizes this class of homologous vector fields.

4.1. Perfect matchings

Definition 4.1. A *perfect matching* on a graph is a matching such that every vertex belongs to an edge of the matching.

As a corollary of Theorem 3.1 we obtain:

Corollary 4.2. *Let M be a closed smooth manifold of dimension 3. Then there exists a C^1 -triangulation of M such that a perfect matching on its Hasse diagram exists.*

Proof. Since M is a closed smooth manifold of dimension 3 we have $\chi(M) = 0$ where χ denotes the Euler characteristic. Take a pointed Heegaard splitting of M $(\Sigma_g; \underline{\alpha} = (\alpha_1, \dots, \alpha_g), \underline{\beta} = (\beta_1, \dots, \beta_g); z)$ of genus g so that there is an n -uplets of intersection points x between the α 's and the β 's which defines a bijection between the sets $\underline{\alpha}$ and $\underline{\beta}$. It is always possible to find such a pointed Heegaard splitting after a finite number of isotopies of the α 's and β 's curves (see [7]). The Morse function f corresponding to this Heegaard splitting has one critical point of index 0 and 3 and g critical points of index 1 and 2. Denote the set of index 1 (respectively 2) critical points by $\{q_i\}_{i=1}^g$ (respectively $\{p_i\}_{i=1}^g$) where q_i (respectively p_i) corresponds to α_i (respectively β_i) for all $i \in \{1, \dots, g\}$. The n -uplet of intersection points $x = (x_{1,i_1}, \dots, x_{n,i_n})$ gives for each $j \in \{1, \dots, g\}$ an integral curve connecting q_j to p_{i_j} . The point z gives an integral curve connecting the index 0 critical point to the index 3 critical point.

Take a combinatorial realization (T, V) as given by Theorem 3.1 of (M, f) . Then to each point $x_{i,j}$ correspond now a V -path γ from an hyperface of the critical cell $\sigma_{p_{i_j}}$ to σ_{q_i} : we change the matching along this path so that both $\tau_{p_{i_j}}$ and σ_{q_i} are no more critical cells. If $\gamma : \sigma_0, \dots, \sigma_r = \sigma_{q_i}$, then do the following:

- match σ_0 with τ ,
- for every $i \in \{1, \dots, r\}$ match σ_i with $V(\sigma_{i-1})$.

Now suppose that z belongs to the interior of a 2-cell τ_z (if not, subdivide T). Denote by ζ the critical cell of dimension 3 and by ν the critical cell of dimension 0. There is by construction a V -path $\gamma : \tau_0, \dots, \tau_r = \tau_z$ from an hyperface of ζ to τ_z since z is in the stable manifold of the index 3 critical point. We modify the matching along γ this way:

- match τ_0 with ζ ,
- for every $i \in \{1, \dots, r\}$ match τ_i with $V(\tau_{i-1})$.

In fact, it is no more a matching since τ_z belongs to two edges of the matching.

Nevertheless, τ_z belongs to the unstable manifold of ν the critical cell of dimension 0. By the construction done in Theorem 3.1, the tubular neighbourhood of the critical point of index 0 is equal to $D^3 = \partial D^3 * \{0\}$. The triangulation of this tubular neighbourhood is given by making the cone over $\{0\} = \nu$ of a triangulation of ∂D^3 . We modify the matching as follows. Let ν denote the critical 0-cell and suppose $\bar{\tau}_z * \nu$ is the tetrahedron $ABCD$ where A corresponds to ν . The

collapsing $\partial D^3 * \{0\} \searrow \{0\}$ gives in particular the following matching on $ABCD$: $(BCD, ABCD)$, (BC, ABC) and (B, AB) (A is critical). Modify the matching by (A, AB) , (B, BC) and $(ABC, ABCD)$. Then, BCD (which is τ_z) is critical. This gives a perfect matching over T . \square

4.2. Euler structures and homologous vector fields

Throughout this subsection we use conventions of Turaev [20].

4.2.1. Combinatorial Euler structures

Perfect matchings have an interpretation as Euler chains. First, we recall Euler structures as defined by Turaev [20]. Let $(M, \partial M)$ be a smooth manifold of dimension n and T be a C^1 -triangulation of M .

Suppose $\partial M = \partial_0 M \sqcup \partial_1 M$ be such that $\chi(M, \partial_0 M) = 0$ and let T_i be equal to $T|_{\partial_i M}$ for $i \in \{0, 1\}$.

Denote K the set of cells of T and K_i the set of cells of T_i for $i \in \{0, 1\}$. For each cell $\sigma \in K$, let $\text{sgn}(\sigma)$ be equal to $(-1)^{\dim(\sigma)}$ and pick a_σ a point in the interior of σ . An *Euler chain* in (T, T_0) is a one-dimensional singular chain ξ in T with the boundary of the form $\sum_{\sigma \in K - K_0} \text{sgn}(\sigma) a_\sigma$. Since $\chi(M, \partial_0 M) = 0$, the set of Euler chains is non-empty. Given two Euler chains ξ and η , the difference $\xi - \eta$ is a cycle. If $\xi - \eta = 0 \in H_1(M)$ then we say that ξ and η are *homologous*. A class of homologous Euler chains in (T, T_0) is called a *combinatorial Euler structure* on (T, T_0) . Let $\text{Eul}(T, T_0)$ be the set of Euler structures on (T, T_0) . If ξ is an Euler chain, denote by $[\xi]$ its class as a combinatorial Euler structure. Euler chains behave well with respect to the subdivision of a triangulation: this allows us to consider the set $\text{Eul}(M, \partial_0 M)$ of Euler structures on $(M, \partial_0 M)$. Taking ξ an element of this set means choosing a triangulation (T, T_0) of $(M, \partial_0 M)$ and considering an Euler chain on (T, T_0) .

Remark 4.3. Let \mathcal{C} be a perfect matching on a C^1 -triangulation (T, T_0) of $(M, \partial_0 M)$. Then it defines an Euler chain $[\xi_{\mathcal{C}}] \in \text{Eul}(T, T_0)$: orient every edge of the matching from odd dimensional cells to even dimensional cells. Perfect matchings are special Euler chains that do not pass through a cell more than one time.

4.2.2. Homologous vector fields

By a *vector field* on $(M, \partial_0 M)$ we mean (except in clearly mentioned case) a non-singular continuous vector field of tangent vectors on M directed into M on $\partial_0 M$ and directed outwards on $\partial_1 M$. Since $\chi(M, \partial_0 M) = 0$, there exists such vector fields on $(M, \partial_0 M)$.

Vector fields u and v on $(M, \partial_0 M)$ are called *homologous* if for some closed ball $B \subset \text{Int}(M)$ the restriction of the fields u and v are homotopic in the class of non-singular vector fields on $M - \text{Int}(B)$ directed into M on $\partial_0 M$, outwards on $\partial_1 M$, and arbitrarily on ∂B . Denote by $\text{vect}(M, \partial_0 M)$ the set of homologous vector fields on $(M, \partial_0 M)$ and the class of a vector field u is denoted by $[u]$.

4.2.3. The canonical bijection

Turaev proved the following:

Theorem 4.4 (Turaev [20]). *Let $(M, \partial_0 M)$ be a smooth pair such that $\dim(M) \geq 2$. For each C^1 -triangulation (T, T_0) of the pair $(M, \partial_0 M)$ there exists a bijection*

$$\rho : \text{Eul}(T, T_0) \rightarrow \text{vect}(M, \partial_0 M)$$

Let us recall the construction of Turaev in the case $\partial M = \emptyset$. Let T be a C^1 -triangulation of M and T' be the first barycentric subdivision of T . We recall the definition of the vector field F_1 with singularities on M . For a simplex a of the triangulation T , let \underline{a} denote its barycenter. If $A = \langle \underline{a}_0, \underline{a}_1, \dots, \underline{a}_p \rangle$ is a simplex of the triangulation T' , where $a_0 < a_1 < \dots < a_p$ are simplexes of T , then, at a point $x \in \text{Int}(A)$,

$$F_1(x) = \sum_{0 \leq i < j \leq p} \lambda_i(x) \lambda_j(x) (\underline{a}_j - x).$$

Here $\lambda_0, \lambda_1, \dots, \lambda_p$ are barycentric coordinates in A and $\underline{a}_j - x$ is the image of the tangent vector $\underline{a}_j - x \in T_x A$ via the homeomorphism between T and M .

Every barycenter of each cell of T is a singular point of F_1 . Let \mathcal{M} be a matching on the Hasse diagram of T : in particular, every edge of the matching connects two singular points. Turaev proved that the index of F_1 in a neighbourhood of every edge of the Hasse diagram (thought as embedded in M in the obvious way) is equal to zero. Thus, if we think about combinatorial (not necessarily Morse) vector field on T as a matching on its Hasse diagram, it encodes a desingularization of the vector field F_1 (where critical points of F_1 remain if the corresponding cell is critical). In the case of ξ an Euler chain, let F_ξ denote an extension of F_1 to a non-singular vector field. Turaev proved that the homotopy $[F_\xi] \in \text{Vect}(M)$ depends only on $[\xi] \in \text{Eul}(T)$. The map ρ is defined by $\rho([\xi]) = [F_\xi]$.

In the general case where ξ corresponds to a matching (instead of a perfect matching), we still denote by ρ the map which assigns to ξ the vector field F_ξ given by the construction above.

We refer to [20] in the case $\partial M \neq \emptyset$.

4.3. \mathcal{M} -realization of vector fields

Definition 4.5. Let $[u] \in \text{Vect}(M, \partial_0 M)$. A \mathcal{M} -realization of $[u]$ is a matching η on the corresponding Hasse diagram of a C^1 -triangulation T of $(M, \partial M)$ such that $[\eta] = \rho([u])$.

Theorem 4.6. *Any class of homologous vector fields on a smooth oriented closed Riemannian 3-manifold has an \mathcal{M} -realization.*

This theorem is a step toward Heegaard–Floer homology [14, 15]. Recall the first steps of the construction of Heegaard–Floer homology for closed, oriented 3-manifolds. Choose a pointed Heegaard splitting $M = U_0 \cup_{\Sigma} U_1$ (that is choose a Morse function) and fix a Spin^c -structure on M given by an n -uplets of intersection points between the α 's and the β 's. Turaev proved that Spin^c -structures are in bijection with Euler structures (and so the set $\text{vect}(M)$). Thus, Theorem 4.6 together with Theorem 3.1 give for a closed oriented 3-manifold a combinatorial realization of both the Spin^c -structure and the pointed Heegaard splitting. The hard part remaining is to understand how holomorphic disks can be combinatorially realized.

The following two lemmas will be useful to prove Theorem 4.6.

Lemma 4.7. *Let $(M \times [0, 1]; M \times \{0\}, M \times \{1\})$ be a smooth $n + 1$ product cobordism such that T_i is a C^1 -triangulation of $M \times \{i\}$ for $i \in \{0, 1\}$. Let $(T; T_0, T_1)$ be any triangulation given by Lemma 3.11 and $[\xi] \in \text{Eul}(M \times [0, 1], M \times \{0\})$ the Euler structure given by the combinatorial Morse vector field induced by $T \searrow T_0$. Then $\rho([\xi]) \in \text{vect}(M \times [0, 1], M \times \{0\})$ is homologous to the vector field $v : M \times [0, 1] \rightarrow T(M \times [0, 1])$ defined by $v(x, t) = ((x, t), dt)$.*

Proof. The construction of the collapsing $T \searrow T_0$ defines a combinatorial Morse vector field pointing downwards. Let $\xi \in \text{Eul}(T, T_0)$ be the corresponding Euler chain. The map $\rho : \text{Eul}(M \times [0, 1], M \times \{0\}) \rightarrow \text{vect}(M \times [0, 1], M \times \{0\})$ sends ξ to a non-singular vector field on $(M \times [0, 1], M \times \{0\})$ which is, by definition of ρ , by the construction of the triangulation and by the definition of the combinatorial Morse vector field, homologous to the desired vector field. \square

Lemma 4.8. *Let f be a generic Morse function on a cobordism $(M; M_0, M_1)$ with exactly one critical point p of index k and $(T; T_0, T_1)$ be a C^1 -triangulation of the cobordism $(T; T_0, T_1)$ given by Theorem 3.13. Let ξ be the one singular chain corresponding to the matching given by the combinatorial Morse vector field on T . Then $\rho(\xi)$ is homologous to the chosen Morse–Smale gradient of f outside a small ball neighbourhood of the critical point p . Moreover, the index of $\rho(\xi)$ at p is equal to k .*

Proof. We follow notations of Theorem 3.13. The two collapsings $T \searrow T_p^s \cup T_0$ and $T_p^s - \sigma_p \searrow (T_p^s \cap T_0)$ define a combinatorial Morse vector field with only one critical cell. Let ξ be the corresponding one singular chain. Thus, the barycenter of this cell (which is p) must be a critical point of $\rho(\xi)$. It remains to check that outside a small neighbourhood of p , $\rho(\xi)$ is homologous to the Morse–Smale gradient v . Since outside the tubular neighbourhood C of the stable manifold the triangulation is constructed by pushing it along gradient lines of v , we can use Lemma 4.7 to see that $\rho(\xi)$ is homologous to v outside the tubular neighbourhood C of the stable manifold. In a small ball neighbourhood (which is a Morse chart at p) of the critical point p , the vector field F_1 coincides with the Morse–Smale gradient v . Since $T^C \searrow \sigma_p \cup (T_0 \cap T^C)$, $\rho(\xi)$ is homologous to v outside the Morse chart of p . The fact that the index of $\rho(\xi)$ at p is equal to k is a consequence of the definition of F_1 . \square

Proof of Theorem 4.6. We apply Theorem 3.1 to obtain a C^1 -triangulation of the 3-manifold and a combinatorial Morse vector field which realizes combinatorially the Thom-Smale complex. Then, the construction done in Corollary 4.2 defines a matching which in turns defines an Euler chain ξ . The map ρ sends ξ to a non-singular vector field which is by construction homologous to the Morse–Smale gradient of f . Thus, to prove the theorem for any $[v] \in \text{vect}(M)$, it remains to find a pointed Heegaard splitting $(\Sigma_g; \underline{\alpha} = (\alpha_1, \dots, \alpha_g), \underline{\beta} = (\beta_1, \dots, \beta_g); z)$ of the 3-manifold M such that an n -uplet of intersection points x corresponds to a given $[v] \in \text{vect}(M)$. Finally, [15, Lemma 5.2] tells that any $[v] \in \text{vect}(M)$ can be realized in such a way. This concludes the proof. \square

References

- [1] D. BAR-NATAN, *Fast Khovanov homology computations*, J. Knot Theory Ramifications **16** (2007), 243–255.
- [2] M. K. CHARI, *On discrete Morse functions and combinatorial decompositions*, Formal power series and algebraic combinatorics (Vienna, 1997), Discrete Math. **217** (2000), 101–113.
- [3] D. FARLEY and L. SABALKA, *Discrete Morse theory and graph braid groups*, Algebr. Geom. Topol. **5** (2005), 1075–1109.
- [4] R. FORMAN, *Combinatorial vector fields and dynamical systems*, Math. Z. **228** (1998), 629–681.
- [5] R. FORMAN, *Morse theory for cell complexes*, Adv. Math. **134** (1998), 90–145.
- [6] R. FORMAN, *Witten-Morse theory for cell complexes*, Topology **37** (1998), 945–979.
- [7] R. E. GOMPF and A. I. STIPSICZ, “4-Manifolds and Kirby Calculus”, Graduate Studies in Mathematics, Vol. 20, American Mathematical Society, Providence, RI, 1999.
- [8] M. JOSWIG and M. E. PFETSCH, *Computing optimal Morse matchings*, SIAM J. Discrete Math. **20** (2006), 11–25.
- [9] T. LEWINER, H. LOPES and G. TAVARES, *Optimal discrete Morse functions for 2-manifolds*, Comput. Geom. **26** (2003), 221–233.
- [10] T. LEWINER, H. LOPES and G. TAVARES, *Applications of Forman’s discrete Morse theory to topology visualization and mesh compression*, Transactions on Visualization and Computer Graphics **10** (2004), 499–508.
- [11] A. T. LUNDELL and S. WEINGRAM, “The Topology of cw Complexes”, Van Nostrand Reinhold Co., New York, 1969.
- [12] J. MILNOR, “Morse Theory”, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [13] J. MILNOR, “Lectures on the h -Cobordism Theorem”, Notes by L. Siebenmann and J. Sondow, Princeton University Press, Princeton, N.J., 1965.
- [14] P. OZSVÁTH and Z. SZABÓ, *Holomorphic disks and knot invariants*, Adv. Math. **1** (2004), 58–116.
- [15] P. OZSVÁTH and Z. SZABÓ, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), 1027–1158.
- [16] A. V. PAJITNOV, *On the Novikov complex for rational Morse forms*, Ann. Fac. Sci. Toulouse Math. (6) **4** (1995), 297–338.
- [17] C. P. ROURKE and B. J. SANDERSON, “Introduction to Piecewise-Linear Topology”, Springer-Verlag, New York, 1972.
- [18] M. SALVETTI and S. SETTEPANELLA, *Combinatorial Morse theory and minimality of hyperplane arrangements*, Geom. Topol. **11** (2007), 1733–1766.

- [19] EMIL SKÖLDBERG, *Morse theory from an algebraic viewpoint*, Trans. Amer. Math. Soc. **358** (2006), 115–129 (electronic).
- [20] V. G. TURAEV, *Euler structures, nonsingular vector fields, and Reidemeister-type torsions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), 607–643, 672.
- [21] J. H. C. WHITEHEAD, *On C^1 -complexes*, Ann. of Math. (2) **41** (1940), 809–824.
- [22] H. WHITNEY, “Geometric Integration Theory”, Princeton University Press, Princeton, New Jersey, 1957.

Laboratoire de Mathématiques
et Applications des Mathématiques (LMAM)
Université de Bretagne Sud
Campus de Tohannic – BP 573
56017 Vannes, France
etienne.gallais@math.univ-nantes.fr