

# *Astérisque*

BURKHARD KÜLSHAMMER

**Morita equivalent blocks in Clifford theory of finite groups**

*Astérisque*, tome 181-182 (1990), p. 209-215

[http://www.numdam.org/item?id=AST\\_1990\\_\\_181-182\\_\\_209\\_0](http://www.numdam.org/item?id=AST_1990__181-182__209_0)

© Société mathématique de France, 1990, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# Morita Equivalent Blocks in Clifford Theory of Finite Groups

BURKHARD KÜLSHAMMER

Let  $F$  be an algebraically closed field of prime characteristic  $p$ , and let

$$1 \longrightarrow K \longrightarrow H \longrightarrow G \longrightarrow 1$$

be an extension of finite groups. Let  $B$  be a block of  $FK$  (considered as a subalgebra of  $FK$ ), and let  $A$  be a block of  $FH$  covering  $B$  (i. e.  $1_A 1_B \neq 0$ ). Following a suggestion by J. L. Alperin [1] we consider the following

**QUESTION.** *When are  $A$  and  $B$  Morita equivalent?*

Our main results concerning this question are given by theorems 1, 7, 8 and proposition 10 below. Special cases of this question are dealt with in [2] and [7].

**THEOREM 1.** *With notation as above, the map  $B \longrightarrow 1_A B \subset A$ ,  $b \longmapsto 1_A b$ , is an isomorphism of  $F$ -algebras.*

Before proving theorem 1 we introduce some notation and state some preliminary results. Obviously  $K$  is contained in  $H(B) := \{h \in H: hBh^{-1} = B\}$ , the stabilizer of  $B$  in  $H$ , and we set  $G(B) := H(B)/K$ . The following facts are well-known (see [8; theorem 1], for example).

**PROPOSITION 2.** (i)  $FH 1_B FH$  is the sum of all blocks of  $FH$  covering  $B$ .

(ii) If  $h_1, \dots, h_t$  denote a transversal for  $H(B)$  in  $H$  then the map

$$\text{Mat}(t, 1_B FH(B)) \longrightarrow FH 1_B FH, [a_{ij}]_{1,j=1}^t \longrightarrow \sum_{i,j=1}^t h_i a_{ij} h_j^{-1},$$

is an isomorphism of  $F$ -algebras.

(iii) The maps

$$Z(1_B FH(B)) \longrightarrow Z(FH 1_B FH), z \longmapsto \sum_{i,j=1}^c h_i z h_j^{-1}.$$

and

$$Z(FH 1_B FH) \longrightarrow Z(1_B FH(B)), z \longmapsto 1_B z,$$

are isomorphisms of  $F$ -algebras and inverse to each other.

For  $h \in H(B)$ , the map  $B \longrightarrow B, b \longmapsto hbh^{-1}$ , is an  $F$ -algebra automorphism of  $B$ . It is easy to see that the elements  $h \in H$  for which the map  $B \longrightarrow B, b \longmapsto hbh^{-1}$ , is an inner automorphism of  $B$  form a normal subgroup  $H(B)$  of  $H(B)$  containing  $K$  (cf. [3; proposition 2.7]). Define  $G(B) := H(B)/K$ .

Setting  $C := 1_B C_{FH}(K)$  and  $C_g := C \cap hFK$  for  $g = hK \in G$  we obtain  $C = \bigoplus_{g \in G} C_g$  and  $C_g C_{g'} \subset C_{gg'}$ , for  $g, g' \in G$ , i. e.  $C$  is a  $G$ -graded  $F$ -algebra in the sense of [4]. It is easy to see that  $C_g = 0$  for  $g \in G \setminus G(B)$ . Thus  $C = \bigoplus_{g \in G(B)} C_g$  can also be viewed as a  $G(B)$ -graded  $F$ -algebra.

**PROPOSITION 3.** ([3; lemma 3.3])  $I := \bigoplus_{g \in G(B)} (JZB)C_g \oplus \bigoplus_{g \in G(B) \setminus G(B)} C_g$  is an ideal of  $C$  contained in the radical  $JC$  of  $C$ .

Setting  $C(B) := \bigoplus_{g \in G(B)} C_g$  we thus have  $C = C(B) + JC$ . By lifting theorems for idempotents one obtains the following result.

**COROLLARY 4.** ([3; theorem 3.5]) All idempotents of  $ZC$  are contained in  $C(B)$ .

It is easy to see that  $C(B)$  is a crossed product of  $G(B)$  with  $ZB$ , in the sense of [4]; in particular,  $C(B)$  is free as a  $ZB$ -module, and  $\overline{C(B)} := C(B)/(JZB)C(B)$  is a crossed product of  $G(B)$  with  $ZB/(JZB) \cong F$ , i. e. a twisted group algebra of  $G(B)$  over  $F$ . Our next result is [8; theorem C].

**PROPOSITION 5.** *If  $G = G[B]$  then the map  $B \otimes_{ZB} C \longrightarrow 1_B FH$ ,  $b \otimes c \longmapsto bc$ , is an isomorphism of  $F$ -algebras.*

We are now in a position to prove theorem 1.

**Proof of theorem 1.** Obviously the map  $B \longrightarrow 1_A B$ ,  $b \longmapsto 1_A b$ , is an epimorphism of  $F$ -algebras. Hence it suffices to prove injectivity. By proposition 2,  $1_A 1_B$  is the block idempotent of a block of  $FH(B)$  covering  $B$ . Hence we may replace  $H$  by  $H(B)$  and assume  $H = H(B)$ . By corollary 4,  $1_A$  is contained in  $FH(B)$ . Replacing  $A$  by a block of  $1_A FH(B)$  we may assume that  $H = H(B)$ . In this case the map  $B \otimes_{ZB} C \longrightarrow 1_B FH$ ,  $b \otimes c \longmapsto bc$ , is an isomorphism of  $F$ -algebras by proposition 5. Moreover,  $C$  is free over  $ZB$ . This isomorphism maps  $B \otimes_{ZB} 1_A C$  onto  $A$ . Since  $C = 1_A C \oplus (1_B - 1_A)C$ ,  $1_A C$  is projective over  $ZB$ . Since  $ZB$  is local,  $1_A C$  is even free over  $ZB$ . Thus  $A$  is free over  $B$ , and the result follows.  $\square$

In order to prove our next theorem we need a result on the behaviour of defect groups.

**PROPOSITION 6.** ([3; theorem 7.7])  *$1_A + (JZB)C(B)$  is a primitive idempotent in  $C_{\overline{C(B)}}(G(B))$ , and  $A$  has a defect group  $P$  such that  $P \cap K$  is a defect group of  $B$  and  $PK/K$  is a defect group of  $1_A + (JZB)C(B)$  in  $G(B)$ .*

Part of proposition 6 has also been proved in [6; 4.21]. We will say that  $A$  and  $B$  are "naturally" Morita equivalent of degree  $n$  if there exists a simple  $F$ -subalgebra  $S$  of  $A$  of dimension  $n^2$  such that the map  $1_A B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of  $F$ -algebras. In this case  $A$  and  $B$  are Morita equivalent since  $1_A B$  is isomorphic to  $B$  by theorem 1 and  $S$  is a complete matrix algebra of degree  $n$  over  $F$ .

**THEOREM 7.**  *$A$  and  $B$  are "naturally" Morita equivalent if and only if  $G = G[B]$  and  $A$  and  $B$  have the same defect.*

**Proof.** Suppose first that  $G = G[B]$  and that  $A$  and  $B$  have the same defect. By proposition 6, the block  $1_A C + (JZB)C/(JZB)C$  of the twisted group algebra  $C/(JZB)C$  of  $G[B] = G$  over  $F$  has defect 0 in  $G(B) = G$ . It is well-known that this implies that

the block  $1_A C + (JZB)C / (JZB)C$  of  $C / (JZB)C$  is a simple  $F$ -algebra; in particular,  $1_A J C = (JZB)1_A C$ . By the Wedderburn-Malcev theorem there is a simple  $F$ -subalgebra  $S$  of  $1_A C$  such that  $1_A C = S \oplus 1_A J C = S \oplus (JZB)1_A C$ . Then  $1_A C = (ZB)S + (JZB)1_A C$ , and Nakayama's lemma implies that  $1_A C = (ZB)S$ . In the proof of theorem 1 we had shown that  $1_A C$  is free over  $ZB$ . Thus  $1_A C / 1_A J C$  is free of the same rank over  $ZB / JZB \cong F$ . Therefore the rank of  $1_A C$  over  $ZB$  equals the dimension of  $S$  over  $F$ . Comparing dimensions we see that the map  $ZB \otimes_F S \longrightarrow 1_A C, z \otimes s \longmapsto zs$ , is an isomorphism of  $F$ -algebras. By proposition 5, the map  $B \otimes_F S \longrightarrow A, b \otimes s \longmapsto bs$ , is an isomorphism as well.

Suppose now conversely that  $A$  and  $B$  are "naturally" Morita equivalent, and let  $S$  be a simple  $F$ -subalgebra of  $A$  such that the map  $1_A B \otimes_F S \longrightarrow A, b \otimes s \longmapsto bs$ , is an isomorphism of  $F$ -algebras. Then  $1_A = 1_S = 1_A 1_B$ . On the other hand, it follows from proposition 2 that  $1_A = \sum_{i=1}^t 1_A (h_i 1_B h_i^{-1})$  with pairwise orthogonal idempotents  $1_A (h_i 1_B h_i^{-1})$  where  $t = |H : H(B)|$ . Thus  $H(B) = H$  and  $G(B) = G$ .

We know from proposition 3 that  $C = C[B] + J C$ ; in particular,  $1_A C = 1_A C[B] + 1_A J C$ . On the other hand, since  $A$  and  $B$  are "naturally" Morita equivalent the map

$$1_A ZB \otimes_F S \longrightarrow 1_A ZB \cdot S = C_A(B) = 1_A C, z \otimes s \longmapsto zs,$$

is an isomorphism of  $F$ -algebras. By the Wedderburn-Malcev theorem we may find a unit  $u$  in  $1_A C$  such that  $S^u$  is contained in  $1_A C[B]$ . Then the map  $1_A B \otimes_F S^u \longrightarrow A, b \otimes s \longmapsto bs$ , is an isomorphism of  $F$ -algebras as well. Hence we may assume that  $S$  is contained in  $FH[B]$ . Since also  $1_A \in FH[B]$  by corollary 4 we obtain  $A \subset FH[B]$  which clearly implies that  $H[B] = H$ .

Since  $1_A C$  is isomorphic to  $ZB \otimes_F S$ ,  $1_A C + (JZB)C / (JZB)C$  is a simple  $F$ -algebra. It is well-known that this implies that the block  $1_A C + (JZB)C / (JZB)C$  of  $\overline{C[B]}$  has defect 0 in  $G[B] = G$ . By proposition 6,  $A$  and  $B$  have the same defect.  $\square$

In the following we assume that  $G(B) = G$ ; in view of proposition 2, this is not an important restriction. In this case we can reduce the question of whether  $A$  and  $B$  are "naturally" Morita equivalent to their Brauer correspondents. Let  $Q$  be a defect group of  $B$ , and let  $B'$  be the Brauer correspondent of  $B$  in  $N_K(Q)$ . Since  $G(B) = G$  the Frattini argument shows that  $H = N_H(Q)K$ , and we obtain a finite group extension

$$1 \longrightarrow N_K(Q) \longrightarrow N_H(Q) \longrightarrow N_H(Q)/N_K(Q) \longrightarrow 1$$

with  $N_H(Q)/N_K(Q)$  naturally isomorphic to  $G$ . By proposition 6,  $A$  has a defect group  $P$  such that  $Q = P \cap K$ ; in particular,  $N_H(P) \subset N_H(Q)$ . By Brauer's First Main Theorem,  $A$  has a unique Brauer correspondent  $A'$  in  $N_H(Q)$ . By [5; theorem],  $A'$  covers  $B'$ .

**THEOREM 8.** *With notation as above,  $A$  and  $B$  are "naturally" Morita equivalent if and only if  $A'$  and  $B'$  are "naturally" Morita equivalent.*

In order to prove theorem 8, we need the following result which is a consequence of [3; corollary 12.6].

**PROPOSITION 9.** *In the situation above,  $HIBI = (N_H(Q)IB')IK$ .*

**Proof of theorem 8.** Suppose first that  $A$  and  $B$  are "naturally" Morita equivalent. By theorem 7,  $G = GIBI$ , and  $A$  and  $B$  have the same defect. By proposition 6,  $Q$  is a defect group of  $A$ . By Brauer's First Main Theorem,  $Q$  is a defect group of  $A'$  and  $B'$  as well. Moreover, since  $H = HIBI = (N_H(Q)IB')IK$  by proposition 9, we have

$$N_H(Q) = (N_H(Q)IB')N_K(Q) = N_H(Q)IB'I.$$

By theorem 7,  $A'$  and  $B'$  are "naturally" Morita equivalent.

Suppose now conversely that  $A'$  and  $B'$  are "naturally" Morita equivalent. By theorem 7,  $N_H(Q) = N_H(Q)IB'I$ , and  $A'$  and  $B'$  have the same defect. By Brauer's First Main Theorem,  $B'$  has defect group  $Q$ . By proposition 6,  $A'$  has defect group  $Q$  as well. Again by Brauer's First Main Theorem,  $A$  has defect group  $Q$ . Moreover, proposition 9 implies that  $HIBI = (N_H(Q)IB')IK = N_H(Q)K = H$ . By theorem 7,  $A$  and  $B$  are "naturally" Morita equivalent.  $\square$

This result can be strengthened by using additional information from [2]. Suppose that  $A$  and  $B$  are "naturally" Morita equivalent. Let  $S$  be a simple  $F$ -subalgebra of  $A$  such that the map  $l_A B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of  $F$ -algebras. By theorem 8,  $A'$  and  $B'$  are "naturally" Morita equivalent. Thus there is a simple  $F$ -subalgebra  $S'$  of  $A'$  such that the map  $l_{A'} B' \otimes_F S' \longrightarrow A'$ ,  $b' \otimes s' \longmapsto b's'$ , is an

isomorphism of  $F$ -algebras.

We have seen above that  $B$  determines a twisted group algebra  $\overline{CIBJ}$  of  $GIBJ = G$  over  $F$ . In the same way,  $B'$  determines a twisted group algebra  $\overline{C'I'B'J}$  of  $N_{H'}(Q)/N_{K'}(Q)$  over  $F$ . Since  $N_{H'}(Q)/N_{K'}(Q)$  and  $G$  are naturally isomorphic we can view  $\overline{C'I'B'J}$  as a twisted group algebra of  $G$  over  $F$ . Then [3; corollary 12.6] (which is the main result of [3]) tells us that the Brauer homomorphism induces a natural isomorphism between  $\overline{CIBJ}$  and  $\overline{C'I'B'J}$ . This isomorphism maps the block of defect 0 in  $\overline{CIBJ}$  determined by  $A$  onto the block of defect 0 in  $\overline{C'I'B'J}$  determined by  $A'$ . Now the proof of theorem 7 shows that  $S$  and  $S'$  are isomorphic. Hence we may add the following result to theorem 8.

**PROPOSITION 10.** *If, in the situation of theorem 8,  $A$  and  $B$  are "naturally" Morita equivalent of degree  $n$  then so are  $A'$  and  $B'$ .*

Finally, let us interpret our results in the language of [9]. The block  $B$  of  $FK$  corresponds to a pointed group  $K_\beta$  over  $FK$ , and the block  $A$  of  $FH$  corresponds to a pointed group  $H_\alpha$  over  $FH$ . Let  $Q_\beta$  be a maximal local pointed subgroup of  $K_\beta$ . Suppose that  $A$  and  $B$  are "naturally" Morita equivalent, and let  $S$  be a simple  $F$ -subalgebra of  $A$  such that the map  $1_A B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of  $F$ -algebras. Then  $S$  and  $B$  centralize each other; in particular, every element of  $S$  is fixed by  $Q$ . It follows easily that the source algebras of  $H_\alpha$  and  $K_\beta$  are isomorphic (as interior  $Q$ -algebras).

#### ACKNOWLEDGEMENTS

The author is very much indebted to Prof. J. L. Alperin. The ideas in this paper grew out of his talk [1] given at the University of Essen in March 1987 and subsequent discussions. In his talk Prof. Alperin had proved the "degree 1" part of our results. (In this case the question is whether  $A$  and  $B$  are isomorphic.) His visit to Essen was part of the research project on representation theory of finite groups and finite dimensional algebras sponsored by the Deutsche Forschungsgemeinschaft. The author is also grateful to L. Puig for providing him with a copy of [2].

REFERENCES

1. J. L. ALPERIN, Isomorphic blocks, Talk given at the seminar on representation theory of finite groups, Essen, March 1987
2. BIAN W.,  $p'$ -extensions centrales d'un bloc, Thèse, Paris 1986
3. E. C. DADE, Block extensions, *Illinois J. Math.* **17** (1973), 198-272
4. E. C. DADE, Group-graded rings and modules, *Math. Z.* **174** (1980), 241-262
5. M. E. HARRIS AND R. KNÖRR, Brauer correspondence for covering blocks of finite groups, *Comm. Algebra* **13** (1985), 1213-1218
6. R. KNÖRR, Blocks, vertices and normal subgroups, *Math. Z.* **148** (1976), 53-60
7. B. KÜLSHAMMER, Quotients, Cartan matrices and Morita equivalent blocks, *J. Algebra* **90** (1984), 364-371
8. B. KÜLSHAMMER, Crossed products and blocks with normal defect groups, *Comm. Algebra* **13** (1985), 147-168.
9. L. PUIG, Pointed groups and construction of characters, *Math. Z.* **176** (1981), 265-292

Burkhard Külshammer  
Department of Mathematics  
University of Dortmund  
4600 Dortmund 50  
Federal Republic of Germany