BULLETIN DE LA S. M. F.

SAMIR KHABBAZ

On a theorem of Charles and Erdélyi

Bulletin de la S. M. F., tome 89 (1961), p. 103-104

http://www.numdam.org/item?id=BSMF_1961__89__103_0

© Bulletin de la S. M. F., 1961, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON A THEOREM OF CHARLES AND ERDÉLYI ;

BY

SAMIR KHABBAZ.

(Lehigh University).

The original purpose of the following was to give a short proof of a theorem of CHARLES [1]. CHARLES then indicated that the proof resembled the proof of a theorem of ERDÉLYI [2], p. 81, and that if modified slightly, would cover both theorems. The same proof however proves also a theorem of J. IRWIN and E. WALKER [3]. In the following the three theorems are combined together.

Let G be a primary group, and if $x \in G$ let $h(x)$ denote the ordinary height of x in G . Also let a represent either an integer or the first infinite ordinal; and if a is the first infinite ordinal, let $p^a G$ represent any subgroup

of $\bigcap_{n=1}^{\infty} p^n G$. Then :

THEOREM (CHARLES, ERDÉLYI, IRWIN and WALKER). — Let M be a subgroup of G maximal with respect to disjointness from $p^a G$. Then M is pure in G .

PROOF. — Deny the theorem. Then there is a least positive integer $n < a$ for which there is an equation $p^n x = y$, $y \in M$ having a solution x in G but not in M . Then there exists an integer m , $0 \leq m < n$, and $z \in M$ such that $0 \neq p^m x + z \in p^a G$. Then $h(z) = h(p^m x) \geq m$, since $h(p^m x) \geq m$ and $h(p^m x + z) \geq a$. Since $m < n$, there is an element $z_1 \in M$ with $p^m z_1 = z$. However $p^{n-m}(p^m x + z) \in M \cap (p^a G)$, and hence is zero. Thus $p^{n-m}(-z) = p^n x = y$. Thus,

$$p^n(-z_1) = p^{n-m}(p^m(-z_1)) = p^{n-m}(-z) = y, \quad \text{with } -z_1 \in M.$$

The author would like to thank J. IRWIN and E. WALKER for letting him read their manuscripts [3] and [4] which are to appear shortly in the *Pacific Journal of Mathematics*.

BIBLIOGRAPHY.

- [1] CHARLES (Bernard). — Étude sur les sous-groupes d'un groupe abélien, *Bull. Soc. math. France*, t. 88, 1960, p. 217-227.
- [2] FUCHS (L.). — *Abelian groups*. — Budapest, Hungarian Academy of Sciences, 1958.
- [3] IRWIN (J) and WALKER (E). — On N -high subgroups of abelian groups, *Pacific J. of Math.* (to appear shortly).
- [4] IRWIN (J.). — High subgroups of abelian torsion groups, *Pacific J. of Math.* (to appear shortly).

(Manuscript reçu le 25 novembre 1960.)

Samir KHABBAZ,
Mathematics Department,
Lehigh University,
Bethlehem, Penn. (États-Unis).