

COMPOSITIO MATHEMATICA

KENTARO YANO

HITOSI HIRAMATU

On the projective geometry of K -spreads

Compositio Mathematica, tome 10 (1952), p. 286-296

http://www.numdam.org/item?id=CM_1952__10__286_0

© Foundation Compositio Mathematica, 1952, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

On the projective geometry of K -spreads

by

Kentaro Yano and Hitosi Hiramatu

1. An N -dimensional space is called by J. Douglas [1] a space of K -spreads if there is given a system of K -dimensional manifolds such that there exists one and only one member of the system passing through any $K + 1$ points given in general position. He shows that such a system of K -dimensional manifolds, that is, a system of K -spreads is represented by a system of completely integrable partial differential equations of the form

$$(1.1) \quad \frac{\partial^2 x^i}{\partial u^\beta \partial u^\gamma} + H_{\beta\gamma}^i(x, p) = 0, \quad \left(p_\alpha^i = \frac{\partial x^i}{\partial u^\alpha} \right)$$

where (x^i) ($i, j, k, \dots = 1, 2, \dots, N$) are coordinates of a point of the space, and (u^α) ($\alpha, \beta, \gamma, \dots = 1, 2, \dots, K$) parameters of a point on a K -spread. The functions $H_{\beta\gamma}^i$ ($= H_{\gamma\beta}^i$) form a homogeneous function system of p_α^i with respect to the lower indices, and consequently satisfy the relations

$$(1.2) \quad p_\lambda^a (H_{\beta\gamma}^i |_\alpha^a) = \delta_\beta^a H_{\lambda\gamma}^i + \delta_\gamma^a H_{\beta\lambda}^i,$$

where $|_\alpha^a$ denotes partial differentiation with respect to p_α^a .

The components of affine connection and those of projective connection are given respectively by

$$(1.3) \quad \Gamma_{jk}^i = \frac{1}{K(K+1)} H_{\beta\gamma}^i |_{j|k}^\beta |_{j|k}^\gamma$$

and

$$(1.4) \quad \begin{aligned} *P_{jk}^i &= \Gamma_{jk}^i - \frac{1}{N+1} (\delta_j^i \Gamma_{ak}^a + \delta_k^i \Gamma_{aj}^a) - \\ &\quad - \frac{1}{N-K} p_\lambda^i \{ \Gamma_{jk}^i |_\alpha^\lambda - \frac{1}{N+1} (\Gamma_{aj}^a |_\alpha^\lambda + \Gamma_{ak}^a |_\alpha^\lambda) \}. \end{aligned}$$

J. Douglas states, among other things, that the quantities defined by

$$(1.5) \quad W_{jkl}^i = *B_{jkl}^i - \frac{1}{N-1} (*B_{jka}^a \delta_l^i - *B_{jla}^a \delta_k^i)$$

are components of a projective tensor, where

$$(1.6) \quad *B^i_{jkl} = (*\Pi^i_{jk,l} - *\Pi^i_{jk}|^{\alpha} * \Pi^{\alpha}_{bl} p^b_{\alpha}) -$$

$$- (*\Pi^i_{jl,k} - *\Pi^i_{jl}|^{\alpha} * \Pi^{\alpha}_{bk} p^b_{\alpha}) + *\Pi^i_{jk} * \Pi^i_{al} - *\Pi^i_{jl} * \Pi^i_{ak},$$

the comma followed by an index denoting partial differentiation with respect to x^l . But, it is pointed out by R. S. Clark [2] that the above defined W^i_{jkl} are not components of a tensor in the general case $1 < K < N$.

J. Douglas states also that necessary and sufficient conditions that equations (1.1) can be reduced to $\partial^2 x^i / \partial u^j \partial u^k = 0$ by suitable coordinate and parameter transformations, in other words, those that the space be projectively flat are

$$(1.7) \quad *\Pi^i_{jk}|^{\alpha} = 0 \quad \text{and} \quad W^i_{jkl} = 0.$$

But, it is pointed out recently by Chih-Ta Yen [3] and Hsien-Chung Wang [5] that the conditions (1.7) are not independent and in fact the latter is a consequence of the former.

To study the projective geometry of K -spreads, it will be certainly helpful to construct a space of K linear elements with projective connection whose K -dimensional flat subspaces coincide with the K -spreads. This was actually done by S. S. Chern [6] and Chih-Ta Yen [3], [4].

But, all the authors mentioned above use the so-called natural frame of reference. If we use the natural frame of reference, Douglas' $*\Pi^i_{jk}$ appear in the coefficients of the connection, but because of the complexity of its law of transformation, it is very difficult to deduce any tensor from it. It seems to the present authors that it is rather desirable to use the so-called semi-natural frame of reference instead of natural one. The purpose of the present Note is to show that if we take up the semi-natural frame of reference, we can find out curvature tensors and their laws of transformation with respect to a projective change of Γ^i_{jk} in a very natural way and to discuss the above topics by the use of these results.

2. We consider a space with projective connection whose elements are points (x^i) and K linearly independent contra-variant vectors (p^i_{α}) and whose projective connection is expressed by

$$(2.1) \quad dA_o = \omega^o_o A_o + \omega^i_o A_i, \quad dA_j = \omega^o_j A_o + \omega^i_j A_i,$$

where A_o is a point coinciding with (x^i) , A_j N points taken in the tangent projective space in such a way that $[A_o, A_j]$ forms a frame

of reference and ω 's are Pfaffian forms with respect to x^i and p_α^i . If $dx^i = 0$, then the point A_o having the same position, we must have $\omega_o^i = 0$, from which we can see that ω_o^i do not contain the differentials dp_α^i . Hence, by a suitable change of frames of reference, we can put (2.1) in the form

$$(2.2) \quad \begin{cases} dA_o = (\varphi_i dx^i + \varphi_i^\alpha dp_\alpha^i) A_o + dx^i A_i, \\ dA_j = (\omega_{jk}^o dx^k + \omega_{jk}^{\alpha\alpha} dp_\alpha^k) A_o + (\omega_{jk}^i dx^k + \omega_{jk}^{i\alpha} dp_\alpha^k) A_i, \end{cases}$$

where $\varphi_i, \omega_{jk}^o, \omega_{jk}^i, \varphi_i^\alpha, \omega_{jk}^{\alpha\alpha}, \omega_{jk}^{i\alpha}$ are homogeneous function systems of p_α^i with respect to Greek index and satisfy

$$\varphi_i^\alpha p_\beta^i = 0, \quad \omega_{jk}^{\alpha\alpha} p_\beta^k = 0, \quad \omega_{jk}^{i\alpha} p_\beta^k = 0.$$

Since the projective connection is completely determined by ω_j^o and $\omega_j^i - \delta_j^i \omega_o^o$, we put

$$(2.3) \quad \omega_j^o = \Pi_{jk}^o dx^k + \Pi_{jk}^{\alpha\alpha} dp_\alpha^k, \quad \omega_j^i - \delta_j^i \omega_o^o = \Pi_{jk}^i dx^k + \Pi_{jk}^{i\alpha} dp_\alpha^k,$$

and call Π 's coefficients of the projective connection. The Π 's have the same properties as ω 's, and satisfy

$$(2.4) \quad \Pi_{jk}^{\alpha\alpha} p_\beta^k = 0, \quad \Pi_{jk}^{i\alpha} p_\beta^k = 0.$$

If we calculate $ddA_o - ddA_o = \Omega_{21}^o A_o + \Omega_o^i A_i$, then the torsion

of the space is given by Ω_o^i . In what follows, we assume that our space has no torsion, from which we have

$$(2.5) \quad \Pi_{jk}^i = \Pi_{kj}^i, \quad \Pi_{jk}^{i\alpha} = 0.$$

To express the projective connection analytically, we have adopted the so-called semi-natural frame of reference. It will be easily seen that the transformation between two semi-natural frames of reference must be of the form

$$(2.6) \quad \bar{A}_o = A_o, \quad \bar{A}_j = \lambda_j A_o + A_j,$$

where λ_j are generalized homogeneous functions of degree zero of the set p_α^i . We call such a transformation of semi-natural frame of reference a transformation of hyperplane at infinity. During a transformation of hyperplane at infinity, the coefficients of the projective connection will be transformed into

$$(2.7) \quad \begin{cases} \bar{\Pi}_{jk}^o = \Pi_{jk}^o + \lambda_{j,k} - \lambda_j \Pi_{ik}^i - \lambda_j \lambda_k, & \bar{\Pi}_{jk}^{\alpha\alpha} = \Pi_{jk}^{\alpha\alpha} + \lambda_j \lambda_k^\alpha, \\ \bar{\Pi}_{jk}^i = \Pi_{jk}^i + \delta_j^i \lambda_k + \delta_k^i \lambda_j. \end{cases}$$

If we change the coordinate system from (x^i) to (\bar{x}^i) , the semi-

natural frame of reference will be transformed into

$$(2.8) \quad \bar{A}_o = A_o, \quad \bar{A}_i = \frac{\partial x^a}{\partial \bar{x}^i} A_a,$$

the hyperplane at infinity being assumed to be invariant. During this transformation, the coefficients of the projective connection will be transformed into

$$(2.9) \quad \begin{cases} \bar{\Pi}_{jk}^o = \frac{\partial x^b}{\partial \bar{x}^j} \frac{\partial x^c}{\partial \bar{x}^k} \Pi_{bc}^o + \frac{\partial x^b}{\partial \bar{x}^j} \frac{\partial^2 x^c}{\partial \bar{x}^k \partial \bar{x}^i} \frac{\partial \bar{x}^i}{\partial x^a} p_\alpha^a \Pi_{bc}^{\alpha}, & \bar{\Pi}_{jk}^{\alpha} = \frac{\partial x^b}{\partial \bar{x}^j} \frac{\partial x^c}{\partial \bar{x}^k} \Pi_{bc}^{\alpha}, \\ \bar{\Pi}_{jk}^i = \frac{\partial \bar{x}^i}{\partial x^a} \left(\frac{\partial x^b}{\partial \bar{x}^j} \frac{\partial x^c}{\partial \bar{x}^k} \Pi_{bc}^a + \frac{\partial^2 x^a}{\partial \bar{x}^j \partial \bar{x}^k} \right). \end{cases}$$

Thus we can see that Π_{jk}^{α} are components of a tensor and Π_{jk}^i are those of an affine connection under the transformation of coordinates.

3. Now let us consider a K -dimensional subspace defined by $x^i = x^i(u)$, $p_\alpha^i = \partial x^i / \partial u^\alpha$, and put

$$(3.1) \quad A_o = A_o, \quad A_\alpha = p_\alpha^o A_o + p_\alpha^i A_i, \quad A_B = p_B^o A_o + p_B^i A_i,$$

where the matrix (p_α^i, p_A^i) ($A, B, C, \dots = K + 1, \dots, N$) is of rank N , p_α^o and p_B^o being arbitrary, then the points A_α are on the K -dimensional plane tangent to the subspace and the points A_o, A_α, A_B are linearly independent. Equation (3.1) can be solved with respect to A_o and A_i and gives the equations of the form

$$(3.2) \quad A_o = A_o, \quad A_i = p_i^o A_o + p_i^\alpha A_\alpha + p_i^B A_B,$$

the matrix (p_α^i, p_B^i) and (p_i^α, p_i^B) being inverse to each other.

Now, the fact that the subspace $x^i = x^i(u)$ is flat is expressed by the equations

$$(3.3) \quad dA_o = \omega_o^o A_o + du^\alpha A_\alpha, \quad dA_B = \omega_B^o A_o + \omega_B^\alpha A_\alpha,$$

from which we have

$$(3.4) \quad \frac{\partial^2 x^i}{\partial u^\beta \partial u^\gamma} + \Pi_{jk}^i p_\beta^j p_\gamma^k = p_\alpha^i G_{\beta\gamma}^\alpha,$$

as equations of the flat subspace, where

$$(3.5) \quad G_{\beta\gamma}^\alpha = p_i^\alpha \left(\frac{\partial^2 x^i}{\partial u^\beta \partial u^\gamma} + \Pi_{jk}^i p_\beta^j p_\gamma^k \right).$$

It will be easily seen that the form of the equations (3.4) is

invariant under the change of hyperplane at infinity, change of coordinates and change of parameters and is independent of the choice of p_B^i .

Now, we shall consider the problem to determine a projective connection with respect to which the system of K -spreads (1.1) or

$$(3.6) \quad \frac{\partial^2 x^i}{\partial u^\beta \partial u^\gamma} + \Gamma_{jk}^i p_\beta^j p_\gamma^k = 0$$

will be exactly the system of K -dimensional flat subspaces. It was shown by J. Douglas that the so-called projective change is given by

$$(3.7) \quad \bar{H}_{\beta\gamma}^i = H_{\beta\gamma}^i + p_\alpha^i C_{\beta\gamma}^\alpha$$

or

$$(3.8) \quad \bar{\Gamma}_{jk}^i = \Gamma_{jk}^i + \delta_j^i \lambda_k + \delta_k^i \lambda_j + p_\alpha^i B_{jk}^\alpha,$$

where

$$\lambda_k = \frac{1}{K(K+1)} C_{\alpha\gamma}^\alpha |_{k}^\gamma, \quad B_{jk}^\alpha = \frac{1}{K(K+1)} C_{\beta\gamma}^\alpha |_{j|k}^\beta |_{k}^\gamma,$$

and, $C_{\beta\gamma}^\alpha$ being a homogeneous function system, these quantities satisfy the relations

$$(3.9) \quad p_\alpha^k (\lambda_k |_{i}^\alpha) = 0, \quad p_\alpha^j B_{jk}^\alpha = 0.$$

For an ordinary space with projective connection and an ordinary space of paths, the change of hyperplane at infinity and the projective change of affine connection correspond to each other [7]. While, in our case, the transformation law of Π_{jk}^i and that of Γ_{jk}^i are not the same. But as we can obtain, from (3.8),

$$\begin{aligned} \frac{1}{N-K} [\bar{\Gamma}_{jk}^a |_{a}^\alpha - \frac{1}{N+1} (\bar{\Gamma}_{aj}^a |_{k}^\alpha + \bar{\Gamma}_{ak}^a |_{j}^\alpha)] = \\ \frac{1}{N-K} [\Gamma_{jk}^a |_{a}^\alpha - \frac{1}{N+1} (\Gamma_{aj}^a |_{k}^\alpha + \Gamma_{ak}^a |_{j}^\alpha)] + B_{jk}^\alpha, \end{aligned}$$

we can see that the transformation law of Π_{jk}^i coincides with that of

$$\Gamma_{jk}^i - \frac{1}{N-K} p_\alpha^i [\Gamma_{jk}^a |_{a}^\alpha - \frac{1}{N+1} (\Gamma_{aj}^a |_{k}^\alpha + \Gamma_{ak}^a |_{j}^\alpha)].$$

Consequently we put

$$(3.10) \quad \Pi_{jk}^i = \Gamma_{jk}^i - \frac{1}{N-K} p_\alpha^i [\Gamma_{jk}^a |_{a}^\alpha - \frac{1}{N+1} (\Gamma_{aj}^a |_{k}^\alpha + \Gamma_{ak}^a |_{j}^\alpha)].$$

To determine the other coefficients Π_{jk}^o and Π_{jk}^{α} , we must calculate the curvature tensors.

4. If we calculate $ddA_j - ddA_j = \Omega_j^o A_o + \Omega_j^i A_i$, the curvatures will be given by

$$(4.1) \quad \Omega_j^o = P_{jkl}^o dx^k dx^l + P_{jkl}^o{}^\alpha (dx^k \delta p_\alpha^l - dx^l \delta p_\alpha^k) + P_{jkl}^o{}^{\alpha\beta} \delta p_\alpha^k \delta p_\beta^l,$$

$$(4.2) \quad \Omega_j^i - \delta_j^i \Omega_o^o = P_{jkl}^i dx^k dx^l + P_{jkl}^i{}^\alpha (dx^k \delta p_\alpha^l - dx^l \delta p_\alpha^k),$$

where δp_α^i defined by

$$\delta p_\alpha^i = dp_\alpha^i + \Pi_{jk}^i p_\alpha^j dx^k$$

are contravariant vectors and

$$(4.3) \quad P_{jkl}^o = (\Pi_{jk,l}^o - \Pi_{jk}^o|_a{}^\alpha \Pi_{bl}^a p_\alpha^b) - (\Pi_{jl,k}^o - \Pi_{jl}^o|_a{}^\alpha \Pi_{bk}^a p_\alpha^b) + \Pi_{jk}^a (\Pi_{al}^o - \Pi_{ab}^{\alpha\beta} \Pi_{cl}^b p_\alpha^c) - \Pi_{jl}^a (\Pi_{ak}^o - \Pi_{ab}^{\alpha\beta} \Pi_{ck}^b p_\alpha^c) + (\Pi_{ja,k}^{\alpha\beta} - \Pi_{ja}^{\alpha\beta}|_b{}^\beta \Pi_{ck}^b p_\beta^c) \Pi_{al}^a p_\alpha^d - (\Pi_{ja,l}^{\alpha\beta} - \Pi_{ja}^{\alpha\beta}|_b{}^\beta \Pi_{cl}^b p_\beta^c) \Pi_{ak}^a p_\alpha^d,$$

$$(4.4) \quad P_{jkl}^o{}^\alpha = \Pi_{jk,l}^o|_i{}^\alpha - (\Pi_{jk,l}^{\alpha\beta} - \Pi_{jk}^{\alpha\beta}|_a{}^\beta \Pi_{bl}^a p_\beta^b) - \Pi_{ja}^{\alpha\beta}|_l{}^\beta \Pi_{bk}^a p_\beta^b + \Pi_{jk}^a \Pi_{al}^{\alpha\beta},$$

$$(4.5) \quad P_{jkl}^o{}^{\alpha\beta} = \Pi_{jk,l}^{\alpha\beta}|_i{}^\beta - \Pi_{jl}^{\alpha\beta}|_k{}^\alpha,$$

$$(4.6) \quad P_{jkl}^i = (\Pi_{jk,l}^i - \Pi_{jk}^i|_a{}^\alpha \Pi_{bl}^a p_\alpha^b) - (\Pi_{jl,k}^i - \Pi_{jl}^i|_a{}^\alpha \Pi_{bk}^a p_\alpha^b) + \Pi_{jk}^i \Pi_{al}^i - \Pi_{jl}^i \Pi_{ak}^i + (\Pi_{jk}^o - \Pi_{ja}^{\alpha\beta} \Pi_{bk}^a p_\alpha^b) \delta_l^i - (\Pi_{jl}^o - \Pi_{ja}^{\alpha\beta} \Pi_{bl}^a p_\alpha^b) \delta_k^i - \delta_j^i \{ (\Pi_{kl}^o - \Pi_{ka}^{\alpha\beta} \Pi_{bl}^a p_\alpha^b) - (\Pi_{lk}^o - \Pi_{la}^{\alpha\beta} \Pi_{bk}^a p_\alpha^b) \},$$

$$(4.7) \quad P_{jkl}^i{}^\alpha = \Pi_{jk,l}^i|_i{}^\alpha - \delta_j^i \Pi_{kl}^{\alpha\beta} - \delta_k^i \Pi_{jl}^{\alpha\beta}.$$

The P_{jkl}^o and P_{jkl}^i are generalized homogeneous functions of degree zero of the set p_α^i and $P_{jkl}^o{}^\alpha$, $P_{jkl}^o{}^{\alpha\beta}$ and $P_{jkl}^i{}^\alpha$ are homogeneous function systems with respect to the upper Greek indices and satisfy

$$(4.8) \quad P_{jkl}^o{}^\alpha p_\alpha^j = 0, P_{jkl}^o{}^{\alpha\beta} p_\alpha^k = 0, P_{jkl}^o{}^{\alpha\beta} p_\alpha^l = 0, P_{jkl}^i{}^\alpha p_\alpha^l = 0.$$

During a change of hyperplane at infinity, the P 's are transformed respectively into \bar{P} 's by the following transformation laws:

$$(4.9) \quad \bar{P}_{jkl}^o = P_{jkl}^o - (P_{jkl}^o{}^\alpha - P_{ilk}^o{}^\alpha) \lambda_\alpha + P_{jkl}^o{}^{\alpha\beta} \lambda_\alpha \lambda_\beta - \lambda_i \{ P_{jkl}^i - (P_{jkl}^i{}^\alpha - P_{ilk}^i{}^\alpha) \lambda_\alpha \},$$

$$(4.10) \quad \bar{P}_{jkl}^o{}^\alpha = P_{jkl}^o{}^\alpha - \lambda_i P_{jkl}^i{}^\alpha - P_{jkl}^o{}^{\beta\alpha} \lambda_\beta,$$

$$(4.11) \quad \bar{P}_{jkl}^o{}^{\alpha\beta} = P_{jkl}^o{}^{\alpha\beta},$$

$$(4.12) \quad \bar{P}_{jkl}^i = P_{jkl}^i - (P_{jkl}^i{}^\alpha - P_{ilk}^i{}^\alpha) \lambda_\alpha,$$

$$(4.13) \quad \bar{P}^i{}_{jkl}{}^\alpha = P^i{}_{jkl}{}^\alpha,$$

where we have put $\lambda_\alpha = \lambda_i p_\alpha^i$.

During a change of coordinates, the P 's, as is easily seen, behave as components of tensors.

To determine $\Pi_{jk}^{\alpha\alpha}$, we put the condition $P_{akl}^a{}^\alpha = 0$, which is invariant one as we can see from (4.13). Thus we obtain

$$(4.14) \quad \Pi_{kl}^{\alpha\alpha} = \frac{1}{N+1} \Pi_{ak}^a |_\alpha^\alpha = \frac{1}{N+1} \Gamma_{ak}^a |_\alpha^\alpha,$$

from which $P_{jkl}^{\alpha\alpha} = 0$ and

$$(4.15) \quad P^i{}_{jkl}{}^\alpha = \left[\Gamma_{jk}^i - \frac{1}{N+1} (\delta_j^i \Gamma_{ak}^a + \delta_k^i \Gamma_{aj}^a) - \right. \\ \left. - \frac{1}{N-K} P_\lambda^i \{ \Gamma_{jk}^a |_\alpha^\lambda - \frac{1}{N+1} (\Gamma_{aj}^a |_\lambda^k + \Gamma_{ak}^a |_\lambda^j) \} \right] |_\alpha^\alpha.$$

Thus we can see that the $P^i{}_{jkl}{}^\alpha$ satisfy

$$(4.16) \quad P_{akl}^a{}^\alpha = P_{jal}^a{}^\alpha = P_{ka}^a{}^\alpha = 0.$$

To determine the Π_{jk}^o , we put the condition $P_{jka}^a = 0$, which is invariant one as we can see from (4.12) and (4.16). Thus we obtain

$$(4.17) \quad \Pi_{jk}^o - \Pi_{ja}^{\alpha\alpha} \Pi_{bk}^a p_\alpha^b = - \frac{1}{N^2-1} (N \Pi_{jka}^a + \Pi_{kja}^a)$$

and consequently

$$(4.18) \quad P^i{}_{jkl} = \Pi_{jkl}^i - \frac{1}{N^2-1} (N \Pi_{jka}^a + \Pi_{kja}^a) \delta_l^i + \\ + \frac{1}{N^2-1} (N \Pi_{jla}^a + \Pi_{lja}^a) \delta_k^i + \frac{1}{N+1} \delta_j^i (\Pi_{kla}^a - \Pi_{lka}^a),$$

where

$$(4.19) \quad \Pi_{jkl}^i = (\Pi_{jk,l}^i - \Pi_{jk}^i |_\alpha^\alpha \Pi_{bl}^a p_\alpha^b) - \\ - (\Pi_{jl,k}^i - \Pi_{jl}^i |_\alpha^\alpha \Pi_{bk}^a p_\alpha^b) + \Pi_{jk}^a \Pi_{al}^i - \Pi_{jl}^a \Pi_{ak}^i.$$

Thus all the coefficients of the projective connection are determined from Γ_{jk}^i by projectively invariant conditions.

5. Among the semi-natural frames of reference, we can select a special one for which $*\Pi_{ak}^a = 0$. We shall call natural frame of reference the semi-natural frame of reference for which these relations hold good. As we can see from (2.7), to obtain the natural frame of reference from an arbitrary semi-natural frame

of reference, we must choose λ_j as

$$\lambda_j = -\frac{1}{N+1} \Pi_{aj}^a = -\frac{1}{N+1} \Gamma_{aj}^a$$

and effect the transformation

$$*A_o = A_o, \quad *A_j = -\frac{1}{N+1} \Gamma_{aj}^a A_o + A_j.$$

Thus, as we can see from (2.7), the coefficients of the projective connection with respect to the natural frame of reference are given by

$$(5.1) \quad \left\{ \begin{array}{l} * \Pi_{jk}^o = \Pi_{jk}^o - \frac{1}{N+1} (\Gamma_{aj, k}^a - \Gamma_{ab}^a \Pi_{jk}^b + \frac{1}{N+1} \Gamma_{aj}^a \Gamma_{bk}^b), \quad * \Pi_{jk}^{o\alpha} = 0, \\ * \Pi_{jk}^i = \Gamma_{jk}^i - \frac{1}{N+1} (\delta_j^i \Gamma_{ak}^a + \delta_k^i \Gamma_{aj}^a) - \\ \quad - \frac{1}{N-K} p_\lambda^i \{ \Gamma_{jk}^a |_\alpha^\lambda - \frac{1}{N+1} (\Gamma_{aj}^a |_\alpha^\lambda + \Gamma_{ak}^a |_\alpha^\lambda) \}. \end{array} \right.$$

The coefficients $* \Pi_{jk}^i$ coincide with those used by the other writers.

If we denote by $*P$ the components of the curvatures with respect to the natural frame of reference, then we have

$$(5.2) \quad *P_{jkl}^o = P_{jkl}^o + \frac{1}{N+1} (P_{jkl}^{o\alpha} - P_{jlk}^{o\alpha}) \Gamma_{ab}^a p_\alpha^b + \\ + \frac{1}{N+1} \Gamma_{ai}^a \{ P_{jkl}^i + \frac{1}{N+1} (P_{jkl}^{i\alpha} - P_{jlk}^{i\alpha}) \Gamma_{bc}^b p_\alpha^c \},$$

$$(5.3) \quad *P_{jkl}^{o\alpha} = P_{jkl}^{o\alpha} + \frac{1}{N+1} \Gamma_{ai}^a P_{jkl}^{i\alpha},$$

$$(5.4) \quad *P_{jkl}^i = P_{jkl}^i + \frac{1}{N+1} (P_{jkl}^{i\alpha} - P_{jlk}^{i\alpha}) \Gamma_{ab}^a p_\alpha^b,$$

$$(5.5) \quad *P_{jkl}^{i\alpha} = P_{jkl}^{i\alpha} = * \Pi_{jk}^i |_\alpha^\alpha.$$

Thus, $*P_{jkl}^o$, $*P_{jkl}^{o\alpha}$ and $*P_{jkl}^i$ are not components of tensors while $*P_{jkl}^{i\alpha}$ are components of a projective tensor. The $*P_{jkl}^i$ coincide with the W_{jkl}^i of J. Douglas. But, its transformation law is, as is easily seen from (5.4),

$$(5.6) \quad * \bar{P}_{jkl}^i = \frac{\partial \bar{x}^i}{\partial x^a} \frac{\partial x^b}{\partial \bar{x}^j} \frac{\partial x^c}{\partial \bar{x}^k} \frac{\partial x^d}{\partial \bar{x}^l} * P_{bcd}^a + * \bar{\Pi}_{jk}^i |_\alpha^\alpha \bar{\theta}_m \bar{p}_\alpha^m - * \bar{\Pi}_{jl}^i |_\alpha^\alpha \bar{\theta}_m \bar{p}_\alpha^m,$$

where

$$\bar{\theta}_m = \frac{1}{N+1} \frac{\partial \log \Delta}{\partial \bar{x}^m}, \quad \bar{p}_\alpha^m = \frac{\partial \bar{x}^m}{\partial u^\alpha} \quad \left(\Delta = \left| \frac{\partial x}{\partial \bar{x}} \right| \right).$$

This is the transformation law of W_{jki}^i obtained by R. S. Clark [2].

It will be remarked here that, when $K = 1$, the above discussion will be reduced to that given already by one of the present authors [8].

6. Now, returning to the semi-natural frame of reference, we have (3.3) along a K -spread. If we put

$$(6.1) \quad \underset{21}{dd}A_\alpha - \underset{12}{dd}A_\alpha = \Omega_\alpha^o A_\alpha, \quad \underset{21}{dd}A_\beta - \underset{12}{dd}A_\beta = \Omega_\beta^o A_\alpha + \Omega_\beta^\alpha A_\alpha.$$

the curvature tensors of the projective connection induced on a K -spread are given by

$$(6.2) \quad \Omega_\beta^o = P_{\beta\gamma\delta}^o du^\gamma du^\delta, \quad \Omega_\beta^\alpha - \delta_\beta^\alpha \Omega_\alpha^o = P_{\beta\gamma\delta}^\alpha du^\gamma du^\delta.$$

On the other hand, as we have

$$A_\alpha = A_o, \quad A_\beta = p_\beta^o A_o + p_\beta^i A_i,$$

we find

$$\begin{cases} \underset{21}{dd}A_\alpha - \underset{12}{dd}A_\alpha = \underset{21}{dd}A_o - \underset{12}{dd}A_o, \\ \underset{21}{dd}A_\beta - \underset{12}{dd}A_\beta = p_\beta^o (\underset{21}{dd}A_o - \underset{12}{dd}A_o) + p_\beta^j (\underset{21}{dd}A_j - \underset{12}{dd}A_j), \end{cases}$$

from which

$$(6.3) \quad \begin{cases} \Omega_\alpha^o A_o = \Omega_o^o A_o, \\ \Omega_\beta^o A_o + \Omega_\beta^\alpha A_\alpha = p_\beta^o \Omega_o^o A_o + p_\beta^j (\Omega_j^o A_o + \Omega_j^\alpha A_\alpha). \end{cases}$$

Thus, remembering that $A_\alpha = p_\alpha^o A_o + p_\alpha^i A_i$, we find

$$p_\alpha^i \Omega_\beta^\alpha = \Omega_j^i p_\beta^j,$$

or

$$p_\alpha^i (\Omega_\beta^\alpha - \delta_\beta^\alpha \Omega_\alpha^o) = (\Omega_j^i - \delta_j^i \Omega_o^o) p_\beta^j,$$

from which, δp_α^i being linear combinations of p_α^i along a K -spread, we find

$$(6.4) \quad p_\alpha^i P_{\beta\gamma\delta}^\alpha = P_{jki}^i p_\beta^j p_\gamma^k p_\delta^l,$$

where $P_{\beta\gamma\delta}^\alpha$ are components of the projective curvature tensor induced on the K -spreads. In a space of K -spreads, the equations of K -spreads being completely integrable, the above equations must be satisfied identically for any K -spread.

Now, we shall seek for the condition that the equations of the K -spreads can be reduced to

$$\frac{\partial^2 x^i}{\partial u^\beta \partial u^\gamma} = 0$$

by suitable coordinate and parameter transformations.

If the equations are reduced to the above form, then we must have

$$(6.5) \quad P_{jkl}^i \alpha \equiv *II_{jk}^i | \alpha = 0.$$

As $P_{jkl}^i \alpha$ is a projective tensor, the vanishing of $P_{jkl}^i \alpha$ is a condition invariant under coordinate and parameter transformations.

Conversely, if $P_{jkl}^i \alpha \equiv *II_{jk}^i | \alpha = 0$, then $*II_{jk}^i$ and consequently

$$*P_{jkl}^i = P_{jkl}^i$$

are the functions of the point only. Thus applying the operator $| \alpha$ to (6.4), we obtain

$$(N - K) P_{\beta\gamma\delta}^{\lambda} = 0,$$

from which

$$P_{jkl}^i p_\beta^j p_\gamma^k p_\delta^l = 0 \quad \text{and consequently} \quad P_{jkl}^i = 0.$$

Thus it is proved that the necessary and sufficient condition that the space of K -spreads be projectively flat is that $P_{jkl}^i \alpha = 0$, that is to say, $*II_{jk}^i$ be independent of p_α^i .

REFERENCES.

J. DOUGLAS

- [1] Systems of K -dimensional manifolds in an N -dimensional space. *Math. Ann.*, 105 (1931), 707—733.

R. S. CLARK

- [2] Projective collineations in a space of K -spreads. *Proc. Cambridge Philosophical Soc.*, 41 (1945), 210—223.

CHIH-TA YEN

- [3] Sur une connexion projective normale associée à un système de variétés à k dimensions. *C.R.*, 227 (1948), 461—462.
 [4] Sur les systèmes de variétés à k dimensions et la connexion projective normale associée. *C.R.*, 228 (1949), 1844—1846.

HSIEN-CHUNG WANG

- [5] Axiom of the plane in a general space of paths. *Annals of Math.*, 49 (1948), 731—737.

S. S. CHERN

- [6] A generalization of the projective geometry of linear spaces. *Proc. Nat. Acad. Sci., U.S.A.*, 29 (1943), 38—43.

K. YANO

- [7] Les espaces à connexion projective et la géométrie projective des paths. *Annales Scientifiques de l'Université de Jassy*, 24 (1938), 395—464.
- [8] Les espaces d'éléments linéaires à connexion projective normale et la géométrie projective générale des paths. *Proc. Physico-Math. Soc. of Japan*, 24 (1942), 9—25.

The Institute for Advanced Study, N.J., U.S.A.
and Yamanashi University, Kofu, Japan.

(Oblatum 19-6-51).