

# COMPOSITIO MATHEMATICA

C. DE VROEDT

## **Metrical problems concerning continued fractions**

*Compositio Mathematica*, tome 16 (1964), p. 191-195

[http://www.numdam.org/item?id=CM\\_1964\\_\\_16\\_\\_191\\_0](http://www.numdam.org/item?id=CM_1964__16__191_0)

© Foundation Compositio Mathematica, 1964, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# Metrical problems concerning continued fractions \*

by

C. de Vroedt

## § 1.

A) Let  $\theta$  be a real number ( $0 \leq \theta \leq 1$ ).

We expand  $\theta$  in the continued fraction:

$$(1) \quad \theta = \{a_1, a_2, a_3, \dots\} = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

with integer partial quotients  $a_n \geq 1$ .

For irrational  $\theta$  we define the sequence of functions  $\theta_n$  as follows:

$$(2) \quad \begin{cases} \theta_0 = \theta \\ \theta_{n+1} = \frac{1}{\theta_n} - \left[ \frac{1}{\theta_n} \right] \end{cases}$$

( $[\alpha]$  means the largest integer  $\leq \alpha$ ).

From the definition of the function  $\theta_n$  it follows that:

$$(3) \quad a_{n+1} = \left[ \frac{1}{\theta_n} \right].$$

B) The first known metrical problem concerning continued fractions is due to Gauss. If  $m_n(x)$  ( $0 \leq x \leq 1$ ;  $n \geq 0$ ) denotes the measure of the set of those  $\theta$  for which  $\theta_n \leq x$ , Gauss asserts:

$$(4) \quad \lim_{n \rightarrow \infty} m_n(x) = \frac{\log(1+x)}{\log 2}.$$

In 1928, R. Kuzmin [3] has given a proof of (4) including an estimate of the remainder which is important for further develop-

\* Nijenrode lecture.

ments. It is to be remarked that in 1929 P. Lévy, independent of Kuzmin's proof, published another one.

In later years A. Khintchine [2], using Kuzmin's result, proved among other things the following theorem:

Let  $f(p)$  be a nonnegative function of the positive integer  $p$  such that  $f(p) \leq K p^{\frac{1}{2}-\beta}$  ( $K$  and  $\beta$  being positive constants). Then for almost all  $\theta$  we have:

$$(5) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(a_n) = \sum_{p=1}^{\infty} f(p) \frac{\log \left\{ 1 + \frac{1}{p(p+2)} \right\}}{\log 2}.$$

Putting in this theorem  $f(x) = \log x$ , Khintchine proved:

$$(6) \quad \lim_{N \rightarrow \infty} \sqrt[N]{a_1 \cdot a_2 \cdot \dots \cdot a_N} = \prod_{p=1}^{\infty} \left\{ 1 + \frac{1}{p(p+2)} \right\}^{\log p / \log 2}$$

for almost all  $\theta$ .

## § 2.

A. Let  $f(x)$  be a real function on  $0 \leq x \leq 1$ . Let  $D$  be a division of the interval  $\langle 0, 1 \rangle$  with dividing points  $x_0 (= 0), x_1, x_2, \dots, x_p (= 1)$ .

Further put

$$\sigma_p = \sup f(x) - \inf f(x)$$

where both,  $\sup$  and  $\inf$ , are to be extended over the closed interval  $x_n \leq x \leq x_{p+1}$ .

Then

$$(7) \quad \varphi(D) = \sum_{p=0}^{P-1} \sigma_p (x_{p+1} - x_p)$$

denotes the "oscillation of  $f(x)$  for the division  $D$ ".

In our thesis [4] we proved the following theorem:

Let  $D_k$  be the division of  $\langle 0, 1 \rangle$  with dividing points

$$h_0^k (= 0), h_1^k \left( = \frac{1}{2^k} \right), h_2^k \left( = \frac{2}{2^k} \right), \dots, h_{2^k}^k (= 1).$$

Let  $f(x)$  be a real function on  $0 \leq x \leq 1$ , such that for the sequence  $\{D_k\}$  of divisions  $D_k$  of  $\langle 0, 1 \rangle$  the corresponding oscillations satisfy

$$(8) \quad \sum_{k=1}^{\infty} \varphi(D_k) < \infty.$$

Then for any couple of integers  $M \geq 0, N \geq 1$  we have:

$$(9) \quad I \stackrel{\text{def}}{=} \int_0^1 \left\{ \sum_{n=M+1}^{M+N} (f(\theta_n) - \int_0^1 \frac{f(x)}{(1+x) \log 2} dx) \right\}^2 d\theta \leq K_1 N,$$

$K_1$  being a constant depending only on  $f(x)$ .

B) We give a sketch of the proof. From the conditions imposed upon  $f(x)$  it follows that  $f(x)$  is bounded and that, without loss of generality, we may limit ourselves to the case  $f(x) > 0$ . Therefore:

$$(10) \quad 0 < f(x) < B$$

and if we put:

$$\int_0^1 \frac{f(x)}{(1+x) \log 2} dx = P$$

we have:

$$(11) \quad 0 < P < B.$$

Working out (9) we get:

$$(12) \quad I = \sum_{n=M+1}^{M+N} \int_0^1 (f(\theta_n) - P)^2 d\theta + 2 \sum_{n=M+1}^{M+N-1} \sum_{\substack{m=M+2 \\ m>n}}^{M+N} \int_0^1 (f(\theta_n) - P)(f(\theta_m) - P) d\theta.$$

Now:

$$(13) \quad \sum_{n=M+1}^{M+N} \int_0^1 (f(\theta_n) - P)^2 d\theta \leq B^2 N$$

and

$$(14) \quad \sum_{\substack{n=M+1 \\ m=n+1}}^{M+N} \int_0^1 (f(\theta_n) - P)(f(\theta_m) - P) d\theta \leq B^2 N.$$

Further we have:

$$(15) \quad \int_0^1 (f(\theta_n) - P)(f(\theta_m) - P) d\theta = \int_0^1 f(\theta_n) f(\theta_m) d\theta - P \int_0^1 f(\theta_n) d\theta - P \int_0^1 f(\theta_m) d\theta + P^2.$$

If

$$m_n(x) \stackrel{\text{def}}{=} mE\{\theta; \theta_n \leq x\},$$

R. Kuzmin [3] proved:

$$(16) \quad \frac{dm_n(x)}{dx} = \frac{1}{(1+x) \log 2} + \delta A e^{-\alpha \sqrt{n}}$$

with  $|\delta| < 1$ , whereas  $A$  and  $\alpha$  are positive constants.

Now it is easily shown that

$$(17) \quad \int_0^1 f(\theta_n) d\theta = \int_0^1 f(x) \cdot \frac{dm_n(x)}{dx} dx = P + \delta_1 A B e^{-\alpha\sqrt{n}}$$

with  $|\delta_1| < 1$ .

As the series  $\sum_{n=1}^{\infty} e^{-\alpha\sqrt{n}}$  converges we get:

$$(18) \quad \sum_{n=M+1}^{M+N-2} \sum_{\substack{m=M+3 \\ m>n}}^{M+N} \left( -P \int_0^1 f(\theta_n) d\theta - P \int_0^1 f(\theta_m) d\theta + P^2 \right) = \\ \sum_{n=M+1}^{M+N-2} \sum_{\substack{m=M+3 \\ m>n}}^{M+N} (-P^2) + \delta_2 K_2 N$$

with  $|\delta_2| < 1$ , whereas  $K_2$  is a positive constant.

In order to estimate

$$\sum_{n=M+1}^{M+N-2} \sum_{\substack{m=M+3 \\ m>n}}^{M+N} \int_0^1 f(\theta_n) f(\theta_m) d\theta$$

we define:

$$(19) \quad k = \left[ \frac{1}{2}(m-n) \right].$$

Then we can prove:

$$(20) \quad \int_0^1 f(\theta_n) f(\theta_m) d\theta = (P + \delta_3 A B e^{-\alpha\sqrt{m-n-k}}) (P + \delta_4 A B e^{-\alpha\sqrt{n}} + 2\delta_5 \varphi(k))$$

where  $\varphi(k)$  means the oscillation of  $f(x)$  for the division  $D_k$ .

Because of our choice of  $k$ (19) and of the condition (8) imposed upon  $f(x)$  we get:

$$(21) \quad \sum_{n=M+1}^{M+N-2} \sum_{\substack{m=M+3 \\ m>n}}^{M+N} \int_0^1 f(\theta_n) f(\theta_m) d\theta = \sum_{n=M+1}^{M+N-2} \sum_{\substack{m=M+3 \\ m>n}}^{M+N} (P^2) + \delta_6 K_3 N$$

with  $|\delta_6| < 1$ , whereas  $K_3$  is a positive constant.

Combination of (12), (13), (14), (15), (18) and (21) proves our theorem.

C) On the result (9) we apply a theorem of I. S. Gál and J. F. Koksma [1] (Theorem III with  $S \equiv$  the set  $0 \leq \theta \leq 1$ ,  $\psi(N) = N$  and  $p = 2$ ). Then we get:

Let  $f(x)$  be a real function on  $0 \leq x \leq 1$ , satisfying the condition (8). Then for almost all  $\theta$  we have:

$$(22) \quad \sum_{n=1}^N f(\theta_n) - N \int_0^1 \frac{f(x)}{(1+x) \log 2} dx = o(N^{\frac{1}{2}} \log^{(3+\varepsilon)/2} N).$$

## § 3.

Shortly, in *Indagationes Mathematicae*, we will publish a note in which we shall prove the results (9) and (22) for functions satisfying somewhat weaker conditions.

As applications then we will get the following refinements of the results (5) and (6) of A. Khintchine.

1) Let  $f(p)$  be a nonnegative function of the positive integer  $p$  such that  $f(p) \leq Kp^{\frac{1}{2}-\beta}$  ( $K$  and  $\beta$  being positive constants). Then for almost all  $\theta$  we have:

$$(23) \quad \sum_{n=1}^N f(a_n) - N \sum_{p=1}^{\infty} f(p) \frac{\log \left\{ 1 + \frac{1}{p(p+2)} \right\}}{\log 2} = o(N^{\frac{1}{2}} \log^{(3+\varepsilon)/2} N).$$

2) For almost all  $\theta$  we have:

$$(24) \quad \sqrt[N]{a_1 \cdot a_2 \cdot \dots \cdot a_N} - \prod_{p=1}^{\infty} \left\{ 1 + \frac{1}{p(p+2)} \right\}^{\log p / \log 2} = o(N^{-\frac{1}{2}} \log^{(3+\varepsilon)/2} N).$$

## REFERENCES

- [1] I. S. GAL et J. F. KOKSMA  
Sur l'ordre de grandeur des fonctions sommables. *Indag. Math.* (1950; Vol. XII; p. 638–653).
- [2] A. KHINTCHINE  
*Kettenbrüche*. B. G. Teubner, Leipzig, 1956.
- [3] R. KUZMIN  
Sur un problème de Gauss, *Atti del congr. intern. Bologna*. (1928; Vol. 6; p. 83).
- [4] C. DE VROEDT  
Metrical problems concerning continued fractions. Thesis. Amsterdam 1960.
- [5] C. DE VROEDT  
Measure-theoretical investigations concerning continued fractions. *Indag. Math.* (1962; Vol. XXIV; p. 583–591).

(Oblatum 29-5-63).