

COMPOSITIO MATHEMATICA

YUNG-MING CHEN

Note on approximation theorems

Compositio Mathematica, tome 17 (1965-1966), p. 36-39

http://www.numdam.org/item?id=CM_1965-1966__17__36_0

© Foundation Compositio Mathematica, 1965-1966, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Note on approximation theorems

by

Yung-ming Chen

This note is concerned with three approximation theorems for a class of definite integrals and sine series with monotone decreasing coefficients. S. Aljančić, R. Bojanić and M. Tomić [1] have considered previously the asymptotic value of the definite integral

$$\Phi(\lambda) = \int_a^b f(t)L(\lambda t)dt, \quad 0 \leq a < b < \infty, \quad \lambda \rightarrow \infty,$$

where $f(t) \in L(a, b)$ and $L(t)$ is a slowly increasing function in the sense of Karamata. In this note, we shall consider the asymptotic behaviour of definite integrals of a more general type:

$$\Phi(\lambda) = \int_0^\infty f(t)\phi(\lambda t)dt,$$

where $\phi(t)$ is such that $\phi(t)/t^{p_1}$ is non-decreasing and $\phi(t)/t^{p_2}$ is non-increasing as t is increasing in $(0, \infty)$ for $0 \leq p_1 \leq p_2 < \infty$. In another paper [2], Aljančić, Bojanić and Tomić have also considered the asymptotic behaviour of $g(x) = \sum_1^\infty \lambda_n \sin nx$, $\lambda_n \searrow 0$, as $x \rightarrow +0$, under the condition that $\lambda_n \sim An^{-\alpha}L(n)$, as $n \rightarrow \infty$. We shall now consider the asymptotic behaviour of the same series when λ_n is asymptotically defined by a more general class of functions.

NOTATION. By $\phi(x) \sim [p_1, p_2]$, $0 \leq p_1 \leq p_2 < \infty$ or $-\infty < p_1 \leq p_2 \leq 0$, we denote the non-negative even function $\phi(x)$, such that $\phi(x)x^{-p_1}$ is non-decreasing and $\phi(x)x^{-p_2}$ is non-increasing, as x is increasing in $(0, \infty)$. By $\phi(x) \sim \langle p_1, p_2 \rangle$, we mean $\phi(x) \sim [p_1 + \varepsilon, p_2 - \varepsilon]$ for some $\varepsilon > 0$. We define $\phi(x) \sim [p_1, p_2 \rangle$, $\phi(x) \sim \langle p_1, p_2]$ in a similar way. By $a(x) \simeq b(x)$ and $a(x) \succ b(x)$, $x \rightarrow c$, we mean $a(x)/b(x) \rightarrow 1$ and $K_1 < a(x)/b(x) < K_2$, respectively, as $x \rightarrow c$.

THEOREM 1. If $\phi(x) \sim [p_1, p_2]$, $0 \leq p_1 \leq p_2 < \infty$, and if $f(t) \geq 0$, $t^{p_1}f(t) \in L(0, 1)$, $t^{p_2}f(t) \in L(1, \infty)$, then $\Phi(\lambda) \asymp \phi(\lambda)$ as $\lambda \rightarrow \infty$. More precisely, we have

$$(1) \quad K_1(f, p_1, p_2)\phi(\lambda) \leq \Phi(\lambda) = \int_0^\infty f(t)\phi(\lambda t)dt \leq K_2(f, p_1, p_2)\phi(\lambda),$$

uniformly for $0 < \lambda < \infty$, where

$$(2) \quad K_1 = \int_0^1 t^{p_2} f(t) dt + \int_1^\infty t^{p_1} f(t) dt, \quad K_2 = \int_0^1 t^{p_1} f(t) dt + \int_1^\infty t^{p_2} f(t) dt.$$

In fact, if $t > 1$, then $\phi(t\lambda)(t\lambda)^{-p_2} \leq \phi(\lambda)\lambda^{-p_2}$, and $\phi(\lambda)\lambda^{-p_1} \leq \phi(t\lambda)(t\lambda)^{-p_1}$, i.e. $t^{p_1}\phi(\lambda) \leq \phi(t\lambda) \leq t^{p_2}\phi(\lambda)$. Similarly, if $0 < t < 1$, then $t^{p_2}\phi(\lambda) \leq \phi(t\lambda) \leq t^{p_1}\phi(\lambda)$. It follows that

$$(3) \quad \begin{aligned} \phi(\lambda) \left\{ \int_0^1 t^{p_2} f(t) dt + \int_1^\infty t^{p_1} f(t) dt \right\} &\leq \int_0^\infty f(t) \phi(\lambda t) dt \\ &\leq \phi(\lambda) \left\{ \int_0^1 t^{p_1} f(t) dt + \int_1^\infty t^{p_2} f(t) dt \right\}. \end{aligned}$$

THEOREM 2. Let

$$(4) \quad g(x) = \sum_1^\infty \lambda_n \sin nx,$$

where λ_n decreases steadily to zero. If $\phi(x) \sim \langle -1, 0 \rangle$ and if $\lambda_n \simeq \phi(n)$, as $n \rightarrow \infty$, then $g(x) \asymp 1/x \phi(1/x)$, as $x \rightarrow +0$.

PROOF. Since $\phi(x) \sim \langle -1, 0 \rangle$, $\varepsilon\phi(x)/x \leq -\phi'(x) \leq (1-\varepsilon)\phi(x)/x$ for some $\varepsilon > 0$. It follows that

$$g(x) = \sum_{n=1}^\infty \lambda_n \sin nx = \sum_{1 \leq n \leq [1/x]} + \sum_{n > 1/x} = T_1 + T_2,$$

where

$$(5) \quad \begin{cases} |T_1| \leq K \int_1^{1/x} \phi(t) dt \leq K \left(\frac{1}{x}\right) \phi\left(\frac{1}{x}\right) - K \int_1^{1/x} t\phi'(t) dt \\ \leq K \left(\frac{1}{x}\right) \phi\left(\frac{1}{x}\right) + (1-\varepsilon) \int_1^{1/x} \phi(t) dt \leq K \left(\frac{1}{x}\right) \phi\left(\frac{1}{x}\right), \end{cases}$$

as $x \rightarrow +0$.

From Abel's transformation, it is easy to see that

$$(6) \quad |T_2| = \left| \sum_{n > 1/x} \lambda_n \sin nx \right| \leq K \phi\left(\frac{1}{x}\right) x^{-1}.$$

It remains to show that $|g(x)| > K(1/x)\phi(1/x)$, as $x \rightarrow +0$. In fact,

$$(7) \quad \begin{cases} g(x) = \sum_1^\infty \Delta\lambda_n \frac{\sin^2(n+\frac{1}{2})\frac{x}{2}}{\sin \frac{x}{2}} - \frac{\lambda_1}{2} \operatorname{tg} \frac{x}{4} \\ = \sum_1^\infty \Delta\lambda_n \frac{\sin^2(n+\frac{1}{2})\frac{x}{2}}{\sin \frac{x}{2}} + o\left\{\frac{1}{x} \phi\left(\frac{1}{x}\right)\right\} = I + II, \end{cases}$$

where

$$(8) \quad \left\{ \begin{aligned} I &> \frac{K}{x} \sum_{\pi/2x \leq n \leq 3\pi/2x} \Delta\phi(n) \geq \frac{K}{x} \left\{ \phi\left(\frac{\pi}{2x}\right) - \phi\left(\frac{3\pi}{2x}\right) \right\} \\ &= \frac{K}{x} \left\{ \phi\left(\frac{\pi}{2x}\right) - \phi\left(\frac{\lambda\pi}{2x}\right) \right\} \quad (\lambda = \frac{3}{2}) \\ &\geq \frac{K}{x} (1 - \lambda^{-\varepsilon}) \phi\left(\frac{\pi}{2x}\right) \geq K \left(\frac{1}{x}\right) \phi\left(\frac{1}{x}\right), \end{aligned} \right.$$

and $II = o\{1/x\phi(1/x)\}$, as $x \rightarrow +0$. Hence $|g(x)| > K(1/x)\phi(1/x)$, as $x \rightarrow +0$.

THEOREM 3. If $\phi(x) \sim \langle -2, -1 \rangle$, $\lambda_n \searrow 0$, $\lambda_n \cong \phi(n)$, as $n \rightarrow \infty$, then $g(x) \asymp (1/x)\phi(1/x)$, as $x \rightarrow +0$.

PROOF. From (7), together with $\phi(x) \sim \langle -2, -1 \rangle$, when $x \simeq 1/n$ we also have

$$(9) \quad \left\{ \begin{aligned} |g(x)| &= \left| \sum_1^\infty \Delta\lambda_n \frac{\sin^2(n+1) \frac{x}{2}}{\sin \frac{x}{2}} \right| + o\left\{ \frac{1}{x} \phi\left(\frac{1}{x}\right) \right\} \\ &\leq \frac{1}{x} \sum_1^n \Delta\lambda_n n^2 x^2 + \frac{\lambda_n}{x} \\ &\leq x \sum_1^n k^2 \Delta\lambda_k + \frac{K}{x} \phi\left(\frac{1}{x}\right) \\ &\leq x \sum_1^n (2k-1)\lambda_k + \frac{K}{x} \phi\left(\frac{1}{x}\right) \\ &\leq Kx \sum_1^n (2k-1)\phi(k) + \frac{K}{x} \phi\left(\frac{1}{x}\right). \end{aligned} \right.$$

Write $\tau(x) = x\phi(x)$, then $\tau(x) \sim \langle -1, 0 \rangle$, since $\tau(x)x^{1-\varepsilon}$ is non-decreasing. It follows that

$$(10) \quad \int_1^n \tau(x) dx = n\tau(n) - \tau(1) - \int_1^n x\tau'(x) dx.$$

But since $x^{1-\varepsilon}\tau(x)$ is non-decreasing, $(1-\varepsilon)x^{-\varepsilon}\tau(x) + x^{1-\varepsilon}\tau'(x) \geq 0$, and $-x\tau'(x) \leq (1-\varepsilon)\tau(x)$. Substituting this in (10) and it follows then from (9) that

$$(11) \quad \int_1^n \tau(x) dx \leq Kn\tau(n), \quad |g(x)| \leq Kx \cdot n^2 \phi(n) \leq K \left(\frac{1}{x}\right) \phi\left(\frac{1}{x}\right),$$

as $x \simeq 1/n \rightarrow 0$, $n \rightarrow \infty$. By slight modification in (7) and (8), it may be readily shown that

$$(12) \quad |g(x)| > K \left(\frac{1}{x}\right) \phi \left(\frac{1}{x}\right).$$

Hence $g(x) \asymp (1/x)\phi(1/x)$, as $x \rightarrow +0$.

Added in proof. The author has just observed that the argument in the proof of Theorem 3 actually gives a more general result, viz. with $\phi(x) \sim \langle -2, -1 \rangle$ replaced by $\phi(x) \sim \langle -2, 0 \rangle$. This includes both Theorem 2 and Theorem 3 as particular cases.

REFERENCES

S. ALJANČIĆ, R. BOJANIĆ and M. TOMIĆ,

- [1] Sur la valeur asymptotique d'une classe des intégrales définies. Acad. Serbe Sci. Publ. Inst. Math. 7, 81–94 (1954).

S. ALJANČIĆ, R. BOJANIĆ and M. TOMIĆ,

- [2] Sur le comportement asymptotique au voisinage de zéro des séries trigonométriques de sinus à coefficients monotones. Acad. Serbe Sci. Publ. Inst. Math. 10, 101–120 (1956).

(Oblatum 4-1-62).