

COMPOSITIO MATHEMATICA

V. M. BHISE

Certain properties of Meijer-Laplace transform

Compositio Mathematica, tome 18, n° 1-2 (1967), p. 1-6

http://www.numdam.org/item?id=CM_1967__18_1-2_1_0

© Foundation Compositio Mathematica, 1967, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

'Certain properties of Meijer-Laplace transform'

by

V. M. Bhise

1

A generalization of the classical Laplace transform

$$(1.1) \quad \phi(p) = p \int_0^{\infty} e^{-pt} f(t) dt$$

has been introduced by the author [1, p. 57] in the form

$$(1.2) \quad \phi(p) = p \int_0^{\infty} G_{m, m+1}^{m+1, 0} \left(pt / \begin{matrix} n_1 + a_1, \dots, n_m + a_m \\ n_1, \dots, n_m, \sigma \end{matrix} \right) f(t) dt$$

and denoted symbolically by $\phi(p) = G[f(t); a_m, n_m, \sigma]$. Certain rules and recurrence relations have also been established [2] for the transform (1.2), which we shall call as Meijer-Laplace transform. Setting $a_j = 0, j = 1, 2, \dots, m-1$.

(i) and $a_m = 0, \sigma = 0$, (1.2) reduces to (1.1)

(ii) and $a_m = -m' - k, n_m = m' - k, \sigma = -m' - k$ we get (1.2) reduced to Meijer's transform [3, p. 730],

$$(1.3) \quad \phi(p) = p \int_0^{\infty} (pt)^{-k-\frac{1}{2}} e^{-\frac{1}{2}pt} W_{k+\frac{1}{2}, m'}(pt) f(t) dt$$

(iii) and $a_m = \frac{1}{2} - m' - k, n_m = 2m', \sigma = 0$ we get (1.2) reduced to Verma's transform [4, p. 209]

$$(1.4) \quad \phi(p) = p \int_0^{\infty} (pt)^{m'-\frac{1}{2}} e^{-\frac{1}{2}pt} W_{k, m'}(pt) f(t) dt$$

We denote (1.1), (1.3) and (1.4) symbolically by

$$\phi(p) \doteq f(t), f(t) \xrightarrow{\frac{k+\frac{1}{2}}{m'}} \phi(p) \text{ and } \phi(p) = W[f(t); k, m']$$

respectively.

In this paper we have obtained a theorem for the Meijer-Laplace transform (1.2), which is a generalization of the well-known Tricomi's theorem [5, p. 564]. Various particular cases have been given and the theorem is illustrated by an example.

2

THEOREM: If

$$\phi(p) = G[f(y); a_m, n_m, \sigma]$$

then,

$$(2.1) \quad p^{n/s-\lambda} \phi(p^{-n/s}) = G \left[t^\lambda \int_0^\infty F(t, y) f(y) dy; \alpha_\mu, \eta_\mu, \rho \right]$$

where ¹

$$F(t, y) = \frac{(2\pi)^{(n-s)/2} \cdot s^{\sigma-\sum a_i + \frac{1}{2}}}{n^{\rho-\sum \alpha_j + \frac{1}{2} + \lambda}} G_{n\mu+sm, sm+s+n\mu+n}^{sm+s, n\mu} \\ \times \left(\frac{y^s t^n}{s^n n^n} \left/ \begin{array}{c} \frac{-\eta_j - \alpha_j - \lambda + l}{n}, \frac{n_i + a_i + \nu}{s} \\ \frac{n_i + \nu}{s}, \frac{\sigma + \nu}{s}, \frac{-\eta_j - \lambda + l}{n}, \frac{-\rho - \lambda + l}{n} \end{array} \right. \right)$$

provided n, s are positive integers $s > n$; $|\arg y^s/p^n| < (s-n)\pi/2$; $\min(\operatorname{Re} \eta_j, \operatorname{Re} \rho) + n \min(\operatorname{Re} n_i/s, \operatorname{Re} \sigma/s) > -1 - \operatorname{Re} \lambda > -\operatorname{Re}(\eta_j + \alpha_j + \lambda) - 2$ for $i = 1, 2, \dots, m$; $j = 1, 2, \dots, \mu$; $l = 0, 1, \dots, n-1$; $\nu = 0, 1, \dots, s-1$. the integral in (2.1) is absolutely and uniformly convergent and the Meijer-Laplace transform of $|t^\lambda \int_0^\infty F(t, y) dy|$ exists.

PROOF: From (1.2), replacing p by $p^{-n/s}$ and multiplying by $p^{n/s-\lambda}$, we obtain

$$(2.2) \quad p^{n/s-\lambda} \phi(p^{-n/s}) = \int_0^\infty p^{-\lambda} G_{m, m+1}^{m+1, 0} \left(yp^{-n/s} \left/ \frac{n_i + a_i}{n_i, \sigma} \right. \right) f(y) dy; \\ i = 1, 2, \dots, m.$$

But we have from [6]

¹ For the sake of brevity the symbol

$$G_{\beta n+m, 2\beta+sm}^{2\beta, \beta n} \left(x \left/ \begin{array}{c} \frac{\alpha_j + \nu}{\beta}, c_i + d_i \\ \frac{\pm b + \nu}{\beta}, \frac{\alpha_i + l}{s} \end{array} \right. \right); \quad i = 1, 2, \dots, m; j = 1, 2, \dots, n; \\ \nu = 0, 1, \dots, \beta-1; l = 0, 1, \dots, s-1$$

has been employed to denote

$$G_{\beta n+m, 2\beta+sm}^{2\beta, \beta n} \left(x \left/ \begin{array}{c} \frac{\alpha_1}{\beta}, \frac{\alpha_1+1}{\beta}, \dots, \frac{\alpha_1+\beta-1}{\beta}, \dots, \frac{\alpha_n}{\beta}, \dots, \frac{\alpha_n+\beta-1}{\beta}, c_1+d_1, \dots, c_m+d_m \\ \frac{b}{\beta}, \frac{b+1}{\beta}, \dots, \frac{b+\beta-1}{\beta}, \frac{-b}{\beta}, \dots, \frac{-b+\beta-1}{\beta}, \frac{\alpha_1}{s}, \dots, \frac{\alpha_m+s-1}{s} \end{array} \right. \right).$$

$$\begin{aligned}
 (2.3) \quad & p^{-\lambda} G_{m, m+1}^{m+1, 0} \left(yp^{-n/s} / \begin{matrix} n_i + a_i \\ n_i, \sigma \end{matrix} \right) \\
 & = p \int_0^\infty G_{\mu, \mu+1}^{\mu+1, 0} \left(pt / \begin{matrix} \eta_j + \alpha_j \\ \eta_j, \rho \end{matrix} \right) t^\lambda F(t, y) dt \\
 & \qquad \qquad \qquad i = 1, 2, \dots, m; j = 1, 2, \dots, \mu.
 \end{aligned}$$

Hence substituting from (2.3) in the integrand on the R.H.S. of (2.2), and changing the order of integration we get the result (2.1).

To justify the change of order of integration we see that the t -integral in (2.3) is absolutely and uniformly convergent as n and s are positive integers,

$$\begin{aligned}
 s > n, \quad \left| \arg \frac{y^s}{p^n} \right| &< \left(\frac{s-n}{2} \right) \pi; \\
 \min (\operatorname{Re} \eta_j, \operatorname{Re} \rho) + n \min \left(\frac{\operatorname{Re} n_i}{s}, \frac{\operatorname{Re} \sigma}{s} \right) \\
 &> -1 - \operatorname{Re} \lambda > -\operatorname{Re} (\eta_j + \alpha_j + \lambda) - 2 \\
 &\qquad \qquad \qquad \text{for } i = 1, 2, \dots, m; j = 1, 2, \dots, \mu.
 \end{aligned}$$

the y -integral in (2.1) is absolutely and uniformly convergent and the repeated integral is absolutely convergent as the Meijer-Laplace transform of $|t^\lambda \int_0^\infty F(t, y) dy|$ exists. Hence by De La Vallée Poussin's Theorem [7, p. 504] the inversion of the order of integration is justified.

3. Particular cases

In all the following cases we assume n, s as positive integers with $s > n$, $|\arg y^s/p^n| < (s-n)\pi/2$ and the integrals involved are absolutely and uniformly convergent.

(i) With $a_j = 0$, $j = 1, 2, \dots, m-1$; $a_m = -m' - k$; $\sigma = -m' - k$, $n_m = m' - k$ we have;

If

$$f(y) \xrightarrow[m']{k+\frac{1}{2}} \phi(p)$$

then

$$p^{n/s-\lambda} \phi(p^{-n/s}) = \frac{(2\pi)^{(n-s)/2} s^{k+m'}}{n^{\rho-\sum \alpha_j + \frac{1}{2} + \lambda}} G \left[t^\lambda \int_0^\infty F(t, y) f(y) dy; \alpha_\mu, \eta_\mu, \rho \right]$$

where

$$F(t, y) = G_{n\mu+s, 2s+n\mu+n}^{2s, n\mu} \left(\frac{y^s t^n}{s^s n^n} \left/ \begin{array}{l} \frac{-\eta_j - \alpha_j - \lambda + l}{n}, \frac{-2k + \nu}{s} \\ \frac{+m' - k + \nu}{s}, \frac{-\eta_j - \lambda + l}{n}, \frac{-\rho - \lambda + l}{n} \end{array} \right. \right)$$

$$\begin{aligned} \min(\operatorname{Re} \eta_j, \operatorname{Re} \rho) + \frac{n}{s} \operatorname{Re}(-k \pm m') &> -1 - \operatorname{Re} \lambda \\ &> -\operatorname{Re}(\eta_j + \alpha_j + \lambda) - 2; \end{aligned}$$

for $j = 1, 2, \dots, \mu$; $l = 0, 1, \dots, n-1$; $\nu = 0, 1, \dots, s-1$.

(ia) Further with $k = \pm m'$ we get

If

$$\phi(p) \doteq f(y)$$

then

$$(3.1) \quad p^{n/s-\lambda} \phi(p^{-n/s}) = \frac{(2\pi)^{(n-s)/2} \cdot s^{\frac{1}{2}}}{n^{\rho-\sum \alpha_j + \frac{1}{2} + \lambda}} G \left[t^\lambda \int_0^\infty F(t, y) f(y) dy; \alpha_\mu, \eta_\mu, \rho \right]$$

where

$$(3.2) \quad F(t, y) = G_{n\mu, s+n\mu+n}^{s, n\mu} \left(\frac{y^s t^n}{s^s n^n} \left/ \begin{array}{l} \frac{-\eta_j - \alpha_j - \lambda + l}{n} \\ \frac{\nu}{s}, \frac{-\eta_j - \lambda + l}{n}, \frac{-\rho - \lambda + l}{n} \end{array} \right. \right)$$

$$\begin{aligned} \min(\operatorname{Re} \eta_j, \operatorname{Re} \rho) + \operatorname{Re} \lambda + 1 &> 0, \operatorname{Re}(\eta_j + \alpha_j) > -1 \\ \text{for } j = 1, 2, \dots, \mu; l = 0, 1, \dots, n-1; \nu = 0, 1, \dots, s-1. \end{aligned}$$

(ib) Further with $k = \pm m'$ and $\alpha_j = 0$, $j = 1, 2, \dots, \mu$; $\rho = 0$ we obtain

If

$$\phi(p) \doteq f(y)$$

then

$$p^{n/s-\lambda} \phi(p^{-n/s}) \doteq \frac{(2\pi)^{(n-s)/2} \cdot s^{\frac{1}{2}}}{n^{\frac{1}{2} + \lambda}} t^\lambda \int_0^\infty G_{0, s+n}^{s, 0} \left(\frac{y^s t^n}{s^s n^n} \left/ \begin{array}{l} \frac{\nu}{s}, \frac{-\lambda + l}{n} \end{array} \right. \right) f(y) dy$$

$$\operatorname{Re} \lambda + 1 > 0; l = 0, 1, \dots, n-1; \nu = 0, 1, \dots, s-1.$$

With $n = s = 1$ we get the well-known Tricomi's theorem [5, p. 564].

(ii) Setting $\alpha_j = 0$, $j = 1, 2, \dots, m-1$; $a_m = \frac{1}{2} - k - m'$; $n_m = 2m'$, $\sigma = 0$ we have

If

$$\phi(p) = W[f(y); k, m']$$

then

$$(3.3) \quad p^{n/s-\lambda} \phi(p^{-n/s}) = \frac{(2\pi)^{(n-s)/2} \cdot s^{\frac{1}{2}}}{n^{\rho-\sum \alpha_j + \frac{1}{2} + \lambda}} G \left[t^\lambda \int_0^\infty F(t, y) f(y) dy; \alpha_\mu, \eta_\mu, \rho \right]$$

where

$$(3.4) \quad F(t, y) = G_{n\mu+s, 2s+n\mu+n}^{2s, n\mu} \left(\frac{y^s t^n}{s^s n^n} \left/ \begin{array}{c} \frac{-\eta_j - \alpha_j - \lambda + l}{n}, \frac{\frac{1}{2} - k + m' + \nu}{s} \\ \frac{m' \pm m' + \nu}{s}, \frac{-\eta_j - \lambda + l}{n}, \frac{-\rho - \lambda + l}{n} \end{array} \right. \right)$$

$$\min (\operatorname{Re} \eta_j, \operatorname{Re} \rho) + \frac{n}{s} \operatorname{Re} (m' \pm m')$$

$$> -1 - \operatorname{Re} \lambda > -\operatorname{Re} (\eta_j + \alpha_j + \lambda) - 2$$

for $j = 1, 2, \dots, \mu$; $l = 0, 1, \dots, n-1$; $\nu = 0, 1, \dots, s-1$.

With $\alpha_j = 0$, $j = 1, 2, \dots, \mu-1$; $\alpha_\mu = \frac{1}{2} - \mu' - \lambda'$; $\eta_\mu = 2\mu'$; $\rho = 0$ and $n = s = 1$ we get after some simplification the result by R. Narain [8, p. 316].

EXAMPLE: The function $K(z)$ of Boersma J. [9] is defined by

$$K \left[\begin{array}{c} a_r, \mu_r \\ b_t, \nu_t \end{array}; z \right] = \frac{1}{2\pi i} \int \frac{\Gamma(-s) \prod_{k=1}^r \Gamma(a_k + \mu_k s)}{\prod_{j=1}^t \Gamma(b_k + \nu_k s)} (-z)^s ds$$

$$= (\text{const.}) G_{\sum \mu_j, 1 + \sum \nu_k}^1 \left(-z \left/ \begin{array}{c} \frac{1 - a_j + i}{\mu_j} \\ 0, \frac{1 - b_k + l}{\nu_k} \end{array} \right. \right)$$

where μ_j, ν_k are natural numbers; $i = 0, 1, \dots, \mu_j - 1$; $j = 1, 2, \dots, r$; $l = 0, 1, \dots, \nu_k - 1$; $k = 1, 2, \dots, t$ and const.

$$= \frac{(2\pi)^{\frac{1}{2}(r-t+\sum \nu_k - \sum \mu_j)} \prod_{j=1}^r \mu_j^{\alpha_j + \mu_j s - \frac{1}{2}}}{\prod_{k=1}^t \nu_k^{b_k + \nu_k s - \frac{1}{2}}}$$

Let

$$f(y) = y^{-h} K \left[\begin{array}{c} a_r, \mu_r \\ b_t, \nu_t \end{array}; -y \right] \doteq p^h K \left[\begin{array}{c} 1, 1; a_r, \mu_r \\ b_t, \nu_t \end{array}; -\frac{1}{p} \right] = \phi(p)$$

$$\operatorname{Re} h < 1; \operatorname{Re} p > 0; \sum \mu_j \leq 1 + \sum \nu_k,$$

in case (ia) of the theorem.

Then (3.1) gives,

$$p^{n/s(1-h)-\lambda} K \left[\begin{matrix} 1, 1; a_r, \mu_r; -p^{n/s} \\ b_t, \nu_t \end{matrix} \right] \\ = \frac{(2\pi)^{(n-s)/2} \cdot s^{\frac{1}{2}}}{n^{\rho-\sum \alpha_j + \frac{1}{2} + \lambda}} G \left[t^\lambda \int_0^\infty F(t, y) K(y) dy; \alpha_\mu, \eta_\mu, \rho \right]$$

where $F(t, y)$ is given by (3.2), provided n, s are positive integers,

$$\operatorname{Re} h < 1; \operatorname{Re} p > 0; \min (\operatorname{Re} \eta_j, \operatorname{Re} \rho) + \operatorname{Re} \lambda + 1 > 0;$$

$$\operatorname{Re} (\eta_j + \alpha_j) > -1$$

for $j = 1, 2, \dots, \mu; l = 0, 1, \dots, n-1; \nu = 0, 1, \dots, s-1; \mu_r, \nu_t$ are natural numbers, $\sum \mu_j \leq 1 + \sum \nu_k; s > n$.

I am thankful to Dr. B. R. Bhonsle for his help in the preparation of this paper. I am also thankful to the referee for his valuable suggestions in improving the paper.

REFERENCES

V. M. BHISE

- [1] Journl. Vikram Univ. 3, No. 3, (1959), p. 57-63.
 [2] Proc. Nat. Acad. Sci., India, 32A (1962), p. 389-404.
 [6] Communicated for publication.

J. BOERSMA

- [9] Compositio Math., 15 (1961) p. 34-63.

BROMWICH T. J. I. 'A.

- [7] Introduction to the theory of Infinite Series, London (1931).

C. S. MEIJER

- [3] Proc. Ned. Acad. V. Weten. Amsterdam, 44 (1941), p. 727-737.

R. NARAIN

- [8] Rendiconti del Seminario Matematico dell'Universita e del Politecnico di Torino, 15 (1955-56), p. 311-328.

F. TRICOMI

- [5] Rendic. Acc. dei Lincei, 22 (1935), p. 564-571.

R. S. VERMA

- [4] Proc. Nat. Acad. Sci, India, 20A (1951), p. 209-216.

(Oblatum 20-5-66)

Department of Mathematics,
 G. S. Technological Institute,
 Indore (India).