

COMPOSITIO MATHEMATICA

RAM SINGH

On a class of star-like functions

Compositio Mathematica, tome 19, n° 1 (1968), p. 78-82

<http://www.numdam.org/item?id=CM_1968__19_1_78_0>

© Foundation Compositio Mathematica, 1968, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On a class of star-like functions¹

by

Ram Singh

1. Introduction

A function $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$, regular and univalent in $|z| < 1$, is said to be star-like if $\operatorname{Re}\{zf'(z)/f(z)\} \geq 0$ in $|z| < 1$. Various sub classes of the class of star-like functions have been considered by different authors. The class S_α of star-like functions of order α ($0 \leq \alpha \leq 1$) i.e. the class of star-like functions $f(z)$ for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq \alpha$, was first introduced by Robertson [1] and has been studied among others by Merkes [5] and Schild [4]. The class $S_{\frac{1}{2}}$ is particularly important. Marx [2] and Stroh  cker [6] proved that if $f(z)$ maps $|z| < 1$ on to a convex domain, then $f(z) \in S_{\frac{1}{2}}$. Somewhat later Gabriel [3] showed that the functions of the class $S_{\frac{1}{2}}$ played an important role in the solution of certain differential equations. In the present paper we investigate the class \bar{S} of star-like functions $f(z)$ such that $\operatorname{Re}\{zf'(z)/f(z)\}$ lies in the circle with centre at $(1, 0)$ and radius unity i.e. the class of functions $f(z)$ such that $|zf'(z)/f(z) - 1| \leq 1$ in $|z| < 1$.

2. A representation formula

THEOREM 1: $f(z) \in \bar{S}$ if and only if

$$(2.1) \quad f(z) = z \exp \left\{ \int_0^z \phi(t) dt \right\},$$

where $\phi(z)$ is regular and $|\phi(z)| \leq 1$ in $|z| < 1$.

PROOF. Suppose that $f(z) \in \bar{S}$. Then $|zf'(z)/f(z) - 1| \leq 1$. Also $\{zf'(z)/f(z) - 1\} = 0$ when z is 0. Therefore, by Schwarz' lemma it follows that

$$(2.2) \quad \frac{zf'(z)}{f(z)} - 1 = z\phi(z),$$

where $\phi(z)$ is regular and $|\phi(z)| \leq 1$ in $|z| \leq 1$.

¹ The author wishes to express his thanks to Prof. V. Singh for helpful guidance.

From (2.2), by integration, it easily follows that $f(z)$ has the representation (2.1).

Conversely, if $f(z)$ has the representation (2.1), then by differentiation etc., we find that $|zf'(z)/f(z)-1| \leq 1$ and therefore $f(z) \in \bar{S}$.

3. Distortion theorems

THEOREM 2. *For all $f(z) \in \bar{S}$, we have*

$$(3.1) \quad |z| e^{-|z|} \leq |f(z)| \leq |z| e^{|z|},$$

and

$$(3.2) \quad (1-|z|) e^{-|z|} \leq |f'(z)| \leq (1+|z|) e^{|z|}.$$

PROOF: Since any function $f(z) \in \bar{S}$ has the representation (2.1), where $\phi(z)$ is regular and $|\phi(z)| \leq 1$ in $|z| < 1$, (3.1) follows immediately.

Inequalities (3.1) are sharp. Equality being attained for the function $f(z) = ze^z$.

Again differentiating (2.1) we get

$$(3.3) \quad f'(z) = [1+z\phi(z)] \exp \left\{ \int_0^z \phi(t) dt \right\}.$$

From the above representation of $f'(z)$, (3.2) at once follows.

The function $f(z) = ze^z$ shows that (3.2) is sharp.

From (3.2) we have the following:

COROLLARY 1: The conformal map of $|z| < 1$ as yielded by functions of class \bar{S} contains all the points of circle $|w| < 1/e$.

4. Coefficient regions

THEOREM 3: *If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in \bar{S}$, then*

$$(4.1) \quad |a_n| \leq \frac{1}{(n-1)}.$$

PROOF. The proof is based on a method of Clunie. If $f(z) \in \bar{S}$, then we know that

$$(4.2) \quad zf'(z) - f(z) = f(z) \cdot z\phi(z).$$

If we put $\psi(z) = z\phi(z) = \sum_{n=1}^{\infty} b_n z^n$, then $|\psi(z)| < 1$ in $|z| < 1$, and (4.2) may be written as

$$(4.3) \quad \sum_{n=2}^{\infty} (n-1)a_n z^n = [z + \sum_{n=2}^{\infty} a_n z^n] \sum_{n=1}^{\infty} b_n z^n.$$

Comparing the coefficients of z^n on both the sides of (4.3) we get

$$(n-1)a_n = b_1 a_{n-1} + b_2 a_{n-2} + b_3 a_{n-3} + \dots + b_{n-1}, \quad n \geq 2.$$

This shows that the coefficient a_n on the left side of (4.3) depends only on the coefficients a_2, a_3, \dots, a_{n-1} on the right hand side of (4.3). Hence for $n \geq 2$, we may write

$$(4.4) \quad \sum_{k=2}^n (k-1)a_k z^k + \sum_{k=n+1}^{\infty} c_k z^k = [z + \sum_{k=2}^{n-1} a_k z^k] \psi(z)$$

say. Squaring the moduli of both sides of (4.4) and integrating round $|z| = r < 1$, we get, using the fact that $|\psi(z)| < 1$ in $|z| < 1$,

$$\sum_{k=2}^n (k-1)^2 |a_k|^2 r^{2k} + \sum |c_k|^2 r^{2k} < 1 + \sum_{k=2}^{n-1} |a_k|^2$$

making $r \rightarrow 1$, we find that

$$\sum_2^n (k-1)^2 |a_k|^2 \leq 1 + \sum_2^{n-1} |a_k|^2$$

or

$$(n-1)^2 |a_n|^2 \leq 1 - \sum_2^{n-1} (k-2)k |a_k|^2.$$

It, therefore follows that for all $n \geq 2$

$$|a_n| \leq \frac{1}{(n-1)}.$$

Equality is attained for $f(z) = ze^{z^{n-1}/(n-1)}$, a function of \bar{S} for which $a_2 = a_3 = \dots a_{n-1} = 0$.

5. Radius of convexity

THEOREM 4. Each function $f(z) \in \bar{S}$ maps

$$(5.1) \quad |z| \leq \frac{\sqrt{13}-3}{2}$$

onto a convex domain.

PROOF. We know that if $f(z) \in \bar{S}$, then

$$\frac{zf''(z)}{f'(z)} = 1 + z\phi(z).$$

Logarithmic differentiation of the above equation yields

$$\begin{aligned}\frac{zf''(z)}{f'(z)} &= \frac{zf''(z)}{f'(z)} - 1 + \frac{z\phi(z) + z^2\phi'(z)}{1 + z\phi(z)} \\ &= z\phi(z) + \frac{z\phi(z) + z^2\phi'(z)}{1 + z\phi(z)}.\end{aligned}$$

Now making use of the fact that $|\phi'(z)| \leq 1 - |\phi(z)|^2/1 - |z|^2$, we get

$$(5.2) \quad \left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{|z|[2|\phi| - 2|\phi| \cdot |z|^2 + |z| - |z| \cdot |z\phi|^2]}{(1 - |z\phi|)(1 - |z|^2)}.$$

$f(z)$ will map $|z| \leq r$ onto a convex domain if the right hand side of (5.2) is less than or equal to 1 for $|z| \leq r$, i.e. if

$$(5.3) \quad (1 - |z\phi|)(1 - |z|^2) \geq |z|[2|\phi| - 2|\phi| \cdot |z|^2 + |z| - |z| \cdot |z\phi|^2].$$

Putting $|z| = p$, $|\phi| = q$ and $pq = x$, where $0 \leq p \leq 1$, $0 \leq q \leq 1$, $0 \leq x \leq p$, (5.3) may be written as

$$(5.4) \quad p^2x^2 - 3(1 - p^2)x + (1 - 2p^2) \geq 0.$$

We consider $f(x) = p^2x^2 - 3(1 - p^2)x + (1 - 2p^2)$. Clearly $f(x)$ decreases for $0 \leq x \leq p$, if

$$2p^2x - 3(1 - p^2) < 0$$

or

$$2p^2 \cdot p - 3(1 - p^2) < 0$$

or

$$(5.5) \quad p < \frac{1}{2}[(5 - 2\sqrt{6})^{\frac{1}{2}} + (5 + 2\sqrt{6})^{\frac{1}{2}} - 1]$$

The value of p as given by (5.5) is easily seen to be less than $\frac{13}{16}$.

The least value of $f(x)$ is $f(p)$, when p is given by (5.5). Therefore, (5.4) will hold if

$$f(p) \geq 0$$

or

$$p^4 + 3p^3 - 2p^2 - 3p + 1 \geq 0$$

or

$$(1 - p^2)(p^2 + 3p - 1) \leq 0$$

or

$$p^2 + 3p - 1 \leq 0$$

or

$$(5.6) \quad p \leq \frac{\sqrt{13}-3}{2}.$$

Clearly the value of p as given by (5.6) is less than that given by (5.5). Hence every function $f(z) \in \bar{S}$ maps $|z| \leq \sqrt{13}-3/2$ onto a convex domain.

The function $f_0(z) = ze^z$ shows that this bound for the radius of convexity cannot be improved.

REFERENCES

ROBERTSON, M. S.

- [1] "On the theory of univalent functions", *Annals of Mathematics*, Vol. 37 (1936), pp. 374-408.

MARX, A.

- [2] "Untersuchungen uber schlichte Appildgen", *Mathematische Annalen*, Vol. 107 (1932-33) pp. 40-67.

GABRIEL, R. F.

- [3] "The Schwarzian derivative and convex functions", *Proc. Amer. Math. Soc.*, Vol. 6 (1955), pp. 58-66.

SCHILD, A.

- [4] "On a class of univalent starshaped functions," *Amer. Math. Soc.* Vol. 9 (1958), pp. 751-757.

MERKES, E. P.

- [5] "On products of Starlike functions", *Proc. Amer. Math. Soc.* Vol. 13 (1962), pp. 960-964.

STROHACKER, E.

- [6] "Beitrage Zur Theorie Der Schlichten Funktionen", *Mathematische Zeitschrift*, Vol. 37 (1933) pp. 356-380.

(Oblatum 21-2-1966)

Punjabi University, Patiala, India