

COMPOSITIO MATHEMATICA

W. L. BYNUM

A class of spaces lacking normal structure

Compositio Mathematica, tome 25, n° 3 (1972), p. 233-236

http://www.numdam.org/item?id=CM_1972__25_3_233_0

© Foundation Compositio Mathematica, 1972, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A CLASS OF SPACES LACKING NORMAL STRUCTURE

by

W. L. Bynum ¹

In [5, Lemma 5], Goebel showed that if B is a Banach space with coefficient of convexity less than 1, then B has normal structure. This paper gives an example (Theorem 1) of a class of spaces which lack normal structure and have coefficient of convexity 1. Thus, Goebel's result is the best result possible. In Theorem 2 of this paper, it is shown that the duals of this class of spaces *have* normal structure and that their coefficients of convexity are between 1 and 2, so that normal structure is not self-dual.

We shall use the following definitions and notation. Let B be a Banach space with norm $\| \cdot \|$ and let K denote the unit sphere of B . The *modulus of convexity* of B is the function δ defined for t in $[0, 2]$ as follows:

$$2\delta(t) = \inf \{2 - \|x+y\| : x, y \in K, \|x-y\| \geq t\}$$

(see [4]). A space B is *uniformly convex* provided that its modulus of convexity is positive on $(0, 2]$ ([3], [4]). The *coefficient of convexity* of B , $\varepsilon_0 = \varepsilon_0(B)$, is $\sup \{t \in [0, 2] : \delta(t) = 0\}$ ([5]).

Let C be a bounded subset of B . The *diameter of C* , $\text{diam } C$, is $\sup \{\|x-y\| : x, y \in C\}$. A member x of C is a *non-diametral point* provided that $\text{diam } C > \sup \{\|x-u\| : u \in C\}$ and a *diametral point of C* is a point x for which the previous inequality is replaced by equality. For y in B , the *distance from y to C* , $d(y, C)$, is

$$\inf \{\|y-u\| : u \in C\}.$$

A space B has *normal structure* if each bounded convex subset of B with positive diameter has a non-diametral point ([2]). The convex hull of a subset A of B will be denoted by $\text{co } A$.

A space B is *uniformly non-square* if there is an $r > 0$ such that for x and y in X having norm 1, $\|x+y\| + \|x-y\| \leq 4-r$ ([7]).

The examples in this paper are based on the l_p spaces. For $1 < p < \infty$ and x in l_p , define sequences x^+ and x^- as follows:

$$\begin{aligned} (x^+)_n &= \sup (x_n, 0) = (|x_n| + x_n)/2 \\ (x^-)_n &= \sup (-x_n, 0) = (|x_n| - x_n)/2. \end{aligned}$$

¹ Research supported by a Faculty Research Grant of the College of William and Mary.

For x in l_p , x^+ and x^- are in l_p and $x = x^+ - x^-$. Denote the l_p norm by $\| \cdot \|$. For $1 \leq q < \infty$, let $l_{p,q}$ denote the set of elements of l_p with the norm:

$$\|x\| = (\|x^+\|^q + \|x^-\|^q)^{1/q}.$$

Let $l_{p,\infty}$ denote the set of elements of l_p with the norm:

$$\|x\| = \sup \{ \|x^+\|, \|x^-\| \}.$$

It is easy to show that for $1 \leq q \leq \infty$, the function $\| \cdot \|$ is indeed a norm for l_p which is equivalent to the l_p norm. We shall show in Theorem 1 that for $1 < p < \infty$, $l_{p,\infty}$ lacks normal structure and $\varepsilon_0(l_{p,\infty}) = 1$ and in Theorem 2 that $l_{p,1}$ has normal structure and $2^{1/p} \leq \varepsilon_0(l_{p,1}) < 2$. Using an argument which is considerably more complicated than the proof of Theorem 2, one can show that $\varepsilon_0(l_{p,1})$ is actually equal to $2^{1/p}$.

If $1 < p < \infty$ and $1 \leq q \leq \infty$ and p^* and q^* are the conjugate indices of p and q , then a straightforward argument shows that $(l_{p,q})^*$ is isometrically isomorphic to l_{p^*,q^*} . Thus, by combining Theorems 1 and 2, we obtain a class of reflexive spaces such that each space lacks (has) normal structure while its dual has (lacks) normal structure, so that normal structure is not self-dual.

A lack of connection between normal structure and reflexivity has been shown previously. In [1, p. 439] Belluce, Kirk and Steiner gave an example of a reflexive space lacking normal structure (the coefficient of convexity of this example is 2); consequently, reflexivity does not imply normal structure. On the other hand, Zizler has shown [8, proposition 2] that each separable Banach space B has an equivalent norm with respect to which B has normal structure; thus, normal structure does not imply reflexivity.

Using the methods of this paper, one can show that for $1 < p, q < \infty$, $l_{p,q}$ is uniformly convex and therefore has normal structure. We shall not analyze these spaces here.

Incidentally, it is not difficult to prove that in an arbitrary Banach space B with modulus of convexity δ , the limit from the left of δ at 2, $\delta(2^-)$, is equal to $1 - (\varepsilon_0/2)$. In [5], Goebel has noted that B is uniformly convex if and only if $\varepsilon_0 = 0$. Consequently, we have that B is uniformly convex if and only if $\delta(2^-) = 1$. It is interesting to compare this with Goebel's observation that B is strictly convex if and only if $\delta(2) = 1$.

THEOREM (1). For $1 < p < \infty$, $l_{p,\infty}$ lacks normal structure and its coefficient of convexity is 1.

PROOF. The following theorem of Brodskii and Mil'man [2] is used to show that $B = l_{p,\infty}$ lacks normal structure:

A space X does not have normal structure if and only if there is a bounded sequence $\{x_n\}$ of elements of X such that the distance from x_{n+1} to $\text{co}\{x_1, \dots, x_n\}$ tends to the diameter of the set of all x_i as $n \rightarrow \infty$ (such a sequence is called a *diametral sequence*).

We shall show that the sequence, $\{e_n\}$, of unit vectors of B (i.e., e_n is that member of B whose n -th coordinate is 1 with all other coordinates 0) is a diametral sequence. If m is a positive integer and $\alpha_1 + \dots + \alpha_m = 1$ and $0 \leq \alpha_i \leq 1$ for $1 \leq i \leq m$, then

$$|e_{m+1} - \sum \alpha_i e_i|^p = \sup \{1, \sum \alpha_i^p\} = 1.$$

Thus, the distance from e_{m+1} to $\text{co}\{e_1, \dots, e_m\}$ is 1. The above equation also shows that the diameter of the sequence $\{e_n\}$ is 1, so B lacks normal structure.

By Goebel's lemma 5 and the previous paragraph, $\varepsilon_0(B) \geq 1$. We shall show that $\varepsilon_0 \leq 1$. Suppose that $\delta(t) = 0$ for some t in $[0, 2]$. Then, there are sequences $\{x_n\}$ and $\{y_n\}$ in K , the unit sphere of B , such that for each n , $|x_n - y_n| \geq t$ and $|x_n + y_n| \rightarrow 2$ as $n \rightarrow \infty$. For each u in B , $(-u)^+ = u^-$ and $(-u)^- = u^+$, so we may assume that for each n , $|x_n + y_n| = \|(x_n + y_n)^+\|$ and $|x_n - y_n| = \|(x_n - y_n)^+\|$. But, $2 \geq \|x_n^+ + y_n^+\| \geq |x_n + y_n| \rightarrow 2$; consequently, the uniform convexity of l_p implies that $\|x_n^+ - y_n^+\| \rightarrow 0$. Therefore,

$$\begin{aligned} t &\leq |x_n - y_n| \leq \|(x_n^+ - y_n^+)^+\| + \|(y_n^- - x_n^-)^+\| \\ &\leq \|x_n^+ - y_n^+\| + 1 \rightarrow 1. \end{aligned}$$

Thus, $\varepsilon_0 = 1$, and the proof is complete.

THEOREM (2). For $1 < p < \infty$, $l_{p,1}$ has normal structure and $2^{1/p} \leq \varepsilon_0(l_{p,1}) < 2$.

PROOF. First, we establish the inequality for ε_0 . Let $x = e_1$ and $y = -e_2$. Then $|x + y| = 2$ and $|x - y| = 2^{1/p}$, so that $\varepsilon_0 \geq 2^{1/p}$. In [5], Goebel notes that a space B is uniformly non-square if and only if $\varepsilon_0(B) < 2$. It is easy to see that if B is uniformly non-square then B^* is also. By Theorem 1, $\varepsilon_0(l_{p^*, \infty}) = 1$, and thus, $\varepsilon_0(l_{p,1}) < 2$.

To show that $l_{p,1}$ has normal structure, we shall use the following theorem of Gossez and Lami Dozo [6]:

Let B be a Banach space with Schauder basis $\{e_n\}$. For each positive integer k and each x in B , let $U_k(x) = \sum_1^k x_n e_n$ and let $V_k(x) = x - U_k(x)$. Suppose that $\{k_n\}$ is a strictly increasing sequence of positive integers with the following property:

If $c > 0$, there is an $r > 0$ with the property that if x is in B and n is an integer such that $\|U_{k_n}(x)\| = 1$ and $\|V_{k_n}(x)\| \geq c$, then $\|x\| \geq 1 + r$.

Then, each convex weakly relatively compact subset of B of at least two points has a non-diametral point.

The sequence $\{e_n\}$ of unit coordinate vectors is a Schauder basis for $l_{p,1}$. It follows from the Minkowski inequality that for each positive integer k and each x in $l_{p,1}$,

$$|x|^p \geq |U_k(x)|^p + |V_k(x)|^p.$$

Therefore, the above theorem is applicable. Since this space is reflexive, each bounded convex subset is weakly relatively compact. Thus, $l_{p,1}$ has normal structure. I want to thank the referee for calling my attention to the paper of Gossez and Lami Dozo.

REFERENCES

- L. P. BELLUCE, W. A. KIRK AND E. F. STEINER
 [1] Normal structure in Banach spaces, *Pacific J. Math.* 26 (1968), 433–440.
- M. S. BRODSKII AND D. P. MIL'MAN
 [2] On the center of a convex set, *Dokl. Akad. Nauk. SSSR N.S.* 59 (1948), 837–840.
- J. A. CLARKSON
 [3] Uniformly convex spaces, *Trans. Amer. Math. Soc.* 40 (1936), 396–414.
- M. M. DAY
 [4] Uniform convexity in factor and conjugate spaces, *Annals of Math.* 45 (1944), 375–385.
- K. GOEBEL
 [5] Convexity of balls and fixed point theorems for mappings with nonexpansive square, *Compositio Math.* 22 (1970), 269–274.
- J. P. GOSSEZ AND E. LAMI DOZO
 [6] Structure normale et base de Schauder, *Bull. de l'Acad. Royale de Belgique (5e Ser.)* 55 (1969), 673–681.
- R. C. JAMES
 [7] Uniformly non-square Banach spaces, *Annals of Math.* 80 (1964), 542–550.
- V. ZIZLER
 [8] Some notes on various rotundity and smoothness properties of separable Banach spaces, *Comment. Math. Univ. Carolinae* 10 (1969), 195–206.

(Oblatum 30–VIII–1971 & 24–IV–1972)

Department of Mathematics
 College of William and Mary
 Williamsburg, Virginia 23185