

COMPOSITIO MATHEMATICA

FLORIS TAKENS

Derivations of vector fields

Compositio Mathematica, tome 26, n° 2 (1973), p. 151-158

http://www.numdam.org/item?id=CM_1973__26_2_151_0

© Foundation Compositio Mathematica, 1973, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

DERIVATIONS OF VECTOR FIELDS

by

Floris Takens*

1. Statement of the result

Let M be a differentiable, i.e. C^∞ , manifold. We denote the Lie-algebra of C^∞ vectorfields on M by $\chi(M)$. A map $D : \chi(M) \rightarrow \chi(M)$ is called a *derivation* if D is \mathbf{R} -linear and if $D([X, Y]) = [D(X), Y] + [X, D(Y)]$ for all $X, Y \in \chi(M)$. It is clear that every $X \in \chi(M)$ defines a derivation $DX : DX(Y) = [X, Y]$. In this note we want to show that every derivation can be obtained in this way.

THEOREM. *For each derivation $D : \chi(M) \rightarrow \chi(M)$ there is a vectorfield $Z \in \chi(M)$, such that for each $X \in \chi(M)$, $D(X) = [Z, X]$.*

This theorem has a certain relation with recent work of M. Gel'fand, D. B. Fuks, and others [1] on the cohomology of Lie-algebras of smooth vectorfields, because it implies that $H^1(\chi(M); \chi(M)) = 0$; $H^1(\chi(M); \chi(M))$ being the first cohomology group of $\chi(M)$ with coefficient in $\chi(M)$ with the adjoined representation (this was pointed out to me by M. Hazewinkel). There is however one difference in their approach: in defining their cohomology they only use cochains which are continuous mappings (with respect to the C^∞ topology). It is however not difficult to show that the nullity of $H^1(\chi(M); \chi(M))$ follows from our theorem in either case.

The theorem will follow from the following lemmas:

LEMMA 1. *Let $D : \chi(M) \rightarrow \chi(M)$ be a derivation and let $X \in \chi(M)$ be zero on some open subset $U \subset M$. Then $D(X)|U \equiv 0$.*

LEMMA 2. *Let $X \in \chi(\mathbf{R}^n)$ be a vectorfield on \mathbf{R}^n with $j^3(X)(0) = 0$, i.e. the 3-jet of each of the component functions of X is zero in the origin. Then there are vectorfields $Y_1, \dots, Y_q, Z_1, \dots, Z_q$ and there is a neighbourhood U of the origin in \mathbf{R}^n such that:*

$$X|U = \sum_i [Y_i, Z_i]|U \quad \text{and}$$

$$j^1(Y_i)(0) = 0, \quad j^1(Z_i)(0) = 0 \quad \text{for all } i = 1, \dots, q.$$

* During the preparation of this paper, the author was a visiting member of the Mathematical Institute of the University of Strasbourg.

LEMMA 3. Let $D : \chi(M) \rightarrow \chi(M)$ be a derivation and let $X \in \chi(M)$ and $p \in M$ be such that $j^3(X)(p) = 0$. Then $D(X)(p) = 0$. In other words, $DX(p)$ is determined by $j^3(X)(p)$, also if $j^3(X)(p) \neq 0$.

Lemma 3 will be derived from the lemmas 1 and 2. Finally we shall use lemma 3 to derive:

LEMMA 4. Let $U \subset \mathbf{R}^n$ be an open connected and simply connected set and let $D_U : \chi(U) \rightarrow \chi(U)$ be a derivation. Then there is a unique vectorfield $Z \in \chi(U)$ such that $D_U(X) = [Z, X]$ for all $X \in \chi(U)$.

Finally, we shall see that the theorem follows from lemma 1 and lemma 4.

2. The Proofs

PROOF OF LEMMA 1. Suppose $X|_U \equiv 0$ and $D(X)(q) \neq 0$ for some point $q \in U$. We take a vectorfield $Y \in \chi(M)$ such that $\text{supp}(Y) \subset U$ and $[D(X), Y](q) \neq 0$. By definition we have $D[X, Y] = [DX, Y] + [X, DY]$; evaluating this in q we get $0 = [DX, Y](q) \neq 0$, which contracts our assumption. Hence the lemma is proved.

PROOF OF LEMMA 2. It is clearly enough to show the lemma for the case

$$X = X(x_1, \dots, x_n) \frac{\partial}{\partial x_1},$$

with $j^3(X)(0) = 0$. Such vectorfields can be written as a finite sum of vectorfields of the following two types:

type I:

$$\tilde{X} = x_1^{m_1} \cdot \dots \cdot x_n^{m_n} \cdot \alpha(x_2, \dots, x_n) \frac{\partial}{\partial x_1},$$

with $\sum m_i \geq 4$ and α a C^∞ function;

type II:

$$\tilde{X} = x_1^4 \cdot g(x_1, \dots, x_n) \frac{\partial}{\partial x_1}$$

with g a C^∞ function.

To prove the lemma we show that each vectorfield, which is either of type I or of type II, can be written as the Lie-product of two vectorfields with zero 1-jet in the origin. For \tilde{X} of type I as above, we observe that

$$\tilde{X} = \left[\frac{1}{k_1 - h_1} \cdot x_1^{h_2} \cdot \dots \cdot x_n^{h_n} \frac{\partial}{\partial x_1}, x_1^{k_2} \cdot \dots \cdot x_n^{k_n}, \alpha(x_2, \dots, x_n) \frac{\partial}{\partial x_1} \right],$$

$$h_1 + k_1 = m_1 + 1 \text{ and } h_1 \neq k_1$$

$$h_2 + k_2 = m_2$$

$$\vdots$$

$$h_n + k_n = m_n.$$

Using the fact that $\sum m_i \geq 4$, we see that we can choose $h_1, \dots, h_n, k_1, \dots, k_n$ so that $\sum h_i \geq 2$ and $\sum k_i \geq 2$; hence for type one we have the required Lie-product.

Suppose that that

$$\tilde{X} = x_1^4 \cdot g(x_1, \dots, x_n) \frac{\partial}{\partial x_1}$$

is of type II. We want to show that there is a function H , defined on a neighbourhood of the origin in \mathbf{R}^n such that

$$\tilde{X} = \left[x_1^2 H(x_1, \dots, x_n) \frac{\partial}{\partial x_1}, x_1^2 \frac{\partial}{\partial x_1} \right]$$

in a neighbourhood of the origin. The existence of such H follows from

SUB-LEMMA (2.1) *Let Z, X be vectorfields on \mathbf{R}^1 , which depend on real variables μ_1, \dots, μ_r , and which can be written in the form*

$$Z = x^k \cdot f(x, \mu_1, \dots, \mu_r) \frac{\partial}{\partial x}, \quad X = x^l \cdot g(x, \mu_1, \dots, \mu_r) \frac{\partial}{\partial x}$$

where f, g are C^∞ functions on \mathbf{R}^{r+1} (at least on a neighbourhood of the origin), $l \geq 2k$ and $f(0, 0, \dots, 0) \neq 0$.

Then there is a vectorfield Y , also depending on μ_1, \dots, μ_r , of the form

$$Y = x^k \cdot H(x, \mu_1, \dots, \mu_r) \frac{\partial}{\partial x},$$

such that

$$[Y, Z] = X$$

for all (x, μ_1, \dots, μ_r) in a small neighbourhood of the origin in \mathbf{R}^{r+1} .

PROOF of (2.1).

$$[Y, Z] = X \text{ or } \left[x^k \cdot H(x, \mu) \frac{\partial}{\partial x}, x^k \cdot f(x, \mu) \frac{\partial}{\partial x} \right] = x^l \cdot g(x, \mu) \frac{\partial}{\partial x}$$

is equivalent with

$$\begin{aligned} x^k \cdot H(x, \mu) \cdot \left[k \cdot x^{k-1} \cdot f(x, \mu) + x^k \frac{\partial f}{\partial x}(x, \mu) \right] \\ - x^k \cdot f(x, \mu) \left[k \cdot x^{k-1} \cdot H(x, \mu) + x^k \frac{\partial H}{\partial x}(x, \mu) \right] = x^l \cdot g(x, \mu). \end{aligned}$$

The terms with x^{2k-1} cancel and $l \geq 2k$, so we can divide by x^{2k} and obtain:

$$H(x, \mu) \cdot \frac{\partial f}{\partial x}(x, \mu) - f(x, \mu) \frac{\partial H}{\partial x}(x, \mu) = x^{l-2k} \cdot g(x, \mu)$$

Restricting ourselves to a small neighbourhood of the origin in the (x, μ) space, we may divide by f and obtain:

$$\frac{\partial H}{\partial x}(x, \mu) = \frac{\frac{\partial f}{\partial x}(x, \mu)}{f(x, \mu)} \cdot H(x, \mu) - x^{l-2k} \cdot \frac{g(x, \mu)}{f(x, \mu)}$$

This is an ordinary differential equation depending on the parameters $\mu = (\mu_1, \dots, \mu_r)$. Hence, by the existence and smoothness of solutions of differential equations depending on parameters, it follows that there is a function H which has the required properties.

PROOF OF LEMMA 3. For X and p as in the statement of the lemma (i.e. $j^3(X)(p) = 0$) we can find, using local coordinates and lemma 2, a neighbourhood U of p in M and vectorfields on M Y_1, \dots, Y_q and Z_1, \dots, Z_q such that

$$X|U = \sum_i [Y_i, Z_i]|U$$

and

$$j^1(Y_i)(p) = 0, \quad j^1(Z_i)(p) = 0 \text{ for all } i = 1, \dots, q.$$

Let $D : \chi(M) \rightarrow \chi(M)$ be any derivation. It follows from Lemma 1 that

$$D(X)(p) = D\left(\sum_i [Y_i, Z_i]\right)(p).$$

By the definition of derivation, this last expression equals

$$\sum_i [D(Y_i), Z_i](p) + \sum_i [Y_i, D(Z_i)](p)$$

which is zero because the 1-jets of Y_i and Z_i are zero in p . This proves lemma 3.

PROOF OF LEMMA 4. For D_U and $U \subset \mathbb{R}^n$ as in the statement of Lemma 4 and x_1, \dots, x_n coordinates on \mathbb{R}^n , we define the functions $D_{ij} : U \rightarrow \mathbb{R}$, $i, j = 1, \dots, n$ by

$$D_U \left(\frac{\partial}{\partial x_i} \right) = \sum D_{ij} \frac{\partial}{\partial x_j}.$$

We know that

$$\left[\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j} \right] \equiv 0,$$

for all i, j so

$$\begin{aligned} 0 \equiv D_U \left[\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j} \right] &= \left[D_U \left(\frac{\partial}{\partial x_i} \right), \frac{\partial}{\partial x_j} \right] + \left[\frac{\partial}{\partial x_i}, D_U \left(\frac{\partial}{\partial x_j} \right) \right] \\ &= - \sum_h \frac{\partial D_{ih}}{\partial x_j} \frac{\partial}{\partial x_h} + \sum_h \frac{\partial D_{jh}}{\partial x_i} \frac{\partial}{\partial x_h}. \end{aligned}$$

Hence, for all i, j, h , we have

$$\frac{\partial D_{ih}}{\partial x_j} = \frac{\partial D_{jh}}{\partial x_i};$$

as U is 1-connected, there are functions $\bar{D}_h : Y \rightarrow \mathbf{R}$, $h = 1, \dots, n$ such that

$$\frac{\partial \bar{D}_h}{\partial x_i} = -D_{ih}.$$

Now we define $\bar{Z} \in \chi(U)$

$$\bar{Z} = \sum \bar{D}_h \frac{\partial}{\partial x_i}.$$

From the above construction it follows that we have for each

$$i = 1, \dots, n : D_U \left(\frac{\partial}{\partial x_i} \right) = \left[\bar{Z}, \frac{\partial}{\partial x_i} \right].$$

Now we define the derivation

$$D_U^1 : \chi(U) \rightarrow \chi(U) \text{ by } D_U^1(X) = D_U(X) - [\bar{Z}, X];$$

clearly

$$D_U^1 \left(\frac{\partial}{\partial x_i} \right) \equiv 0$$

for all i .

Next we define the functions $D_{ijk} : U \rightarrow \mathbf{R}$, $i, j, k = 1, \dots, n$ by

$$D_U^1 \left(x_i \frac{\partial}{\partial x_j} \right) = \sum D_{ijk} \frac{\partial}{\partial x_k}.$$

First we show that all these functions are constant: as

$$\left[\frac{\partial}{\partial x_i}, x_j \frac{\partial}{\partial x_k} \right] = \delta_{ij} \frac{\partial}{\partial x_k}$$

we have

$$\begin{aligned} 0 \equiv D_U^1 \left[\frac{\partial}{\partial x_i}, x_j \frac{\partial}{\partial x_k} \right] &= \left[D_U^1 \left(\frac{\partial}{\partial x_i} \right), x_j \frac{\partial}{\partial x_k} \right] \\ &\quad + \left[\frac{\partial}{\partial x_i}, D_U^1 \left(x_j \frac{\partial}{\partial x_k} \right) \right] = \sum \frac{\partial D_{jkh}}{\partial x_i} \frac{\partial}{\partial x_h}. \end{aligned}$$

Hence, for all i, j, k, h ,

$$\frac{\partial D_{jkh}}{\partial x_i} \equiv 0,$$

so the functions D_{jkh} must be constant (because U is connected). We de-

note these constants by c_{jkh} . Next we want to show that

- a) $c_{ijk} = 0$ whenever $j \neq k$ and
- b) $c_{ijj} = c_{ikk}$ for all i, j, k .

To prove this we observe that

$$\left[x_i \frac{\partial}{\partial x_j}, x_k \frac{\partial}{\partial x_l} \right] = \delta_{jk} x_i \frac{\partial}{\partial x_l} - \delta_{li} x_k \frac{\partial}{\partial x_j}.$$

Applying D_U^1 to this, we obtain

$$\delta_{jk} \sum_h c_{ilh} \frac{\partial}{\partial x_h} - \delta_{li} \sum_h c_{kjh} \frac{\partial}{\partial x_h} = c_{ijk} \frac{\partial}{\partial x_l} - c_{kli} \frac{\partial}{\partial x_j} \dots^*$$

If we take in $^* k \neq l = j = i$ (this assumes that the dimension $n \neq 1$ because for $n = 1$ we cannot take $k \neq l$; if $n = 1$ however a) and b) above are trivially true) we obtain:

$$- \sum_h c_{kjh} \frac{\partial}{\partial x_h} = c_{ilk} \frac{\partial}{\partial x_l} - c_{kli} \frac{\partial}{\partial x_l}$$

from which it follows that $c_{kjh} = 0$ if $l \neq h$ which proves a) above.

Next we take in $^* k \neq j$ and $l = i$ and obtain (using the above result):

$$- c_{kjj} \frac{\partial}{\partial x_j} = - c_{kli} \frac{\partial}{\partial x_j} \text{ and hence:}$$

$$c_{kjj} = c_{kli} \text{ if } k \neq j, \text{ which implies b).}$$

From the above calculations it follows that for all i, j ,

$$D_U^1 \left(x_i \frac{\partial}{\partial x_j} \right) = \left[\sum_h c_{hhh} \frac{\partial}{\partial x_h}, x_i \frac{\partial}{\partial x_j} \right].$$

We now define $Z \in \chi(U)$ by

$$Z = \bar{Z} + \sum_h c_{hhh} \frac{\partial}{\partial x_h}$$

and observe that for all i, j ,

$$D_U \left(\frac{\partial}{\partial x_i} \right) = \left[Z, \frac{\partial}{\partial x_i} \right] \text{ and } D_U \left(x_i \frac{\partial}{\partial x_j} \right) = \left[Z, x_i \frac{\partial}{\partial x_j} \right];$$

it is not hard to see that Z is uniquely determined by these properties. In order to complete the proof of this Lemma we have to show that the derivation D_U^2 , defined by $D_U^2(X) = D_U(X) - [Z, X]$ is identically zero:

SUB-LEMMA (4.1). Let D_U and $U \subset \mathbf{R}^n$ be as in Lemma 4. If, for all i, j ,

$$D_U \left(\frac{\partial}{\partial x_i} \right) \equiv 0 \text{ and } D_U \left(x_i \frac{\partial}{\partial x_j} \right) \equiv 0$$

then $D_U(X) \equiv 0$ for all $X \in \chi(U)$.

PROOF of (4.1) We define the functions $D_{ijkl} : U \rightarrow \mathbf{R}$ by

$$D_U \left(x_i x_j \frac{\partial}{\partial x_k} \right) = \sum D_{ijkl} \frac{\partial}{\partial x_l}.$$

To prove that these functions are all constant one can proceed just as in the case with D_{ijk} above, but now we use the fact that

$$D_U \left(\left[\frac{\partial}{\partial x_i}, x_j x_k \frac{\partial}{\partial x_l} \right] \right) \equiv 0;$$

we omit the computation. We denote the corresponding constants again by c_{ijkl} . Next we observe that

$$\left[\sum_h x_h \frac{\partial}{\partial x_h}, x_i x_j \frac{\partial}{\partial x_k} \right] = x_i x_j \frac{\partial}{\partial x_k};$$

applying D_U to this we obtain:

$$\left[\sum_h x_h \frac{\partial}{\partial x_h}, \sum_l c_{ijkl} \frac{\partial}{\partial x_l} \right] = \sum c_{ijkl} \frac{\partial}{\partial x_l}, \text{ or } - \sum_l c_{ijkl} \frac{\partial}{\partial x_l} = \sum c_{ijkl} \frac{\partial}{\partial x_l};$$

hence all the constants c_{ijkl} are zero. In the same way one can show that

$$D_U \left(x_i x_j x_k \frac{\partial}{\partial x_l} \right) \equiv 0$$

for all i, j, k, l . Finally, we apply lemma 3 to obtain the proof: Let $X \in \chi(U)$ and $p \in U$, we want to show $D_U(X)(p) = 0$. There is a vectorfield $\hat{X} \in \chi(U)$ such that the coefficient functions of \hat{X} are polynomials of degree ≤ 3 and such that $j^3(X)(p) = j^3(\hat{X})(p)$. By our previous computations we have $D_U(X) \equiv 0$ and by lemma 3 we have $D(X)(p) = D(\hat{X})(p)$; hence $D_U(X)(p) = 0$, this proves (4.1).

PROOF OF THE THEOREM. For a given derivation $D : \chi(M) \rightarrow \chi(M)$ and an open $U \subset M$, we get an induced derivation $D_U : \chi(U) \rightarrow \chi(U)$. This D_U is constructed as follows:

For $X \in \chi(U)$ an $p \in U$ one defines $D_U(X)(p)$ to be $D(X)(p)$, where $\tilde{X} \in \chi(M)$ is some vectorfield which equals X on some open neighbourhood of p . Clearly $D_U(X)(p)$ is well defined (by Lemma 1) and D_U is a derivation on $\chi(U)$.

Now we take an atlas $\{U_i, \varphi_i(U_i) \rightarrow \mathbf{R}^n\}$ of M such that each U_i is connected and simply connected. Using the coordinates $x_j \varphi_i$ on each U_i we can apply Lemma 4 to each D_{U_i} and obtain on each U_i a vectorfield $Z_i \in \chi(U_i)$ such that $D_{U_i}(X) = [Z_i, X]$ for each $X \in \chi(U_i)$.

As D_{U_i} and D_{U_j} both restricted to $U_i \cap U_j$ are equal, Z_i and Z_j both restricted to $U_i \cap U_j$ also have to be equal. Hence there is a vectorfield $Z \in \chi(M)$ such that for each i , $Z_i = Z|_{U_i}$. It follows easily that, for each $X \in \chi(M)$, $D(X) = [Z, X]$.

3. Remark

One can also take, instead of $\chi(M)$, the set of vectorfields which respect a certain given structure. To be more explicit, let ω be a differential form on M defining a symplectic structure or a volume structure, and let $\chi_\omega(M)$ be the Lie-algebra of those vectorfields X for which $L_X \omega \equiv 0$ (L_X means : Lie derivative with respect to X). Now one can ask again whether every derivation $D : \chi_\omega(M) \rightarrow \chi_\omega(M)$ is induced by a vectorfield $Z \in \chi_\omega(M)$. This is in general not the case. Take for example $M = \mathbb{R}^n$ and $\omega = dx_1 \wedge \cdots \wedge dx_n$ the usual volume form and

$$Z = \sum_{i=1}^n x_i \frac{\partial}{\partial x_i}.$$

Then $Z \notin \chi_\omega(\mathbf{R}^n)$ but for each $X \in \chi_\omega(\mathbf{R}^n)$, $[Z, X] \in \chi_\omega(\mathbf{R}^n)$; so $[\cdot, -]$ is a derivation on $\chi_\omega(\mathbf{R}^n)$. This derivation cannot be induced by any $Z' \in \chi_\omega(\mathbf{R}^n)$.

REFERENCES

M. GEL'FAND and D. B. FUKS

- [1] Cohomologies of Lie algebra of tangential vectorfields of a smooth manifold, *Funktional'nyi Analiz i Ego Prilozheniya* 3 (1969) pp. 32-52. (translation: *Functional Analysis and its applications* 3 (1969), pp. 194-210.

(Oblatum 3-VIII-1972)

Mathematisch Instituut
Postbus 800
GRONINGEN (The Netherlands).