

COMPOSITIO MATHEMATICA

C. H. HOUGHTON

Ends of groups and baseless subgroups of wreath products

Compositio Mathematica, tome 27, n° 2 (1973), p. 205-211

http://www.numdam.org/item?id=CM_1973__27_2_205_0

© Foundation Compositio Mathematica, 1973, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ENDS OF GROUPS AND BASELESS SUBGROUPS OF WREATH PRODUCTS

C. H. Houghton

Introduction

In this paper we investigate the relationship between Specker's theory [4] of ends of groups and Hartley's results [2] on baseless subgroups of wreath products. The wreath product $W = A \wr B$ of groups A and B is the split extension of the group $K = A^B$ of functions from B to A , by B , with $f^b(x) = f(xb^{-1})$, for $f \in A^B$, $b, x \in B$. We shall assume throughout that A and B are non-trivial. Let α be an infinite cardinal and, for $f \in K$, let $\text{supp}(f)$ denote the support of f . We define K_α to consist of those $f \in K$ with $|\text{supp}(f)| < \alpha$ and let $W_\alpha = BK_\alpha \leq W$. We note that in the case $\alpha = \aleph_0$, K_α consists of those functions with finite support and W_α is the restricted wreath product $A \wr B$ of A and B .

Hartley [2] investigates baseless subgroups of W_α , that is, subgroups which have trivial intersection with the base group K_α . He gives conditions for baseless subgroups of W_α to be conjugate to subgroups of B and conditions for the existence of baseless subgroups which are maximal in some class of subgroups of W_α . He applies these results to construct locally finite groups with certain given sets of locally finite p -groups as Sylow p -subgroups. Our aim is to establish a connection between Hartley's results and the theory of ends, which we now consider.

For a group G , we define $Q_\alpha(G)$ to consist of those $S \subseteq G$ such that $|Sg \cap S'| < \alpha$, for all $g \in G$, where $S' = G \setminus S$. The set of α -ends of G consists of the ultrafilters of sets S in $Q_\alpha(G)$ with $|S| \geq \alpha$. We denote the number of α -ends of G by $e_\alpha(G)$; in the case $\alpha = \aleph_0$, we omit mention of the cardinal. From now on we assume $|G| \geq \alpha$, which is equivalent to the condition $e_\alpha(G) > 0$. Specker [4] shows that $e_\alpha(G) = 1, 2$, or is infinite and that $e_\alpha(G) = 2$ if and only if $\alpha = \aleph_0$ and G is infinite cyclic by finite. We note that $e_\alpha(G) = 1$ if and only if, for $S \subseteq G$, the condition $|Sg \cap S'| < \alpha$, for all $g \in G$, implies $|S| < \alpha$ or $|S'| < \alpha$.

The theory of ends of groups, with $\alpha = \aleph_0$, has been studied extensively (see Cohen [1], Stallings [5]); in particular, finitely generated groups with more than one end have been characterised. Various sufficient con-

ditions for G to have 1 end are known and some may be extended to the case of a general cardinal α . We do not pursue this point, as the results in the case $\alpha = \aleph_0$ are far from complete. We require the following result.

THEOREM (1): (Specker [4, p. 173 (a)]). *If either $|G| = \alpha > \aleph_0$ or $|G| = \alpha = \aleph_0$ and G is locally finite, then $e_\alpha(G)$ is infinite.*

Baseless subgroups and ends

Let L be a baseless subgroup of $W = A Wr B$. Then, for a unique $C \leq B, LA^B = CA^B$. Hartley shows [2, Lemma 3.2. (i)] that there is an element $g \in A^B$ such that $L = C^g$. It follows that the baseless subgroups of W_α are all the subgroups of W of the form C^g with $C \leq B, g \in A^B$ and $C^g \leq W_\alpha$. Throughout this section, let $S' = B \setminus S$, for $S \subseteq B$.

LEMMA (2): *Suppose $C \leq B$ and $g \in A^B$. Let $g(B) = \{a_i : i \in I\}$ and $B_i = \{b \in B : g(b) = a_i\}$. Then*

(i) $C^g \leq W_\alpha$ if and only if

$$|\bigcup_{i \in I} B_i c \cap B'_i| < \alpha,$$

for all $c \in C$,

(ii) if $|C| \geq \alpha, C^g$ is conjugate to C in W_α if and only if there is a subset J of I and a partition $\{D_j : j \in J\}$ of B with $D_j C = D_j$ such that

$$|\bigcup_{j \in J} D_j \cap B'_j| < \alpha.$$

PROOF: (i) If $c \in C, c^g = g^{-1} c g = c(g^{-1})^c g$ and so $c^g \in W_\alpha$ if and only if $|\text{supp } ((g^{-1})^c g)| < \alpha$. Now $((g^{-1})^c g)(x) = (g(xc^{-1}))^{-1} g(x)$, so

$$\text{supp } ((g^{-1})^c g) = \bigcup_{i \in I} B_i c \cap B'_i.$$

Thus $C^g \leq W_\alpha$ if and only if

$$|\bigcup_{i \in I} B_i c \cap B'_i| < \alpha, \text{ for all } c \in C.$$

(ii) Suppose $C^g = C^h$ with $h \in K_\alpha$. Then $z = gh^{-1}$ is in the centraliser of C and hence is constant on each left coset bC of C in B . Since $|\text{supp } (z^{-1}g)| = |\text{supp } (h)| < \alpha$ and $|C| \geq \alpha, z(B) \subseteq g(B)$. Let $J = \{i \in I : a_i \in z(B)\}$ and let $D_j = \{b \in B : z(b) = a_j\}$, for $j \in J$. Then $\{D_j : j \in J\}$ is a partition of B with $D_j C = D_j$ and

$$\bigcup_{j \in J} D_j \cap B'_j = \text{supp } (z^{-1}g) = \text{supp } (h)$$

so

$$|\bigcup_{j \in J} D_j \cap B'_j| < \alpha.$$

Conversely, suppose the given condition is satisfied and define $z \in A^B$ by $z(D_j) = (g(B_j))^{-1}$. Then $C^g = C^{zg}$ and

$$\text{supp}(zg) = \bigcup_{j \in J} D_j \cap B'_j,$$

so $|\text{supp}(zg)| < \alpha$.

THEOREM (3): *Suppose L is a baseless subgroup of W_α with $LA^B = CA^B$, $C \leq B$. If $e_\alpha(C) = 1$, then L is conjugate to C in W_α .*

PROOF: We have $L = C^g$ where the partition associated with g satisfies

$$|\bigcup_{i \in I} B_i c \cap B'_i| < \alpha,$$

for $c \in C$. For $i \in I$, $b \in B$ we have, for all $c \in C$,

$$B_i c \cap B'_i \cong (bC \cap B_i)c \cap (bC \cap B'_i) = b((C \cap b^{-1}B_i)c \cap (C \cap b^{-1}B'_i))$$

and hence $|(C \cap b^{-1}B_i)c \cap (C \setminus (C \cap b^{-1}B_i))| < \alpha$. Since $e_\alpha(C) = 1$, $|C \cap b^{-1}B_i| < \alpha$ or $|C \cap b^{-1}B'_i| < \alpha$ and so $|bC \cap B_i| < \alpha$ or $|bC \cap B'_i| < \alpha$. Let $M = \{(bC, i) : 0 < |bC \cap B_i| < \alpha\}$.

We first suppose $|C| > \alpha$ and take $N \subseteq M$, with $|N| \leq \alpha$. Then

$$|\bigcup_{(bC, i) \in N} (bC \cap B_i)^{-1}(bC \cap B_i)| \leq \alpha,$$

so there is an element $c \in C$ such that, for $(bC, i) \in N$, $c \notin (bC \cap B_i)^{-1}(bC \cap B_i)$ and $(bC \cap B_i)c \subseteq B'_i$. Then

$$\begin{aligned} |\bigcup_{(bC, i) \in N} bC \cap B_i| &= |\bigcup_{(bC, i) \in N} (bC \cap B_i)c| \\ &= |\bigcup_{(bC, i) \in N} (bC \cap B_i)c \cap B'_i| \\ &\leq |\bigcup_{i \in I} B_i c \cap B'_i| < \alpha. \end{aligned}$$

So $|N| < \alpha$ and we deduce that $|M| < \alpha$ and

$$|\bigcup_{(bC, i) \in M} bC \cap B_i| < \alpha.$$

In the other case, $|C| = \alpha$. Since we are assuming $e_\alpha(C) = 1$, Theorem 1 implies that $\alpha = \aleph_0$ and C is not locally finite. Let $D = \langle d_1, \dots, d_n \rangle$ be a finitely generated infinite subgroup of C . Since, for $j = 1, \dots, n$,

$$\bigcup_{i \in I} B_i d_j \cap B'_i$$

is finite, almost all B_i are fixed under right multiplication by d_1, \dots, d_n . So $H = \{i : B_i \neq B_i D\}$ is finite. If $(bC, i) \in M$ then $bC \cap B_i$ is finite and non-empty. Thus $bC \cap B_i \neq (bC \cap B_i)D = bC \cap B_i D$, so $B_i \neq B_i D$

and $i \in H$. Also, for some d_j , $(bC \cap B_i)d_j \cap B'_i \neq \emptyset$ so, for some $c \in C$, $bc \in B_i \cap B'_i d_j^{-1}$ and $b \in (B_i \cap B'_i d_j^{-1})C$. Thus if $(bc, i) \in M$, then $i \in H$ and $bC \subseteq FC$, where F is the finite set

$$\bigcup_{j=1}^n \left(\bigcup_{i \in I} B_i d_j \cap B'_i \right) d_j^{-1}.$$

So M is finite and hence

$$\bigcup_{(bc, i) \in M} bC \cap B_i$$

is also finite.

In each case, we now have

$$\left| \bigcup_{(bc, i) \in M} bC \cap B_i \right| < \alpha.$$

Furthermore,

$$|C| \geq \alpha \text{ and } bC = \bigcup_{i \in I} bC \cap B_i.$$

Thus, for $b \in B$, $|bC \cap B_i| \geq \alpha$ for some i and then $|bC \cap B_j| < \alpha$ for $j \neq i$. For $i \in I$, let $T_i = \{bc : |bc \cap B_i| \geq \alpha\}$ and let

$$D_i = \bigcup_{bc \in T_i} bc.$$

Put $J = \{i \in I : D_i \neq \emptyset\}$. Then $\{D_j : j \in J\}$ is a partition of B with $D_j = D_j C$. For $j \in J$,

$$D_j \cap B'_j \subseteq \bigcup_{(bc, i) \in M} bC \cap B_i$$

so

$$\left| \bigcup_{j \in J} D_j \cap B'_j \right| \leq \left| \bigcup_{(bc, i) \in M} bC \cap B_i \right| < \alpha.$$

Then Lemma 2 (ii) implies L is conjugate to C in W_α .

Theorem 1 describes a class of groups C with $e_\alpha(C)$ infinite. We now consider these groups in more detail.

LEMMA (4): *Suppose $|C| = \alpha > \aleph_0$ or $|C| = \alpha = \aleph_0$ and C is locally finite. Then there is a partition $\{C_1, C_2\}$ of C such that $|C_1| = |C_2| = \alpha$ and $|sC_1 t \cap C_2| < \alpha$, for all $s, t \in C$.*

PROOF: Consider α as an ordinal equivalent to none of its predecessors. Under the conditions of the theorem,

$$C = \bigcup_{\lambda < \alpha} H_\lambda$$

for some system of subgroups such that $|H_\lambda| < \alpha$ and $H_\lambda < H_\mu$ if $\lambda < \mu$.

Let

$$C_1 = \bigcup_{\lambda < \alpha} H_{2\lambda+1} \setminus H_{2\lambda}.$$

Then

$$C_2 = C \setminus C_1 = H_0 \cup \bigcup_{\lambda < \alpha} H_{2\lambda+2} \setminus H_{2\lambda+1}$$

and $|C_1| = |C_2| = \alpha$. Suppose $s, t \in C$. For some ordinal $\mu < \alpha$, we have $s, t \in H_\lambda$, for $\lambda \geq \mu$, and so $sH_\lambda t = H_\lambda$ and $s(H_{\lambda+1} \setminus H_\lambda)t = H_{\lambda+1} \setminus H_\lambda$. Thus $sC_1 t \cap C_2 \subseteq sH_\mu t$ and $|sC_1 t \cap C_2| < \alpha$.

A refinement of this argument gives a proof of Theorem 1. For a positive integer n ,

$$C_i = \bigcup_{\lambda < \alpha} H_{n\lambda+i} \setminus H_{n\lambda+i-1}$$

is in $Q_\alpha(C)$ and $\{C_1, \dots, C_{n-1}, C_n \cup H_0\}$ is a partition of C . Thus the number of α -ends of C is unbounded.

Suppose $e_\alpha(C) = 2$. Then $\alpha = \aleph_0$ and C has an infinite cyclic normal subgroup $E = \langle e \rangle$ of finite index. The centraliser H of E in C has index 1 or 2 and so $C = H \cup Hd = E(F \cup Fd)$, where $d = 1$ or $d \in C \setminus H$ and the finite set F is a set of coset representatives for E in H . Let $P = \{e^i : i > 0\}$, $C_1 = P(F \cup Fd)$ and $C_2 = C \setminus C_1$. For $s \in H$, $t \in C$, $s(F \cup Fd)t$ is a set of coset representatives for E in C and $sC_1 t = Ps(F \cup Fd)t = P\{e^{i(f)}f : f \in F \cup Fd\}$, for some finite set of integers $\{i(f)\}$. Then $sC_1 t \cap C_2$ is finite, and similarly $sC_2 t \cap C_1$ is finite, for $s \in H$, $t \in C$. Thus we have proved the following result.

LEMMA (5): *Suppose C has 2 ends. Then there is a partition $\{C_1, C_2\}$ of C , with $C_1 \in Q(C)$, such that the sets $\{c \in C_2 : cC_1 \cap C_2 \text{ is finite}\}$ and $\{c \in C_1 : cC_2 \cap C_1 \text{ is finite}\}$ are infinite.*

Once again we consider C as a subgroup of B in W_α , but now assuming $e_\alpha(C) > 1$. We recall that this implies that either $e_\alpha(C)$ is infinite or $\alpha = \aleph_0$ and $e_\alpha(C) = e(C) = 2$.

THEOREM (6): *Suppose $C \leq B$ with $e_\alpha(C) > 1$. Then there is a baseless subgroup L of W_α satisfying the following conditions:*

- (i) $LA^B = CA^B$,
- (ii) L is not conjugate to C in W_α ,
- (iii) if M is a baseless subgroup of W_α with $M > L$ then $C = N_D(C)$ where $MA^B = DA^B$ with $D \leq B$. Furthermore, if either $|C| = \alpha > \aleph_0$ or otherwise $|C| = \alpha = \aleph_0$ and C is locally finite or has 2 ends, then the assumptions of (iii) imply the following stronger result:
- (iv) $|C \cap uCv^{-1}| < \alpha$ for all $u, v \in D \setminus C$.

PROOF: Since $e_\alpha(C) > 1$, there is a partition $\{C_1, C_2\}$ of C with $|C_i| \geq \alpha$ and $|C_1 c \cap C_2| < \alpha$, for all $c \in C$. Then $|C_2 c \cap C_1| = |C_2 \cap C_1 c^{-1}| < \alpha$, for all $c \in C$. Take $a \in A$, $a \neq 1$, and define $g \in A^B$ by $g(C_1) = a$, $g(C'_1) = 1$, where C'_1 denotes $B \setminus C_1$. For $c \in C$, $|(C_1 c \cap C'_1) \cup (C'_1 c \cap C_1)| = |(C_1 c \cap C_2) \cup (C_2 c \cap C_1)| < \alpha$, so Lemma 2(i) implies $C^g \leq W_\alpha$. Let $\{D_1, D_2\}$ be a partition of B with $D_i C = D_i$ and suppose $C \subseteq D_1$ (here we allow $D_2 = \emptyset$). Then $D_1 \cap C'_1 \supseteq C_2$ and $D_1 \cap C_1 = C_1$ so, from Lemma 2 (ii), C^g is not conjugate to C in W_α .

Under the assumptions of (iii) we have $C^g \leq D^h \leq W_\alpha$ for some $h \in A^B$. Then $C^g = C^h$ so hg^{-1} centralises C and the parts of the partition of B corresponding to hg^{-1} consist of unions of left cosets bC . Suppose $hg^{-1}(C) = a_1$ and so $h(C_1) = a_1 a$, $h(C_2) = a_1$, and $h(b) = hg^{-1}(b)$ for $b \notin C$. Let $B_1 = \{b : h(b) = a_1 a\}$, $B_2 = \{b : h(b) = a_1\}$. Then

$$B_1 = C_1 \cup \bigcup_{bC \in T_1} bC, \quad B_2 = C_2 \cup \bigcup_{bC \in T_2} bC,$$

where T_1, T_2 are disjoint sets of cosets distinct from C . From Lemma 2(i), $|B_i d \cap B'_i| < \alpha$, for $d \in D$ and $i = 1, 2$. If $d \in N_D(C) \setminus C$, then $|C_i d \cap dC| \geq \alpha$ so $dC \cap B_1 \neq \emptyset \neq dC \cap B_2$ and $dC \in T_1 \cap T_2 = \emptyset$. Thus $N_D(C) = C$.

We now suppose that either $|C| = \alpha > \aleph_0$ or otherwise $|C| = \alpha = \aleph_0$ and C is locally finite or has 2 ends. We can then assume that the partition $\{C_1, C_2\}$ has been chosen as described in Lemma 4 or 5. Given $u, v \in D \setminus C$, we have $|C_i u \cap B'_i| < \alpha$, $|C_i v \cap B'_i| < \alpha$. Thus for some $G_i \subseteq C_i$ with $|C_i \setminus G_i| < \alpha$, we have $G_i u \cup G_i v \subseteq B_i$. Then

$$G_1 u \cup G_1 v \subseteq \bigcup_{bC \in T_1} bC.$$

Thus, for $c_1 \in G_1$, $c_2 \in G_2$, $c_1 u C \neq c_2 v C$ and $c_1 v C \neq c_2 u C$ and so $c_1^{-1} c_2$, $c_2^{-1} c_1 \notin u C v^{-1}$. From Lemmas 4 and 5, since $|C_1 \setminus G_1| < \alpha$, we may choose $c_1 \in G_1$ such that $|c_1 C_2 \cap C_1| < \alpha$ and hence $|c_1^{-1} C_1 \cap C_2| < \alpha$. Now $\{c_2 \in C_2 : c_2 \notin u C v^{-1}\} \supseteq c_1^{-1} G_2 \cap C_2$ so $C_2 \cap u C v^{-1} \subseteq C_2 \cap c_1^{-1} (C \setminus G_2) \subseteq (C_2 \cap c_1^{-1} C_1) \cup c_1^{-1} (C_2 \setminus G_2)$ and hence $|C_2 \cap u C v^{-1}| < \alpha$. Similarly, $|C_1 \cap u C v^{-1}| < \alpha$ and so $|C \cap u C v^{-1}| < \alpha$.

We note a consequence of Theorems 3 and 6. This generalises a result in [3] that if $A^{(B)}$ is the group of functions from B to A with finite support and B acts as in the wreath product, then the first cohomology set $H^1(B, A^{(B)})$ is trivial if and only if B has 1 end. However $H^1(B, A^{(B)})$ is precisely the set of conjugacy classes of complements of the base group $A^{(B)}$ in $A \text{ wr } B = W_{\aleph_0}$.

COROLLARY (7): Suppose $C \leq B$ with $|C| \geq \alpha$. Every baseless subgroup L of W_α such that $LA^B = CA^B$ is conjugate to C in W_α if and only if $e_\alpha(C) = 1$.

For certain subgroups C of B , Theorem 6 gives sufficient conditions for the existence of baseless subgroups L , with $LA^B = CA^B$, which are maximal in certain classes. Hartley's Theorem B, the first part of Theorem A and the second part of Theorem D [2] may be deduced. Using Corollary 7, the other parts of his Theorems A and D lead to new sufficient conditions for a group to have 1 end. Thus, unless they are infinite cyclic by finite, radical groups with non-periodic Hirsch-Plotkin radical have 1 end and so also do uncountable locally finite groups satisfying the normaliser condition. In this connection we mention the conjecture that uncountable locally finite groups have 1 end.

I would like to thank Dr. Hartley for sending me a preprint of [2].

REFERENCES

- [1] D. E. COHEN: Groups of cohomological dimension one. *Lecture Notes in Mathematics* 245. (Springer-Verlag, Berlin, 1972).
- [2] B. HARTLEY: Complements, baseless subgroups and Sylow subgroups of infinite wreath products. *Compositio Mathematica*, 26 (1973) 3–30.
- [3] C. H. HOUGHTON: Ends of groups and the associated first cohomology groups. *J. London Math. Soc.*, 6 (1972) 81–92.
- [4] E. SPECKER: Endenverbände von Räumen und Gruppen. *Math. Ann.* 122 (1950) 167–174.
- [5] J. STALLINGS: Group theory and three-dimensional manifolds. *Yale Mathematical Monographs*, 4. (Yale University Press, New Haven, 1971).

(Oblatum: 23–I–1973)

Department of Pure Mathematics
University College
Cardiff,
Wales, U.K.