

COMPOSITIO MATHEMATICA

KNUD LØNSTED

On the trivialization of line bundles over schemes

Compositio Mathematica, tome 27, n° 2 (1973), p. 217-221

http://www.numdam.org/item?id=CM_1973__27_2_217_0

© Foundation Compositio Mathematica, 1973, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE TRIVIALIZATION OF LINE BUNDLES OVER SCHEMES

Knud Lønsted

1. Introduction

Consider a line bundle L over a scheme S , and assume that $L \oplus \cdots \oplus L$ is trivial. We show the existence of a finite flat covering $\rho : T \rightarrow S$ such that $\rho^*(L)$ is trivial over T , and give a condition which ensures ρ to be étale. This condition is verified e.g. when S is of finite type over a field of characteristic zero. In particular, if S is connected and simply connected in the algebraic sense, then L is trivial.

The corresponding theorem for complex *topological* line bundles over, say, a 1-connected CW-complex S is essentially trivial, as pointed out to us by J. Dupont and V. Lundsgaard Hansen: Note that the first Chern class yields a bijection from the isomorphism classes of line bundles to $H^2(S, \mathbb{Z})$ (see [4, Chap. 1]). By the universal coefficient theorem one gets $H^2(S, \mathbb{Z})$ torsionfree as $H_1(S, \mathbb{Z}) = 0$. Since in this case we have

$$c_1(L \oplus \cdots \oplus L) = c_1(L^{\otimes n}) = n \cdot c_1(L)$$

then $n \cdot c_1(L) = 0$ implies $c_1(L) = 0$, whence L is trivial.

With a minor change of vocabulary in our proof of the algebraic case below, it becomes a proof of the complex analytic version.

In the remarks at the end of the paper we have inserted a translation into commutative algebra, and there are a few comments on ‘universality’ of the covering ρ in the cases where S is either the spectrum of a Dedekind ring or a smooth projective curve.

This paper grew out of an unsuccessful attempt to solve Serre’s problem on algebraic vector bundles over affine spaces, and we are indebted to Serre for giving a counterexample to the naive generalization of the theorem below to vector bundles of arbitrary rank.

The notations will follow [2]. Note specifically that μ is used to denote the multiplicative group scheme, instead of the usual G_m . All schemes and morphisms of schemes are over some fixed base scheme that never occurs in the notation, and *sheaves* are taken to be sheaves in the *fffp*-topology (faithfully flat finitely presented) relative to the base scheme.

We denote the group scheme of the n ’th roots of unity by ${}_n\mu$ and ${}_n\mu_S$

equals ${}_n\mu \times S$ for any scheme S . Finally, if S is noetherian and connected, $\pi_1(S)$ denotes the algebraic fundamental group of S [3], and for any prime p , $\pi_1(S)^{(p)}$ denotes the non- p -part of $\pi_1(S)$. We set $\pi_1(S)^{(1)} = \pi_1(S)$.

2. Results

THEOREM: *Let S be any scheme and \mathcal{L} a quasi-coherent locally free \mathcal{O}_S -Module of rank 1 such that $\mathcal{L} \oplus \cdots \oplus \mathcal{L}$ (n summands) is free, for some integer n . Then there exists a finite flat morphism $\rho : T \rightarrow S$ of degree at most n such that $\rho^*(\mathcal{L})$ is a free \mathcal{O}_T -Module.*

Furthermore, if ${}_n\mu_S$ is étale over S , then ρ may be chosen étale.

COROLLARY: *If, in addition to all the hypothesis of the theorem, S is connected and of finite type over a field of exponent p , so that $(p, n) = 1$ and $\pi_1(S)^{(p)} = 0$, then \mathcal{L} is a free \mathcal{O}_S -Module.*

This conclusion holds in particular for $p = 1$ and S connected and simply connected.

PROOFS: We may assume that $n > 1$. Let \mathfrak{G} be a *ffffp*-sheaf of groups over the base scheme, and $\mathfrak{H} \subset \mathfrak{G}$ an invariant subgroup. To the exact sequence

$$1 \rightarrow \mathfrak{H} \rightarrow \mathfrak{G} \rightarrow \mathfrak{G}/\mathfrak{H} \rightarrow 1$$

belongs an exact sequence of pointed sets (see [3, III.4] for this and the sequel) of which we shall need the following piece

$$(1) \quad \tilde{H}^0(S, \mathfrak{G}) \rightarrow \tilde{H}^0(S, \mathfrak{G}/\mathfrak{H}) \xrightarrow{\circ} \tilde{H}^1(S, \mathfrak{H}) \rightarrow \tilde{H}^1(S, \mathfrak{G})$$

Recall that one defines $\tilde{H}^0(S, \mathfrak{G}) = \text{Hom}(S, \mathfrak{G})$ and that $\tilde{H}^1(S, \mathfrak{G})$ is the set of isomorphism classes of *ffffp*-torsors over S under \mathfrak{G} . It is an immediate consequence of the definitions of the maps in (1) that the sequence is functorial in S . Hence for any morphism $\rho : T \rightarrow S$ we have a commutative diagram:

$$(2) \quad \begin{array}{ccccccc} \tilde{H}^0(S, \mathfrak{G}) & \rightarrow & \tilde{H}^0(S, \mathfrak{G}/\mathfrak{H}) & \xrightarrow{\circ} & \tilde{H}^1(S, \mathfrak{H}) & \rightarrow & \tilde{H}^1(S, \mathfrak{G}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \tilde{H}^0(T, \mathfrak{G}) & \rightarrow & \tilde{H}^0(T, \mathfrak{G}/\mathfrak{H}) & \xrightarrow{\circ} & \tilde{H}^1(T, \mathfrak{H}) & \rightarrow & \tilde{H}^1(T, \mathfrak{G}) \end{array}$$

We apply these generalities to $\mathfrak{G} = \mathfrak{GL}_n$, $\mathfrak{H} = \mu$, and $\mathfrak{G}/\mathfrak{H} = \mathfrak{PGL}_{n-1}$. Then $\tilde{H}^1(S, \mu) = \text{Pic}(S)$ and the map $\beta : \text{Pic}(S) \rightarrow \tilde{H}^1(S, \mathfrak{GL}_n)$ sends the isomorphism class of an invertible sheaf \mathcal{M} of \mathcal{O}_S -Modules onto the isomorphism class of $\mathcal{M} \oplus \cdots \oplus \mathcal{M}$ (n summands). Denote the isomorphism class of \mathcal{L} in $\text{Pic}(S)$ by l . By assumption $\beta(l) = e$, where e is the marked point in $\tilde{H}^1(S, \mathfrak{GL}_n)$, whence $l = \partial(x)$ for some $x \in \tilde{H}^0(S, \mathfrak{PGL}_{n-1})$. We set $l' = \tilde{H}^1(\rho, \mu)(l)$, and have $l' = \partial(x')$ with $x' = \tilde{H}^0(\rho,$

$\mathfrak{P}\mathcal{G}\mathcal{L}_{n-1})(x)$. In order to get $l' = 0$ we must have $x' = \alpha(y)$, where $y \in \tilde{H}^0(T, \mathcal{G}\mathcal{L}_n)$ and $\alpha : \tilde{H}^0(T, \mathcal{G}\mathcal{L}_n) \rightarrow \tilde{H}^0(T, \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1})$ is the canonical map, coming from the projection $\pi : \mathcal{G}\mathcal{L}_n \rightarrow \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1}$. This means that we want to lift the morphism $x' : T \rightarrow \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1}$ through π to some morphism $y : T \rightarrow \mathcal{G}\mathcal{L}_n$.

Let $i : \mathcal{E}\mathcal{L}_n \rightarrow \mathcal{G}\mathcal{L}_n$ denote the inclusion of the unimodular group. Then we have an exact sequence

$$1 \rightarrow {}_n\mu \rightarrow \mathcal{E}\mathcal{L}_n \xrightarrow{p} \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1} \rightarrow 1$$

where $p = \pi \circ i$. Now define the morphism $\rho : T \rightarrow S$ as the pull-back of p by $x : S \rightarrow \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1}$,

$$(3) \quad \begin{array}{ccc} T & \longrightarrow & \mathcal{E}\mathcal{L}_n \\ \rho \downarrow & & \downarrow p \\ S & \xrightarrow{x} & \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1} \end{array}$$

Since p is finite and flat, so is ρ , and by construction the morphism $x' = x \circ \rho$ factors through $\mathcal{E}\mathcal{L}_n$, hence through $\mathcal{G}\mathcal{L}_n$. Furthermore, if we pull the cartesian diagram (3) back over S , we get

$$\begin{array}{ccc} T & \longrightarrow & \mathcal{E}\mathcal{L}_{n,S} \\ \downarrow & & \downarrow p_S \\ S & \xrightarrow{(x, 1_S)} & \mathfrak{P}\mathcal{G}\mathcal{L}_{n-1,S} \end{array}$$

where ${}_n\mu_S = \text{Ker}(p_S)$, and $p_S = (p, 1_S)$, thus p_S is étale if and only if ${}_n\mu_S$ is an étale S -group scheme. This proves the theorem.

The corollary follows from the observation that under the mentioned hypothesis the morphism $\rho : T \rightarrow S$ constructed above has a section, which provides a lifting $S \rightarrow \mathcal{G}\mathcal{L}_n$ of x , i.e. $\partial(x) = 0$.

3. Remarks

In terms of commutative algebra the theorem states: Given a commutative ring A and an invertible A -module P such that $P \oplus \cdots \oplus P$ (n summands) is free. Then there exists a finite flat (possibly unramified) extension $A \rightarrow B$ such that $P \otimes_A B$ is a free B -module.

Suppose that A is a Dedekind ring. It follows from the isomorphism $K_0(A) = \mathbb{Z} \oplus \text{Pic}(A)$ (see e.g. [1, Chap. XIII]) and the cancellation law, that for an invertible A -module P the conditions $P \oplus \cdots \oplus P$ (n summands) free, and $P^{\otimes n}$ free, are equivalent. If A is of arithmetic type, e.g. the ring of integers in a number field, then $\text{Pic}(A)$ is finite, and we conclude that every invertible P becomes trivial after some finite flat ex-

tension of A (thus this is true for every projective A -module, in view of their structure), and the knowledge of the order of P in $\text{Pic}(A)$ enables one to predict where this extension might ramify. Furthermore, one may even take a fixed ‘universal’ extension of A that works for all projective A -modules: choose an extension for each element in $\text{Pic}(A)$ and take their tensor product over A .

Assume finally that S is a smooth projective connected curve defined over an algebraically closed field k of exponent p , and let n be an integer for which $(p, n) = 1$. In this case there exists a universal finite étale covering $\rho_n : S_n \rightarrow S$ of degree n^{2g} , where g is the genus of S , and S_n is a smooth projective connected curve, such that $\rho_n^*(\mathcal{L})$ is trivial for every invertible sheaf \mathcal{L} on S for which $\mathcal{L} \oplus \cdots \oplus \mathcal{L}$ (n summands) is trivial. This is seen as follows: Let \mathcal{L} be such a sheaf. Then the étale covering $\rho : T \rightarrow S$ constructed in the proof of the theorem is an abelian Galois covering of S with Galois group ${}_n\mu(k)$. We replace T by one of its connected components, which again is an abelian Galois covering of S , the Galois group of which is a subgroup of ${}_n\mu(k)$. This new covering is also denoted by $\rho : T \rightarrow S$. By a theorem due to Serre [5, Chap. VI p. 128] every connected abelian covering of S is the pull-back of some separable isogeny of its Jacobian J , via one of the canonical immersions $S \rightarrow J$. Let $A \rightarrow J$ be an isogeny that gives the covering ρ . Then the kernel is obviously killed by n , and one readily sees that there exists an isogeny $J \rightarrow A$ such that the composition $J \rightarrow A \rightarrow J$ is multiplication by n . Now define $\rho_n : S_n \rightarrow S$ to be the pull-back of $J \xrightarrow{\times n} J$. Since $\text{Ker}(J \xrightarrow{\times n} J) = {}_nJ$ is a finite constant subgroup of J of rank n^{2g} , the morphism ρ_n has the properties claimed above, and we clearly have a morphism $\sigma : S_n \rightarrow T$ for which $\rho \circ \sigma = \rho_n$. Thus $\rho_n^*(\mathcal{L})$ is trivial.

The hypothesis k algebraically closed may be weakened to the assumptions that S has a k -rational point, and that ${}_n\mu$ and ${}_nJ$ are constant group schemes over k .

Added in proof

R. Fossum has called our attention to a paper by G. Garfinkel: Generic Splitting Algebras for Pic, Pacific J. Math. 35 (1970), 369–380, which deals with torsion elements of the Picard group in the affine case, also treated in [2, p. 376]. It contains our theorem in the case of a Dedekind domain.

REFERENCES

- [1] H. BASS: *Algebraic K-theory*. (W. A. Benjamin 1968).
- [2] M. DEMAZURE et P. GABRIEL: *Groupes algébriques*, tome 1. (Masson et Cie – North Holland 1970).
- [3] A. GROTHENDIECK: *SGA 1: Revêtements Étales et Groupe Fondamental*. Lecture Notes in Mathematics 224. (Springer Verlag 1971).
- [4] F. HIRZEBRUCH: *Topological Methods in Algebraic Geometry*. 3rd Edition. (Springer Verlag 1966).
- [5] J. -P. SERRE: *Groupes algébriques et corps de classes*. (Hermann 1959).

(Oblatum 30–IV–1973)

Matematisk Institut
Universitetsparken 5
2100 Copenhagen
Denmark