

# COMPOSITIO MATHEMATICA

JAMES AUSTIN FRENCH

**A condition equivalent to covering dimension  
for normal spaces**

*Compositio Mathematica*, tome 28, n° 3 (1974), p. 223-227

[http://www.numdam.org/item?id=CM\\_1974\\_\\_28\\_3\\_223\\_0](http://www.numdam.org/item?id=CM_1974__28_3_223_0)

© Foundation Compositio Mathematica, 1974, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## A CONDITION EQUIVALENT TO COVERING DIMENSION FOR NORMAL SPACES

James Austin French<sup>1</sup>

In this paper a concept called *boundary covering dimension* is defined. Boundary covering dimension is proven to be equivalent to covering dimension for normal spaces. Also included is a definition of complete boundary covering dimension. Complete boundary covering dimension is proven to be equivalent to complete covering dimension for paracompact  $T_2$ -spaces (complete covering dimension is equivalent to covering dimension for paracompact  $T_2$ -spaces).

NOTATIONS: If  $X$  is a space and  $V \subset X$ , then  $B(V)$  denotes the boundary of  $V$ . If  $X$  is a space,  $M \subset X$ , and  $H \subset M$ , then  $B(M, H)$  denotes the boundary in the subspace  $M$  of  $H$ .

DEFINITIONS: The collection  $G$  of subsets of the space  $X$  is *discrete* means every point of  $X$  is contained in an open set that intersects at most one element of  $G$ .

*Covering dimension* is denoted by *dim*.  $\dim X \leq n$  means if  $G$  is a *finite* open cover of  $X$ , then there exists an open cover  $R$  of  $X$  such that  $R$  refines  $G$  and  $\text{ord } R \leq n + 1$ .

*Boundary covering dimension* is denoted by *bcd*. For  $n \geq 1$ ,  $\text{bcd } X \leq n$  means if  $H$  is a closed set,  $W$  is an open set,  $H \subset W$ , and  $G$  is a *finite* open cover of  $X$ , then there are an open set  $V$  and discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $H \subset V \subset W$ ,  $\bigcup_{j=1}^n G_j$  refines  $G$ , and  $B(V) = \bigcup (\bigcup_{j=1}^n G_j)$ . Now  $\text{bcd } X = n$  means  $\text{bcd } X \leq n$  and  $\text{bcd } X \not\leq n - 1$ .

*Complete covering dimension* is denoted by *complete dim*. Complete  $\dim X \leq n$  means if  $G$  is an open cover of  $X$ , then there exists an open cover  $R$  of  $X$  such that  $R$  refines  $G$  and  $\text{ord } R \leq n + 1$ .

*Complete boundary covering dimension* is denoted by *complete bcd*. For  $n \geq 1$ , complete  $\text{bcd } X \leq n$  means if  $H$  is a closed set,  $W$  is an open set,  $H \subset W$ , and  $G$  is an open cover of  $X$ , then there exist an open set  $V$  and discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $H \subset V \subset W$ ,  $\bigcup_{j=1}^n G_j$  refines  $G$ , and  $B(V) = \bigcup (\bigcup_{j=1}^n G_j)$ .

<sup>1</sup> The work for that paper was done while the author was on a Cottrell College Science Grant for Research Corporation.

REMARK: What is meant by  $\text{bcd } X \leq 0$ ? Let us note that the definition for  $\text{bcd}$  can be written another way.  $\text{bcd } X \leq n$  means if  $H$  is a closed set,  $W$  is an open set,  $H \subset W$ , and  $G$  is a finite open cover of  $X$ , then there are an open set  $V$  and a collection  $T$  of at most  $n$  elements such that  $H \subset V \subset W$ , each element of  $T$  is a discrete collection of closed sets,  $\bigcup T$  refines  $G$ , and  $B(V) = \bigcup (\bigcup T)$ . So when  $n = 0$ ,  $T = \phi$  and  $B(V) = \phi$ . Thus,  $\text{bcd } X \leq 0$  means  $\text{Ind } X \leq 0$ . In our proofs we will not be considering the case where  $n = 0$  since it will be evident what the proof would be for  $n = 0$ .

LEMMA 1: *If  $X$  is a topological space,  $\text{bcd } X \leq n$ , and  $M$  is a closed subset of  $X$ , then  $\text{bcd } M \leq n$ . (The proof is straight-forward and will not be given.)*

LEMMA 2: *If  $X$  is a topological space and  $\text{bcd } X \leq n$ , then if  $H$  is a closed set,  $W$  is an open set,  $H \subset W$ , and  $G$  is a finite open cover of  $X$ , then there are an open set  $V$  and finite discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $H \subset V \subset W$ ,  $\bigcup_{i=1}^n G_i$  refines  $G$ , and  $B(V) = \bigcup (\bigcup_{i=1}^n G_i)$ . (The proof is straight-forward and will not be given.)*

LEMMA 3: *If each of  $G_1, G_2, \dots, G_n$  is a finite open cover of the topological space  $X$ , then there is a finite open cover  $G$  of  $X$  such that for every  $i \in \{1, \dots, n\}$ ,  $G$  refines  $G_i$ .*

PROOF: For every  $p \in X$ , let  $T(p) = \{g | \exists i \in \{1, \dots, n\} \text{ such that } g \in G_i \text{ and } p \in g\}$ . Let  $G = \{\bigcap T(p) | p \in X\}$ .  $G$  is a finite open cover of  $X$  such that for every  $i \in \{1, \dots, n\}$ ,  $G$  refines  $G_i$ .

LEMMA 4: *If  $X$  is a paracompact  $T_2$ -space,  $M \subset X$ ,  $M$  is closed,  $n$  is a positive integer,  $G$  is a collection of open sets of  $X$  covering  $M$ , and no point of  $M$  belongs to  $n+1$  elements of  $G$ , then there exist discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $\bigcup_{j=1}^n G_j$  refines  $G$  and  $\bigcup (\bigcup_{j=1}^n G_j) = M$ .*

PROOF: Since every paracompact  $T_2$ -space is collectionwise normal, Theorem 2 of [1] can be applied to prove the Lemma.

LEMMA 5: *If  $X$  is a normal topological space,  $M \subset X$ ,  $M$  is closed,  $n$  is a positive integer,  $G$  is a finite collection of open sets of  $X$  covering  $M$ , and no point of  $M$  belongs to  $n+1$  elements of  $G$ , then there exist discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $\bigcup_{j=1}^n G_j$  refines  $G$  and  $\bigcup (\bigcup_{j=1}^n G_j) = M$ .*

PROOF: The proof is similar to the proofs of Theorem 1 and Theorem 2 of [1]. Only normality is needed instead of collectionwise normality since the open cover  $G$  is finite.

**THEOREM 1:** *If  $X$  is a normal topological space, then  $\text{bcd } X = \text{dim } X$ .*

**PROOF:**

*Part I: Show  $\text{dim } X \leq \text{bcd } X$ .* Assume  $n$  is a positive integer and  $\text{bcd } X \leq n$ . Assume  $G$  is a finite open cover of  $X$ . Let  $G = \{g_1, \dots, g_m\}$ . Let  $H_1 = g_1 - (\bigcup_{j=2}^m g_j) = X - \bigcup_{j=2}^m g_j$ . Now  $g_1$  is an open set containing the closed set  $H_1$ . Since  $\text{bcd } X \leq n$ , by Lemma 2, there exist an open set  $V_1$ , and finite discrete collections  $L_1, L_2, \dots, L_n$  of closed sets such that  $H_1 \subset V_1 \subset g_1$ ,  $\bigcup_{j=1}^n L_j$  refines  $G$ , and  $B(V_1) = \bigcup (\bigcup_{j=1}^n L_j)$ . For every  $j \in \{1, \dots, n\}$ , let  $S(1, j) = L_j$ . Let  $X_1 = X$ .

Assume  $k$  is a positive integer such that  $1 \leq k \leq m$  and for every  $i \in \{1, \dots, k\}$ ,

(a)  $X_i = X - \bigcup_{j=1}^{i-1} V_j = X_{i-1} - V_{i-1}$

(b)  $H_i = X_i - \bigcup_{j=i+1}^m g_j$

(c)  $H_i \subset V_i \subset g_i$ ,  $V_i \subset X_i$ ,  $V_i$  open in  $X_i$  (Hence  $X_i$  and  $H_i$  are closed in  $X$ )

(d)  $\forall j \in \{1, \dots, n\}$ ,  $S(i, j)$  is a finite discrete collection of closed sets and  $S(i, j)$  refines  $G$ , and

(e)  $\bigcup_{j=1}^n (\bigcup_{i=1}^k S(i, j)) = \bigcup_{j=1}^n B(X_j, V_j)$ .

Now let  $X_{k+1} = X - \bigcup_{j=1}^k V_k = X_k - V_k$  and let  $H_{k+1} = X_{k+1} - \bigcup_{j=k+2}^m g_j$

Now  $H_{k+1} \subset g_{k+1}$ . For every  $j \in \{1, \dots, n\}$ , let  $E_j = \{e(j, w) | w \in S(k, j)\}$  and  $F_j = \{f(j, w) | w \in S(k, j)\}$  be finite discrete collections of open sets such that  $F_j$  refines  $G$ ,  $\forall w \in S(k, j)$   $w \subset e(j, w) \subset \overline{e(j, w)} \subset f(j, w)$  and  $f(j, w)$  intersects only one element of  $S(k, j)$ , and let  $T_j = \{f(j, w) \cap X_{k+1} | w \in S(k, j)\} \cup \{g - (\bigcup E_j) \cap X_{k+1} | g \in G\}$ . By Lemma 3, there is a finite cover  $T$  of  $X_{k+1}$  such that each element of  $T$  is open in  $X_{k+1}$ , and for every  $j \in \{1, \dots, n\}$ ,  $T$  refines  $T_j$ . By Lemma 1,  $\text{bcd } X_{k+1} \leq n$  so by Lemma 2 there exist a set  $V_{k+1}$ , open in  $X_{k+1}$ , and finite discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $H_{k+1} \subset V_{k+1} \subset g_{k+1}$ ,  $\bigcup_{j=1}^n G_j$  refines  $T$ , and  $B(X_{k+1}, V_{k+1}) = \bigcup (\bigcup_{j=1}^n G_j)$ .  $\forall j \in \{1, \dots, n\}$ ,  $\forall w \in S(k, j)$ , let  $b(j, w) = \{w\} \cup \{h | h \in G_j \text{ and } h \subset f(j, w)\}$ .  $\forall j \in \{1, \dots, n\}$ , let  $M_j = \{h | h \in G_j \text{ and } \forall w \in S(k, j), h \not\subset b(j, w)\}$  and let  $S(k+1, j) = \{\bigcup b(j, w) | w \in S(k, j)\} \cup M_j$ .  $\forall j \in \{1, \dots, n\}$ ,  $S(k+1, j)$  is a finite collection of closed sets and  $S(k+1, j)$  refines  $G$ .

Assume  $j \in \{1, \dots, n\}$ . It will now be shown that  $S(k+1, j)$  is discrete. Since  $S(k+1, j)$  is finite, we need only to show that no two elements of  $S(k+1, j)$  intersect. It should be clear that no two elements of  $M_j$  intersect and no two elements of  $\{\bigcup b(j, w) | w \in S(k, j)\}$  intersect. Assume  $\exists w_0 \in S(k, j)$  and  $h_0 \in M_j$  such that  $\bigcup b(j, w_0)$  intersects  $h_0$ .

*Case 1:  $\exists h_1 \in G_j$  such that  $h_1 \subset f(j, w_0)$  and  $h_1$  intersects  $h_0$ .* Since no two elements of  $G_j$  intersect,  $h_0 = h_1$ .  $\forall w \in S(k, j)$ ,  $h_0 \notin b(j, w)$  since

$h_0 \in M_j$ . But  $h_0$ , which is  $h_1$ , is an element of  $b(j, w_0)$ . Contradiction.

*Case 2:  $h_0$  intersects  $w_0$ .* Since  $G_j$  refines  $T$  which refines  $T_j$ , there is an element  $g_0$  of  $T_j$  such that  $h_0 \subset g_0$ . Thus  $g_0$  intersects  $w_0$ , and  $w_0 \subset \bigcup E_j$ . No element of  $\{[g - (\bigcup E_j)] \cap X_{k+1} | g \in G\}$  intersects  $\bigcup E_j$  so  $g_0 \in \{f(j, w) \cap X_{k+1} | w \in S(k, j)\}$ . Thus  $\exists w_1 \in S(k, j)$  such that  $g_0 = f(j, w_1) \cap X_{k+1}$ . This means  $h_0 \subset f(j, w_1)$ . So  $h_0 \in b(j, w_1)$ . Since  $h_0 \in M_j$ , we know that  $\forall w \in S(k, j)$ ,  $h_0 \notin b(j, w)$ . This means  $h_0 \notin b(j, w_1)$ , but  $h_0 \in b(j, w_1)$ . Contradiction. Therefore, no two elements of  $S(k+1, j)$  intersect.

It follows that  $\bigcup (\bigcup_{j=1}^n S(k+1, j)) = \bigcup_{j=1}^{k+1} B(X_j, V_j)$ . We have now completed our inductive definition. Thus each of  $S(m, 1), S(m, 2), \dots, S(m, n)$  is a finite discrete collection of closed sets that refines  $G$ .  $\forall j \in \{1, \dots, n\}$ , let  $Z_j$  be a finite discrete collection of open sets such that  $S(m, j)$  refines  $Z_j$  and  $Z_j$  refines  $G$ .  $\forall i \in \{1, \dots, m\}$ , let  $V'_i = V_i - [\bigcup (\bigcup_{j=1}^n Z_j)]$ . Now  $\{V'_1, V'_2, \dots, V'_m\}$  is a finite collection of mutually exclusive closed sets such that  $\forall i \in \{1, \dots, m\}$ ,  $V'_i \subset g_i$ . Let  $Z_{n+1} = \{a_1, \dots, a_m\}$  be a finite discrete collection of open sets such that  $\forall i \in \{1, \dots, m\}$ ,  $V'_i \subset a_i \subset g_i$ . Let  $Z = \bigcup_{j=1}^{n+1} Z_j$ .  $Z$  is an open cover of  $X$  such that  $Z$  refines  $G$  and  $\text{ord } Z \leq n+1$ . Thus  $\dim X \leq n$ .

*Part II: Show  $\text{bcd } X \leq \dim X$ .* Assume  $n$  is a positive integer and  $\dim X \leq n$ . Assume  $H$  is a closed set,  $W$  is an open set  $H \subset W$ , and  $G$  is a finite open cover of  $X$ . Let  $F$  be a finite open cover of  $X$  such that  $F$  refines  $G$  and every element of  $F$  that intersects  $H$  is a subset of  $W$ . Let  $T = \{t_i | i = 1, \dots, k\}$  be a finite open cover of  $X$  such that  $T$  refines  $F$ ,  $\text{ord } T \leq n+1$ , and if  $i \neq j$ , then  $t_i \neq t_j$ . Let  $R = \{r_i | i = 1, \dots, k\}$  be an open cover of  $X$  such that  $\forall i \in \{1, \dots, k\}$ ,  $r_i \subset t_i$ . Let  $V = \bigcup \{r_i | i \in \{1, \dots, k\} \text{ and } r_i \text{ intersects } H\}$ . Assume  $p \in B(V)$  and  $n+1$  elements of  $R$  contain  $p$ . There exist positive integers  $j_1 < j_2 < \dots < j_{n+1} \leq k$  such that  $\forall i \in \{1, \dots, n+1\}$ ,  $p \in r_{j_i}$ . Since  $R$  is finite,  $\exists j_{n+2} \in \{1, \dots, k\}$  such that  $p \in B(r_{j_{n+2}})$ .  $\forall i \in \{1, \dots, n+2\}$ ,  $p \in t_{j_i}$  since  $r_{j_i} \subset t_{j_i}$ . Thus,  $n+2$  elements of  $T$  contain  $p$ , which is a contradiction. Therefore no point of  $B(V)$  is contained by  $n+1$  elements of  $R$ . By Lemma 5, there exist discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $\bigcup_{j=1}^n G_j$  refines  $G$  and  $B(V) = \bigcup (\bigcup_{j=1}^n G_j)$ . So  $\text{bcd } X \leq n$ .

**THEOREM 2:** *If  $X$  is a paracompact  $T_2$ -space, then  $\text{bcd } X = \dim X = \text{complete bcd } X = \text{complete dim } X$ .*

**PROOF:** Assume  $X$  is a paracompact  $T_2$ -space. Theorem II.6 page 22 of [2] makes it clear  $\dim X = \text{complete dim } X$ , Theorem 1 gives us  $\text{bcd } X = \dim X$ . It is trivial that  $\text{bcd } X \leq \text{complete bcd } X$ . It will now be shown that  $\text{complete bcd } X \leq \text{bcd } X$ . Assume  $n$  is positive integer and  $\text{bcd } X \leq n$ . Thus  $\dim X \leq n$ , and hence  $\text{complete dim } X \leq n$ .

Assume  $H$  is a closed set,  $W$  is an open set,  $H \subset W$ , and  $G$  is an open cover of  $X$ . Let  $F$  be an open cover of  $X$  such that  $F$  refines  $G$  and every element of  $F$  that intersects  $H$  is a subset of  $W$ . Let  $T = \{t_b | b \in B\}$  be a locally finite open cover of  $X$  such that  $T$  refines  $F$ ,  $\text{ord } T \leq n+1$ , and if  $b_1, b_2 \in B$  and  $b_1 \neq b_2$  then  $t_{b_1} \neq t_{b_2}$  (Theorem 3 of [1] assures the existence of such a  $T$ ). Let  $R = \{r_b | b \in B\}$  be an open cover of  $X$  such that  $\forall b \in B, \overline{r_b} \subset t_b$ . Let  $V = \bigcup \{r_b | b \in B \text{ and } r_b \text{ intersects } H\}$ . Assume  $p \in B(V)$  and  $n+1$  elements of  $R$  contain  $p$ . There exist  $n+1$  elements  $b_1, b_2, \dots, b_{n+1}$  of  $B$  such that  $\forall i \in \{1, \dots, n+1\}, p \in r_{b_i}$ . Since  $R$  is locally finite, there exists  $b_{n+2} \in B$  such that  $p \in B(r_{b_{n+2}})$ .  $\forall i \in \{1, \dots, n+2\}, p \in t_{b_i}$ , since  $\overline{r_{b_i}} \subset t_{b_i}$ . Thus  $n+2$  elements of  $R$  contain  $p$ , which is a contradiction. Therefore, no point of  $B(V)$  is contained by  $n+1$  elements of  $R$ . By Lemma 4, there exist discrete collections  $G_1, G_2, \dots, G_n$  of closed sets such that  $\bigcup_{j=1}^n G_j$  refines  $G$  and  $B(V) = \bigcup (\bigcup_{j=1}^n G_j)$ . Thus  $\text{complete bcd } X \leq n$ . Therefore  $\text{bcd } X = \dim X = \text{complete bcd } X = \text{complete dim } X$ .

**COROLLARY:** *Assume  $X$  is a normal topological space. Then  $\dim X \leq n$  if and only if for all mutually exclusive closed sets  $H$  and  $K$ , for every finite (the word 'finite' can be deleted for  $X$  a paracompact  $T_2$ -space) open cover  $G$  of  $X$ , there exist mutually exclusive open sets  $D_H$  and  $D_K$  and a collection  $T$  of at most  $n$  elements such that  $H \subset D_H, K \subset D_K$ , every element of  $T$  is a discrete collection of closed sets,  $\bigcup T$  refines  $G$ , and  $X - (D_H \cup D_K) = \bigcup (\bigcup T)$ .*

**PROOF:** The proof follows from Theorem 1 (If  $X$  is  $T_2$ -paracompact and the open cover  $G$  is not necessarily finite, then the proof follows from Theorem 2).

**REMARK:** Note the similarity between the above Corollary and the following familiar theorem on large inductive dimension (denoted  $\text{Ind}$ ): For  $X$  normal,  $\text{Ind } X \leq n$  if and only if for all mutually exclusive open sets  $H$  and  $K$ , there exist mutually exclusive open sets  $D_H$  and  $D_K$  and a closed set  $T$  such that  $H \subset D_H, K \subset D_K, \text{Ind } T \leq n-1$ , and  $X - (D_H \cup D_K) = T$ . The similarity of the Corollary and this theorem on  $\text{Ind}$  enable one to pattern some  $\dim$  proofs after some  $\text{Ind}$  proofs.

#### REFERENCES

- [1] JAMES AUSTIN FRENCH: A characterization of covering dimension for collectionwise normal spaces. *Proceedings of the American Mathematical Society*, Vol. 25, 3, (1970) 646-649.
- [2] J. NAGATA: Modern dimension theory. *Bibliotheca Math.*, vol. 6, Interscience, New York, 1965, MR 34 #8380.

(Oblatum 1-X-1973)

David Lipscomb College  
Nashville  
Tenn. 37203