

COMPOSITIO MATHEMATICA

M. VAN DER PUT

**A problem on coefficient fields and equations
over local rings**

Compositio Mathematica, tome 30, n° 3 (1975), p. 235-258

<http://www.numdam.org/item?id=CM_1975__30_3_235_0>

© Foundation Compositio Mathematica, 1975, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A PROBLEM ON COEFFICIENT FIELDS AND EQUATIONS OVER LOCAL RINGS

M. van der Put

Introduction

Let R be a noetherian local ring, $\mathfrak{m}$ its maximal ideal and $\pi : R \rightarrow K$ the natural map of R onto its residue field K . Given a subfield k of R (hence R has equal characteristic) does there exist a coefficient field of R containing k ?

Stated in a more general way: Given subfields $k \subset l$ of K and a ring-homomorphism $\phi : k \rightarrow R$ such that $\pi\phi = \text{id}_k$, does ϕ extend to a ringhomomorphism $\Phi : l \rightarrow R$ with $\pi\Phi = \text{id}_l$?

As is well known, the answer is “yes” when R is complete and l/k is separable (See [3]).

In this paper we consider the case when l/k is inseparable. A necessary condition for the existence of Φ is the existence for all $n \geq 1$ of a ring-homomorphism $\Phi_n : l \rightarrow R/\mathfrak{m}^n$ with $\pi\Phi_n = \text{id}_l$ and $\Phi_n|_k = \phi$ (For convenience all the natural maps $R/\mathfrak{m}^a \rightarrow R/\mathfrak{m}^b$, $\infty \leq a \leq b \leq 1$, are denoted by π). Assume that this condition is satisfied and let H_t denote the set of all $\Phi : l \rightarrow R/\mathfrak{m}^t$ with $\pi\Phi = \text{id}_l$ and $\Phi|_k = \phi$. By assumption $H_t \neq \emptyset$ for all t and clearly $\varprojlim H_t = \{\Phi : l \rightarrow \hat{R} \mid \pi\Phi = \text{id}_l \text{ and } \Phi|_k = \phi\}$. The problem splits in two parts:

- (i) Is $\varprojlim H_t \neq \emptyset$?
- (ii) If $\varprojlim H_t \neq \emptyset$ does there exist a $\Phi : l \rightarrow R$ with $\Phi|_k = \phi$ and $\pi\Phi = \text{id}_l$?

Results

In Section 1 it is shown that (i) and (ii) have a positive answer for l/k finitely generated and R an s -ring, i.e. R has the following property: For every ideal F in $R[X_1, \dots, X_N]$ there exists a function $s : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $x = (x_1, \dots, x_N) \in R^N$ with $F(x) \in \mathfrak{m}^{s(n)}$ there exists a $x' \in R^N$ with $x' \equiv x(\mathfrak{m}^n)$ and $F(x') = 0$.

Further a list of s -rings is given. In Sections 2, 3 it is shown that $\varprojlim H_t \neq \emptyset$ if $\dim_l \Omega_{l/k} < \infty$. In Sections 4, 5, 6 a proof is given of the statement: A complete local ring of equal characteristic is an s -ring. In Section 7 it is shown that some complete local rings of unequal characteristic are s -rings.

1. l/k finitely generated

DEFINITION: A local ring R is called an s -ring if for any set $F = (F_1, \dots, F_k)$ of elements in $R[X] = R[X_1, \dots, X_N]$ there exists a function $s: \mathbb{N} \rightarrow \mathbb{N}$, $s(n) \geq n$ for all n , such that: For every $x \in R^N$ with $F(x) \equiv 0(\mathfrak{m}^{s(n)})$ there exists $x' \in R^N$ with $x' \equiv x(\mathfrak{m}^n)$ and $F(x') = 0$.

EXAMPLE:

(1) Any Henselian discrete valuation ring R , such that the quotient field of $\hat{R}$ is separable over the quotient field of R (equivalently R is Henselian and excellent) is an s -ring (see M. Greenberg [4]).

(2) Any complete local ring of equal characteristic is an s -ring. This statement is close to an approximation theorem of M. Artin (see [2], Theorem (6.1)). Since there seems to be no proof available we will give a proof in Sections 4, 5 and 6.

(3) If R is the Henselization of a local ring R_0 which is of essentially finite type over R_1 and R_1 is a field or an excellent discrete valuation ring of equal characteristic, then R is an s -ring. This follows from (2) and M. Artin's approximation theorem ([2], Theorem 1.10).

(4) Any analytic local ring over a complete valued field k ($[k:k^p] < \infty$ if $\text{char } k = p \neq 0$) is an s -ring. We will discuss this in Section 8.

(1.1) **THEOREM:** Let R be an s -ring with residue field K , let $k \subset l \subset K$ be subfields such that l/k is finitely generated and $\phi: k \rightarrow R$ a ringhomomorphism with $\pi\phi = \text{id}_k$. There exists a positive integer v such that $H_v \neq \emptyset$ implies $\varprojlim H_t \neq \emptyset$ and ϕ extends to $\Phi: l \rightarrow R$.

PROOF: The field l can be considered as the quotient field of $A = k[X_1, \dots, X_N]/(F_1, \dots, F_k)$, where the images of $X_1, \dots, X_t$ in l form a transcendence base of l/k . The map $\phi: k \rightarrow R$ extends to

$$\phi^*: k[X_1, \dots, X_N] \rightarrow R[X_1, \dots, X_N]$$

in the obvious way and we obtain a set of polynomials $\phi^*(F)$ in $R[X_1, \dots, X_N]$. Let s be its s -function and put $v = s(1)$. The condition

$H_v \neq \emptyset$ is equivalent to the existence of $x \in R^N$ with $\phi^*(F)(x) \equiv 0(\mathfrak{m}^{s(n)})$ and $\pi(x_i) = \bar{X}_i$ where $\bar{X}_i$ is the image of X_i in l .

There exists $x' \in R^N$ with $x' \equiv x(\mathfrak{m})$, $\phi^*(F)(x') = 0$. Consequently we have a map $\Phi : A = k[X_1, \dots, X_N]/(F) \rightarrow R$ such that $\pi\Phi = \text{id}_A$. This map extends to l , the quotient field of A .

REMARK: (1.1) solves both problems (i) and (ii) for finitely generated field extension l/k and s -rings R .

2. Complete regular local rings

In this section we assume that R is a complete regular local ring with residue field K and $\text{chc } R = \text{chc } K = p > 0$. We assume $d = \dim R$ and denote by $t_1, \dots, t_d \in R$ a base for the maximal ideal. Further we always take $l = K$.

(2.1) THEOREM: Let $k \subset K$ and $\phi : k \rightarrow R$ be given such that $\pi\phi = \text{id}_k$. Assume that $H_t \neq \emptyset$ for all t . If $\dim_K \Omega_{K/k} < \infty$ then $\varprojlim H_t \neq \emptyset$ and ϕ extends to $\Phi : K \rightarrow R$.

PROOF: This is divided in some lemmata.

DEFINITION: Let G^t be the group of all k -automorphisms γ of $K[[T]]/(T)^t$ satisfying: $\gamma \equiv 1(\mathfrak{m})$; $\gamma(T_i) = T_i$ ($i = 1, \dots, d$). Let G_n^t ($n \leq t$) denote the subgroup of G^t consisting of the γ 's with $\gamma \equiv 1(\mathfrak{m}^n)$.

(2.2) LEMMA:

- (1) Let $\psi_0 \in H_t$ be given then $H_t = \psi_0 G^t$.
- (2) $\psi_0 G_n^t = \{\psi \in H_t \mid \psi \equiv \psi_0(\mathfrak{m}^n)\}$.

PROOF (1): For $\psi \in H_t$ we make the extension $\psi^e : K[[T]]/(T)^t \rightarrow R/\mathfrak{m}^t$ given by $\psi^e(T_i) = t_i$ ($i = 1, \dots, d$). This is an isomorphism. Then $\psi_0^{-1}\psi^e \in G^t$. Conversely for $\gamma \in G^t$ we have $\psi = \psi_0 \gamma \in H_t$.

- (2) If $\psi = \psi_0 \gamma$ then $\psi \equiv \psi_0(\mathfrak{m}^n)$ if and only if $\gamma \equiv 1(\mathfrak{m}^n)$.

DEFINITION: $\chi : G_n^t \rightarrow \text{Der}_k(K, (T)^n/(T)^{n+1})$ is the map given by $\chi(\gamma)(\lambda) = \gamma(\lambda) - \lambda$ (where $\lambda \in K$; $\gamma \in G_n^t$).

(2.3) LEMMA: The image V of χ satisfies:

- (1) $V + V \subset V$
- (2) $a^n V \subset V$ for all $a \in K$

- (3) V is a constructible subset of the finite-dimensional vectorspace $\text{Der}_k(K, (T)^n/(T)^{n+1})$.

PROOF: $\gamma \in G_n^t$ can explicitly be described by $\gamma(\lambda) = \sum_{|\alpha| < t} \gamma_\alpha(\lambda) T^\alpha$, where: each γ_α is a k -linear map of $K \rightarrow K$, $\gamma_0 = \text{id}_K$, $\gamma_\alpha = 0$ if $0 < |\alpha| < n$, and for all $a, b \in K$ and

$$\alpha : \gamma_\alpha(ab) = \sum_{\alpha_1 + \alpha_2 = \alpha} \gamma_{\alpha_1}(a) \gamma_{\alpha_2}(b).$$

Further

$$\chi(\gamma)(\lambda) = \sum_{|\alpha|=n} \gamma_\alpha(\lambda) T^\alpha \bmod (T)^{n+1}.$$

Clearly $\chi(\gamma\gamma^*) = \chi(\gamma) + \chi(\gamma^*)$, hence (1). Further for $a \in K$, $\gamma \in G_n^t$ we define $\gamma^a \in G_n^t$ by $\gamma^a(\lambda) = \sum_{|\alpha| < t} a^{|\alpha|} \gamma_\alpha(\lambda) T^\alpha$. So we proved (2).

(3) Let $\gamma : K \rightarrow K[[T]]/(T)^t$ be a homomorphism such that $\gamma \equiv 1(\mathfrak{m})$ and γ is k -linear. Then for any β with $p^\beta \geq t$ we find that $\gamma|_{K^{p^\beta}(k)}$ is the ordinary inclusion map, or what amounts to the same γ is $K^{p^\beta}(k)$ -linear. Let $a_1, \dots, a_d$ be a p -base of K/k (i.e. $\Omega_{K/k}$ has base $da_1, \dots, da_d$) then $K = K^{p^\beta}(k)[a_1, \dots, a_d] = K^{p^\beta}(k)[X_1, \dots, X_d]/(F)$ where F is some set of polynomials.

Consider F as a set of polynomials with coefficients in $K[[T]]/(T)^t$ then there exists a natural bijection between G_n^t and

A is the set of elements $(x_1, \dots, x_d) \in (K[[T]]/(T)^t)^d$ such

$$\text{that } F(x_1, \dots, x_d) = 0 \quad \text{and} \quad (x_1, \dots, x_d) \equiv (a_1, \dots, a_d)(m^n)$$

Consider the map

$$\begin{aligned} x^* : G_n^t &\xrightarrow{x} \text{Der}_k(K, (T)^n/(T)^{n+1}) \simeq \text{Hom}_K(\Omega_{K/k}, (T)^n/(T)^{n+1}) \\ &\simeq \text{Hom}_K(Kda_1 + \dots + Kda_d, (T)^n/(T)^{n+1}) \simeq ((T)^n/(T)^{n+1})^d. \end{aligned}$$

The image of χ^* is the same as the image of $A - (a_1, \dots, a_d)$ in $((T)^n/(T)^{n+1})^d$. Since A is an algebraic set $/K$ this image is constructible. Hence also W is constructible.

(2.4) LEMMA: Let $(n, p) = 1$ and let $W \neq \{0\}$ be a subset of K satisfying $W + W \subseteq W$ and $a^n W \subseteq W$ for all $a \in K$. Then $W = K$. (provided that K is infinite).

PROOF: We may suppose that $1 \in W$. Let W_0 be the smallest subset of K which satisfies $1 \in W_0$, $W_0 + W_0 \subseteq W_0$, $a^n W_0 \subseteq W_0$ for all n . Then any element of W_0 has the form $\sum_i a_i^n$. Hence W_0 is a subring of K . For $f, g \in W_0$, $g \neq 0$ we have $f/g = g^{-n} f \cdot g^{n-1} \in W_0$. So W_0 is a subfield.

Since K is infinite also W_0 is infinite. Take $x \in K$ and let T be an indeterminate. Consider the polynomial

$$p(T) = \frac{(x+T)^n - x^n - T^n}{nT} = x^{n-1} + \dots + xT^{n-2}.$$

For every $\lambda \in W_0^*$, $p(\lambda) \in W_0$. Take distinct elements $\lambda_1, \dots, \lambda_{n-2} \in W_0^*$ and let $p(\lambda_i) = a_i \in W_0$. Then

$$p(T) = \sum_{i=1}^{n-2} a_i \prod_{j \neq i} \left(\frac{T - \lambda_j}{\lambda_i - \lambda_j} \right)$$

and belongs to $W_0[T]$. Hence the coefficient x in $p(T)$ belongs to W_0 . So $W = W_0 = K$.

(2.5) LEMMA: *The image of $\chi : G_n^t \rightarrow \text{Der}_k(K, (T)^n/(T)^{n+1})$ is a K -linear subspace.*

PROOF: If $(n, p) = 1$ this follows from (2.3) part (1) and (2) and (2.4). If $p|n$ we have to use that the image W is a constructible subset. Take $z \neq 0$, $z \in \text{Der}_k(K, (T)^n/(T)^{n+1})$ then $W \cap Kz$ is a constructible set, hence is finite or cofinite in Kz . Property (2) of (2.3) implies that either $Kz \subset W$ or $Kz \cap W = \{0\}$. So W is a K -linear subspace.

Conclusion of the proof (2.1).

Let $H_n^* = \bigcap_{m \geq n} \text{im}(H_m \rightarrow H_n)$. It suffices to show that $H_{n+1}^* \rightarrow H_n^*$ is surjective since it follows that $\emptyset \neq \varprojlim H_n^* \subseteq \varprojlim H_n$. Choose $\phi_0 \in H_n^*$ and for $t > n$ let $\tilde{H}_t$ be the preimage of ϕ_0 in H_t . If we can show that $\bigcap_{t > n} \text{im}(\tilde{H}_t \rightarrow \tilde{H}_{n+1}) \neq \emptyset$ then any $\phi_1 \in \bigcap_{t > n} \text{im}(\tilde{H}_t \rightarrow \tilde{H}_{n+1})$ satisfies $\phi_1 \in H_{n+1}^*$ and ϕ_1 is mapped onto $\phi_0 \in H_n^*$.

Take some $\alpha \in \tilde{H}_t$ and consider the map $[\alpha] : \tilde{H}_{n+1} \rightarrow G_{n+1}^{n+1}$ given by $[\alpha](\alpha\gamma) = \gamma$ for all $\gamma \in G_{n+1}^{n+1}$. Then we have an induced map

$$\tilde{H}_t \rightarrow \tilde{H}_{n+1} \xrightarrow{[\alpha]} G_{n+1}^{n+1} \xrightarrow{\chi} \text{Der}_k(K, (T)^n/(T)^{n+1})$$

which depends on the choice of $\alpha \in \tilde{H}_t$ but for which the image is independent of $\alpha \in \tilde{H}_t$. According to (2.5) the image is a finite dimensional

vectorspace over K . Hence $\text{im}(\tilde{H}_t \rightarrow \tilde{H}_{n+1})$ is constant for $t \gg n$ and $\bigcap \text{im}(\tilde{H}_t \rightarrow \tilde{H}_{n+1}) \neq \emptyset$.

3. Complete local rings

In this section we extend (2.1) to a more general case:

(3.1) THEOREM: *Let R be a complete local ring with residue field K and let subfields $k \subset l \subset K$ and a homomorphism $\tau: k \rightarrow R$ with $\pi\tau = \text{id}_k$ be given. Suppose that $H_t \neq \emptyset$ for all t . Then if $\dim_l \Omega_{l/k} < \infty$ then $\varprojlim H_t \neq \emptyset$ and τ extends to a $\phi: l \rightarrow R$ with $\pi\phi = \text{id}_l$.*

PROOF: First we introduce some notation. For any local ring R of characteristic $p > 0$ we define $\mathfrak{m}^{[n]} =$ the ideal generated by $\{x^{p^n} | x \in \mathfrak{m}\}$. If $\mathfrak{m}$ is generated by e elements then $\mathfrak{m}^{p^n \cdot e} \subseteq \mathfrak{m}^{[n]} \subseteq \mathfrak{m}^{p^n}$.

Instead of working with the powers of $\mathfrak{m}$ (as in Sections 1, 2) we can also work with the sequence of ideals $\mathfrak{m}^{[n]}$. Let $H_{[n]}$ denote the set of ringhomomorphisms $\phi: l \rightarrow R/\mathfrak{m}^{[n]}$ such that $\pi\phi = \text{id}_l$ and $\phi|_k = \tau$. By assumption $H_{[n]} \neq \emptyset$. For each $\phi \in H_{[n]}$ we form $\phi|^{p^n}(k) \rightarrow R/\mathfrak{m}^{[n]}$. This map is independent of the choice of ϕ and we will denote it by τ_n . Further $l^{p^n}(k)$ will be abbreviated with l_n .

Indeed, $x \in l_n$ has the form $\sum a_i x_i^{p^n} (a_i \in k, x_i \in l)$ and for $\phi, \phi^* \in H_{[n]}$ we have

$$\phi(x) - \phi^*(x) = \sum \tau(a_i)(\phi(x_i) - \phi^*(x_i))^{p^n}.$$

This is 0 since $\phi(x_i) - \phi^*(x_i) \in \mathfrak{m}$.

We define $A_n = R/\mathfrak{m}^{[n]} \otimes_{i_n} l$. In the next lemma we enumerate some properties of A_n .

- (3.2) LEMMA: (1) *Each A_n is a local ring and noetherian if $\dim \Omega_{l/k} < \infty$.*
 (2) *The natural map $A_{n+1} \rightarrow A_n$ is surjective and has kernel $\mathfrak{m}(A_{n+1})^{[n]}$.*
 (3) *$A = \varprojlim A_n$ is a complete local ring and noetherian if $\dim \Omega_{l/k} < \infty$.*
 (4) *$A/\mathfrak{m}(A)^{[n]} = A_n$.*
 (5) *There is a natural bijection $\chi_n: \text{Hom}_R(A, R/\mathfrak{m}^{[n]}) \rightarrow H_{[n]}$ and all diagrams*

$$\begin{array}{ccc} \text{Hom}_R(A, R/\mathfrak{m}^{[n]}) & \xrightarrow{\chi_n} & H_{[n]} \\ \uparrow & & \uparrow \\ \text{Hom}_R(A, R/\mathfrak{m}^{[n+1]}) & \xrightarrow{\chi_{n+1}} & H_{[n+1]} \end{array} \quad \text{are commutative.}$$

PROOF: (1) A_n is clearly local. If $\dim \Omega_{l/k} < \infty$ and $a_1, \dots, a_s$ is a p -base of l/k then for all n , $l = l_n[a_1, \dots, a_s]$. Hence A_n is a finite $R/\mathfrak{m}^{[n]}$ -module and thus noetherian.

(2) The map $\rho : A_{n+1} \rightarrow A_n$ decomposes as follows:

$$R/\mathfrak{m}^{[n+1]} \otimes_{l_{n+1}} l \xrightarrow{\alpha} R/\mathfrak{m}^{[n]} \otimes_{l_{n+1}} l \xrightarrow{\beta} R/\mathfrak{m}^{[n]} \otimes_{l_n} l,$$

where α and β are the obvious maps. Clearly $\ker \rho \supseteq \mathfrak{m}(A_{n+1})^{[n]}$. The kernel of β is generated by $\{\tau_n(x^{p^n}) \otimes 1 - 1 \otimes x^{p^n} | x \in l\}$. Take $\phi \in H_{[n+1]}$ then $\ker \rho$ is generated by $\mathfrak{m}^{[n]}/\mathfrak{m}^{[n+1]} \otimes_{l_{n+1}} l$ and

$$\{(\phi(x) \otimes 1 - 1 \otimes x)^{p^n} | x \in l\}.$$

Hence $\ker \rho \subseteq \mathfrak{m}(A_{n+1})^{[n]}$.

(3) That A is a complete local ring (possibly not noetherian) follows from its definition. Let $\dim_l \Omega_{l/k} < \infty$ and let $a_1, \dots, a_s$ be a p -base of l/k . Choose elements $b_1, \dots, b_s \in R$ with $\pi(b_i) = a_i$ ($i = 1, \dots, s$). Consider the sequence of maps $\phi_n : R[[y_1, \dots, y_s]] \rightarrow A_n$ given by $y_i \mapsto b_i \otimes 1 - 1 \otimes a_i$. This sequence of R -homomorphisms is coherent and each ϕ_n is surjective. Hence $\phi = \varprojlim \phi_n : R[[y_1, \dots, y_s]] \rightarrow A$ is a surjective R -homomorphism and A is noetherian.

(4) $A/\mathfrak{m}(A)^{[n]} = \varprojlim A_k/\mathfrak{m}(A_k)^{[n]} = A_n$ according to (2).

(5) For every n we have a map $l \rightarrow A_n = R/\mathfrak{m}^{[n]} \otimes_{l_n} l$ by $x \mapsto 1 \otimes x$. This induces a map $l \xrightarrow{i} A$. Define χ_n by $\chi_n(\phi) = \phi \circ i$. This makes the diagrams commutative. Further $\text{Hom}_R(A, R/\mathfrak{m}^{[n]}) = \text{Hom}_R(A_n, R/\mathfrak{m}^{[n]}) = \text{Hom}_R(R/\mathfrak{m}^{[n]} \otimes_{l_n} l, R/\mathfrak{m}^{[n]}) =$ the set of l_n -linear homomorphisms $\phi : l \rightarrow R/\mathfrak{m}^{[n]} = H_{[n]}$.

Conclusion of the proof of (3.1)

According to the lemma $\varprojlim H_t \simeq \text{Hom}_R(A, R)$ and A has the form $R[[y_1, \dots, y_s]]/G$ where G is some ideal.

Given is $\text{Hom}_R(A, R/\mathfrak{m}^t) \neq \emptyset$ for all t . Then by a theorem on the existence of an s -function for ideals in $R[[y_1, \dots, y_s]]$ (see Section 4, Theorem (4.1)) we can conclude $\text{Hom}_R(A, R) \neq \emptyset$.

4. Equations over complete local rings

Let R be a ring and let $X = (X_1, \dots, X_h; X_{h+1}, \dots, X_N)$ denote a set of indeterminates. The ring $R[[X_1, \dots, X_h]][X_{h+1}, \dots, X_N]$ will be denoted by $R[[X_1, \dots, X_h; X_{h+1}, \dots, X_N]]$ or by $R[[X]]$. We consider a complete local ring R and sets of elements $F = (F_1, \dots, F_s)$ in $R[[X]]$. A solution x

modulo $\mathfrak{m}^t$ of F is a set of elements $x = (x_1, \dots, x_N)$ with $x_1, \dots, x_h \in \mathfrak{m}$ and $x_{h+1}, \dots, x_N \in R$ such that $F_i(x_1, \dots, x_N) \in \mathfrak{m}^t$ for all i . We abbreviate this by $F(x) \equiv 0(\mathfrak{m}^t)$. The ideal in $R[[X]]$ generated by $\{F_1, \dots, F_s\}$ is also denoted by F . Solutions of F modulo $\mathfrak{m}^t$ are into one-one correspondence with $\text{Hom}_R(R[[X]]/F, R/\mathfrak{m}^t)$.

A local noetherian ring R is called a *strong s-ring* if for every F in $R[[X]]$ there exists a function $s: \mathbb{N} \rightarrow \mathbb{N}$, $s(n) \geq n$ for all n , such that:

If $F(x) \equiv 0(\mathfrak{m}^{s(n)})$ then there exists x' with $x' \equiv x(\mathfrak{m}^n)$ and $F(x') = 0$. We note that a strong s-ring is necessarily complete. In trying to prove the converse we have encountered some difficulties in the mixed characteristic case and we cannot show much more than:

(4.1) THEOREM: *Every noetherian complete local ring of equal characteristic is a strong s-ring.*

Our proof of (4.1) follows closely proofs of M. Greenberg [4] and M. Artin [2] where special cases of (4.1) are treated.

(4.2) PROPOSITION: (Descent). *Let R_0 and R be complete local noetherian rings and let $R_0 \rightarrow R$ be a finite map. If R_0 is a strong s-ring then so is R .*

PROOF: Let $e_1, \dots, e_a$ be a base of the R_0 -module R and let $r_1, \dots, r_b \in R_0^a$ be a base of the relations between $e_1, \dots, e_a$. Let $\mathfrak{m}_0$ denote the maximal ideal of R_0 and e an integer satisfying $\mathfrak{m}^e \subseteq \mathfrak{m}_0 R \subseteq \mathfrak{m}$. Let the set of equations $F = (F_1, \dots, F_s)$ in $R[[X_1, \dots, X_h; X_{h+1}, \dots, X_N]]$ be given.

We introduce new variables

$$\begin{aligned} \tilde{X}_{ij} (i = 1, \dots, h; j = 1, \dots, a); & \quad X_{ij} (i = 1, \dots, N; j = 1, \dots, a); \\ Y_{il} (i = 1, \dots, s; l = 1, \dots, b); & \quad Z_{il} (i = 1, \dots, h; l = 1, \dots, b). \end{aligned}$$

F_i can be written as $\tilde{F}_i(x_1^e, \dots, x_h^e; x_1, \dots, x_N)$ where $\tilde{F}_i$ is a formal power series in the first h variables and a polynomial in the last N variables. Substitute in $\tilde{F}_i: X_i^e = \sum_{j=1}^a \tilde{X}_{ij} e_j$; $X_i = \sum_{j=1}^a x_{ij} e_i$. Then $\tilde{F}_i$ becomes $\sum_{j=1}^a G_{ij}(\tilde{X}_{..}, X_{..}) e_j$ where $G_{ij} \in R_0[[\tilde{X}_{..}; X_{..}]]$. Further

$$\left(\sum_{j=1}^a X_{ij} e_j \right)^e = \sum_{j=1}^a H_{ij}(X_{..}) e_j$$

for some $H_{ij} \in R_0[[X_{..}]]$. We consider over R_0 the system of equations F^* in $R_0[[\tilde{X}_{..}; X_{..}, Y_{..}, Z_{..}]]$ given by $G_{ij}(\tilde{X}_{..}, X_{..}) + \sum_{l=1}^b Y_{il} r_{lj}$ and $H_{ij}(X_{..}) - \tilde{X}_{ij} + \sum_{l=1}^b Z_{il} r_{lj}$ where $r_l = (r_{l1}, \dots, r_{la}) \in R_0^a (l = 1, \dots, b)$.

By assumption the system F^* has a function s^* . Then $s = e \cdot s^*$ is an

s -function for F . Indeed let $F(x) \equiv 0(m^{es*(n)})$. Write $x_i = \sum_{j=1}^b x_{ij}e_j$ ($x_{ij} \in R_0$; $i = 1, \dots, N$) and $x_i^e = \sum_{j=1}^a \tilde{x}_{ij}e_j$ ($\tilde{x}_{ij} \in R_0$; $i = 1, \dots, h$).

Then $(\sum_j x_{ij}e_j)^e = \sum_j \tilde{x}_{ij}e_j$ and so for suitable $z_{il} \in R_0$ we have $H_{ij}(x_{..}) - \tilde{x}_{ij} + \sum_{l=1}^b z_{il}r_{lj} = 0$. Further $\sum_{j=1}^a G_{ij}(\tilde{x}_{..}, x_{..})e_j = \sum \tau_{ij}e_j$ with $\tau_{ij} \in m_0^{s*(n)}$ since $m^{es*(n)} \subseteq m_0^{s*(n)}R$. Hence for suitable $y_{il} \in R_0$ we have $G_{ij}(\tilde{x}_{..}, x_{..}) + \sum y_{il}r_{lj} \in m_0^{s*(n)}$. So we found a solution modulo $m_0^{s*(n)}$ of F^* namely $(\tilde{x}_{..}, x_{..}, y_{..}, z_{..})$. Let $(\underline{\tilde{x}}_{..}, \underline{x}_{..}, \underline{y}_{..}, \underline{z}_{..})$ be a solution of F^* which is equivalent modulo m_0^n with $(\tilde{x}_{..}, x_{..}, y_{..}, z_{..})$. Put $x_i = \sum \underline{x}_{ij}e_j$. Then $(\sum \underline{x}_{ij}e_j)^e = \sum \underline{\tilde{x}}_{ij}e_j$ and it follows that $\underline{x} \equiv x(m^n)$ and $F(\underline{x}) = 0$.

(4.3) LEMMA: *Let R be a regular complete local ring. If there exists an s -function for every prime ideal in $R[[X]]$ then there exists an s -function for every ideal in $R[[X]]$.*

PROOF: Let F be an ideal in $R[[X]]$. The radical of F is the intersection of prime ideals $p_1, \dots, p_t$ which have s -functions $s_1, \dots, s_t$. For some number d we have $F \supset p_1^d \dots p_t^d$. Define $s = dt \max \{s_1, \dots, s_t\}$. If $F(x) \equiv 0(m^{s(n)})$ then for some i , $p_i(x) \equiv 0(m^{s_i(n)})$. Hence there exists $x' \equiv x(m^n)$ with $p_i(x') = 0$ and in particular $F(x') = 0$.

REMARK: (4.2) and (4.3) reduce the general statement to proving the existence of an s -function for prime ideals F in $R[[X]]$ where R is a complete regular local ring and $F \cap R = 0$. In the rest of the proof of (4.1) we apply induction on $\dim R$ and on $\dim R[[X]]/F$. According to the next lemma we may further assume that the quotient field of $R[[X]]/F$ is separable over the quotient field of R .

(4.4) LEMMA: *Suppose that F is a prime ideal of $R[[X]]$, R a regular complete local ring with $F \cap R = 0$, such that the quotient field of $A = R[[X]]/F$ is inseparable (i.e. not separable) over that of R . Then there exists an ideal $G \not\supseteq F$ of $R[[X]]$ and a function $\tau: \mathbb{N} \rightarrow \mathbb{N}$ ($\tau(n) \geq n$ for all n) such that $F(x) \equiv 0(m^{\tau(n)})$ implies $G(x) \equiv 0(m^n)$.*

PROOF: Let $f_1, \dots, f_s \in A$ be linearly independent over R such that $f_1^p, \dots, f_s^p$ are dependent ($p = \text{char of } R > 0$). Hence $\alpha_1 f_1^p + \dots + \alpha_s f_s^p = 0$ for some $\alpha_1, \dots, \alpha_s \in R$ not all zero. Let $\{\alpha_1, \dots, \alpha_t\}$ be a maximal p -independent subset over R^p . After multiplying with β^p , $\beta \neq 0$, $\beta \in R$ we may suppose $\alpha_i \in R^p[\alpha_1, \dots, \alpha_t]$ for all $i > t$. The equation $\alpha_1 f_1^p + \dots + \alpha_s f_s^p = 0$ becomes $\sum_{0 \leq \beta_i < p} g_\beta^p \alpha_1^{\beta_1} \dots \alpha_t^{\beta_t} = 0$ and not all $g_\beta \in F$. (Otherwise the $f_1, \dots, f_s$ are linearly dependent over R). Put

$G = (F, g_\beta) \not\supseteq F$. The local ring $B = R^p[\alpha_1, \dots, \alpha_t]$ has the free base $\{\alpha_1^{\beta_1} \dots \alpha_t^{\beta_t} | 0 \leq \beta_i < p\}$ over R^p . Hence for some e we have

$$\mathfrak{m}(B)^e \subseteq \mathfrak{m}(R^p)B \subseteq \mathfrak{m}(B).$$

Further since B is complete there exists a function $\tau : \mathbb{N} \rightarrow \mathbb{N}$, $\tau(n) \geq n$ for all n , such that $\mathfrak{m}(R)^{\tau(n)} \cap B \subseteq \mathfrak{m}(R^p)^n B$. (See Nagata [5] Theorem (30.1) on page 103.)

If now $F(x) \equiv 0(\mathfrak{m}^{\tau(n)})$ then $\sum g^\beta(x) \alpha_1^{\beta_1} \dots \alpha_t^{\beta_t} \equiv 0(\mathfrak{m}(R)^{\tau(n)})$ and all $g_\beta(x) \equiv 0(\mathfrak{m}^n)$. Hence $G(x) \equiv 0(\mathfrak{m}^n)$.

(4.5) REMARK: It suffices to prove (4.1) in the following situation: R is a complete regular local ring, F is a prime ideal of $R[[X]]$ such that

- (1) The quotient field of $R[[X]]/F$ is separable over the quotient field of R
- (2) For all $n \geq 1$ there exists a solution of $F(x) \equiv 0(\mathfrak{m}^n)$.

If the second condition were not satisfied then F has clearly an s -function, namely $s(n) = n + \max \{k | \text{there exists } x \text{ with } F(x) \equiv 0(\mathfrak{m}^k)\}$.

Our next step in proving (4.1) will be to show that the conditions above imply that the Jacobian ideal of $(F_1, \dots, F_s)$ with respect to the variables $X_1, \dots, X_N$ is not contained in F . This will be done in Section 5.

5. Modules of differentials

Let R be a complete regular local ring and let $A = R[[X]]/F$ satisfy the condition (4.5). Let s denote the height of the ideal F . We want to show that the ideal generated by the $s \times s$ -minors of the Jacobian matrix

$$\begin{pmatrix} \frac{\partial F_1}{\partial X_1}, \dots, \frac{\partial F_m}{\partial X_1} \\ \vdots \\ \frac{\partial F_1}{\partial X_N}, \dots, \frac{\partial F_m}{\partial X_N} \end{pmatrix}$$

is not contained in F . We consider separately the cases $\text{char } R = p > 0$ and $\text{char } R = 0$.

(5.1) THEOREM: Suppose that $\text{char } R = p > 0$ and let $A = R[[X]]/F$ satisfy

- (1) F is a prime ideal and the quotient field L of A is separable over the quotient field K of R .
- (2) $\text{Hom}_R(A, R/\mathfrak{m}) \neq \emptyset$.

Then $\text{rank}_A \Omega_{A/R} = \dim A - \dim R$ and the ideal of the $s \times s$ -minors of $\partial F / \partial X$ is not contained in F .

PROOF: Let k be a coefficient field of R and consider the exact sequence

$$\Omega_{R/R^p[k]} \otimes A \xrightarrow{\alpha} \Omega_{A/R^p[k]} \rightarrow \Omega_{A/R} \rightarrow 0.$$

We note that $R^p[k]$ is a noetherian local in between $R^p = k^p[[T_1^p, \dots, T_d^p]]$ and $R = k[[T_1, \dots, T_d]]$. It's completion $R_1 = k[[T_1^p, \dots, T_d^p]]$. Hence $\Omega_{R/R_1} \otimes A$ is a free A -module of rank $= \dim R$. Likewise the other modules in the sequence are finitely generated. The map α is injective since $\alpha \otimes 1_L : \Omega_{K/l} \otimes L \rightarrow \Omega_{L/l}$ is injective ($l =$ the quotient field of R_1 and L/K is separable).

Hence $\text{rank } \Omega_{A/R} = \text{rank } \Omega_{A/R_1} - \dim R$ and we have to show that $\text{rank } \Omega_{A/R_1} = \dim A$.

Let $\rho : A \rightarrow k$ be an R -homomorphism (exists, since (2)) and let p be its kernel. Then $B = \hat{A}_p$ has the properties (see [3] EGA IV, Ch. 0, (7.8.2) and (7.8.3))

(a) B has no nilpotents.

(b) every minimal prime q of B satisfies $\dim B/q = \dim B (= \dim A_p = \dim A)$.

(c) the quotient field of B/q is separable over L (and hence over k). Further since $A \subset B$ have no zero divisors $\text{rank}_A \Omega_{A/R_1} = \text{rank}_B \Omega_{A/R_1} \otimes B$. It is easily seen that $\Omega_{B/k} = \Omega_{B/R_1} \simeq \Omega_{A/R_1} \otimes B$. Hence the statement $\text{rank}_A \Omega_{A/R} = \dim A - \dim R$ will follow from lemma (5.2).

The last statement of (5.1) follows directly from the exact sequence:

$$A^m \xrightarrow{\alpha} \Omega_{R[[X]]/R} \otimes A \rightarrow \Omega_{A/R} \rightarrow 0,$$

in which $\Omega_{R[[X]]/R} \otimes A$ is the free A -module on generators $dX_1, \dots, dX_N$ and α is the map given by

$$\alpha(a_1, \dots, a_m) = \sum_{i=1}^m a_i dF_i = \sum_{i,j} a_i \frac{\partial F_i}{\partial X_j} dX_j.$$

Indeed

$$\dim A - \dim R = \text{rank } \Omega_{A/R} = N - \text{rank} \left(\frac{\partial F}{\partial X} \right) \text{ modulo } F$$

and $\dim A = \dim R + N - \text{height } F$.

DEFINITION: Let $A \rightarrow B$ be a ringhomomorphism. By $\Omega_{B/A}^f$ we denote the *universal finite module of differentials* i.e.

- (i) $\Omega_{B/A}^f$ is a finite B -module and $d: B \rightarrow \Omega_{B/A}^f$ is an A -derivation.
- (ii) The natural map $\text{Hom}_B(\Omega_{B/A}^f, M) \rightarrow \text{Der}_A(B, M)$ is an isomorphism for all finitely generated B -modules M .

REMARK: (a) If B is of essentially finite type over A then $\Omega_{B/A}^f = \Omega_{B/A}$.

(b) If B is a complete local noetherian ring with coefficient ring or field A then $\Omega_{B/A}^f$ exists.

(c) If the noetherian local ring has a coefficient field k of characteristic $p \neq 0$ then $\Omega_{B/k}^f = \Omega_{B/k}$.

(d) If $A = k[[X]]$ where k is a field of characteristic 0, then $\Omega_{A/k}^f \neq \Omega_{A/k}$.

(e) If $A = k[[X]][[Y]]$ then $\Omega_{A/k}^f$ does not exist.

(5.2) LEMMA: Let B be a complete local ring such that

- (i) $A \subset B$ is a coefficient ring (or field) consisting of non-zero divisors.
- (ii) B has no nilpotents and for every minimal prime q of B , $\dim B = \dim B/q$.
- (iii) For every minimal prime q of B , the quotient field of B/q is separable over that of A .

Then $\text{rank}_B \Omega_{B/A}^f = \dim B - \dim A$.

PROOF: (a) $\dim A = 1$ (i.e. A is a discrete valuation ring with maximal ideal pA). The ring B has the form $A[[X_1, \dots, X_N]]/F$. Since p is a non-zero divisor on B we find that $F \notin pA[[X_1, \dots, X_N]]$. Take an element $f \in F$ with non-zero image $\bar{f}$ in $k[[X_1, \dots, X_N]]$ where $k = A/pA$. After a change of coordinates, $\bar{f}$ is general in X_N of say order d . The Weierstrasz theorem for $k[[X_1, \dots, X_N]]$ implies that for every $g \in A[[X_1, \dots, X_N]]$ one has $g = q_0 f + r_0 + pg_1$ where $r_0 \in A[[X_1, \dots, X_{N-1}]] [X_N]$ has $\text{degree}_{X_N}(r_0) < d$. By induction we find $g_1 = q_1 f + r_1 + pg_2, \dots, g_n = q_n f + r_n + pg_{n+1}, \dots$. Hence $g = (q_0 + pq_1 + \dots)f + (r_0 + pr_1 + \dots)$. So we proved that for any $g \in A[[X_1, \dots, X_N]]$ we can write $g = qf + r$ where $r \in A[[X_1, \dots, X_{N-1}]] [X_N]$ has $\text{degree}_{X_N}(r) < d$. In particular $f = (\text{unit})(X_N^d + a_{d-1}X_N^{d-1} + \dots + a_0)$ with all $a_i \in A[[X_1, \dots, X_{N-1}]]$. So $A[[X_1, \dots, X_N]]/F$ is a finite extension of $A[[X_1, \dots, X_{N-1}]]/G$ where $G = F \cap A[[X_1, \dots, X_{N-1}]]$. Repeating this process we find that B is finite over $A[[X_1, \dots, X_l]]$. Since all the minimal primes q of B satisfy $\dim B/q = \dim B$ we have $q \cap A[[X_1, \dots, X_l]] = 0$. The total quotientring $Q_t(B) = K_1 \times \dots \times K_t$ of B is a product of fields $K_i = B/q_i$ where $q_1, \dots, q_t$ are the minimal primes of B . Each K_i contains the quotient field K of $A[[X_1, \dots, X_l]]$.

The natural map $\alpha: \Omega_{A[[X_1, \dots, X_l]]/A}^f \otimes B \rightarrow \Omega_{B/A}^f$ has the property that $\alpha \otimes 1_{Q_t(B)}: \Omega_{A[[X_1, \dots, X_l]]/A}^f \otimes Q_t(B) \rightarrow \Omega_{B/A}^f \otimes Q_t(B)$ is an isomorphism.

Indeed for any Λ -derivation $D: A[[X_1, \dots, X_l]] \rightarrow M$, M a finitely generated B -module, we have a unique extension $D_i: K_i \rightarrow M \otimes_B K_i$ since K_i is a finite separable extension of K . So we have a unique extension

$$D_1 \times \dots \times D_l: Qt(B) \rightarrow M \otimes Qt(B) = (M \otimes K_1) \oplus \dots \oplus (M \otimes K_l).$$

Since $\Omega_{A[[X_1, \dots, X_k]]/A}^f$ is a free module of rank $= \dim B - \dim A$ also $\text{rank } \Omega_{B/A}^f = \dim B - \dim A$.

(b) $\Lambda = k$ is a field of characteristic zero. Same proof as in case (a)

(c) $\Lambda = k$ is a field of characteristic $p \neq 0$. A refined version of the Weierstrasz-theorems yields that B is a finite extension of $k[[X_1, \dots, X_d]]$ such that $q \cap k[[X, \dots, X_d]] = 0$ for all minimal primes and such that the quotient field of B/q is separable over $k((X_1, \dots, X_d))$ for all minimal primes q of B . After this we can finish the proof as in case (a).

The characteristic zero case of (5.1) is more complicated. Let $A = R[[X]]/F$ satisfy (4.5) and let A_0 be the image in A of $R[[X_1, \dots, X_h]]$, hence $A_0 = R[[X_1, \dots, X_h]]/G$ with $G = F \cap R[[X_1, \dots, X_h]]$. Further $A = A_0[[X_{h+1}, \dots, X_N]]/H$ where $H = F/G$. Complete local rings satisfy the universal chain condition, so $\text{height } F = \text{height } H + \text{height } G$.

Let K_0 be the quotient field of A_0 then $A \otimes_{A_0} K_0 = K_0[[X_{h+1}, \dots, X_N]]/L$ where L is the ideal generated by the image of F .

The usual 'Jacobian criterium for simple points' yields some $\text{height } H \times \text{height } H$ -minor δ of the matrix

$$\frac{\partial F}{\partial X_{h+1}, \dots, \partial X_N}$$

is not contained in L (and hence not in F).

If we can find a $\text{height } G \times \text{height } G$ -minor of

$$\frac{\partial G}{\partial X_1, \dots, \partial X_h}$$

which is not contained in G then we can combine this with δ to produce a $\text{height } F \times \text{height } F$ -minor of

$$\frac{\partial F}{\partial X_1 \dots \partial X_n}$$

which is not contained in F . Hence we showed that it suffices to prove:

(5.3) THEOREM: If $A = R[[X_1, \dots, X_N]]/F$ satisfies (4.5) then some height $F \times \text{height } F$ -minor of

$$\frac{\partial F}{\partial X}$$

is not contained in F .

PROOF: Suppose that there exists a $\rho \in \text{Hom}_R(A, R)$; after changing the coordinates we may suppose that $\rho(X_i) = 0$ for all i . So $F \subset (X_1, \dots, X_N) = p$. The ring $B = \hat{A}_p$ has the properties: (i) B has no nilpotents and (ii) For every minimal prime q of B , $\dim B/q = \dim B = \dim A_p = \text{height } p/F = N - \text{height } F$. Further clearly

$$B = K[[X_1, \dots, X_N]]/FK[[X_1, \dots, X_N]]$$

where $K = Q_t(B)$. From (5.2) it follows that

$$\frac{\partial F}{\partial X}$$

has an height $F \times \text{height } F$ -minor which is not contained in $FK[[X_1, \dots, X_N]]$ (and hence not contained in F).

(5.4) PROPOSITION: Let A be a coefficient ring or field (according to $\text{char } R/\mathfrak{m} > 0$ or $= 0$) of R . The assumptions (4.5) and $\text{Hom}_R(A, R) \neq \emptyset$ for $A = R[[X]]/F$ imply that the sequence $0 \rightarrow \Omega_{R/A}^f \otimes A \xrightarrow{\alpha} \Omega_{A/A}^f \rightarrow \Omega_{A/R}^f \rightarrow 0$ is exact.

PROOF: The only thing to show is the injectivity of α . Now $\Omega_{A/R}^f$ is equal to the free A -module on generators $dX_1, \dots, dX_N$ divided by the submodule AdF . Since some height $F \times \text{height } F$ -minor is not contained in F we have $\text{rank}_A \Omega_{A/R}^f \leq N - \text{height } F = \dim A - \dim R$. By (5.2) $\text{rank}_A \Omega_{A/A}^f = \dim A - \dim A$ and $\Omega_{R/A}^f$ is a free-module of rank $\dim R - \dim A$. Let K denote the quotient field of A then for dimension reasons

$$0 \rightarrow \Omega_{R/A}^f \otimes K \rightarrow \Omega_{A/A}^f \otimes K \rightarrow \Omega_{A/R}^f \otimes K \rightarrow 0$$

is exact. Since $\Omega_{R/A}^f \otimes A$ is a free A -module, also α must be injective.

(5.5) LEMMA: Let R be a complete local ring with a residue field k

which is algebraically closed and uncountable. Let $A = R[[X]]/F$ satisfy $\text{Hom}_R(A, R/\mathfrak{m}^n) \neq \emptyset$ for all n . Then $\text{Hom}_R(A, R) \neq \emptyset$.

PROOF: Fix a coefficient field of R or in the unequal characteristic case a map $W(k) \rightarrow R$ where $W(k)$ denotes the ring of Witt-vectors over k . Then each $R/\mathfrak{m}^n$ has the structure of a finite-dimensional vector space over k in which addition and multiplication are morphisms. Then $\text{Hom}_R(A, R/\mathfrak{m}^n)$ is an algebraic subset of $(R/\mathfrak{m}^n)^N$ (we identify a map ρ with $(\rho(X_1), \dots, \rho(X_N)) \in (R/\mathfrak{m}^n)^N$).

The intersection of a descending sequence of non-empty constructible sets is non-empty (see F. Oort [6], Lemma 2 on page 221). Hence

$$\bigcap_{m \geq n} \text{im}(\text{Hom}_R(A, R/\mathfrak{m}^m) \rightarrow \text{Hom}_R(A, R/\mathfrak{m}^n)) \neq \emptyset$$

and with the usual compactness-argument it follows that

$$\text{Hom}_R(A, R) = \varprojlim \text{Hom}_R(A, R/\mathfrak{m}^n) \neq \emptyset.$$

Continuation of the proof of (5.3). Let A be a coefficient ring (or field) of R and denote by A' a flat extension such that (i) $\mathfrak{m}(A)A' = \mathfrak{m}(A')$; (ii) $A'/\mathfrak{m}(A')$ is algebraically closed and uncountable. We use the following notations $R' = R \hat{\otimes} A'$ or if $R = A[[T_1, \dots, T_d]]$ then $R' = A'[[T_1, \dots, T_d]]$ and let $A' = R'[[X]]/FR'[[X]]$.

Consider the exact sequence

$$\Omega_{R/A}^f \otimes_A A \xrightarrow{\alpha} \Omega_{A/A}^f \rightarrow \Omega_{A/R}^f \rightarrow 0.$$

As shown before $\Omega_{R/A}^f \otimes_A A$ is a free module of rank $= \dim R - \dim A$ and $\text{rank}_A \Omega_{A/A}^f = \dim A - \dim A$. If we can show that α is injective then it follows that $\text{rank} \Omega_{A/R}^f = \dim A - \dim R$. The module $\Omega_{A/R}^f$ is equal to the free A -module on generators $dX_1, \dots, dX_N$ modulo the submodule generated by dF . As in the proof of (5.1) one concludes that the rank of the matrix

$$\frac{\partial F}{\partial X}$$

modulo F is equal to height F .

So we want to show that $\Omega_{R/A}^f \otimes_A A \xrightarrow{\alpha} \Omega_{A/A}^f$ is injective. Consider $S = R[[X]] \setminus F$. Every $s \in S$ is a non-zero divisor on $A = R[[X]]/F$ and since A'/A is flat, S consists of non-zero divisors on A' . In $R[[X]]_S$ the ideal F is the regular maximal ideal, hence generated by a regular sequence

$F_1, \dots, F_s$ ($s = \text{height } F$). By flatness $\{F_1, \dots, F_s\}$ is a regular sequence on $R'[[X]]_S$ and all the associated ideals of $(F_1, \dots, F_s)$ in $R'[[X]]_S$ have height s . Take a minimal prime q of $FR'[[X]]$ such that

$$\text{Hom}_{R'}(R'[[X]]/q, R') \neq \emptyset$$

((5.5) guarantees the existence of q). Put $A_1 = R'[[X]]/q$.

Then we have a commutative diagram

$$\begin{array}{ccccccc} \Omega_{R/A}^f \otimes A & \xrightarrow{\alpha} & \Omega_{A/A}^f & & & & \\ \downarrow \gamma & & \downarrow & & & & \\ 0 \longrightarrow & \Omega_{R'/A'}^f \otimes A' & \xrightarrow{\beta} & \Omega_{A'/A'}^f & \longrightarrow & \Omega_{A'/R'}^f & \longrightarrow 0 \end{array}$$

in which the row is exact according to (5.4). Clearly also γ is injective. Hence α is injective and we are done.

6. Inductionsteps

In this section we finally give a proof of (4.1). Let $F \subset R[[X]]$ satisfy (4.5). Let A be the ideal generated by the $s \times s$ -minors of

$$\frac{\partial F}{\partial X}$$

(where $s = \text{height } F$). According to Section 5, $F \not\subseteq (F, A)$. By induction on $\dim R[[X]]/F$ there exists an s -function for (F, A) . Hence it suffices to show (6.1) in the equal characteristic case. In the unequal characteristic case we also have to consider elements x with $F(x) \equiv 0(\mathfrak{m}^{\succ})$, $A(x) \not\equiv 0(\mathfrak{m}^b)$ and $A(x) \equiv 0(p, \mathfrak{m}^{\succ})$.

(6.1) **PROPOSITION:** *Suppose that F satisfies (4.5). Let p denote the characteristic of $R/\mathfrak{m}$ considered as an element of R .*

For all n and b there exists an $a \in \mathbb{N}$ such that $F(x) \equiv 0(\mathfrak{m}^a)$ and $A(x) \not\equiv 0(p, \mathfrak{m}^b)$ imply the existence of $x' \equiv x(\mathfrak{m}^n)$ with $F(x') = 0$.

The proof of (6.1) requires a string of lemmata.

(6.2) **LEMMA:** *Let R be a complete regular local ring (unramified in the unequal characteristic case) with infinite residue field. There is a finite set of subrings $R_1, \dots, R_s$ of R and $T \in R$ such that*

- (i) each R_i is regular and $R_i[[T]] = R$
(ii) for any $g \in R$, $g \not\equiv 0(p, \mathfrak{m}^b)$ there exists an i such that

$$g = (\text{unit})(T^d + a_{d-1}T^{d-1} + \dots + a_0)$$

with $d < b$ and $a_1, \dots, a_{d-1} \in R_i$.

PROOF: The image $\bar{g}$ of g in R/pR has order c , $c < b$; let h be its homogeneous part of order c (with respect to a presentation $A[[X_1, \dots, X_n]]$ of R). Let A_0 be a finite subset of A such that the set of residues in A/pA is of cardinal $> b$. There are $\lambda_1, \dots, \lambda_n \in A_0$ such that $\lambda_n \not\equiv 0(p)$ and $h(\bar{\lambda}_1, \dots, \bar{\lambda}_n) \neq 0$. Put $Y_i = X_i - \lambda_i \lambda_n^{-1} X_n$ for $i = 1, \dots, n-1$ and $Y_n = X_n$. Then $h(X_1, \dots, X_n) = k(Y_1, \dots, Y_n)$ for some homogeneous polynomial k . Then $k(0, \dots, 0, Y_n) = \bar{\lambda}_n^{-c} Y_n^c h(\bar{\lambda}_1, \dots, \bar{\lambda}_n) \neq 0$. Hence $\bar{g}$ is general in $T = Y_n = X_n$. By the Weierstrasz-preparation theorem

$$g = \text{unit} (T^c + a_{c-1} T^{c-1} + \dots + a_0)$$

with $a_i \in R' = A[[Y_1, \dots, Y_{n-1}]]$.

(6.3) PROPOSITION: (Induction on $\dim R$). Let $F = (F_1, \dots, F_m) \in R[[X]]$ and $G \in R[[X]]$. For all n and b there exists $a \in \mathbb{N}$ such that

$$\left. \begin{array}{l} F(x) \equiv 0(\mathfrak{m}^a) \\ G(x) \not\equiv 0(p, \mathfrak{m}^b) \end{array} \right\} \text{ imply the existence of } x' \equiv x(\mathfrak{m}^n)$$

with $F(x') \equiv 0G(x')$.

PROOF: If x satisfies $G(x) \not\equiv 0(p, \mathfrak{m}^b)$ then according to (6.2) there is a presentation $R = R'[[T]]$ and an integer $d < b$ such that

$$G(x) = a \text{ unit times } (T^d + a_{d-1} T^{d-1} + \dots + a_0)$$

with all $a_i \in \mathfrak{m}(R')$. Since we have a finite choice for R' and d we can restrict ourselves to a fixed choice for R' and d .

Introduce new variables $A_0, \dots, A_{d-1}$; $Y_1, \dots, Y_N$; Y_{ij} ($i = 1, \dots, N$; $j = 0, \dots, d-1$); Z_i ($i = 1, \dots, h$); Z_{ij} ($i = 1, \dots, h$; $j = 0, \dots, d-1$). Then $C = R'[[A_0, \dots, A_{d-1}]] \hookrightarrow R[[A_0, \dots, A_{d-1}]]/(T^d + A_{d-1}T^{d-1} + \dots + A_0) = D$ is a finite extension and there is a number e with $\mathfrak{m}(D)^e \subseteq \mathfrak{m}(C)D$.

Make the substitutions:

$$X_i = Y_i(T^d + A_{d-1}T^{d-1} + \dots + A_0) + \sum_{j=0}^{d-1} Y_{ij}T^j$$

$$X_i^e = Z_i(T^d + A_{d-1}T^{d-1} + \dots + A_0) + \sum_{j=0}^{d-1} Z_{ij}T^j$$

and consider Weierstrasz-division by $W = T^d + A_{d-1}T^{d-1} + \dots + A_0$. Then

$$G = Q(Z., Z., A., Y., Y.)W + \sum_{j=0}^{d-1} G_j(Z., A., Y.)T^j$$

$$F_i = Q_j(Z., Z., A., Y., Y.)W + \sum_{j=0}^{d-1} F_{ij}(Z., A., Y.)T^j$$

where G_j, F_{ij} belong to $R'[[Z., A.]][Y.]$. Consider also the equations:

$$(Y_i W + Y_{ij} T^j)^e - (Z_i W + Z_{ij} T^j)$$

which amounts to the equations

$$Z_{ij} - H_{ij}(Y.) \in R'[[Z., A.]][Y.].$$

The system of equations $F^* = \{G_j, F_{ij}, H_{ij} - Z_{ij}\}$ over R' has an almost solution with $A_i = a_i$ as given above. Further by Weierstrasz-division

$$x_i = y_i(T^d + a_{d-1}T^{d-1} + \dots + a_0) + \sum y_{ij}T^j, \quad \text{all } y_{ij} \in R'$$

$$x_i^e = z_i(T^d + a_{d-1}T^{d-1} + \dots + a_0) + \sum z_{ij}T^j, \quad \text{all } z_{ij} \in \mathfrak{m}(R').$$

$$\text{These elements satisfy } \begin{cases} z_{ij} - H_{ij}(y.) = 0 \\ G_j(z., a., y.) = 0 \\ F_{ij}(z., a., y.) \equiv 0(\mathfrak{m}_0^{a-d}) \end{cases}$$

where $\mathfrak{m}_0$ is the maximal ideal of R' .

Since F^* has an s -function, we find for sufficiently high $a \in \mathbb{N}$ a solution $(z'., a'., y'.) \equiv (z., a., y.)(\mathfrak{m}_0^n)$ of F . Define

$$x'_i = y'_i(T^d + a'_{d-1}T^{d-1} + \dots + a'_0) + \sum_{j=0}^{d-1} y'_{ij}T^j.$$

Then $x \equiv x'(\mathfrak{m}^n)$ and

$$F_i(x') \equiv 0(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$$

for all i and

$$G(x') = \text{unit } (T^d + a'_{d-1} T^{d-1} + \dots + a'_0).$$

Hence $F(x') \equiv 0(G(x'))$.

(6.4) LEMMA: Let $F_1, \dots, F_s \in R[[X]]$ and let δ be an $s \times s$ -minor of

$$\frac{\partial F}{\partial X}$$

and let $a \neq 0$ be an element of R and x such that $F(x) \equiv 0(a\delta(x)^2)$. Then there exists $x' \equiv x(a\delta(x))$ with $F(x') = 0$.

PROOF: We may suppose $x = 0$ and we may replace $R[[X]]$ by $R[[X]]$. Then we are reduced to a well known case of this lemma. See [1] lemma (5.10) and (5.11).

Conclusion of the proof of (4.1)

Let (i) resp. (j) denote subsets of s elements from $\{1, \dots, m\}$ resp. $\{1, \dots, N\}$ and let $\Delta_{(i),(j)}$ denote the corresponding $s \times s$ -minor $(\partial F / \partial X)$.

For any (i) let $F_{(i)}$ denote the ideal generated by $\{F_\alpha | \alpha \in (i)\}$. The radical $\sqrt{F_{(i)}}$ of $F_{(i)}$ equals $p_{(i),1} \cap \dots \cap p_{(i),t_i}$ = the intersection of prime ideals. Let $G_{(i)} = \bigcap \{p_{(i),a} | p_{(i),a} \not\subseteq F\}$. By induction $(F, G_{(i)})$ has an s -function $s_{(i)}$ and (F, Δ) has an s -function s .

Let $F(x) \equiv 0(\mathfrak{m}^\tau)$ with τ sufficiently high, then:

- (a) If $\Delta(x) \equiv 0(\mathfrak{m}^{s_0(n)})$ then there exists $x' \not\equiv x(\mathfrak{m}^n)$ with $F(x') = \Delta(x') = 0$.
- (b) If $\Delta(x) \not\equiv 0(\mathfrak{m}^{s_0(n)})$ then for some (i) and (j) we have

$$\Delta_{(i)(j)}(x) \not\equiv 0(\mathfrak{m}^{s_0(n)}).$$

Choose $u \in R$, $u \neq 0$ of order τ' , then by (6.3) there $x' \equiv x(\mathfrak{m}^{\tau'})$ with $F(x') \equiv 0(u\Delta_{(i)(j)}(x')^2)$. By lemma (6.4) there exists $x'' \equiv x'(\mathfrak{m}^{\tau'})$ with $F_{(i)}(x'') = 0$ and $F(x'') \equiv 0(\mathfrak{m}^{\tau'})$ and $\Delta_{(i)(j)}(x'') \equiv 0(\mathfrak{m}^{s_0(n)})$ where τ' is sufficiently high.

(c) For some minimal prime $p_{(i),a}$ of $F_{(i)}$ we have $p_{(i),a}(x'') = 0$. Since $\Delta_{(i)(j)}(x'') \neq 0$ it follows that height $p_{(i),a}(x'') = s^{(i),a}$. If $p_{(i),a} = F$ we are finished.

If $p_{(i),a} \neq F$ then $p_{(i),a} \not\subseteq F$ and $G_{(i)}(x'') = 0$. So we find

$$(F(x''), G(x'')) \equiv 0(\mathfrak{m}^{\tau'}).$$

From the existence of $s_{(i)}$ we conclude that there is an element $x''' \equiv x''(\mathfrak{m}^n)$ such that $F(x''') = G(x''') = 0$.

This concludes the proof of (4.1).

7. The mixed characteristic case

In this section we give the results that we could obtain in the mixed characteristic case.

(a) If the residue field k of R is finite then R is clearly a strong s -ring since every $\text{Hom}_R(R[[X]]/F, R/\mathfrak{m}^n)$ is a finite set.

(b) If $\dim R = 1$ (i.e. R is a discrete valuation ring of mixed characteristic) then R is a strong s -ring. In this case we don't need (6.3) and the hypothesis of (6.4) is automatically satisfied.

(c) For general R we would have proved that R is a strong s -ring if we could prove a more general version of (6.3), for instance: 'For all b and n there exists $a \in \mathbb{N}$ such that $F(x) \equiv 0(\mathfrak{m}^a)$ and $G(x) \not\equiv 0(\mathfrak{m}^b)$ imply the existence of $x' \equiv x(\mathfrak{m}^n)$ with $F(x') \equiv 0(G(x'))'$ '.

If $\dim R = 2$ and $[k : k^p] < \infty$ we will prove this more general version. But first another result.

(7.1) PROPOSITION: *Let R be a complete local ring with residue characteristic $p \neq 0$. Suppose that $k = R/\mathfrak{m}$ is finite over k^p and that for some l , $p^{l+1}R = 0$. Then R is a strong s -ring.*

(7.2) COROLLARY: *Let R be a complete local ring of residue characteristic $p \neq 0$. Let $k = R/\mathfrak{m}$ be finite over k^p . Given $F \in R[[X]]$ there exists a function $\tau : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that for all x with $F(x) \equiv 0(\mathfrak{m}^{\tau(a,b)})$ there exists $x' \equiv x(\mathfrak{m}^b)$ and $F(x') \equiv 0(p^a R)$.*

PROOF: Replace R by $R/p^a R$ and apply (7.1).

PROOF OF (7.1): (a) Suppose that we have shown the existence of a local ring homomorphism $R_0 = W_{l+1}(k_0[[T_1, \dots, T_d]]) \rightarrow R$ where k_0 is a subfield of k which makes R into a finite R -module. With (4.2) it suffices to show that R_0 is a strong s -ring. Let $F \in R_0[[X]]$ be given. Replace each variable X_i by a Witt-vector $(Y_{i,0}, \dots, Y_{i,l})$. Then the system F is equivalent to a set of equations over $k_0[[T_1, \dots, T_d]]$. From (4.1) the assertion (7.1) would follow.

(b) The structure theorem for complete local rings yields the existence of a finite map $R_1 = V/p^{l+1}V[[T_1, \dots, T_d]] \rightarrow R$, where V is a complete discrete valuation ring with $V/pV = k$. Let K be a perfect field containing

k , then $V/p^{l+1}V \hookrightarrow W_{l+1}(K)$ and $R_1 \hookrightarrow W_{l+1}(K[[S_1, \dots, S_d]])$ where $T_i \mapsto (S_i, 0, \dots, 0)$ ($i = 1, \dots, d$).

The image of R_1 contains $W_{l+1}(k[[S_1, \dots, S_d]]p^l)$, since for any $f \in k[[S_1, \dots, S_d]]$ there exists $f^* = (f, f_1, \dots, f_l) \in R_1$ and hence

$$(f^*)^{p^l} = (f^{p^l}, 0, \dots, 0)$$

belongs to R_1 . Further

$$p(f^*)^{p^{l-1}} = (0, f^{p^l}, 0, \dots, 0), \dots, p^l f^* = (0, \dots, 0, f^{p^l})$$

all belong to R .

So we found a finite map $W_{l+1}(k^{p^l}[[S_1^{p^l}, \dots, S_d^{p^l}]] \rightarrow R_1 \rightarrow R$ and the proof is completed.

(7.3) THEOREM: *Let R be a complete regular local ring of mixed characteristic. Suppose that $k = R/\mathfrak{m}$ is finite over k^p . If $\dim R = 2$ then R is a strong s -ring.*

PROOF: As remarked above we have to show that for

$$G, F = (F_1, \dots, F_m) \in R[[X]]$$

and all b and n there exists $a \in \mathbb{N}$ such that $F(x) \equiv 0(\mathfrak{m}^a)$, $G(x) \not\equiv 0(\mathfrak{m}^b)$ implies that there exists $x' \equiv x(\mathfrak{m}^n)$ with $F(x') \equiv 0(G(x'))$.

(1) If $G(x)$ has order c ($c < b$) and $G(x) \equiv 0(p^c)$ then $G(x) = \text{unit } p^c$ and we can apply (7.2).

(2) If $G(x) \equiv 0(p^c, \mathfrak{m}^{\geq})$ then applying (7.2) we are reduced to case (1), etc.

We see that we have only to do the case $G(x) = p^a$ with $a \in R$ satisfying $a \not\equiv 0(p, \mathfrak{m}^d)$ where d is some fixed number. Using (6.2) it is enough to consider the case

$$G(x) = \text{unit} \cdot p^a (T^d + a_{d-1} T^{d-1} + \dots + a_0)$$

where $R = V[[T]]$, V a valuation-ring, and all $a_i \in V$ and moreover all $a_i \in \mathfrak{m}(V)$.

Let I be the ideal generated by p^a and $T^d + a'_{d-1} T^{d-1} + \dots + a'_0$ with all $a'_i \in \mathfrak{m}(V)$. Then $\mathfrak{m}(R)^{2d\alpha} \subseteq I$. Further there is a number $\varepsilon \geq 1$, independent of the choice of $a'_0, \dots, a'_{d-1} \in \mathfrak{m}(V)$, such that

$$ap^a + b(T^d + a'_{d-1} T^{d-1} + \dots + a'_0) \equiv 0(\mathfrak{m}(R)^{\varepsilon n})$$

implies $ap^x \equiv 0(\mathfrak{m}^n)$.

Choose n such that $n > 2d\alpha$. Now we proceed as in the proof of (6.3). Choose new variables $A_0, \dots, A_{d-1}; Y_i; Y_{ij}; Z_i; Z_{ij}$ and substitute

$$\begin{aligned} X_i &= Y_i(T^d + A_{d-1}T^{d-1} + \dots + A_0) + \sum Y_{ij}T^j \\ X_i^e &= Z_i(T^d + A_{d-1}T^{d-1} + \dots + A_0) + \sum Z_{ij}T^j \end{aligned}$$

Then

$$\begin{aligned} G &= Q(Z_., Z_., A_., Y_., Y_..)(T^d + A_{d-1}T^{d-1} + \dots + A_0) \\ &\quad + \sum G_j(Z_., A_., Y_..)T^j \\ F &= Q_f(Z_., Z_., A_., Y_., Y_..)(T^d + A_{d-1}T^{d-1} + \dots + A_0) \\ &\quad + \sum F_{ij}(Z_., A_., Y_..)T^j Z_{ij} - H_{ij}(Y_..) \end{aligned}$$

We find a system of equations F^* over V namely $\{Z_{ij} - H_{ij}, G_j, F_{ij}\}$ and we are given an almost solution of F^* .

So there is (for $a \gg 0$) an $x' \equiv x(\mathfrak{m}^{en})$ with $F(x')$, $G(x') \equiv 0$ modulo $(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$.

According to (7.2) there is also an $x'' \equiv x(\mathfrak{m}^{en})$ with $F(x')$, $G(x'') \equiv 0(p^x)$. Hence $x'' - x' \equiv 0(\mathfrak{m}^{en})$. Since $n > 2d\alpha$ we find a and b with

$$x'' - x' = ap^x + b(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$$

and $ap^x \equiv 0(\mathfrak{m}^n)$.

Put $z = x'' - ap^x = x' + b(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$ then $z \equiv x(\mathfrak{m}^n)$ and $F(z)$, $G(z)$ are divisible by p^x and $(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$. So $F(z)$ and $G(z)$ are divisible by $p^x(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$. Since order $G(z) =$ order $G(x)$ we must have $G(z) = \text{unit } p^x(T^d + a'_{d-1}T^{d-1} + \dots + a'_0)$. It follows that $F(z) \equiv 0(G(z))$. End of the proof.

REMARKS: (1) F. Oort's theorem 1: 'Every complete local domain R is an f -ring' will follow from the statement: ' R is an s -ring'.

PROOF: Consider the polynomial $F = XY \in R[X, Y]$; by assumption it has an s -function. Define $f(i, j) = s(\max(i, j))$ for all $i, j \in \mathbb{N}$. Then $x \in R \setminus \mathfrak{m}^i$ and $y \in R \setminus \mathfrak{m}^j$ implies $xy \in \mathfrak{m}^{f(i, j)}$. Indeed, $F(x, y) \equiv 0(\mathfrak{m}^{s(\max(i, j))})$ implies the existence of $(x', y') \equiv (x, y)(\mathfrak{m}^{\max(i, j)})$ and $x'y' = 0$. Since R has no zero divisors $x' = 0$ or $y' = 0$ and one finds a contradiction.

(2) Using Oort's theorem 1 one can conversely prove that an s -function exists in some cases e.g.: If R is a complete local domain with quotient field K . Then an s -function exists for every ideal $F \subset R[X_1, \dots, X_n]$ such that $K[X_1, \dots, X_n]/FK[X_1, \dots, X_n]$ has Krull-dimension zero.

PROOF: As in (4.3), using the f -function of R one reduces to the case where F is a prime ideal and F has a zero in every $R/\mathfrak{m}^s$. Let $P_i = P_i(X_i)$ be a minimal polynomial for $X_i \bmod FK[X_1, \dots, X_n]$ over K . The polynomials P_i are irreducible over K and are normed such that all coefficients belong to R .

Since P_i has a zero in every $R/\mathfrak{m}^s$, it has a zero in R according to [6] Theorem 2. Hence $F = (x_1 - a_1, \dots, x_n - a_n)$ for suitable $a_1, \dots, a_n \in R$. Clearly an s -function exists for F .

(3) It might be possible to extend the reasoning of (2) to more general cases.

8. Analytic local rings

In this section we want to show that analytic local rings R over a complete valued field k (with $[k : k^p] < \infty$ if $\text{char } k = p \neq 0$) are s -rings. Let $F \subset R[X_1, \dots, X_n]$ be some ideal. According to (4.1) it suffices to show that every formal solution of F can be approximated by solutions in R . This is again a theorem of M. Artin [1] theorem (1.2) in the case $\text{char } k = 0$. The only instance in Artin's proof where $\text{char } k = 0$ is used is lemma (2.2) [1] page 283. It suffices to show the following:

(8.1) PROPOSITION: *Let k be a (pseudo-)complete valued field of $\text{char } p \neq 0$ with $[k : k^p] < \infty$, let $X = (X_1, \dots, X_n)$, $Y = (Y_1, \dots, Y_n)$; $k\{X, Y\}$ the ring of convergent power series over k and $F \subset k\{X, Y\}$ a prime ideal such that (i) $F \cap k\{X\} = 0$ and (ii) F has a solution in $k[[X]]$.*

Then the ideal Δ in $k\{X, Y\}$ generated by the height $F \times \text{height } F$ -minors of

$$\frac{\partial F}{\partial Y_1, \dots, \partial Y_n}$$

is not contained in F .

PROOF: (Analogous to (5.1)). We are given $k\{X\} \hookrightarrow k\{X, Y\}/F = A \hookrightarrow k[[X]]$. Hence the quotient field L of A is separable over the quotient field K of $k\{X\}$. So $\Omega_{K/k} \otimes L \xrightarrow{\beta} \Omega_{L/k}$ is injective. Consider the exact sequence

$$\Omega_{k\{X\}/k} \otimes A \xrightarrow{\alpha} \Omega_{A/k} \rightarrow \Omega_{A/k\{X\}} \rightarrow 0$$

with $\beta = \alpha \otimes_A 1$. Since $\Omega_{k\{X\}/k} \otimes A$ is a free A -module this implies that

α is injective and hence $\text{rank } \Omega_{A/k\{X\}} = \text{rank } \Omega_{A/k} - n$.

Weierstrasz-preparation theorem yields $k\{T_1, \dots, T_a\} \hookrightarrow A$ such that A is finite and separable over $k\{T_1, \dots, T_a\}$ and $a = \dim A$. The map $\gamma: \Omega_{k\{T_1, \dots, T_a\}} \otimes A \rightarrow \Omega_{A/k}$ has the property $\gamma \otimes 1_L$ is bijective. So $\text{rank } \Omega_{A/k} = a = \dim A$.

Further we have an exact sequence:

$$A^m \xrightarrow{\delta} \Omega_{k\{X, Y\}/k\{X\}} \otimes A \rightarrow \Omega_{A/k\{X\}} \rightarrow 0$$

where δ is given by

$$\delta(a_1, \dots, a_m) = \sum_i a_i dF_i = \sum_{i,j} a_i \frac{\partial F_i}{\partial Y_j} dY_j;$$

and $F = (F_1, \dots, F_m)$. The middle term is a free module of rank N , and the term on the right has rank $a - n$. Hence some $(N + n - a) \times (N + n - a)$ -minor of

$$\frac{\partial F_1 \dots F_m}{\partial Y_1 \dots Y_N}$$

is non-zero modulo F . Note further that $N + n - a = \text{height } F$.

REFERENCES

- [1] M. ARTIN: *On the solutions of Analytic Equations*. Inventiones Math. 5, 277–291. (1968).
- [2] M. ARTIN: *Algebraic Approximation of structures over complete local rings*. IHES no. 36, 23–58.
- [3] J. DIEUDONNÉ and A. GROTHENDIECK: *Eléments de géométrie algébrique*. IHES no. 20, 24,
- [4] M. J. GREENBERG: *Rational points in Henselian discrete valuation rings*. IHES no. 31, 59–64.
- [5] M. NAGATA: *Local rings*. Interscience Publishers 1962.
- F. OORT: *Hensel's lemma and rational points over local rings*. Symposia Mathematica, vol. III, 1970, 217–232.

(Oblatum 23-IX-1974)

Mathematisch Instituut
Budapestlaan
Utrecht, The Netherlands