

COMPOSITIO MATHEMATICA

TAMMO TOM DIECK

**A finiteness theorem for the burnside ring
of a compact Lie group**

Compositio Mathematica, tome 35, n° 1 (1977), p. 91-97

http://www.numdam.org/item?id=CM_1977__35_1_91_0

© Foundation Compositio Mathematica, 1977, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A FINITENESS THEOREM FOR THE BURNSIDE RING OF A COMPACT LIE GROUP

Tammo tom Dieck

Let G be a compact Lie group and let $A(G)$ be its Burnside ring [6]. We show that after inverting a finite number of primes the ring $A(G)$ is generated by idempotent elements. The following result (Theorem 1) about compact Lie groups is basic for our investigations.

Let H be a subgroup of G (subgroups will always be closed), let NH be its normalizer in G and denote NH/H by WH . If K is a compact Lie group let K_0 be its component of the unit element.

THEOREM 1: *There exists an integer b such that for each closed subgroups H of G the index $|WH : (WH)_0|$ is less than b .*

Let $A_c(G)$ be the integral closure of $A(G)$ in its total quotient ring.

THEOREM 2: *There exists an integer $n > 0$ such that $nA_c(G) \subset A(G)$. The minimal such n is the least common multiple of the numbers WH where H runs through all subgroups such that WH is finite.*

The minimal integer $n(G)$ provided by Theorem 1 replaces the order of the finite group if one extends the general Artin induction theorem (see Dress [7], Theorem 2, p. 204) to compact Lie groups, whence its importance.

1. Normalizers

1.1. We prove in this section Theorem 1. The proof proceeds in three steps: We first reduce to the case that WH is finite; then we reduce to the case that H is finite; and finally show that for finite H

with finite WH the order of WH is uniformly bounded.

Let H be a closed subgroup of G (notation: $H < G$). Let $\text{Aut}(H)$ be the automorphism group of H and $\text{In}(H)$ the closed normal subgroup of inner automorphisms. The group $\text{Aut } H/\text{In } H$ is discrete. Mapping $n \in NH$ to the conjugation automorphism $c(n): h \mapsto nhn^{-1}$ of H induces a homomorphism $NH \rightarrow \text{Aut } H/\text{In } H$ with kernel $ZH \cdot H$, where ZH denotes the centralizer of H . Hence $NH/ZH \cdot H$ being a compact subgroup of the discrete group $\text{Aut } H/\text{In } H$ is finite. We conclude

LEMMA 1: *WH is finite if and only if $ZH/(ZH \cap H)$ is finite.*

LEMMA 2: *A compact Lie group contains only a finite number of conjugacy classes (K) where K is the centralizer of a closed subgroup.*

PROOF: Let G act on $M = G$ via conjugation $G \times M \rightarrow M: (g, m) \mapsto gmg^{-1}$. If $H < G$ then the fixed point set M^H is the centralizer ZH . A compact differentiable G -manifold has finite orbit type. Hence there exist finitely many conjugacy classes $(H_1), \dots, (H_k)$ such that for any closed subgroup H $M^H = M^K$ and $(K) = (H_i)$ for a suitable i .

LEMMA 3: *For any $H < G$ the group $ZH \cdot H$ has finite index in its normalizer.*

PROOF: We have $Z(ZH \cdot H) < ZH < ZH \cdot H$; hence the assertion follows from Lemma 1.

If $n \in G$ normalizes H then also ZH and hence $ZH \cdot H$. We therefore have

$$NH/ZH \cdot H < N(ZH \cdot H)/ZH \cdot H.$$

Using Lemma 3 and the existence of an upper bound for the set

$$\{|WH| \mid H < G, WH \text{ finite}\} =: F(G)$$

we obtain

LEMMA 4: *There exists an integer c such that for all $H < G$ we have*

$$|NH/ZH \cdot H| < c.$$

Now we are able to obtain the first reduction of our problem. From the exact sequence $1 \rightarrow ZH/ZH \cap H \rightarrow WH \rightarrow NH/ZH \cdot H \rightarrow 1$ we see

that $WH/(WH)_0 \rightarrow NH/ZH \cdot H$ has a kernel which is a quotient of $ZH/(ZH)_0$. Lemmas 2 and 4 then show that

$$\{|WH/(WH)_0|H < G\}$$

is bounded. But note that Lemma 4 requires a bound for the set $F(G)$.

1.2. We show by induction over $|G/G_0|$ and $\dim G$ that $F(G)$ has an upper bound $a = a(G/G_0, \dim G)$. For finite G we can take $a = |G|$.

Suppose that an upper bound $a(K/K_0, \dim K)$ is given for all K with $\dim K < \dim G$. Let $\Sigma(G) = \{H < G \mid WH \text{ finite}\}$. Suppose $H \in \Sigma(G)$ is not finite. We consider the projection $p: NH_0 \rightarrow NH_0/H_0 =: U$. Let V be the normalizer of H/H_0 in U . Then $WH = V/(H/H_0)$ and therefore $H/H_0 \in \Sigma(U)$. Since $\dim U < \dim G$ we obtain by induction hypothesis

$$|WH| \leq a(U/U_0, \dim U).$$

We show that the possible values for $|U/U_0|$ are finite in number. The group NH_0 is the normalizer of a connected subgroup. By [8], Ch. VII, Lemma 3.2, there are only a finite number of conjugacy classes of such subgroups. Hence for a given G the possible $|U/U_0|$ are bounded, say $|U/U_0| \leq m(G)$. We have inequalities

$$\begin{aligned} |U/U_0| &\leq |NH_0/N_0H_0| |N_0H_0/(NH_0)_0| \\ &\leq |G/G_0| m(G), \end{aligned}$$

where N_0 means normalizer in G_0 . By the classification theory of compact connected Lie groups there are only a finite number in each dimension. Hence there exists a bound for $|U/U_0|$ depending only on $|G/G_0|$ and $\dim G$. This proves the induction step as far as the non-finite H in $\Sigma(G)$ are concerned.

1.3. Let $H \in \Sigma(G)$ be finite. Let $\sigma(G)$ be the set of finite subgroups of G . We use the following classical theorem of Jordan.

LEMMA 5: *There exists an integer $j = j(|G/G_0|, \dim G)$ with the following properties: To each $H \in \sigma(G)$ there exists an abelian normal subgroup A_H of H such that $|H/A_H| \leq j$. Moreover the A_H can be chosen such that $H < K$ implies $A_H < A_K$.*

PROOF: Boothby and Wang [2]. Wolf [9]. In these references only connected groups are considered. The straightforward extension to non-connected groups we leave to the reader.

If $H \in \Sigma(G)$ is finite then also $K := NH$ is finite and by Lemma 1 $K \in \Sigma(G)$. We choose $j = j(|G/G_0|, \dim G)$ and A_H, A_K according to Lemma 5. We have

$$|K/H| \leq |K/A_K| \cdot |A_K/H \cap A_K| \leq j|A_K/H \cap A_K|.$$

Hence it is sufficient to find a bound for $|A_K/H \cap A_K|$. Consider the exact sequence $1 \leftarrow S \leftarrow H \leftarrow A_H \leftarrow 1$. The conjugation $c(a)$ with $a \in A_K$ is trivial on A_H , because $A_K > A_H$, and hence $c(a)$ induces an automorphism of S . Since $|S| \leq j$ this automorphism has order at most $J = j!$, i.e. $c(a^r)$ is the identity on S and A_H for a suitable $r \leq J$. The group of such automorphisms modulo the subgroups of inner automorphisms by elements of A_H is isomorphic to $H^1(S; A_H)$, with S acting on A_H by conjugation. Since this group is annihilated by $|S|$ we see that $c(a^s)$ is an inner automorphism by an element of A_H for a suitable $s \leq J|S| \leq jJ$. In other words: $a^s h^{-1} \in ZH$. Hence it is sufficient to find a bound for the order of $A_K \cap ZH/H \cap A_K \cap ZH$.

Let $U_1 = A_K \cap ZH$. By [3], Théorème 1, U_1 is contained in the normalizer NT of a maximal torus of G . Put $U = U_1 \cap T$. Then $|U_1/U| \leq |G/G_0| |w(G_0)|$ where $w(G_0)$ denotes the Weyl group of G_0 . We estimate the order of U . Since U is abelian we have $U < ZU$. Moreover $H < ZU$ by definition of ZH . Since U is contained in ZU it is contained in the center $C := CZU$ of ZU . The inclusion $H < ZU$ implies $C < NH$. Hence C is finite.

We proceed to show that for the order of a finite center $C(G)$ of G there exists a bound depending only on $|G/G_0|$ and $\dim G$. We let G/G_0 act by conjugation on $C(G_0)$. Then $C(G) \cap G_0$ is the fixed point set of this action. We have $C(G_0) = A \times T_1$, where A is a finite abelian group and T_1 a torus. The group A is the center of a semisimple group and therefore, by the classification theory of these groups, $|A|$ is bounded by a constant c depending only on $\dim G$. The exact cohomology sequence associated to the universal covering $0 \rightarrow \pi_1 T_1 \rightarrow V \rightarrow T_1 \rightarrow 0$ shows, that the fixed point set of the action of G/G_0 on $T_1 = C(G_0)_0$ is isomorphic to $H^1(G/G_0, \pi_1 T_1)$, hence its order is bounded by a constant d depending only on $|G/G_0|$ and the rank of T_1 . Hence $|C(G)| \leq |G/G_0|cd$.

Finally we have to show that for the possible groups ZU the order $|ZU/(ZU)_0|$ is bounded.

U is contained in a maximal torus of G . Therefore ZU is a subgroup of maximal rank and $(ZU)_0$ a connected subgroup of maximal rank. By [4] there exist only finitely many conjugacy classes of connected subgroups of maximal rank. We have

$$|ZU/(ZU)_0| \leq |N(ZU)_0/(ZU)_0| \leq |G/G_0| |N_0(ZU)_0/(ZU)_0|.$$

There are only finitely many possibilities for normalizers $N_0(ZU)$ in G_0 of $(ZU)_0$.

This finishes the proof of Theorem 1.

2. The integral closure of $A(G)$

Let $\phi = \text{Spec}(A(G) \otimes Q)$ be the prime ideal spectrum of $A(G) \otimes_Z Q =: A_0$ with Zariski topology. By [6], Theorem 4, the prime ideals of A_0 correspond bijectively to kernels of ring homomorphisms $A_0 \rightarrow Q$. Therefore ϕ is a totally disconnected compact Hausdorff space ([5], §4. Ex. 16; [1], Ch. 3. Ex. 11).

LEMMA 6: (1) For each $a \in A_0$ the map $\varphi_a: \phi \rightarrow Q: \varphi \mapsto \varphi(a)$ is locally constant. If $a \in A(G)$ then $\varphi(a) \in Z$. (2) Let $C(\phi, Q)$ be the ring of locally constant functions $\phi \rightarrow Q$. The ring homomorphism $\alpha: A_0 \rightarrow C(\phi, Q): a \mapsto \varphi_a$ is an isomorphism. The image $\alpha A(G)$ is contained in $C(\phi, Z)$. (3) The map $A(G) \rightarrow A_0: a \mapsto a \otimes 1$ is the inclusion of A into its total quotient ring. The map $\alpha: A(G) \rightarrow C(\phi, Z)$ is the inclusion of $A(G)$ into the integral closure of $A(G)$ in $A_0 \cong C(\phi, Q)$.

PROOF: (1) For $k \in Q$ the set $\varphi_a^{-1}(k)$ is closed in ϕ , by definition of the Zariski topology. Since A_0 is an absolutely flat ring, this set is also open by [5], §4. Ex. 16.b. Hence φ_a is continuous and there exist only a finite number of non-empty sets $\varphi_a^{-1}(k)$, because ϕ is compact. A homomorphism $A_0 \rightarrow Q$ is induced from a homomorphism $A(G) \rightarrow Z$, by [6] Theorem 4.

(2) Since the localizations of A_0 at its prime ideals are canonically isomorphic to Q , we can identify $C(\phi, Q)$ with the ring of sections of the structure sheaf of A_0 . Then α corresponds to the canonical map of A_0 into this ring, hence α is an isomorphism.

(3) To form the total quotient ring we have to invert the elements which are not zero divisors. Hence A_0 is contained in the total quotient ring. If $x \in C(\phi, Q) \cong A_0$ is not a zero divisor then it is a locally constant function without zeros, hence a unit. Therefore $C(\phi, Q)$ is its own total quotient ring. Since a locally constant function $\phi \rightarrow Z$ takes only finitely many values the ring $C(\phi, Z)$ is generated by idempotent elements hence integral over any subring. Under the isomorphism α the ring $C(\phi, Z)$ corresponds to $\{a \in A_0 \mid \varphi \in \phi \Rightarrow \varphi(a) \in Z\}$. If $x \in A_0$ is integral over $A(G)$ then $\varphi(x)$ is integral over Z , hence $\varphi(x) \in Z$ and $\alpha(x) \in C(\phi, Z)$.

Let $A_c(G)$ denote the pre-image of $C(\phi, Z)$ under α . We recall that $A(G)$ is additively the free abelian group on homogeneous spaces G/H where H runs through a complete system of non-conjugate subgroups H of G with finite index in their normalizer ([6], Theorem 1). Let $\Sigma(G)$ denote the set of conjugacy classes (H) of subgroups H of G with finite $WH = NH/H$.

LEMMA 7: $A_c(G)$ is additively the free abelian group with basis $x_{(H)} := |WH|^{-1}G/H$, $(H) \in \Sigma(G)$.

PROOF: The elements $x_{(H)}$ are contained in $A_c(G)$: Suppose $\varphi \in \phi$ is given. Then there exists $(K) \in \Sigma(G)$ such that $\varphi(x \otimes r) = r\chi(x^K)$ where χ denotes the Euler characteristic and x^K the K fixed point set of any manifold representing x . But WH acts freely as a G -automorphisms group on G/H , hence also on G/H^K . Therefore $\chi(G/H^K)$ is divisible by $|WH|$ and we see that $\varphi(x_{(H)}) \in Z$ for all φ . By definition of $A_c(G)$ this means $x_{(H)} \in A_c(G)$.

The elements $x_{(H)}$ are obviously linearly independent over Z . We have to show that any $x \in A_c(G)$ is an integral linear combination of the $x_{(H)}$. In any case we have an expression $x = \sum r_H x_{(H)}$ with rational r_H . Take $(L) \in \Sigma(G)$ maximal with respect to inclusion such that $r_L \neq 0$. Then $\sum r_H |WH|^{-1} \chi(G/H^L) \in Z$. But $G/H^L = \phi$ for $(H) \neq (L)$, $r_H \neq 0$; and $\chi(G/L^L) = |WH|$. Therefore $r_L \in Z$. We apply the same argument to $x - r_L x_{(L)}$ and complete the proof by induction.

Lemma 7 and Theorem 1 give a proof of Theorem 2.

REFERENCES

- [1] M. F. ATIYAH and I. G. MACDONALD: *Introduction to Commutative Algebra*. Addison-Wesley Publ. Comp. 1969.
- [2] W. BOOTHBY and H.-C. WANG: On the finite subgroups of connected Lie groups. *Comment. Math. Helv.* 39 (1964) 281-294.
- [3] A. BOREL et J.-P. SERRE: Sur certain sous groupes des groupes de Lie compacts. *Comment. Math. Helv.* 27 (1953) 128-139.
- [4] A. BOREL et J. DE SIEBENTHAL: Les sous-groupes fermés de rang maximum des groupes de Lie clos. *Comment. Math. Helv.* 23 (1949) 200-221.
- [5] N. BOURBAKI: *Algèbre commutative, Chapitre 2*. Hermann, Paris 1961.
- [6] T. TOM DIECK: The Burnside Ring of a Compact Lie Group I. *Math. Ann.* 215 (1975) 235-250.
- [7] A. DRESS: Contributions to the theory of induced representations. *Springer Lecture Notes* 342, (1973) 183-240.

[8] *Seminar on Transformation Groups*. Ed. A. Borel et al. Princeton University Press 1960.

[9] J. A. WOLF: *Spaces of Constant Curvature*. McGraw-Hill, New York 1967.

(Oblatum 26-V-1976)

Math. Institut
34 Göttingen
Bunsenstrasse 3-5
BRD