

# COMPOSITIO MATHEMATICA

ANTONIO CASSA

**A theorem on complete intersection curves and a  
consequence for the runge problem for analytic sets**

*Compositio Mathematica*, tome 36, n° 2 (1978), p. 189-202

<[http://www.numdam.org/item?id=CM\\_1978\\_\\_36\\_2\\_189\\_0](http://www.numdam.org/item?id=CM_1978__36_2_189_0)>

© Foundation Compositio Mathematica, 1978, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**A THEOREM ON COMPLETE INTERSECTION CURVES AND  
 A CONSEQUENCE FOR THE RUNGE PROBLEM FOR  
 ANALYTIC SETS**

Antonio Cassa

**Summary**

The main goal of this article is to prove the following:

**APPROXIMATION THEOREM:** *Let  $X$  be a Stein complex analytic manifold of dimension  $n \geq 2$ ,  $A$  a Runge and Stein open set of  $X$  and  $C$  a curve of  $A$ ; there exists a sequence of curves  $\{C_k\}_{k \geq 1}$  of  $X$  such that:*

$$C = \lim_{k \rightarrow \infty} (C_k|_A)$$

*in the topological space  $Z_1^+(A)$  of positive analytic 1-cycles of  $A$ .*

The proof makes use essentially of the following:

**COMPLETE INTERSECTION THEOREM:** *For each relatively compact open set  $B$  of  $A$  there exist functions  $g_1, \dots, g_{n-1}$  holomorphic on  $B$  such that the positive analytic 1-cycle defined by  $g = (g_1, \dots, g_{n-1})$  in  $B$  is:*

$$V_1(g) = C|_B + m_1 \cdot (D_1|_B) + \dots + m_s \cdot (D_s|_B)$$

*where  $D_1, \dots, D_s$  are curves of  $X$  and  $m_1, \dots, m_s$  positive integers.*

In fact if  $\{g^{(k)}\}_{k \geq 1}$  is a sequence of maps  $g^{(k)}: X \rightarrow \mathbb{C}^{n-1}$  holomorphic on  $X$ , having at least multiplicity  $m_i$  on  $D_i$  for each  $i = 1, \dots, s$  and converging to  $g$ , for  $k$  big enough we have:

$$V_1(g^{(k)}) = C_{k|B} + m_1 \cdot (D_1|_B) + \dots + m_s \cdot (D_s|_B)$$

where the  $C_k$  are curves of  $X$ ; then in  $Z_1^+(B)$ :

$$C = \lim_{k \rightarrow \infty} (C_{k|B})$$

So every curve  $C$  of  $A$  can be approximated by curves of  $X$  on every relatively compact open set  $B$  of  $A$ , that is the restriction map:

$$Z_1(X) \longrightarrow Z_1(A)$$

has dense image in  $Z_1(A)$ .

Moreover if  $C$  is irreducible in  $A$  the curves  $C_k$  can be chosen irreducible in  $X$  and if  $X$  is an open set of  $\mathbb{C}^n$  they can be taken algebraic.

This proves that the so-called Runge problem has always solution for analytic cycles of dimension one. This is no longer true in general for higher dimension; Cornalba and Griffiths show in [7] page 76 there exists a non trivial condition for the approximability of an analytic set.

Under that condition they state a general Runge problem for analytic sets that they solve in the case of codimension one.

In that article (as in [4]) the topology of  $Z_d^+(X)$  is defined through the space of currents  $\mathcal{D}_{2d}(X)$ ; the properties of that topology are described in [11] and in a more geometric way in [3] or in [5].

I take the opportunity of thanking prof. A. Andreotti for all his help and mainly for his precious suggestions; likewise I wish to thank prof. M. Cornalba and prof. Ph. Griffiths for having communicated to me their ideas about the Runge problem.

### List of symbols

$\text{reg } V$  = manifold of all the regular points of the analytic space  $V$ .

$\text{sing } V = V - \text{reg } V$  = subspace of the singular points of  $V$ .

$T_x(V) = \text{Hom}_{\mathbb{C}}(\mathcal{M}_x/\mathcal{M}_x^2, \mathbb{C})$  = Zarinski tangent space at  $x \in V$ .

$\dim t_x(V) = \dim_{\mathbb{C}} T_x(V)$  = embedding dimension = tangential dimension.

$Z_d(W)$  = topological group of the analytic  $d$ -cycles in the manifold  $W$ .

$Z_d^+(W)$  = cone in  $Z_d(W)$  of the positive  $d$ -cycles of  $W$ .

$V_d(f)$  = positive  $d$ -cycle defined by the equation  $f = 0$ , where  $f: W \rightarrow \mathbb{C}^r$  is an holomorphic map.

**§1. The estimate of the rank of a sheaf using the Endromisbündel of Forster and Ramspott**

Let  $\mathcal{F}$  be a coherent sheaf on a complex analytic manifold  $X$ .

For each point  $x \in X$  the least number of generators of the stalk  $\mathcal{F}_x$  is given by the dimension on  $\mathbb{C}$  of the vector space  $L_x(\mathcal{F}) = \mathcal{F}_x / (\mathcal{M}_x \cdot \mathcal{F}_x)$ .

If this number is bounded in  $X$  the sheaf  $\mathcal{F}$  has finite rank, that is there exist sections  $f_1, \dots, f_r \in \Gamma(X, \mathcal{F})$  generating all the stalks  $\mathcal{F}_x$  for every  $x \in X$  (see [6]).

Taken a positive integer  $s \leq r$ , the existence of  $s$  sections  $g_1, \dots, g_s \in \Gamma(X, \mathcal{F})$  with the same property is equivalent to the existence of an holomorphic section of a bundle  $E(\mathcal{F}; f, r)$  on  $X$  called Endromisbündel (see [8]).

The Endromisbündel is an open set of  $X \times \mathbb{C}^{rs}$  obtained subtracting analytic subspaces defined by the sections  $f_1, \dots, f_r$  and by the numbers  $\{\dim_{\mathbb{C}} L_x(\mathcal{F}); x \in X\}$ .

Let's put for each integer  $k \geq 0$ :

$$Y_k(\mathcal{F}) = \{x \in X : \dim_{\mathbb{C}} L_x(\mathcal{F}) \geq k\}$$

the family  $\{Y_k(\mathcal{F})\}_{k \geq 0}$  is a decreasing sequence of analytic subspaces of  $X$  which are surely empty for  $k \geq r + 1$ .

On the analytic space  $X_k(\mathcal{F}) = Y_k(\mathcal{F}) - Y_{k+1}(\mathcal{F})$  the Endromisbündel is a locally trivial holomorphic bundle whose fibre  $F_{r,s,k}$  is homotopic to the manifold  $W_{sk}$  of all the orthonormal  $k$ -frames of  $\mathbb{C}^s$ .

The main result of [8] (satzen 5 and 6) claims that if  $X$  is holomorphically convex the existence of a holomorphic section of the Endromisbündel is equivalent to the existence of a continuous section.

Therefore the evaluation of the rank of  $\mathcal{F}$  is a purely topological problem whose main ingredients are the spaces  $Y_k(\mathcal{F})$  and the fibres  $W_{sk}$ .

The following proposition is a way to make sure the existence of a continuous section of  $E(\mathcal{F}, f, s)$  supposing zero all the cohomology groups containing the obstructions.

**PROPOSITION:** *Let  $X$  be a Stein manifold and  $\mathcal{F}$  a coherent analytic sheaf having his rank bounded by an integer  $s$ .*

*If for each  $k \geq 0$  and  $q \geq 1$ :*

$$H^{q+1}(Y_k(\mathcal{F}), Y_{k+1}(\mathcal{F}); \pi_q(W_{sk})) = 0$$

then there exist  $s$  sections  $g_1, \dots, g_s \in \Gamma(X, \mathcal{F})$  generating all the stalks of  $\mathcal{F}$ .

PROOF: Let  $f_1, \dots, f_r$  be global sections of  $\mathcal{F}$  generating all the stalks of  $\mathcal{F}$ ; proceeding by induction on  $h = r - k$  from 0 to  $r$  we will prove there exists a continuous section of  $E(\mathcal{F}, f, s)$  on  $Y_{r-h}(\mathcal{F})$  for  $h = 0, \dots, r$ .

If  $h = 0$  since  $Y_{r+1} = \phi$  the bundle  $E(\mathcal{F}, f, s)$  is a locally trivial fibre bundle with fibre homotopic to  $W_{sr}$ ; the condition  $H^{q+1}(Y_r; \pi_q(W_{sr})) = 0$  is just the one we need to prove the existence of a continuous section on  $Y_r$  (see [13] page 174).

Let's prove now we can extend a continuous section from  $Y_{r-(h-1)}$  to  $Y_{r-h}$ ; we can find a triangulation of  $Y_{r-h}$  in such a way  $Y_{r-(h-1)}$  is a subpolyhedron furnished of a neighborhood  $U$  which is again a subpolyhedron of  $Y_{r-h}$  and contractible on  $Y_{r-(h-1)}$ .

Since  $E(\mathcal{F}, f, s)$  is an open set of  $C^r \times X$  choosing  $U$  suitably small we can, first of all, extend our continuous section from  $Y_{r-(h-1)}$  to  $U$ ; then we can extend the section from  $U - Y_{r-(h-1)}$  to  $Y_{r-h} - Y_{r-(h-1)}$  because for each  $q \geq 1$  we have:

$$H^{q+1}(Y_{r-h} - Y_{r-(h-1)}, U - Y_{r-(h-1)}; \pi_q(W_{s,r-h})) = 0$$

In fact:

$$\begin{aligned} H^{q+1}(Y_{r-h} - Y_{r-(h-1)}; U - Y_{r-(h-1)}) &\simeq H^{q+1}(Y_{r-h}, U) \\ &\simeq H^{q+1}(Y_{r-h}, Y_{r-(h-1)}) \simeq H^{q+1}(Y_k, Y_{k+1}) = 0. \end{aligned}$$

## §2. Complete intersection curves

Let  $C$  be a curve of an open set of  $C^n$  and  $x_0$  a singular point of  $C$ , if  $\dim t_{x_0}(C) = 2$  then the curve  $C$  is complete intersection at  $x_0$ .

In fact there exist a manifold  $M$  of dimension 2 in  $C^n$  and a neighborhood  $U$  of  $x_0$  such that  $C \cap U \subset M \cap U$ ; restricting, in case,  $U$  we can find a function  $f_n$  holomorphic on  $U$  such that  $\mathcal{F}_{C \cap U, M \cap U} = f_n \cdot \mathcal{O}_{M \cap U}$  and functions  $f_2, \dots, f_{n-1}$  holomorphic on  $U$  such that  $\mathcal{F}_{M \cap U, U} = f_2 \cdot \mathcal{O}_U + \dots + f_{n-1} \cdot \mathcal{O}_U$ ; therefore

$$\mathcal{F}_{C \cap U, U} = f_2 \cdot \mathcal{O}_U + \dots + f_n \cdot \mathcal{O}_U.$$

The following two lemmas prove in most cases that if  $t = \dim t_{x_0}(C)$  is bigger than 2, then adding to  $C$  some lines  $L_1, \dots, L_{t-2}$  through  $x_0$  the curve  $C \cup (L_1 \cup \dots \cup L_{t-2})$  is complete intersection at  $x_0$ .

LEMMA 1: Let  $C$  be a curve of an open set of  $\mathbb{C}^n$  (with  $n \geq 2$ ) and the origin  $0$  a singular point of  $C$ .

Denoted by  $L_1, \dots, L_n$  the coordinate axes of  $\mathbb{C}^n$  and written  $L_0 = \{0\}$ , if the following hypothesis is verified:

- (i) the projection map  $p: \mathbb{C}^n \rightarrow \mathbb{C}^2$  defined by  $p(z_1, \dots, z_n) = (z_{n-1}, z_n)$  is injective on  $C$  in a neighborhood of  $0$ .

then a neighborhood  $V$  of  $0$ , an integer  $s = 0, \dots, n - 2$ , a Stein neighborhood  $U$  of  $(L_0 \cup \dots \cup L_s)$  and functions  $f_1, \dots, f_{n-1}$  holomorphic on  $U$  exist such that:

- (1)  $\{x \in U : f_1(x) = \dots = f_{n-1}(x) = 0\} = (C \cap V \cap U) \cup (L_0 \cup \dots \cup L_s)$
- (2) the germs  $f_{1,x}, \dots, f_{n-1,x}$  generate the stalk  $\mathcal{T}_{C,x}$  for each  $x \in C \cap V \cap U - \{0\}$ .

PROOF: Let's proceed by induction on  $n \geq 2$ . For  $n = 2$  the conclusion is well known. For  $n \geq 3$  let's suppose we have already proved the lemma for all the curves  $C'$  of  $\mathbb{C}^{n'}$  with  $n' < n$  and let's prove it for the curves  $C$  of  $\mathbb{C}^n$ .

Let's denote by  $q: \mathbb{C}^n \rightarrow \mathbb{C}^{n-1}$  the projection along the axis  $L_{n-2}$  defined by  $q(z_1, \dots, z_{n-2}, z_{n-1}, z_n) = (z_1, \dots, 0, z_{n-1}, z_n)$  with values in  $\mathbb{C}^{n-1} = \{z \in \mathbb{C}^n : z_{n-2} = 0\}$ .

For the hypothesis (i) it is possible to find a neighborhood  $V$  of  $0$  where  $q$  is injective on  $C$ . Rechoosing in case  $V$  we can suppose the map  $q: V \cap C \rightarrow q(V)$  proper; therefore  $C' = q(C \cap V)$  is a curve of  $V' = q(V)$  open neighborhood of  $0$  in  $\mathbb{C}^{n-1}$ .

We can choose  $V$  small enough to have also  $\text{sing}(C) \cap V = \{0\} = \text{sing}(C')$ .

The curve  $C'$  of  $\mathbb{C}^{n-1}$  in respect to the coordinates  $z_1, \dots, \hat{z}_{n-2}, z_{n-1}, z_n$  verifies the hypothesis (i); for the induction there exist an integer  $s' = 0, \dots, n - 3$ , a Stein neighborhood  $U'$  of  $L_0 \cup \dots \cup L_{s'}$  and functions  $f'_1, \dots, f'_{n-2}$  holomorphic on  $U'$  verifying the theses (1) and (2).

Using (i) it can be verified that the restriction of  $q$  gives a map  $\hat{q}: (C \cap V) \cup (L_0 \cup \dots \cup L_s) \rightarrow C' \cup (L_0 \cup \dots \cup L_{s'})$  bijective and holomorphic, whose inverse is meromorphic, continuous and bi-holomorphic out of  $0$ . Likewise the function  $m$  in  $C' \cup (L_0 \cup \dots \cup L_{s'})$  defined by  $m(x') = z_{n-2}(\hat{q}^{-1}(x'))$  is meromorphic, continuous, holomorphic out of  $0$  and vanishes on  $(L_0 \cup \dots \cup L_{s'})$ . Therefore  $m(x') = a'(x')/b'(x')$  everywhere  $b'(x') \neq 0$  for two functions  $a', b'$  holomorphic on  $C' \cup (L_0 \cup \dots \cup L_{s'})$  with  $b'$  not identically zero on any irreducible component and  $a' = 0$  on  $L_0 \cup \dots \cup L_{s'}$ .

Solving a  $\mathcal{O}^*$ -cohomological problem we can find two functions  $a$  and  $b$  holomorphic such that  $b(x') \neq 0$  if  $x' \neq 0$  and  $m(x') = a(x')/b(x')$  for each  $x' \neq 0$ .

Since  $U'$  is Stein we can suppose  $a$  and  $b$  defined on  $U'$ ; written  $U = q^{-1}(U')$ ,  $f_1 = f'_1 \circ q, \dots, f_{n-2} = f'_{n-2} \circ q, f_{n-1} = (b \circ q) \cdot z_{n-2} - (a \circ q)$  the theses (1) and (2) are verified for the curve  $C$  together with the lines  $L_0, \dots, L_{s'}, L_{n-2}$  if  $b(0) = 0$  or the lines  $L_0, \dots, L_{s'}$  if  $b(0) \neq 0$ .

LEMMA 2: *Let  $C$  be as in Lemma 1 and  $n \geq 3$ ; there exists a coordinate system in  $\mathbb{C}^n$  verifying (i).*

*Moreover if  $E$  is a measurable subset of  $\mathbb{C}^n - \{0\}$  with Hausdorff measure  $H_r(E) = 0$  for each  $r > 2$ , the coordinate system can be chosen in such a way to have:*

$$(L_1 \cup \dots \cup L_{n-2}) \cap E = \emptyset$$

PROOF: For each  $n$ -uple of lines  $L = (L_1, \dots, L_n)$  in general position and for each  $i = 1, \dots, n - 1$  let's write  $V_{L,i} = L_i + \dots + L_n$  and let's denote  $p_{L,i}: \mathbb{C}^n \rightarrow V_{L,i}$  and  $q_{L,i+1}: V_{L,i} \rightarrow V_{L,i+1}$  the natural projections.

Since  $p_{n-1} = (q_{n-1}) \circ \dots \circ (q_2)$ , if  $p_{n-1}$  is not injective on  $C$  in any neighborhood of  $0$ , then some of the projections  $q_{i+1}$  (where  $i = 1, \dots, n - 2$ ) is not injective on the set  $p_i(C)$  in any neighborhood of  $0$ ; therefore for each integer  $j \geq 1$  there exist two points  $z'_j$  and  $z''_j$  of  $C$  such that  $p_i(z'_j)$  and  $p_i(z''_j)$  are different, non zero,  $|p_i(z'_j)| < 1/j, |p_i(z''_j)| < 1/j$  and  $(p_i(z'_j) - p_i(z''_j)) \in L_i$ .

Then the intersection  $L_i \cap (p_i(C) - p_i(C))$  has interior part not empty in  $L_i$ , this set is in fact the image of the holomorphic map  $d_i: d^{-1}(L_i) \cap (C \times C) \rightarrow L_i$  where  $d: \mathbb{C}^n + \mathbb{C}^n \rightarrow V_i$  is defined by  $d(z', z'') = p_i(z') - p_i(z'')$  which is of rank one at least in some point containing in its image the sequence  $\{(p_i(z'_j) - p_i(z''_j))\}_{j \geq 1}$  infinite and converging to  $0$ .

Written  $G = \{g \in \mathbb{C}^*: g = e^{a+bi} \text{ with } a, b \in \mathbb{Q}\}$ ,  $S_i = p_i(C) - p_i(C)$ ,  $S'_i = \cup_{g \in G} g \cdot S_i$  we have  $L_i = \cup_{g \in G} g \cdot (L_i \cap S_i)$  and therefore  $L_i \subset S'_i + (L_0 + \dots + L_{i-1})$ . Let's prove at this point that for each  $i = 1, \dots, n - 2$  and for each  $L' = (L_0, \dots, L_{i-1}) \in \{L_0\} \times (\mathbb{P}^{n-1})^{i-1}$  (where  $L_0 = \{0\}$ ) the set  $R_{L'} = \{L \in \mathbb{P}^{n-1}: L \subset S'_i + L_0 + \dots + L_{i-1}\}$  has measure zero in  $\mathbb{P}^{n-1}$ .

In fact written  $T_{L'} = \cup_{L \in R_{L'}} L$ , because  $T_{L'} \leq S'_i + L_0 + \dots + L_{i-1}$  and  $H_r(S'_i) = 0$  if  $r > 4$ , it must be  $H_r(T_{L'}) = 0$  for  $r > 4 + 2(i - 1)$  and therefore  $H_r(R_{L'}) = 0$  if  $r > 2 + 2(i - 1) = 2i$  since  $T_{L'} - \{0\} \cong R_{L'} \times \mathbb{C}^*$ , so we can conclude  $\mu(R_{L'}) = H_{2n-2}(R_{L'}) = 0$  because  $2n - 2 > 2i$ .

We are able now to prove that for each  $k = 1, \dots, n - 2$  there exist  $k$  lines  $L_1, \dots, L_k$  in general position such that  $(L_1 \cup \dots \cup L_k) \cap E = \emptyset$  and for each  $l = 1, \dots, k$  written  $L'_l = (L_0, \dots, L_{l-1})$  we have  $L_l \notin R_{L'}$ .

For  $k = n - 2$  the lemma will result proved.

For  $k = 1$  we have to find  $L_1 \in \mathbb{P}^{n-1}$  such that  $L_1 \not\subseteq R_{L_0} \cup E'$  where  $E'$  is the image of  $E$  in  $\mathbb{P}^{n-1}$  for the natural quotient map. It is possible to find  $L_1$  since the set of directions to avoid has measure zero in  $\mathbb{P}^{n-1}$ .

Given the lines  $L_1, \dots, L_{k-1}$  with the properties listed above we have to find a line  $L_k$  in general position in respect with the others and in such a way  $L_k \not\subseteq R_{L'} \cup E'$ .

Again it is possible to choose  $L_k$  since the set of directions to avoid has measure zero.

**THEOREM 1:** *Let  $X$  be a Stein manifold of dimension  $n \geq 3$ ,  $A$  a Runge and Stein open set of  $X$  and  $C$  a curve of  $A$ .*

*For each relatively compact open set  $B$  of  $A$  there exist a curve  $D$  of  $X$  and functions  $g_1, \dots, g_{n-1}$  holomorphic on  $B$  such that:*

- (1)  $\{x \in B : g_1(x) = \dots = g_{n-1}(x) = 0\} = (C \cup D) \cap B$
- (2) *the germs  $g_{1,x}, \dots, g_{n-1,x}$  generate the stalk  $\mathcal{T}_{C,x}$  for each  $x \in C \cap B - S$ , where  $S = \{x \in C \cap B : C \text{ is not complete intersection at } x\}$ .*

**PROOF:** Enlarging  $B$  we can suppose it a Runge and Stein open set yet relatively compact in  $A$ .

The set  $S$  contained in  $\text{sing}(C) \cap B$  is finite; if  $S = \emptyset$  the curve  $C \cap B$  is locally complete intersection (ideal theoretically) and therefore it is complete intersection in  $B$  (see [8] page 162, the Remark (b) to Corollary (2) of Theorem (9)).

If  $S \neq \emptyset$  let's write  $S = \{x_1, \dots, x_p\}$ ; we show that however fixed an integer  $r = 1, \dots, p$  for each  $j = 1, \dots, r$  there exist a curve  $D_j$  of  $X$ , an open neighborhood  $U_j$  of  $D_j$  and functions  $f_{j,1}, \dots, f_{j,n-1}$  holomorphic on  $U_j$  such that:

- (A)  $\{x \in U_j \cap B : f_{j,1}(x) = \dots = f_{j,n-1}(x) = 0\} = (C \cup D_j) \cap U_j \cap B$
- (B) the germs  $f_{j,1,x}, \dots, f_{j,n-1,x}$  generate  $\mathcal{T}_{C,x}$  for each  $x \in C \cap U_j - \{x_j\}$
- (C)  $D_j \cap C \cap B = \{x_j\}$
- (D)  $U_k \cap U_j \cap B = \emptyset$  for each  $k < j \leq r$ .

Let's proceed by induction on  $r$ . Let  $r = 1$ ; it is possible to find a holomorphic map  $R : X \rightarrow \mathbb{C}^n$  regular in  $x_1$  and such that  $R^{-1}(0) = \{x_1\}$  (see [8] page 161, Corollary 1 of Theorem 9). Replacing  $R$  with another map (denoted again by  $R$ ) near enough to  $R$  we can have (see [9] page 168, Theorem 4):

- (1) for all the points  $x$  of a neighborhood  $W$  of  $x_1 : R^{-1}(R(x)) \cap \bar{B} = \{x\}$
- (2)  $R(x_1) = 0$

(3)  $\dim(R^{-1}(R(x))) = 0$  for each  $x \in X$ .

(4)  $R$  establishes a biholomorphism between  $W$  and  $W' = R(W)$  open set of  $C^n$ .

Applying the Lemmas 1 and 2 to the curve  $C' = R(C \cap W)$  of the open set  $W'$  of  $C^n$  and to the set  $E = R(C) - \{0\}$ , it is possible to find a coordinate system  $(z_1, \dots, z_n)$  in  $C^n$  whose coordinate axes we denote by  $L_1, \dots, L_n$  ( $L_0 = \{0\}$ ), a neighborhood  $V'$  of 0 contained in  $W'$ , an integer  $s = 0, \dots, n - 2$ , a neighborhood  $U'$  of  $L_0 \cup \dots \cup L_s$  and functions  $f'_1, \dots, f'_{n-1}$  holomorphic on  $U'$  such that:

(1)  $\{z \in U': f'_i(z) = \dots = f'_{n-1}(z) = 0\}$   
 $= (C' \cap V' \cap U') \cup (L_0 \cup \dots \cup L_s)$

(2) the germs  $f'_{1,z}, \dots, f'_{n-1,z}$  generate  $\mathcal{I}_{C',z}$  for each  $z \in C' \cap V' \cap U' - \{0\}$

(3)  $(L_0 \cup \dots \cup L_s) \cap R(C) = \{0\}$ .

Let's put  $D_1 = R^{-1}(L_0 \cup \dots \cup L_s)$ , since  $D_1 \cap C \cap \bar{B} = \{x_1\}$  we can find a neighborhood  $U_1$  of  $D_1$  contained in  $R^{-1}(U')$  such that  $C \cap (U_1 \cap B) \subset C \cap W$ ; on  $U_1$  let's define the functions  $f_{1,1} = f'_1 \circ R, \dots, f_{1,n-1} = f'_{n-1} \circ R$ .

For these sets and functions the conditions (A) (B) (C) and (D) listed above are verified.

Let's suppose now  $r > 1$  and we have found for each  $j = 1, \dots, r - 1$  a curve  $D_j$  of  $X$ , a neighborhood  $U_j$  of  $D_j$  and functions  $f_{j,1}, \dots, f_{j,n-1}$  satisfying the conditions (A) (B) (C) and (D) and let's show how to add a curve  $D_r$ , a neighborhood  $U_r$  of  $D_r$  and functions  $f_{r,1}, \dots, f_{r,n-1}$  in such a way the properties (A) (B) (C) and (D) are verified for each  $k < j \leq r$ .

Again we consider a holomorphic map  $R_r: X \rightarrow C^n$  such that:

(1) for all the points  $x$  of an open neighborhood  $W_r$  of  $x_r$  we have  $R_r^{-1}(R_r(x)) \cap \bar{B} = \{x\}$

(2)  $R_r(x_r) = 0$

(3)  $\dim R_r^{-1}(R_r(x)) = 0$  for each  $x \in X$ .

As above we apply the Lemmas (1) and (2) to the curve  $C'_r = R_r(C \cap W_r)$  of the open set  $W'_r = R_r(W_r)$  of  $C^n$  and the set  $E_r = R_r(C \cup D_1 \cup \dots \cup D_{r-1}) - \{0\}$ ; written  $D_r = R_r^{-1}(L_0 \cup \dots \cup L_{s_r})$ , again we can find a neighborhood  $U_r$  of  $D_r$  such that  $C \cap U_r \cap B \subset C \cap W_r$  and define  $f_{r,1} = f'_1 \circ R_r, \dots, f_{r,n-1} = f'_{n-1} \circ R_r$ ; moreover since  $D_i \cap D_j \cap B = \emptyset$  if  $i \neq j \leq r$  we can choose  $U_1, \dots, U_r$  in such a way to have  $U_i \cap U_j \cap B = \emptyset$  for each  $i \neq j \leq r$ , and again these sets and functions satisfy the conditions (A) (B) (C) and (D).

Arrived with  $r$  to  $p$ , let's put  $D = D_1 \cup \dots \cup D_p$  and let's define for each  $i = 1, \dots, p$  a coherent sheaf  $\mathcal{F}_i$  on  $U_i \cap B$  putting:

$$\mathcal{F}_i = f_{i,1} \cdot \mathcal{O}_{|U_i \cap B} + \dots + f_{i,n-1} \cdot \mathcal{O}_{|U_i \cap B}$$

For each  $x \in U_i \cap B - D$  we have  $\mathcal{T}_{i,x} = \mathcal{T}_{C,x}$ .  
 We can now define a sheaf  $\mathcal{T}$  on  $B$  writing:

$$\mathcal{T}_x = \begin{cases} \mathcal{T}_{i,x} & \text{if } x \in B \cap U_i \\ \mathcal{T}_{C,x} & \text{if } x \in B - D \end{cases}$$

The sheaf  $\mathcal{T}$  is well defined and coherent; moreover

$$\dim L_x(\mathcal{T}) = \begin{cases} 1 & \text{for each } x \in B - (C \cup D) \\ n - 1 & \text{for each } x \in (C \cup D) \cap B \end{cases}$$

therefore the sheaf  $\mathcal{T}$  has limited rank on  $B$ .

To complete the theorem's proof we have to check that the rank of  $\mathcal{T}$  is just  $n - 1$ .

For what has been reported in §1 since we have:

$$Y_0(\mathcal{T}) = Y_1(\mathcal{T}) = B, \quad Y_2(\mathcal{T}) = \dots = Y_{n-1}(\mathcal{T}) = (C \cup D) \cap B \quad \text{and} \\ Y_r(\mathcal{T}) = \emptyset \text{ for each } r \geq n, \text{ we have to prove that for each } q \geq 1:$$

$$H^{q+1}((C \cup D) \cap B; \pi_q(W_{n-1,n-1})) = 0$$

and

$$H^{q+1}(B, (C \cup D) \cap B; \pi_q(W_{n-1,1})) = 0$$

The first cohomology groups vanish because  $(C \cup D) \cap B$  is a Stein curve; for the second we have  $W_{n-1,1} \approx S^{2n-3}$ , therefore  $\pi_q(W_{n-1,1}) = 0$  for each  $1 \leq q \leq 2n - 4$ .

For  $q \geq 2n - 3 \geq n \geq 3$  from the exact sequence:

$$\dots \longrightarrow H^q((C \cup D) \cap B; G) \longrightarrow H^{q+1}(B, (C \cup D) \cap B; G) \\ \longrightarrow H^{q+1}(B; G) \longrightarrow H^{q+1}((C \cup D) \cap B; G) \longrightarrow \dots$$

where  $G = \pi_q(W_{n-1,1})$ , it follows:

$$H^{q+1}(B, (C \cup D) \cap B; G) \cong H^{q+1}(B; G) \cong 0$$

because  $H^{q+1}((C \cup D) \cap B; G) = 0 = H^{q+1}(B; G)$  for each  $q \geq n \geq 3$  (see [2] and [12]).

When  $X$  is an open set of  $\mathbb{C}^n$  we can prove something more precise:

**THEOREM 1':** *Let  $X$  be a Stein open set of  $\mathbb{C}^n$  ( $n \geq 3$ ),  $A$  a Runge and Stein open set of  $X$  and  $C$  a curve of  $A$ .*

If the set:

$$S = \{x \in C : C \text{ is not complete intersection at } x\}$$

is finite, then for each  $x \in S$  there exists a finite family of lines  $L_{x,0}, \dots, L_{x,s_x}$  through 0 such that the curve:

$$\left( (C \cup \bigcup_{\substack{x \in S \\ i=1, \dots, s_x}} L_{x,i}) \cap A \right)$$

is a set-theoretically complete intersection in  $A$ .

More precisely there exist functions  $g_1, \dots, g_{n-1}$  holomorphic on  $A$  such that:

$$(1) \{x \in A; g_1(x) = \dots = g_{n-1}(x) = 0\} = (C \cup \bigcup_{\substack{x \in S \\ i=1, \dots, s_x}} L_{x,i}) \cap A$$

(2) the germs  $g_{1,x}, \dots, g_{n-1,x}$  generate the stalk  $\mathcal{T}_{C,x}$  for each  $x \in C - S$ .

PROOF: As in the theorem 1 forgetting about  $B$  or  $\bar{B}$  and using as maps  $R_r : X \rightarrow \mathbb{C}^n$  the translations sending the points  $x_r$  in 0.

THEOREM 2: Let  $X$  be a Stein manifold of dimension  $n \geq 2$ ,  $A$  a Runge and Stein open set of  $X$  and  $C$  a curve of  $A$ .

For each relatively compact open set  $B$  of  $A$  there exist a holomorphic map  $g : B \rightarrow \mathbb{C}^{n-1}$  and a positive 1-cycle  $D \in Z_1^+(X)$  such that:

$$V_1(g) = C|_B + D|_B.$$

PROOF: Let's prove first the theorem when  $n \geq 3$ ; enlarging  $B$  we can suppose it Runge and Stein in  $A$ . For the Theorem 1 there exist a map  $g : B \rightarrow \mathbb{C}^{n-1}$  and a curve  $D$  of  $X$  such that:

$$(1) \{x \in B : G(x) = 0\} = (C \cup D) \cap B$$

$$(2) g_{1,x}, \dots, g_{n-1,x} \text{ generate } \mathcal{T}_{C,x} \text{ for each } x \in \text{reg}(C) \cap B.$$

Let's denote by  $D$  the sum of the components of the cycle  $V_1(g)$  not contained in  $C$ ;  $D$  is a cycle of  $X$  and we have:

$$V_1(g) = m_1 \cdot (C|_B) + \dots + m_r \cdot (C_r|_B) + D|_B$$

where  $C_1, \dots, C_r$  are curves contained in  $C \cap B$  decomposing it in its irreducible components, and  $m_1, \dots, m_r$  are positive integers.

We have just to prove that  $m_1 = \dots = m_r = 1$ ; let  $i = 1, \dots, r$  and  $x_i \in \text{reg}(C_i) \cap B$ , at  $x_i$  we can find a coordinate system  $(z_1, \dots, z_n)$  such

that  $g_1 = z_1, \dots, g_{n-1} = z_{n-1}$ ; in this coordinate system  $V_1(g)$  is the  $n$ th axis counted only once.

If  $n = 2$ , enlarging in case the open set  $B$  we can suppose it Runge and Stein in  $X$  and with smooth boundary. Therefore (see [2]) we have  $H_3(X, B; \mathbb{Z}) = 0$  and the group  $H_2(X, B; \mathbb{Z})$  is free of finite rank; then the restriction map:

$$r: H^2(X; \mathbb{Z}) \longrightarrow H^2(B; \mathbb{Z})$$

is surjective.

Therefore there exists a positive divisor  $D$  of  $X$  such that  $r(c(D)) = -c(C|_B)$ , that is  $c(D|_B + C|_B) = 0$ .

Since the divisor has Chern class zero, there exist a holomorphic map  $g: B \rightarrow \mathbb{C}$  such that:  $V_1(g) = C|_B + D|_B$ .

### §3. Approximation of curves

**THEOREM 3:** *Let  $X$  be a Stein manifold of dimension  $n \geq 2$ ,  $A$  a Runge and Stein open set of  $X$  and  $C$  an irreducible curve of  $A$ .*

*There exists a sequence of irreducible curves  $\{C_k\}_{k \geq 1}$  such that:*

$$\lim_{k \rightarrow \infty} (C_k \cap A) = C$$

*in the space of positive 1-cycles  $Z_1^+(A)$ .*

**PROOF:** Let  $\{B_i\}_{i \geq 1}$  be a sequence of relatively compact open sets of  $A$  which are Runge and Stein and invade  $A$ .

For each  $i \geq 1$  for the Theorem 3 we can find irreducible curves  $D_{i1}, \dots, D_{is_i}$  of  $X$  and a map  $g: B_i \rightarrow \mathbb{C}^{n-1}$  such that:

$$V_1(g_i) = (C \cap B_i) + m_{i1} \cdot (D_{i1} \cap B_i) + \dots + m_{is_i} \cdot (D_{is_i} \cap B_i).$$

Let's write  $\mathcal{F}_i = (\mathcal{F}_{D_{i1}})^{m_{i1}} \cap \dots \cap (\mathcal{F}_{D_{is_i}})^{m_{is_i}}$ , since  $g_i \in [\Gamma(B_i, \mathcal{F}_{i|_{B_i}})]^{n-1}$  for theorem 11 at page 241 of [9] there exists a sequence of maps  $\{g_i^{(k)}\}_{k \geq 1} \subset [\Gamma(X, \mathcal{F}_i)]^{n-1}$  converging to  $g_i$  on  $B_i$ ; therefore for the prop. 7 of [5] we have:

$$V_1(g_i) = \lim_{k \rightarrow \infty} (V_1(g_i^{(k)})|_{B_i}).$$

Let's denote by  $T_{ik}$  the sum of the terms of  $V_1(g_i^{(k)})$  whose support

is not in  $D_i = D_{i_1} \cup \dots \cup D_{i_{s_i}}$ ; we can write:

$$V_1(g_i^{(k)}) = T_{ik} + m_{i_1}^{(k)} \cdot D_{i_1} + \dots + m_{i_{s_i}} \cdot D_{i_{s_i}}$$

where  $m_{ij}^{(k)} \geq m_{ij}$  for each  $k \geq 0$  and  $j = 1, \dots, s_i$ .

Let's fix a point  $x_{ij} \in \text{reg}(D_{ij}) \cap B_i$  and choose in a neighborhood of  $x_{ij}$  a coordinate system where  $D_{ij}$  is the first coordinate axis; let's call  $R$  and  $L$  respectively a cube of center  $x_{ij}$  and  $L$  the normal hyperplane to  $D_{ij}$  in  $x_{ij}$ ; for the Bochner–Martinelli formula (see [10]) we have:

$$m_{ij} = \int_{L \cap \partial R} \frac{\lambda(g_i)}{|g_i|^{4n+2}} \quad \text{and} \quad m_{ij}^{(k)} \leq \int_{L \cap \partial R} \frac{\lambda(g_i^{(k)})}{|g_i^{(k)}|^{4n+2}}$$

for  $k$  big enough, where  $\lambda(g)$  is a form whose coefficients are polynomials in  $g$  and its derivatives.

For the integral continuity for  $k$  big enough we have  $m_{ij} \geq m_{ij}^{(k)}$ .

Therefore:

$$V_1(g_i^{(k)}) = T_{ik} + m_{i_1} \cdot D_{i_1} + \dots + m_{i_{s_i}} \cdot D_{i_{s_i}}$$

and then subtracting the common terms between  $V_1(g_i^{(k)})$  and  $V_1(g_i)$ :

$$C \cap B_i = \lim_{k \rightarrow \infty} (T_{ik|B_i}).$$

For the convergence is a local property (see [5]) we have:

$$C = \lim_{i \rightarrow \infty} (T_{ii}).$$

To complete the proof we need only to prove the following:

LEMMA: *Let  $X$  be a manifold of dimension  $n \geq 2$ ,  $A$  an open set of  $X$  and  $C$  an irreducible curve of  $A$ .*

*If there exists a sequence of 1-cycles  $\{T_k\}_{k \geq 1} \subset Z_1^+(X)$  such that:*

$$C = \lim_{k \rightarrow \infty} (T_k|_A)$$

*then there exists a sequence of irreducible curves  $\{C_k\}_{k \geq 1}$  of  $X$  such that:*

$$C = \lim(C_k \cap A).$$

LEMMA'S PROOF: It's enough to prove the lemma for each relatively compact open set  $B$  of  $A$ .

Let  $x$  be a regular point of  $C$ , we can find a coordinate system in a neighborhood of  $x$  making  $C$  a line; let  $P_x$  be a polycylinder with center  $x$  in this coordinate system. For  $k$  big enough the analytic set  $(\text{supp}(T_k)) \cap P_x$  is regular because each normal plane to  $C$  meets, in  $P_x$ , the space  $\text{supp}(T_k)$  in a simple point for the Bochner–Martinelli formula; moreover  $(\text{supp}(T_k)) \cap P_x$  is a connected manifold and there exists an irreducible curve  $C_{kx}$  of  $X$  such that  $T_{k|P_x} = C_{kx|P_x}$  for each  $k$  bigger than a suitable  $k_x$ .

Let's fix in  $\text{reg}(C)$  a sequence of connected compact sets invading  $\text{reg}(C)$  (such a sequence can be constructed using a triangulation of the connected smooth manifold  $\text{reg}(C)$ ); let's call  $U$  a compact neighborhood of  $\text{sing}(C) \cap B$  small enough to be contained in a Stein open set of  $B$ .

Since the set  $(B - U) \cap \text{reg}(C)$  is relatively compact in  $\text{reg}(C)$  there exists a connected compact set  $K$  of  $\text{reg}(C)$  containing the set  $(B - U) \cap \text{reg}(C)$  and it is possible to find a finite number of points  $x_1, \dots, x_m$  of  $K$  and polycylinders  $P_{x_1}, \dots, P_{x_m}$  centered in those points such that  $P = \bigcup_{i=1}^m P_{x_i} \supset K$ ; therefore we have  $C \cap B \subset P \cup U$ .

Moreover whenever  $P_{x_i} \cap P_{x_j} \neq \emptyset$  we can find a point  $x_{ij} \in P_{x_i} \cap P_{x_j}$ , a polycylinder  $P_{ij}$  centered in  $x_{ij}$  contained in  $P_{x_i} \cap P_{x_j}$  and an integer big enough  $k_{ij}$  such that  $(\text{supp}(T_k)) \cap P_{ij}$  is non-empty and irreducible for each  $k \geq k_{ij}$ .

Since  $P$  is connected for  $k \geq \bar{k} = \max\{k_{x_i}, k_{ij}\}$  the irreducible curve representing  $T_k$  in each  $P_{x_i}$  must be the same, that is there exists an irreducible curve  $C_k$  of  $X$  for each  $k \geq \bar{k}$  such that:  $T_{k|P} = C_{k|P}$ .

Moreover for  $k$  big enough we have  $(\text{supp}(T_k)) \cap B \subset (P \cap U) \cap B$  (see the Remark 5 of [5]); then  $T_{k|P \cap B} = C_{k|P \cap B}$ , that is  $T_{k|B-U} = C_{k|B-U}$  and at last  $T_{k|B} = C_{k|B}$ .

**THEOREM 4:** *Let  $X$  be an holomorphically convex open set of  $\mathbb{C}^n$  ( $n \geq 2$ ),  $A$  a Runge and holomorphically convex open set of  $X$  and  $C$  an analytic irreducible curve of  $A$ .*

*There exists a sequence of algebraic curves  $\{C_k\}_{k \geq 1}$  of  $\mathbb{C}^n$  irreducible in  $X$  such that:*

$$\lim_{k \rightarrow \infty} (C_k \cap A) = C$$

*in the space of positive analytic 1-cycles  $Z_1^+(A)$ .*

PROOF: Trivial for  $n = 2$ .

For  $n \geq 3$  following Theorem 3 let's observe that, being  $X$  an open set of  $\mathbb{C}^n$ , we can take as curves  $D_{i_1}, \dots, D_{i_{r_i}}$  some lines of  $\mathbb{C}^n$  as in Lemma 1 and therefore the section of the sheaf  $\mathcal{T}_i = (\mathcal{T}_{D_{i_1}})^{m_{i_1}} \cap \dots \cap (\mathcal{T}_{D_{i_{r_i}}})^{m_{i_{r_i}}}$  are generated by some polynomials  $p_{i_1}, \dots, p_{i_{r_i}}$  of  $\mathbb{C}^n$ ; that is for each  $j = 1, \dots, n - 1$  it holds:

$$(g_i)_j = \sum_{l=1, \dots, r_i} h_{ijl} \cdot p_{il}$$

for some functions  $h_{ijl}$  holomorphic on  $B_i$ .

Moreover we can choose the open sets  $B_i$  to be Runge in  $\mathbb{C}^n$  and then find sequences of polynomials  $\{q_{ijl}^{(k)}\}_{k \geq 1}$  of  $\mathbb{C}^n$  converging to  $h_{ijl}$  on  $B_i$ .

Denoting  $(g_i^{(k)})_j = \sum_{l=1, \dots, r_i} q_{ijl}^{(k)} \cdot p_{il}$ , the positive 1-cycles  $\{T_{ik}\}$  are algebraic and even more so the curves  $\{C_k\}_{k \geq 1}$ .

#### REFERENCES

- [1] A. ANDREOTTI and W. STOLL: Analytic and algebraic dependence of meromorphic functions. *Lecture notes 234* (1971) Springer-Verlag.
- [2] A. ANDREOTTI and R. NARASHIMAN: A topological property of Runge pairs. *Ann. of Math.* 76 (1962) 499–509.
- [3] D. BARLET: Espace analytique réduit des cycles analytiques complexes compacts d'un espace analytique. *Lecture notes 482* (1975) 1–158.
- [4] A. CASSA: A theorem on Runge problem for analytic curves. *Ann. di Matematica (IV)* Vol. CVII (1976) 169–184.
- [5] A. CASSA: The topology of the space of positive analytic cycles. (To appear in: *Annali di Matematica*).
- [6] S. COEN: Sul rango de fasci coerenti. *Boll. U.M.I. Vol. XXII* (3) (1967) 373–382.
- [7] M. CORNALBA and P. GRIFFITHS: Analytic cycles and vector bundles on non-compact algebraic varieties. *Inv. Math.* 28 (1975) 1–106.
- [8] O. FORSTER and K.J. RAMSPOTT: Analytische Modulgarben und Endromisbündel. *Inv. Math.* 2 (1966) 145–170.
- [9] R.C. GUNNING and H. ROSSI: *Analytic functions of several complex variables*. Prentice Hall Series. 1965.
- [10] J.R. KING: A residue formula for complex subvarieties. *Carolina conference proceedings* (1970) 43–56.
- [11] J.R. KING: The currents defined by analytic varieties. *Acta Math.* 127 (1971) 185–220.
- [12] R. NARASHIMAN: On the homology of Stein spaces. *Inv. Math.* 2 (1967) 377–385.
- [13] N. STEENROD: *The topology of fibre bundles*. Princeton University Press 14 (1951).

(Oblatum 2–XII–1976)

Università degli Studi  
Istituto Matematico U. Dini  
Vialé Morgagni 571A  
I-50 134 Firenze, Italia.