

COMPOSITIO MATHEMATICA

C. SOULÉ

**The rank of étale cohomology of varieties
over p -adic or number fields**

Compositio Mathematica, tome 53, n° 1 (1984), p. 113-131

http://www.numdam.org/item?id=CM_1984__53_1_113_0

© Foundation Compositio Mathematica, 1984, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

THE RANK OF ÉTALE COHOMOLOGY OF VARIETIES OVER p -ADIC OR NUMBER FIELDS

C. Soulé

Let K be a number field or its completion at a finite prime, p a rational prime, and V a smooth and proper variety over K . In this paper we study the corank of the groups $H^k(V, \mathbb{Q}_p/\mathbb{Z}_p(i))$, $k \geq 0$, $i \in \mathbb{Z}$, of étale cohomology of V with coefficients $\mathbb{Q}_p/\mathbb{Z}_p$ twisted i -times by the cyclotomic character.

If \bar{V} denotes the extension of V to an algebraic closure \bar{K} of K (imbedded in the field \mathbb{C} of complex numbers), Grothendieck showed that the group $H^k(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i))$ is equal to the ordinary (singular) cohomology group $H^k(\bar{V}(\mathbb{C}), \mathbb{Q}_p/\mathbb{Z}_p)$ of the set of complex points of \bar{V} with constant coefficients $\mathbb{Q}_p/\mathbb{Z}_p$. Therefore the computation of the cohomology of V is a problem of Galois descent, which is of arithmetic nature. We cannot solve this problem in general, but what we show is that *for all but finitely many* integers i , the corank of $H^k(V, \mathbb{Q}_p/\mathbb{Z}_p(i))$ admits a simple (“geometric”) expression.

In the global case this result was suggested by conjectures in algebraic K -theory due to D. Quillen and A. Beilinson (which are shown here to be compatible, cf. I.7.). In the local case the Hodge-Tate decomposition of the cohomology of \bar{V} ([19], [2]) tells us which values of i it is sufficient to avoid.

During this work I was helped by S. Bloch (in particular, Theorem 2(iv) is due to him), J.-L. Colliot-Thélène, K. Ribet, P. Schneider and K. Wingberg. I would like to thank them warmly.

I also thank the University of Princeton for its hospitality when the last version of this paper was written.

1. The global case

1.1. Statement of the result

1.1.1. Notations

Let p be a rational prime, and A an abelian group. We denote by $A[p]$ (resp. A/p) the kernel (resp. cokernel) of the multiplication by p in A .

Let $\text{Div}(A)$ be the maximal divisible subgroup of A . When A is a p -torsion group, we define its dimension (or corank) to be the dimension of the vector space $\text{Div}(A)[p]$ over the field \mathbb{F}_p of order p :

$$\dim A = \dim_{\mathbb{F}_p} \text{Div}(A)[p].$$

We also denote by

$$A^\wedge = \text{Hom}(A, \mathbb{Q}_p/\mathbb{Z}_p)$$

the Pontryagin dual of A .

Given a scheme X such that all its residue fields contain $1/p$ and an integer $n \geq 1$, we denote by μ_{p^n} the étale sheaf over X over p^n -th roots of unity and

$$\mathbb{Z}_p(1) = \varprojlim_n \mu_{p^n}$$

their projective limit. When M is an étale sheaf of p -torsion groups over X we define the Tate twists $M(i)$, $i \in \mathbb{Z}$, of M to be

$$M(0) = M$$

$$M(i) = M(i-1)_{\mathbb{Z}_p} \otimes \mathbb{Z}_p(1) \quad \text{when } i \geq 1,$$

$$M(i) = \text{Hom}_{\mathbb{Z}_p}(\mathbb{Z}_p(-i), M) \quad \text{when } i < 0.$$

For instance, when $i \in \mathbb{Z}$, the sheaf $\mathbb{Q}_p/\mathbb{Z}_p(i)$ is the inductive limit of the finite sheaves $(\mathbb{Z}/p^n)(i)$, the transition maps being induced by the standard inclusions $\mathbb{Z}/p^n \rightarrow \mathbb{Z}/p^{n+1}$. The fact that inductive limits preserve exact sequences leads us to take $\mathbb{Q}_p/\mathbb{Z}_p(i)$ as coefficients (instead of $\mathbb{Z}_p(i)$) in the Theorem 1 below.

1.1.2. Let K be a number field, \bar{K} an algebraic closure of K , G the Galois group $G = \text{Gal}(\bar{K}/K)$. Let V be a proper smooth variety defined over K , and $\bar{V} = V \otimes_K \bar{K}$ the variety over \bar{K} obtained by extending scalars.

For any archimedean prime v of K , let K_v be the completion of K at v , and choose an imbedding $v: K_v \rightarrow \mathbb{C}$ of K_v into the complex numbers. Let $V_v = V \otimes_K K_v$ and $V_v(\mathbb{C})$ be the set of complex points of the K_v -variety V_v . For any integer $k \geq 1$, we denote by $H^{k-1}(V_v(\mathbb{C}), \mathbb{R})$ the group of ordinary (singular) cohomology of the topological space $V_v(\mathbb{C})$ with real coefficients. The Galois group $\text{Gal}(\mathbb{C}/K_v)$ acts upon $H^{k-1}(V_v(\mathbb{C}), \mathbb{R})$ and for any integer $i \in \mathbb{Z}$ we define $H^{k-1}(V_v(\mathbb{C}), \mathbb{R})^{(i-1)}$ to be $H^{k-1}(V_v(\mathbb{C}), \mathbb{R})$ when $K_v = \mathbb{C}$, and the eigenspace of eigenvalue $(-1)^{(i-1)}$ of the generator of $\text{Gal}(\mathbb{C}/K_v)$ when $K_v \simeq \mathbb{R}$.

If S_∞ is the set of archimedean primes of K , $\ell \geq 1$ and $i \in \mathbb{Z}$, we define

$$n_{i,\ell} = \sum_{v \in S_\infty} \dim_{\mathbb{R}} H^{\ell-1}(V_v(\mathbb{C}), \mathbb{R})^{(i-1)}.$$

On the other hand we consider the groups of étale cohomology

$$H^\ell(V, \mathbb{Q}_p/\mathbb{Z}(i)) = \varinjlim_n H^\ell(V, \mathbb{Z}/p^n(i))$$

defined in 1.1.1.

THEOREM 1: *Let V be as above, $\ell \geq 1$ an integer, and p a rational prime.*

- (i) *For any integer $i \in \mathbb{Z}$ we have $\dim H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)) \geq n_{i,\ell}$.*
- (ii) *For almost all integer $i \in \mathbb{Z}$ we have $\dim H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)) = n_{i,\ell}$.*

1.2. To prove Theorem 1 we first show that we can assume $K = \mathbb{Q}$. For let $\pi: V \rightarrow \text{Spec } K$ be the morphism of schemes defining V and \tilde{V} the variety over \mathbb{Q} obtained by composing π with the map

$$\text{Spec } K \rightarrow \text{Spec } \mathbb{Q}.$$

Since the étale cohomology does not depend on the field of definition we have

$$H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)) = H^\ell(\tilde{V}, \mathbb{Q}_p/\mathbb{Z}_p(i)).$$

On the other hand the numbers $n_{i,\ell}$ are the same for V/K and \tilde{V}/\mathbb{Q} . To see this, for any imbedding $\sigma: K \rightarrow \mathbb{C}$, denote by $V \otimes_\sigma \mathbb{C}$ the complex variety obtained from V by extending scalars to \mathbb{C} through σ . We have

$$\tilde{V} \otimes_{\mathbb{Q}} \mathbb{C} = V \otimes_K \left(K \otimes_{\mathbb{Q}} \mathbb{C} \right) \simeq \prod_{\sigma} \left(V \otimes_{\sigma} \mathbb{C} \right)$$

where the product is taken over all imbeddings of K into \mathbb{C} . A complex valuation v of K gives two imbeddings σ_v and $c\sigma_v$ of K_v into \mathbb{C} (where c denotes the complex conjugation) and complex conjugation acting upon $\tilde{V} \otimes_{\mathbb{Q}} \mathbb{C}$ permutes the two factors $V \otimes_{\sigma_v} \mathbb{C}$ and $V \otimes_{c\sigma_v} \mathbb{C}$. When v is a real valuation, there is only one imbedding $\sigma_v: K_v \rightarrow \mathbb{C}$ attached to it, and the complex conjugation acting upon $\tilde{V} \otimes_{\mathbb{Q}} \mathbb{C}$ induces the action of $\text{Gal}(\mathbb{C}/K_v)$ on the factor $V \otimes_{\sigma_v} \mathbb{C}$. Therefore, for any $i \in \mathbb{Z}$, we have

$$H^{\ell-1}(\tilde{V}(\mathbb{C}), \mathbb{R})^{(i-1)} \simeq \sum_{v \in S_\infty} H^{\ell-1}(V_v(\mathbb{C}), \mathbb{R})^{(i-1)}.$$

1.3. From now on we will assume $K = \mathbb{Q}$. Let S be a finite set of rational primes containing p and such that V has good reduction outside S . That is to say, if we call \mathbb{Z}_S the ring of rational numbers which are units outside S , there exist a smooth proper scheme \mathcal{V} over $\text{Spec}(\mathbb{Z}_S)$ with generic fiber $\mathcal{V} \otimes_{\mathbb{Z}_S} \mathbb{Q}$.

We fix an integer $\ell \geq 1$ and let $M = H^{\ell-1}(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p)$. The module M is a discrete module under the Galois group $G = \text{Gal}(K/K)$, and we can also consider M as an étale sheaf over $\text{Spec}(\mathbb{Q})$ [5].

Let $j: \text{Spec}(\mathbb{Q}) \rightarrow \text{Spec}(\mathbb{Z}_S)$ be the canonical inclusion and let $j_* M$ be the direct image of M .

PROPOSITION 1:

- (i) For any $i \in \mathbb{Z}$ we have $\dim H^r(\text{Spec } \mathbb{Q}, M(i)) = 0$ when $r \geq 3$ and $\dim H^1(\text{Spec } \mathbb{Q}, M(i)) \geq \dim H^1(\text{Spec } \mathbb{Z}_S, j_* M(i))$.
- (ii) Let $i \in \mathbb{Z}$ be an integer such that $\ell \neq 2i - 1$ and $H^2(\text{Spec } \mathbb{Z}_S, j_* M(i))$ is finite. Then for any integer $r \geq 0$ we have

$$\dim H^r(\text{Spec } \mathbb{Q}, M(i)) = \dim H^r(\text{Spec } \mathbb{Z}_S, j_* M(i)).$$

PROOF: It follows from [21], Theorem 3.1 c) that $\dim H^r(\text{Spec } \mathbb{Q}, M(i)) = 0$ when $r \geq 3$. Let M_n be the finite module $H^{\ell-1}(\bar{V}, \mathbb{Z}/p^n)$, so that

$$H^\ell(\text{Spec } \mathbb{Q}, M(i)) = \varinjlim_n H^\ell(\text{Spec } \mathbb{Q}, M_n(i)) \quad \text{and}$$

$$H^\ell(\text{Spec } \mathbb{Z}_S, M(i)) = \varinjlim_n H^\ell(\text{Spec } \mathbb{Z}_S, j_* M_n(i)).$$

The Leray spectral sequence for the functor j_* and the sheaf $M_n(i)$ has E_2 -term

$$E_2^{rs} = H^r(\text{Spec } \mathbb{Z}_S, R^s j_* M_n(i))$$

and converges to $H^{r+s}(\text{Spec } \mathbb{Q}, M_n(i))$.

For any prime l not in S , let \mathbb{F}_l be the field with l elements and

$$i_l: \text{Spec } \mathbb{F}_l \rightarrow \text{Spec } \mathbb{Z}_S$$

the canonical inclusion. When $s \geq 1$ the morphism of étale sheaves over $\text{Spec } \mathbb{Z}_S$

$$R^s j_* M_n(i) \rightarrow \bigoplus_{l \notin S} i_{l*} i_l^* R^s j_* M_n(i)$$

is an isomorphism, as can be seen by looking at the geometric fibers.

Since i_l is a finite morphism we get ([5], II.3., Proposition 3.6.) $H^r(\text{Spec } \mathbb{Z}_S, i_{l*} i_l^* R^s j_* M_n(i)) \simeq H^r(\text{Spec } \mathbb{F}_l, i_l^* R^s j_* M_n(i))$. Let \mathbb{Q}_l be the field of l -adic numbers and \mathbb{Q}_l^{nr} its maximal unramified extension. Then the sheaf $i_l^* R^s j_* M_n(i)$, considered as a $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ -module, is isomorphic to $H^s(\mathbb{Q}_l^{nr}, M_n(i))$, where $\text{Gal}(\overline{\mathbb{Q}}_l/\mathbb{Q}_l^{nr})$ acts upon M_n through its inclusion in $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

The smooth and proper base change theorem ([5], .3., Theorem 3) asserts that the action of the inertia group $\text{Gal}(\overline{\mathbb{Q}}_l/\mathbb{Q}_l^{nr})$ on M_n is trivial, and that there is an isomorphism of $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ -modules between M_n and $H^{\ell-1}(\overline{W}_l, \mathbb{Z}/p^n)$, where $W_l = \mathcal{V} \otimes_{\mathbb{Z}_S} \mathbb{F}_l$ is the reduction of \mathcal{V} modulo l and $\overline{W}_l = W_l \otimes_{\mathbb{Z}_S} \overline{\mathbb{F}}_l$ is the extension of W_l to an algebraic closure of \mathbb{F}_l . Therefore we get isomorphisms of $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ -modules

$$\begin{aligned} H^s(\mathbb{Q}_l^{nr}, M_n(i)) &\simeq H^s(\mathbb{Q}_l^{nr}, \mu_{p^n}) \otimes M_n(i-1) \\ &= \begin{cases} H^{\ell-1}(\overline{W}_l, \mathbb{Z}/p^n)(i-1) & \text{when } s=1 \\ 0 & \text{when } s>1 \end{cases} \end{aligned}$$

(compare [16], III.1.3.).

Since $cd_p(\mathbb{F}_l) = 1$, the Leray spectral sequence for j_* and $M_n(i)$ gives an exact sequence

$$\begin{aligned} 0 \rightarrow H^1(\text{Spec } \mathbb{Z}_S, j_* M_n(i)) &\rightarrow H^1(\text{Spec } \mathbb{Q}, M_n(i)) \\ &\rightarrow \bigoplus_{l \notin S} H^0(\mathbb{F}_l, H^{\ell-1}(\overline{W}_l, \mathbb{Z}/p^n)(i-1)) \rightarrow H^2(\text{Spec } \mathbb{Z}_S, j_* M_n(i)) \\ &\rightarrow H^2(\text{Spec } \mathbb{Q}, M_n(i)) \rightarrow \bigoplus_{l \notin S} H^1(\mathbb{F}_l, H^{\ell-1}(\overline{W}_l, \mathbb{Z}/p^n)(i-1)). \end{aligned}$$

Taking the inductive limits over n of these exact sequences we see that to get Proposition 1 it will be enough to show the following

LEMMA 1: *When $2i \neq \ell + 1$ the groups*

$$\begin{aligned} H^0(\mathbb{F}_l, H^{\ell-1}(\overline{W}_l, \mathbb{Q}_p/\mathbb{Z}_p)(i-1)) \quad \text{and} \\ H^1(\mathbb{F}_l, H^{\ell-1}(\overline{W}_l, \mathbb{Q}_p/\mathbb{Z}_p)(i-1)) \end{aligned}$$

are finite.

PROOF OF LEMMA 1: Since $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ is isomorphic to the profinite completion of \mathbb{Z} , when T is a torsion discrete $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ -module, the group $H^0(\mathbb{F}_l, T)$ is equal to the group of invariants $T^{\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)}$ of T by $\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$, and $H^1(\mathbb{F}_l, T)$ is equal to the group of coinvariants $T_{\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)}$.

Since the cohomology of \overline{W}_l with \mathbb{Z}_p -coefficients is finitely generated, we have

$$\text{Div } H^{\ell-1}(\overline{W}_l, \mathbb{Q}_p/\mathbb{Z}_p) = H^{\ell-1}(\overline{W}_l, \mathbb{Z}_p) \otimes \mathbb{Q}_p/\mathbb{Z}_p$$

where $H^{\ell-1}(\overline{W}_l, \mathbb{Z}_p) = \varprojlim_n H^{\ell-1}(\overline{W}_l, \mathbb{Z}/p^n)$. Let $\phi \in \text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$ be the arithmetic Frobenius. It will be enough to show that the endomorphism of $H^{\ell-1}(\overline{W}_l, \mathbb{Q}_p(i-1)) = H^{\ell-1}(\overline{W}_l, \mathbb{Z}_p) \otimes \mathbb{Q}_p(i-1)$ induced by ϕ has no fixed vector. But the Weil conjectures proved by Deligne [4] tell us that any eigenvalue λ of ϕ acting upon $H^{\ell-1}(\overline{W}_l, \mathbb{Q}(i-1))$ is an algebraic number whose archimedean absolute values are equal to $l^{(-\ell+1+2(i-1))/2}$. Therefore λ is different from 1 when $2i \neq \ell + 1$. Q.E.D.

1.4. PROPOSITION 2: *Let M be as above. Then, for any $i \in \mathbb{Z}$,*

$$\sum_{r=0}^2 (-1)^r \dim H^r(\text{Spec } \mathbb{Z}_S, j_* M(i)) = -n_{i,\ell}$$

PROOF: We shall extend to our situation the arguments in [14], 4, Theorem 6. Let $\chi = \sum_{r=0}^2 (-1)^r \dim H^r(\text{Spec } \mathbb{Z}_S, j_* M(i))$. The module M is the direct sum of its divisible subgroup $A = \text{Div}(M)$ with a finite group, and the cohomology groups of $\text{Spec } \mathbb{Z}_S$ with finite coefficients are finite, therefore we have

$$\chi = \sum_{r=0}^2 (-1)^r \dim H^r(\text{Spec } \mathbb{Z}_S, j_* A[i]).$$

Let $\mathbb{Q}_S \subset \overline{\mathbb{Q}}$ be the maximal extension of \mathbb{Q} which is unramified outside S (and infinity), and $G_S = \text{Gal}(\mathbb{Q}_S/\mathbb{Q})$ its Galois group over \mathbb{Q} . Then, given any finite $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -module F , we have

$$H^r(\text{Spec } \mathbb{Z}_S, j_* F) \simeq H^r(G_S, F^{\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}_S)}).$$

As we saw when proving Proposition 1, the module $j_* A(i)$ is unramified outside S , therefore

$$H^r(\text{Spec } \mathbb{Z}_S, j_* A(i)) \simeq H^r(G_S, A(i)).$$

Now, the exact sequence of coefficients

$$0 \rightarrow A(i)[p] \rightarrow A(i) \xrightarrow{\times p} A(i) \rightarrow 0$$

gives a long exact sequence of cohomology groups:

$$\begin{aligned}
0 &\rightarrow H^0(G_S, A(i)[p]) \rightarrow H^0(G_S, A(i)) \xrightarrow{\times p} H^0(G_S, A(i)) \\
&\rightarrow H^1(G_S, A(i)[p]) \rightarrow \dots \rightarrow H^2(G_S, A(i)) \xrightarrow{\times p} H^2(G_S, A(i)) \\
&\rightarrow H^2(G_S, A(i))/p \rightarrow 0.
\end{aligned}$$

From this we deduce:

$$\chi = \sum_{r=0}^2 (-1)^r \dim_{\mathbb{F}_p} H^r(G_S, A(i)[p]) - \dim_{\mathbb{F}_p} H^2(G_S, A(i))/p.$$

By a theorem of Tate ([20], Theorem 2) we get

$$\begin{aligned}
&\sum_{r=0}^2 (-1)^r \dim_{\mathbb{F}_p} H^r(G_S, A(i)[p]) \\
&= \dim_{\mathbb{F}_p} H^0(\mathbb{R}, A(i)[p]) - \dim_{\mathbb{F}_p} A(i)[p].
\end{aligned}$$

Furthermore, since the maps $H^3(G_S, A(i)[p]) \rightarrow H^3(\mathbb{R}, A(i)[p])$ and $H^3(G_S, A(i)) \rightarrow H^3(\mathbb{R}, A(i))$ are isomorphisms ([21], Thm. 3.1.c) we see that the map

$$H^2(G_S, A(i))/p \rightarrow H^2(\mathbb{R}, A(i))/p = H^2(\mathbb{R}, A(i))$$

is an isomorphism. So we get

$$\begin{aligned}
\chi &= \dim_{\mathbb{F}_p} H^0(\mathbb{R}, A(i)[p]) - \dim A - \dim_{\mathbb{F}_p} H^2(\mathbb{R}, A(i)) \\
&= \dim H^0(\mathbb{R}, A(i)) + \dim_{\mathbb{F}_p} H^0(\mathbb{R}, A(i))/p \\
&\quad - \dim A - \dim_{\mathbb{F}_p} H^2(\mathbb{R}, A(i)).
\end{aligned}$$

Since $\text{Gal}(\mathbb{C}/\mathbb{R}) = \mathbb{Z}/2$, the numbers

$$\begin{aligned}
&\dim_{\mathbb{F}_p} H^1(\mathbb{R}, A(i)[p]) \\
&= \dim_{\mathbb{F}_p} H^0(\mathbb{R}, A(i))/p + \dim_{\mathbb{F}_p} H^1(\mathbb{R}, A(i))
\end{aligned}$$

and

$$\dim_{\mathbb{F}_p} H^2(\mathbb{R}, A(i)[p]) = \dim_{\mathbb{F}_p} H^1(\mathbb{R}, A(i)) + \dim_{\mathbb{F}_p} H^2(\mathbb{R}, A(i))$$

are equal. Therefore we get

$$\chi = \dim H^0(\mathbb{R}, A(i)) - \dim A = -\dim H^0(\mathbb{R}, A(i-1)).$$

On the other hand, we have isomorphism of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -modules ([5], V, Cor. 3.3., and IV, Thm. 6.3.):

$$\begin{aligned} A &= \text{Div } H^{\ell-1}(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p) = \text{Div } H^{\ell-1}\left(V \otimes_{\mathbb{Q}} \mathbb{C}, \mathbb{Q}_p/\mathbb{Z}_p\right) \\ &= \text{Div } H^{\ell-1}(V(\mathbb{C}), \mathbb{Q}_p/\mathbb{Z}_p) = H^{\ell-1}(V(\mathbb{C}), \mathbb{Z}) \otimes \mathbb{Q}_p/\mathbb{Z}_p \end{aligned}$$

(since the groups of ordinary cohomology of $V(\mathbb{C})$ with constant coefficients \mathbb{Z} are finitely generated). So we have

$$\begin{aligned} \chi &= -\dim H^0(\text{Gal}(\mathbb{C}/\mathbb{R}), H^{\ell-1}(V(\mathbb{C}), \mathbb{Z})(i-1) \otimes \mathbb{Q}_p/\mathbb{Z}_p) \\ &= -\dim_{\mathbb{R}} H^0(\text{Gal}(\mathbb{C}/\mathbb{R}), H^{\ell-1}(V(\mathbb{C}), \mathbb{R})(i-1)) = -n_{i,\ell}. \end{aligned}$$

Q.E.D.

1.5. PROPOSITION 3: *For all but finitely many $i \in \mathbb{Z}$ the groups $H^0(\text{Spec } \mathbb{Z}_S, j_* M(i))$ and $H^2(\text{Spec } \mathbb{Z}_S, j_* M(i))$ are finite.*

PROOF:

(a) Let l be a rational prime outside S . As we saw in Proposition 1

$$\begin{aligned} H^0(\text{Spec } \mathbb{Z}_S, j_* M(i)) &= H^0(\text{Spec } \mathbb{Q}, M(i)) \subset H^0(\mathbb{Q}_l, M(i)) \\ &= H^0(\mathbb{F}_l, H^{\ell-1}(\bar{W}_l, \mathbb{Q}_p/\mathbb{Z}_p(i))), \end{aligned}$$

and these groups are finite when $2i \neq \ell + 1$.

(b) Let $\mathbb{Q}(\mu_{p^n}) \subset \bar{\mathbb{Q}}$ be the cyclotomic extension of \mathbb{Q} obtained by adding the p^n -th roots of unity, $n \geq 1$, and $\mathbb{Q}(\mu_{p^\infty}) = \bigcup_{n \geq 1} \mathbb{Q}(\mu_{p^n})$ be the maximal p -cyclotomic extension of \mathbb{Q} . The extension $\mathbb{Q}(\mu_{p^n})$ of \mathbb{Q} is unramified outside p , therefore $\mathbb{Q}(\mu_{p^\infty})$ is contained in \mathbb{Q}_S (since $p \in S$). Let

$$H = \text{Gal}(\mathbb{Q}_S/\mathbb{Q}_\infty), \Delta = \text{Gal}(\mathbb{Q}(\mu_{p^2})/\mathbb{Q}),$$

$$\Gamma = \text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}(\mu_{p^2}))$$

and $G'_S = \text{Gal}(\mathbb{Q}_S/\mathbb{Q}(\mu_{p^2}))$.

The order of Δ is $p(p-1)$. The composite of the corestriction map

$$H^2(G_S, M(i)) \rightarrow H^2(G'_S, M(i))$$

with the transfer map

$$H^2(G'_S, M(i)) \rightarrow H^2(G_S, M(i))$$

is the product by the cardinal of Δ . Therefore the kernel of

$$H^2(G_S, M(i)) \rightarrow H^2(G'_S, M(i))$$

is contained in $H^2(G_S, M(i))[p]$, which is a finite group. In fact it is a quotient of

$$H^2(G_S, (M/A)(i)[p]) \oplus H^2(G_S, A(i))[p],$$

where $A = \text{Div } M$, and the cohomology groups of G_S with finite coefficients are finite. Therefore, to prove that the groups

$$H^2(\text{Spec } \mathbf{Z}_S, j_* M(i)) = H^2(G_S, M(i))$$

are finite for almost all i , it will be enough to show that $H^2(G'_S, M(i))$ is finite (for almost all i).

For this let us consider the extension of groups

$$1 \rightarrow H \rightarrow G'_S \rightarrow \Gamma \rightarrow 1.$$

The Hochschild-Serre spectral sequence deduced from it has E_2 -term

$$E_2^{rs} = H^r(\Gamma, H^s(H, M(i)))$$

and converges to $H^{r+s}(G'_S, M(i))$.

The cyclotomic character $\kappa: \text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}) \rightarrow \mathbf{Z}_p^*$ is an isomorphism and induces an isomorphism between Γ and the multiplicative group $1 + p^2\mathbf{Z}_p$, i.e. with the additive group \mathbf{Z}_p . Therefore $cd_p \Gamma = 1$. On the other hand Ferrero and Washington proved that the μ -invariant of $\mathbb{Q}(\mu_{p^\infty})$ is zero [6], and Iwasawa deduced from this that $cd_p H_p = 1$, where H_p is the maximal pro- p quotient group of H ([8], Theorem 2). Therefore $cd_p H = 1$ and $E_2^{rs} = 0$ when $r \geq 2$ or $s \geq 2$. So we have

$$H^2(G'_S, M(i)) = H^1(\Gamma, H^1(H, M(i))) = H^1(H, M)(i)_\Gamma$$

(since the action of H on the roots of unity is trivial).

Let $X = H^1(H, M)^\wedge$ be the Pontryagin dual of $H^1(H, M)$. We want to show that $X(-i)^\Gamma$ is finite for almost all i . Call

$$\Lambda = \mathbf{Z}_p[[\Gamma]] = \varprojlim_n \mathbf{Z}_p[\Gamma/\Gamma^{p^n}]$$

the \mathbb{Z}_p -algebra of the pro- p -group Γ . Choosing a generator γ_0 of Γ gives an isomorphism between Λ and the ring of powers series $\mathbb{Z}_p[[T]]$, which sends γ_0 to $1 + T$. The module X is a Λ -module.

We first notice that X is a noetherian Λ -module. In fact Λ is a local domain, compact for the topology defined by its maximal ideal (p, T) , and X is a compact Λ -module therefore ([7], 1.1.) it is enough, by Nakayama's lemma, to show that $X_\Gamma \otimes \mathbb{Z}/p$ is finite. But

$$(X_\Gamma \otimes \mathbb{Z}/p)^\wedge = \hat{X}^\Gamma[p] = H^1(H, M)^\Gamma[p].$$

Since M is the direct sum of its divisible subgroup $\text{Div}(M)$ with a finite group, we have to show that $H^1(H, \text{Div } M)^\Gamma[p]$ is finite; but this is a quotient of $H^1(G'_S, M[p])$ (by the Hochschild-Serre spectral sequence), therefore it is finite.

The Proposition 3 will then be a consequence of the following

LEMMA 2: *Let X be a noetherian Λ -module. Then $X(-i)^\Gamma$ is finite for almost all integers $i \in \mathbb{Z}$.*

PROOF: It was proved by Iwasawa [7] that X is pseudo-isomorphic to a finite product of Λ -modules of the type $\Lambda/(f(T))$, where $f(T)$ is a polynomial in $\mathbb{Z}_p[T]$. Let $c = \kappa(\gamma_0) \in \mathbb{Z}_p^*$ be the image of γ_0 by the cyclotomic character. Then, when twisting the Λ -module $\Lambda/(f(T))$, we get

$$(\Lambda/(f(t)))(-i) \simeq \Lambda/(f(c^i(1 + T) - 1)).$$

The euclidean algorithm shows that $(\Lambda/(f(T)))(-i)^\Gamma$ is finite whenever $f = 0$ or $f(c^i - 1) \neq 0$ (cf. [12], Lemma 4.2). A nonzero polynomial having only finitely many roots, the Lemma 2 follows.

1.6. PROOF OF THEOREM 1: The Hochschild-Serre spectral sequence attached to the Galois covering $\bar{V} \rightarrow V$ has E_2 -term

$$E_2^{r,s} = H^r(\mathbb{Q}, H^s(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i)))$$

and converges to $H^{r+s}(V, \mathbb{Q}_p/\mathbb{Z}_p(i))$. From Proposition 1 and 3 we know that, for almost all i , we have

$$\begin{aligned} \dim H^r(\mathbb{Q}, H^s(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i))) \\ = \dim H^r(\text{Spec } \mathbb{Z}_S, j_* H^s(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i))) \end{aligned}$$

and that this dimension is zero for $r \neq 1$. From Proposition 2 we get

$$\dim H^1(\mathbb{Q}, H^{\ell-1}(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i))) = n_{i,\ell}.$$

Therefore, for almost all $i \in \mathbb{Z}$,

$$\begin{aligned} \dim H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)) &= \dim E_2^{1,\ell-1} \\ &= \dim H^1(\mathbb{Q}, H^{\ell-1}(\bar{V}, \mathbb{Q}_p/\mathbb{Z}_p(i))) = n_{i,\ell}. \end{aligned}$$

This proves (ii). To prove (i) we notice that for any $i \in \mathbb{Z}$ we have

$$\begin{aligned} \dim H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)) &\geq \dim E_2^{1,\ell-1} \\ &\geq \dim H^1(\text{Spec } \mathbb{Z}_S, j_* M(i)) \geq n_{i,\ell}. \end{aligned}$$

1.7. Connection with algebraic K-theory

The statement of Theorem 1 was motivated by the following conjectures in algebraic K-theory.

Let V be as in Theorem 1, let $K_m(V)$, $m \geq 0$, be the higher algebraic K-groups of the variety V [13], let $K_m(V; \mathbb{Z}/p^n)$ be the groups of algebraic K-theory with coefficients \mathbb{Z}/p^n of V [3], and let

$$K_m(V; \mathbb{Q}_p/\mathbb{Z}_p) = \varinjlim_n K_m(V; \mathbb{Z}/p^n).$$

Let $Gr_\gamma^i K_m(V; \mathbb{Q}_p/\mathbb{Z}_p)$ be the i -th quotient of the γ -filtration of the K-theory of V [11].

The theory of p -adic Chern classes [16] gives morphisms

$$c_{i,\ell}: Gr_\gamma^i K_{2i-\ell}(V, \mathbb{Q}_p/\mathbb{Z}_p) \rightarrow H^\ell(V, \mathbb{Q}_p/\mathbb{Z}_p(i)), \quad i \geq 1,$$

and Quillen's conjecture [13] would imply that the kernel and the cokernel of $c_{i,\ell}$ have dimension zero when i is big enough.

On the other hand Beilinson [1] and Karoubi [9] defined transcendental Chern classes

$$\rho_{i,\ell}: \mathbb{R} \otimes_{\mathbb{Z}} Gr_\gamma^i K_{2i-\ell}(V) \rightarrow \bigoplus_{v \in S_\infty} H^{\ell-1}(V_v(\mathbb{C}), \mathbb{R})^{(i-1)}, \quad i \geq \ell,$$

which Beilinson expects to be isomorphisms when i is big enough [18].

Assuming that $K_m(V)$ does not contain any divisible subgroup, the Bockstein exact sequences

$$0 \rightarrow K_m(V) \otimes \mathbb{Z}/p^n \rightarrow K_m(V; \mathbb{Z}/p^n) \rightarrow K_{m-1}(V)[p^n] \rightarrow 0$$

will show that

$$\dim Gr'_\gamma K_{2i-\ell}(V; \mathbb{Q}_p/\mathbb{Z}_p) = \dim_{\mathbb{R}} \mathbb{R} \otimes_{\mathbb{Z}} Gr'_\gamma K_{2i-\ell}(V).$$

Therefore Theorem 1 expresses the compatibility of Beilinson's conjecture with Quillen's one.

1.9. It is hard to decide for which values of i the equality in Theorem 1 holds. For instance, assume V is a point. Then it is proved in [17], using algebraic K-theory, that

$$\dim H^\ell(K, \mathbb{Q}_p/\mathbb{Z}_p(i)) = n_{i,\ell} \quad \text{when } i \geq 2.$$

When $i = 0$, the Leopoldt's conjecture asserts that

$$\dim H^1(K, \mathbb{Q}_p/\mathbb{Z}_p) = n_{0,1} + 1$$

([14], §7, Lemma 1). The case $i < 0$ is not completely understood [14].

2. The local case

2.1. Statement of the result

2.1.1. Notations

Let K be a finite extension of the field \mathbb{Q}_p of p -adic numbers, $[K:\mathbb{Q}_p]$ its degree, \bar{K} an algebraic closure of K , C the completion of \bar{K} , $G = \text{Gal}(\bar{K}/K)$ the Galois group of \bar{K} over K , O_K the ring of integers of K , and k its residue field.

Assume V is a smooth proper variety over K with good reduction, i.e. such that there exists a smooth proper scheme \mathcal{V} over $\text{Spec } O_K$ whose generic fiber $\mathcal{V} \otimes_{O_K} K$ is isomorphic to V . We call $W = \mathcal{V} \otimes_{O_K} k$ the special fiber of \mathcal{V} .

Recall that when X is a scheme whose residual characteristics are different from p we define

$$H^\ell(X, \mathbb{Z}_p(i)) = \varprojlim_n H^\ell(X, \mathbb{Z}/p^n(i))$$

and

$$H^\ell(X, \mathbb{Q}_p(i)) = H^\ell(X, \mathbb{Z}_p(i)) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p.$$

2.1.2. *Hodge-Tate decomposition*

The action of $G = \text{Gal}(\bar{K}/K)$ upon \bar{K} extends to C by continuity. Let G act upon the module $H^\ell(\bar{V}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C$ diagonally. Call s.s. $(H^\ell(\bar{V}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C)$ the semi-simplification of this $\mathbb{Q}_p[G]$ -module. In [19] (§4.1., Remark), Tate conjectured that there exists a direct sum decomposition (“Hodge-Tate decomposition”)

$$\text{s.s. } H^\ell(\bar{V}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C \cong \bigoplus_{j=0}^{\ell} C(-j)^{m_{j,\ell}}$$

where $m_{j,\ell}$ are positive integers, $C(-j)$ are the Tate twists of C , and the isomorphism above is a G -isomorphism.

This conjecture was recently proved by S. Bloch and K. Kato [2] when V is “ordinary” in the following sense:

DEFINITION: The variety V is called *ordinary* when the De Rham cohomology groups $H^q(W, d\Omega_{W/k}^i)$ of its reduction W with coefficients the image of $\Omega_{W/k}^i$ by the De Rham differential are zero for all positive integers q and j .

2.1.3. THEOREM 2: *Let p be any prime, let V be a smooth proper variety with good reduction over a p -adic field K , and let $\bar{V} = V \otimes_K \bar{K}$.*

(i) *When l is a prime different from p and $\ell \neq 2i - 1, 2i, 2i + 1$, the groups $H^\ell(V, \mathbb{Q}_l(i))$ are zero.*

(ii) *For all $i \in \mathbb{Z}$ we have*

$$\dim_{\mathbb{Q}_p} H^\ell(V, \mathbb{Q}_p(i)) \geq [K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p).$$

Equality holds for almost all $i \in \mathbb{Z}$.

(iii) *When the cohomology groups of \bar{V} in dimensions $\ell, \ell - 1$ and $\ell - 2$ have a Hodge-Tate decomposition (cf. 2.1.2) and $i < 0$ or $i > \ell$, we have*

$$\dim_{\mathbb{Q}_p} H^\ell(V, \mathbb{Q}_p(i)) = [K: \mathbb{Q}_p] \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p).$$

(iv) *When V is projective and ordinary, and $\ell \neq 2i - 1, 2i, 2i + 1$, we have*

$$\dim_{\mathbb{Q}_p} H^\ell(V, \mathbb{Q}_p(i)) = [K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p).$$

2.2. The equalities of Theorem 2 will be deduced from the following:

PROPOSITION 4: *Assume that*

$$\begin{aligned} H^\ell(\bar{V}, \mathbb{Q}_l)(i)^G &= H^{\ell-1}(\bar{V}, \mathbb{Q}_l)(i)^G = H^{\ell-1}(\bar{V}, \mathbb{Q}_l(i-1))_G \\ &= H^{\ell-2}(\bar{V}, \mathbb{Q}_l(i-1))_G = 0. \end{aligned}$$

Then

$$\dim_{\mathbb{Q}_l} H^\ell(V, \mathbb{Q}_l(i)) = \begin{cases} 0 & \text{when } l \neq p \\ [K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p) & \\ & \text{when } l = p. \end{cases}$$

PROOF: The Hochschild-Serre spectral sequence attached to the Galois covering $\bar{V} \rightarrow V$ has E_2 -term

$$E_2^{rs} = H^r(K, H^s(\bar{V}, \mathbb{Z}/l^n(i)))$$

and converges to $H^{r+s}(V, \mathbb{Z}/l^n(i))$. Since all the groups above are finite, one can perform a projective limit of such spectral sequences when n varies (the Mittag-Leffler property holds). We get

$$E_2^{rs} = H^r(K, H^s(\bar{V}, \mathbb{Z}_l)(i)) \rightarrow H^{r+s}(V, \mathbb{Z}_l(i)).$$

The duality theorem for finite G -modules ([15], II.5.2., Theorem 2) gives

$$\begin{aligned} H^2(K, H^s(\bar{V}, \mathbb{Z}/l^n)(i)) &= H^0(K, H^s(\bar{V}, \mathbb{Z}/l^n)(i)^\wedge(1))^\wedge \\ &= H^s(\bar{V}, \mathbb{Z}/l^n)(i-1)_G, \end{aligned}$$

and taking a projective limit we get

$$H^2(K, H^s(\bar{V}, \mathbb{Z}_l)(i)) = H^s(\bar{V}, \mathbb{Z}_l)(i-1)_G.$$

The hypotheses of the Proposition imply that

$$E_2^{0,\ell} \otimes \mathbb{Q}_l = H^0(K, H^\ell(\bar{V}, \mathbb{Z}_l(i))) \otimes \mathbb{Q}_l = 0,$$

$$E_2^{0,\ell-1} \otimes \mathbb{Q}_l = 0, E_2^{2,\ell-1} \otimes \mathbb{Q}_l = H^{\ell-1}(\bar{V}, \mathbb{Q}_l(i-1))_G = 0$$

$$E_2^{2,\ell-2} \otimes \mathbb{Q}_l = H^{\ell-2}(\bar{V}, \mathbb{Q}_l(i-1))_G = 0.$$

On the other hand let M be any $\mathbb{Z}_l[[G]]$ -module which is finitely generated over \mathbb{Z}_l , and F its quotient by its torsion subgroup. We have (as in Proposition 2)

$$\begin{aligned}\chi(M) &= \sum_{r=0}^2 (-1)^r \dim_{\mathbb{Q}_l} H^r(K, M) \otimes \mathbb{Q}_l \\ &= \sum_{r=0}^2 (-1)^r \dim_{\mathbb{Z}_l} H^r(K, F) \\ &= \sum_{r=0}^2 (-1)^r \dim_{\mathbb{F}_l} H^r(K, F/IF),\end{aligned}$$

and by [15] (II.5.7, Theorem 5) we get $\chi(M) = 0$ when $l \neq p$ and

$$\begin{aligned}\chi(M) &= -[K: \mathbb{Q}_p] \times \dim_{\mathbb{F}_l}(F/IF) \\ &= -[K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} M \otimes \mathbb{Q}_p \quad \text{when } p = l.\end{aligned}$$

Here we get

$$\begin{aligned}\dim_{\mathbb{Q}_l} H^\ell(V, \mathbb{Q}_l(i)) &= \dim_{\mathbb{Q}_l} H^1(K, H^{\ell-1}(\bar{V}, \mathbb{Q}_l(i))) \\ &= -\chi(H^{\ell-1}(\bar{V}, \mathbb{Z}_l(i))) = \begin{cases} 0 & \text{when } l \neq p \\ [K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p) & \text{when } \ell = p. \end{cases}\end{aligned}$$

2.3. PROOF OF (i): Assume $l \neq p$. We want to show that the hypotheses of Proposition 4 are satisfied when $i \neq 2k - 1, 2k, 2k + 1$. Since V has good reduction, we get, by the smooth and proper base change theorem [5], an isomorphism of G -modules

$$H^\ell(\bar{V}, \mathbb{Q}_l(i)) \simeq H^\ell(\bar{W}, \mathbb{Q}_l(i))$$

where $\bar{W} = W \otimes \bar{k}$, \bar{k} is an algebraic closure of k , and G acts on \bar{W} through its projection onto $\text{Gal}(\bar{k}/k)$. The Weil conjectures then imply that

$$H^\ell(\bar{W}, \mathbb{Q}_l(i))^{\text{Gal}(\bar{k}/k)} = H^\ell(\bar{W}, \mathbb{Q}_l(i))_{\text{Gal}(\bar{k}/k)} = 0$$

when $\ell \neq 2i$ (cf. Proposition 1 and Lemma 1). Therefore, when $k = 2i - 1, 2i, 2i + 1$, the hypotheses of Proposition 4 hold.

2.4. PROOF OF (ii): From the Hochschild-Serre spectral sequence considered in Proposition 4 we get, for all $i \in \mathbb{Z}$,

$$\begin{aligned} \dim_{\mathbb{Q}_p} H^\ell(V, \mathbb{Q}_p(i)) &\geq \dim_{\mathbb{Q}_p} E_2^{1, \ell-1} \otimes \mathbb{Q}_p \geq \chi(H^{\ell-1}(\bar{V}, \mathbb{Z}_p(i))) \\ &= [K: \mathbb{Q}_p] \times \dim_{\mathbb{Q}_p} H^{\ell-1}(\bar{V}, \mathbb{Q}_p). \end{aligned}$$

To see that equality holds for almost all $i \in \mathbb{Z}$, let $K(\mu_{p^\infty}) = \bigcup_{n \geq 1} K(\mu_{p^n})$ be the maximal p -cyclotomic extension of K , $\Gamma = \text{Gal}(K(\mu_{p^\infty})/K)$ its Galois group over K , and $H = \text{Ker}(G \rightarrow \Gamma)$.

Fixing an integer $\ell \geq 0$, let $X = H^\ell(\bar{V}, \mathbb{Z}_p)^H$. It is a Λ -module, where $\Lambda = \mathbb{Z}_p[[\Gamma]] \simeq \mathbb{Z}_p[[T]]$ (cf. Proposition 3 and Lemma 2). Furthermore X is finitely generated as a \mathbb{Z}_p -module, therefore it is pseudo-isomorphic to a finite product of Λ -modules of the type $\mathbb{Z}_p[[T]]/(f(T))$ where $f(T)$ is a nonzero polynomial. But

$$(\Lambda/(f(T))(i))^\Gamma = (\Lambda/f(c^i(1+T)-1))^\Gamma$$

is finite when $f(c^i - 1) \neq 0$. Therefore

$$X(i)^\Gamma = H^\ell(\bar{V}, \mathbb{Z}_p(i))^G$$

is finite for almost all i . A similar proof gives that $H^\ell(\bar{V}, \mathbb{Z}_p(i))_G$ is finite for almost all i .

2.5. PROOF OF (iii): Assume that $H^\ell(\bar{V}, \mathbb{Q}_p)$ admits a Hodge-Tate decomposition

$$H^\ell(\bar{V}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C \simeq \bigoplus_{j=0}^{\ell} C(-j)^{m_{j,\ell}}$$

i.e.

$$H^\ell(\bar{V}, \mathbb{Q}_p(i)) \otimes_{\mathbb{Q}_p} C \simeq \bigoplus_{j=0}^{\ell} C(i-j)^{m_{j,\ell}}.$$

The group of invariants $H^\ell(\bar{V}, \mathbb{Q}_p(i))^G$ is a \mathbb{Q}_p -vector space contained in $(H^\ell(\bar{V}, \mathbb{Q}_p(i)) \otimes C)^G$, and Tate proved in [13] (Theorem 2) that when $n \neq 0$ we have $C(n)^G = 0$. Therefore, when $i < 0$ or $i > \ell$, we have $H^\ell(\bar{V}, \mathbb{Q}_p(i))^G = 0$.

On the other hand there are isomorphisms of C -vector spaces with

semilinear G -action:

$$\begin{aligned} & \text{s.s. Hom}_{\mathbb{Q}_p} \left(H^\ell(\bar{V}, \mathbb{Q}_p(i)), \mathbb{Q}_p \right) \otimes_{\mathbb{Q}_p} C \\ & \simeq \text{Hom}_C \left(\text{s.s. } H^\ell(\bar{V}, \mathbb{Q}_p(i)) \otimes_{\mathbb{Q}_p} C, C \right) \\ & \simeq \bigoplus_{j=0}^{\ell} \text{Hom}_C(C(i-j), C)^{m_{i,\ell}} \simeq \bigoplus_{j=0}^{\ell} C(j-i)^{m_{i,\ell}}. \end{aligned}$$

So, when $i < 0$ or $i > \ell$, we have

$$\text{Hom}_{\mathbb{Q}_p} \left(H^\ell(\bar{V}, \mathbb{Q}_p(i)), \mathbb{Q}_p \right)^G = \text{Hom}_{\mathbb{Q}_p} \left(H^\ell(\bar{V}, \mathbb{Q}_p(i))_G, \mathbb{Q}_p \right) = 0.$$

From this we deduce that when $i < 0$ or $i > \ell$ the hypotheses of Proposition 4 are satisfied.

2.6. PROOF OF (iv): By Proposition 4 it will be enough to show that

$$H^\ell(\bar{V}, \mathbb{Q}_p(i))^G = H^\ell(\bar{V}, \mathbb{Q}_p(i))_G = 0 \quad \text{when } \ell \neq 2i.$$

But Bloch and Kato proved in [2] that there exists a filtration $F^*H^\ell(\bar{V}, \mathbb{Q}_p)$ of $H^\ell(\bar{V}, \mathbb{Q}_p)$ whose corresponding graded module is $gr^*H^\ell(\bar{V}, \mathbb{Q}_p) = \bigoplus_{j=0}^{\ell} H^{n-j,j}(-j)$, where $(-j)$ is the twisting by the p -cyclotomic character, and $H^{n-j,j}$ is a module under $\text{Gal}(\bar{k}/k)$ upon which G acts via its projection to $\text{Gal}(\bar{k}/k)$. More precisely, writing $\bar{W} = W \otimes_k \bar{k}$, let $H_{\text{crys}}^n(\bar{W}/\mathbb{W}(\bar{k}))$ be its crystalline cohomology and $F: \bar{W} \rightarrow \bar{W}$ be the absolute Frobenius (which raises the coordinates over \bar{k} to the p -th power). Then

$$H^{n-j,j} = \left(H_{\text{crys}}^n(\bar{W}/\mathbb{W}(\bar{k})) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \right)^{[F=p^j]}$$

is the kernel of $F - p^j$ acting upon the crystalline cohomology of \bar{W} .

Let p^s be the order of the residue field k , let $f_g: W \rightarrow W$ be the geometric Frobenius of W (which raises the coordinates over k to the p^s -th power), and let $f_a \in \text{Gal}(\bar{k}/k)$ be the arithmetic Frobenius (sending $x \in \bar{k}$ to x^{p^s}). If we call $\tilde{f}_g = f_g \otimes id: \bar{W} \rightarrow \bar{W}$ (resp. $\tilde{f}_a = id \otimes f_a: \bar{W} \rightarrow \bar{W}$) the tensor product of f_g (resp. f_a) with the identity on \bar{k} (resp. W), we have the following equalities

$$F^s = \tilde{f}_g \circ \tilde{f}_a = \tilde{f}_a \circ \tilde{f}_g$$

and since \bar{f}_a and \bar{f}_g commute with F they act upon $H^{n-1,1}$. Since $H^{n-1,1}$ is contained in the crystalline cohomology of \bar{W} , the Weil conjectures in crystalline cohomology [10] tell us that the eigenvalues of \bar{f}_g acting on $H^{n-1,1}$ are algebraic numbers whose archimedean absolute values are equal to $p^{s_n/2}$.

Since $F = p^j$ on $H^{n-1,1}$, the equality $p^{s_j} = \bar{f}_g \circ \bar{f}_a$ shows that \bar{f}_a can have a fixed vector in $H^{n-1,1}$ only when $n = 2j$. Let $\tau \in G$ be a lifting of $f_a \in \text{Gal}(\bar{k}/k)$ such that τ lies in the kernel of the p -cyclotomic character. Then the endomorphism $\tau - 1$ of $H^{n-1,1}(i-j)$ has trivial kernel and cokernel unless $n = 2j$.

On the other hand, if γ_0 is a generator of $\text{Gal}(K(\mu_{p^\infty})/K)$, the endomorphism $\gamma_0 - 1$ of $H^{j,1}(i-j)$ has trivial kernel and cokernel unless $i = j$.

From this we conclude that $H^\ell(\bar{V}, \mathbb{Q}_p(i))^G = H^\ell(\bar{V}, \mathbb{Q}_p(i))_G = 0$ unless $\ell = 2i$. Q.E.D.

References

- [1] A. BEILINSON: Higher regulators and values of L-functions of curves. *Funk. Analysis 14* (2) (1980) 116–117.
- [2] S. BLOCH and K. KATO: *P-adic étale cohomology*, preprint.
- [3] W. BROWDER: Algebraic K-theory with coefficients \mathbb{Z}/p , in *Springer Lec. Notes in Maths. 657* (1978), 40–84.
- [4] P. DELIGNE: La conjecture de Weil II. *Publ. math. IHES 52* (1980) 137–252.
- [5] P. DELIGNE et al.: SGA IV $\frac{1}{2}$, Cohomologie étale, exp. 1. *Springer Lecture Notes in Maths. 568* (1977).
- [6] B. FERRERO and L. WASHINGTON: The Iwasawa μ_p invariant vanishes for abelian number fields. *Ann. Math. 109* (1979) 377–395.
- [7] K. IWASAWA: On \mathbb{Z}_p -extensions of algebraic number fields. *Ann. Math. 98* (1973) 246–326.
- [8] K. IWASAWA: Riemann-Hurwitz formula and p -adic Galois representations for number fields. *Tohoku Math. Journal 33* (1981) 263–288.
- [9] M. KAROUBI: Connexions, courbures et classes caractéristiques en K -théorie algébrique. in *Current Trends in Algebraic Topology*, CMS Conference Proceedings, Vol. 2, Part 1 (1982) 19–29.
- [10] M. KATZ and W. MESSING: Some consequences of the Riemann hypothesis for varieties for finite fields. *Invent. Math. 23* (1974) 73–77.
- [11] C. KRATZER: λ -structure en K -théorie algébrique. *Comm. Math. Helvetici 55* (1980) 233–254.
- [12] S. LICHTENBAUM: On the values of zeta and L-functions I. *Ann. Math. 96* (1972) 338–360.
- [13] D. QUILLEN: Higher Algebraic K-theory. *Proceedings of the International Congress of Mathematicians*, Vancouver, 1974.
- [14] P. SCHNEIDER; Über gewisse Galois Cohomologie Gruppen. *Math. Zeitschrift 168* (1979) 181–205.
- [15] J.-P. SERRE: Cohomologie galoisienne. *Springer Lecture Notes in Maths. 5* (1964).
- [16] C. SOULÉ: K -théorie des anneaux d'entiers de corps de nombres et cohomologie étale. *Inv. Math. 55* (1979) 251–295.
- [17] C. SOULÉ: On higher p -adic regulators, in *Springer Lectures Notes in Maths. 854* (1980) 372–401.

- [18] C. SOULÉ: On K -theory and values of zeta functions, in *Current Trends in Algebraic Topology*, CMS Conference Proceedings, Vol. 2, Part 1 (1982) 49–59.
- [19] J. TATE: p -divisible groups, in *Proceedings of a Conf. on local fields* (1967) Springer.
- [20] J. TATE: On the conjectures of Birch and Swinnerton-Dyer and a geometric analog., Sem. Bourbaki (1965–1966), exposé 306, in *Dix exposés sur la cohomologie des schémas* (1968), North-Holland and Masson.
- [21] J. TATE: Duality theorems in Galois cohomology over number fields. *Proc. Cong. Stockholm* (1962) 288–295.

(Oblatum 8-XI-1982)

U.E.R. de Mathématiques
Université Paris VII
Tour 45-55, 5^{ème} ét.
2, Place Jussieu
75251 Paris CEDEX 05
France