

COMPOSITIO MATHEMATICA

MICHELE COOK

An improved bound for the degree of smooth surfaces in P^4 not of general type

Compositio Mathematica, tome 102, n° 2 (1996), p. 141-145

http://www.numdam.org/item?id=CM_1996__102_2_141_0

© Foundation Compositio Mathematica, 1996, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

An improved bound for the degree of smooth surfaces in \mathbf{P}^4 not of general type

MICHELE COOK

Department of Mathematics, Oklahoma State University, Stillwater, OK 74078

Received 15 November 1994; accepted in final form 30 March 1995

This is an addendum to the paper of Braun and Fløystad ([BF]) on the bound for the degree of smooth surfaces in \mathbf{P}^4 not of general type. (In fact, any statements made here without comment will be found there.) Using their construction and the regularity of curves in \mathbf{P}^3 , one may lower the bound a little more. We will prove the following:

THEOREM. *Let S be a smooth surface of degree d in \mathbf{P}^4 not of general type. Then either $d \leq 70$ or S lies on a hypersurface in \mathbf{P}^4 of degree 5 and $d \leq 80$. Thus $d \leq 80$.*

(We will also indicate how one may actually get the bound down to 76.)

The main result of [BF] is to bound $\chi\mathcal{O}_S$ from below using generic initial ideal theory and information coming from a generic hyperplane section of S .

Let C be a generic hyperplane section of S , then C has associated to it two pieces of information relevant to this situation; *connected invariants* $\lambda_0 > \lambda_1 > \dots > \lambda_{s-1} > 0$ and *sporadic zeros* which are defined as follows:

DEFINITION. Let $\mathbf{C}[x_0, x_1, x_2, x_3]$ be the ring of polynomials of \mathbf{P}^3 , with reverse lexicographical ordering. Let C be a curve in \mathbf{P}^3 , then the generic initial ideal of C , $\text{gin}(I_C)$, is generated by elements of the form $x_0^i x_1^j x_2^k$. We say that a monomial $x_0^a x_1^b x_2^c$ is a *sporadic zero* if $x_0^a x_1^b x_2^c \notin \text{gin}(I_C)$, but there exists $c' > c$ such that $x_0^a x_1^b x_2^{c'} \in \text{gin}(I_C)$.

Let α_t is the number of sporadic zeros of degree t and m be the maximal degree of the sporadic zeros.

If π is the genus of C , then

$$\pi = 1 + \sum_{i=0}^{s-1} \left(\binom{\lambda_i}{2} + (i-1)\lambda_i \right) - \sum_{t=0}^m \alpha_t \quad (1)$$

and

$$1 + \sum_{i=0}^{s-1} \left(\binom{\lambda_i}{2} + (i-1)\lambda_i \right) \leq \frac{d^2}{2s} + (s-4)\frac{d}{2} + 1 = G(d, s) \tag{2}$$

for $d > (s-1)^2 + 1$.

(This is due to the work of Gruson and Peskine on the numerical invariants of points (see [GP]).)

In [BF] they show that

$$\chi_{\mathcal{O}_S} \geq \sum_{t=0}^{s-1} \left(\binom{\lambda_t + t - 1}{3} - \binom{t-1}{3} \right) - \sum_{t=0}^m \alpha_t(t-1),$$

and if $s \geq 2$ and $d > (s-1)^2 + 1$,

$$\sum_{t=0}^{s-1} \left(\binom{\lambda_t + t - 1}{3} - \binom{t-1}{3} \right) \geq s \binom{\frac{d}{s} + \frac{s-3}{2}}{3} + 1 - \binom{s-1}{4}.$$

They Combine this with the double point formula (see [H, p. 434])

$$d^2 - 5d - 10(\pi - 1) + 2(6\chi - K^2) = 0,$$

and the fact that if the degree of S is $d > 5$ then $K^2 \leq 9$, to get

$$\begin{aligned} 18 \geq 2K^2 &= d^2 - 5d - 10(\pi - 1) + 12\chi \\ &\geq d^2 - 5d - 10 \left(\frac{d^2}{2s} + (s-4)\frac{d}{2} - \sum_{t=0}^m \alpha_t \right) \\ &\quad + 12 \left(s \binom{\frac{d}{s} + \frac{s-3}{2}}{3} + 1 - \binom{s-1}{4} - \sum_{t=0}^m \alpha_t(t-1) \right) \\ &= d^2 - 5d - 10 \left(\frac{d^2}{2s} + (s-4)\frac{d}{2} \right) \\ &\quad + 12 \left(s \binom{\frac{d}{s} + \frac{s-3}{2}}{3} + 1 - \binom{s-1}{4} \right) - \sum_{t=0}^m \alpha_t(12t - 22). \tag{3} \end{aligned}$$

We will use regularity conditions to find an upper bound for

$$A = \sum_{t=0}^m \alpha_t(12t - 22).$$

First we should note that if $s = \min\{k | h^0(\mathcal{I}_S(k)) \neq 0\}$. Then by the work of Ellingsrud and Peskine ([EP]) if S is a surface, of degree d , not of general type then either $d \leq 90$ or $s \leq 5$. Furthermore, if one imitates their argument with $s = 6$, one gets either $d \leq 70$ or $s \leq 6$. We will assume $s \leq 6$. We also know that if $s \leq 3$ then $d \leq 8$. Hence we only need to bound the degree of surfaces not of general type with $s = 4, 5$, or 6 .

By [BS] the regularity of an ideal I is equal to the regularity of $\text{gin}(I)$, which is the highest degree of a minimal generator of $\text{gin}(I)$. Furthermore, by [GLP], the regularity of a smooth curve of degree d in \mathbb{P}^3 is $\leq d - 1$. Hence the largest degree of the minimal generators of $\text{gin}(I_C)$ is $\leq d - 1$ and so all the sporadic zeros of C are in degree $\leq d - 2$.

We will now bound the number of sporadic zeros. Let $\gamma = G(d, s) - \pi$. Any bound on γ will also bound the number of sporadic zeros (see equations (1) and (2)). By [EP], $\gamma \leq \frac{d(s-1)^2}{2s}$. Furthermore, by the double point formula and the fact that if S is a surface not of general type $K^2 < 6\chi$, we get $\pi \geq \frac{d^2-5d+10}{10}$ and thus $\gamma \leq \frac{d^2}{2s} + (s-4)\frac{d}{2} + 1 - \frac{d^2-5d+10}{10}$. Taking the minimum of these bounds for γ , we get, for $s = 4, \gamma \leq \frac{9d}{8}$, for $s = 5, \gamma \leq d$ and for $s = 6, \gamma \leq \frac{d(90-d)}{60}$.

Furthermore, by connectedness of the invariants $\lambda_0 \leq \frac{d}{s} + s - 1$ and $\lambda_1 \leq \frac{d}{s} + s - 2$.

Putting all this together, we have

$$A \leq \sum_{t=\frac{d}{s}+s-1}^{d-2} (12t - 22) + \sum_{t=\frac{d}{s}+s-1}^{\gamma-d+\frac{2d}{s}+2s-2} (12t - 22)$$

and using the bounds on γ , we get

for $s = 4, A \leq \frac{243}{32}d^2 - 9d + 192,$

for $s = 5, A \leq \frac{162}{25}d^2 - 16d + 300.$

Substituting back into the original equation (3) above, we get

for $s = 4, 0 \geq \frac{1}{8}d^3 - \frac{275}{32}d^2 + \frac{7}{2}d - 195$ and hence $d \leq 68,$

for $s = 5, 0 \geq \frac{2}{25}d^3 - \frac{162}{25}d^2 + 4d - 318$ and hence $d \leq 80.$

Now suppose S does not lie on a hypersurface of degree 5. Then the degree of $S = d \leq 90$. By [EP], we know that either $d \leq 70$ or S is contained in a hypersurface of degree 6. Therefore it remains to show that there are no smooth surfaces not of general type with $s = 6$ and $71 \leq d \leq 90$.

Suppose such a surface existed. Then $\gamma \leq \frac{71(90-71)}{60}$, i.e. $\gamma \leq 22$ and so

$$A \leq \sum_{t=\frac{d}{6}+5}^{\frac{d}{6}+5+21} (12t - 22) = 44d + 3608.$$

Substituting as above we get

$$0 \geq \frac{d^3}{18} + \frac{2}{3}d^2 - \frac{119}{2}d - \frac{7321}{2}.$$

But this is a contradiction for $71 \leq d \leq 90$. □

By more careful consideration of the equations involved, it is possible to lower the bound a little more:

Making use of the equation

$$0 \leq \sum_{t=0}^m \alpha_t \leq \sum_{i=0}^{s-1} \left(\binom{\lambda_i}{2} + i(\lambda_i) \right) - \frac{d^2 + 5d}{10}$$

arising from the double point formula, and

$$\begin{aligned} 0 \geq & d^2 - 5d - 18 - 10 \left(\sum_{i=0}^{s-1} \left(\binom{\lambda_i}{2} + (i-1)(\lambda_i) \right) \right) \\ & + 12 \left(\sum_{i=0}^{s-1} \binom{\lambda_i + i - 1}{3} - \binom{s-1}{4} \right) - \sum_{t=0}^m \alpha_t (12t - 22) \end{aligned}$$

it is possible to show that $d \leq 76$ and $s \leq 8$. This is done by writing out all the possible connected invariants of length s and degree d and checking the inequalities on “Mathematica”. (I would like to thank Rich Liebling for sending me the program which calculates the invariants.)

The temptation at this point is to try to lower the bound by considering each degree individually and using ad hoc methods. For example, we know for certain configurations of the invariants the curve is Arithmetically Cohen-Macaulay and hence has no sporadic zeros. But, to make any significant improvements using these methods, one needs to understand which configurations of the sporadic zeros can occur.

Acknowledgements

I would like to thank Mei-Chu Chang for asking me to talk on the paper of Braun and Fløystad, which made me read the details of the paper. I would like to thank Robert Braun and Gunnar Fløystad for answering all my questions and Robert for pointing out the better bound for γ for $s = 6$. Finally, I would like to thank Bruce Crauder and Sheldon Katz for making me feel welcome during my stay here at OSU.

References

- [BS] Bayer, D. and Stillman, M.: “A criterion for detecting m -regularity”, *Invent. Math.* 87 (1987) 1–11.
- [BF] Braun, R. and Fløystad, G.: “A bound for the degree of smooth surfaces in \mathbf{P}^4 not of general type”, *Compositio Mathematica*, Vol. 93, No. 2, September(I) 1994, 211–229.
- [EP] Ellingsrud, G. and Peskine, C.: “Sur les surfaces lisses de \mathbf{P}^4 ”, *Invent. Math.* 95 (1989) 1–11.
- [GLP] Gruson, L., Lazarsfeld, R. and Peskine, C.: “On a Theorem of Castelnuovo, and the equations defining space curves”, *Invent. Math.* 72 (1983) 491–506.
- [GP] Gruson, L. and Peskine, C.: “Genres des courbes de l’espace projectif”, *Lecture Notes in Mathematics, Algebraic Geometry*, Tromsø 1977, 687 (1977) 31–59.
- [H] Hartshorne, R.: *Algebraic Geometry*, Springer-Verlag (1977).