

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

ARMIN FREI

On the preservation of homotopy invariance by Kan extensions

Cahiers de topologie et géométrie différentielle catégoriques, tome 22, n° 3 (1981), p. 329-336

http://www.numdam.org/item?id=CTGDC_1981__22_3_329_0

© Andrée C. Ehresmann et les auteurs, 1981, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE PRESERVATION OF HOMOTOPY INVARIANCE BY KAN EXTENSIONS

by Amin FREI

1. INTRODUCTION.

In the literature (e. g. in [1, 2, 3, 4, 6, 7]) one considers Kan extensions of functors (usually homology or cohomology functors) defined on a category $\mathcal{T}_0$ of spaces and homotopy classes of maps to another such category $\mathcal{T}_1$, containing $\mathcal{T}_0$ as a full subcategory. The following question arises: (in rough terms) does it matter whether one takes the Kan extension at the level of homotopy classes or at the level of maps? In order to lay out the situation we consider the commutative diagram:

$$(1.1) \quad \begin{array}{ccc} \mathcal{T}_0 & \xrightarrow{E} & \mathcal{T}_1 \\ \downarrow H_0 & \searrow F & \downarrow H_1 \\ & \mathcal{K} & \\ \downarrow \tilde{H}_0 & \nearrow \tilde{F} & \\ \tilde{\mathcal{T}}_0 & \xrightarrow{\tilde{E}} & \tilde{\mathcal{T}}_1 \end{array}$$

where $\mathcal{T}_0$ and $\mathcal{T}_1$ are full subcategories of the category of spaces and continuous maps, E is a full embedding and $\tilde{\mathcal{T}}_0$, $\tilde{\mathcal{T}}_1$, $\tilde{E}$ are their quotients modulo homotopy. H_1 and H_0 are the quotient functors and F is any homotopy invariant functor. Throughout the paper we use the notations G^K , ${}^K G$ for the right respectively the left Kan extension of G along K . As we take Kan extensions along full embeddings we may, and do, always choose them so that the universal transformation is the identity. Thus, in more precise terms, our question is:

When is $\tilde{F}^{\tilde{E}} H_1 = F^E$?

It turns out that this is the case precisely when F^E exists and is homotopy invariant (Theorem 3.4). By duality the same holds when F^E , $\tilde{F}^{\tilde{E}}$ are replaced by the left Kan extensions ${}^E F$, ${}^{\tilde{E}} \tilde{F}$, or when F is a cofunctor. In

view of this we investigate when the Kan extension of a homotopy invariant functor is homotopy invariant.

We shall always refer to the situation of diagram (1.1). We assume that $\mathcal{K}$ is sufficiently complete or cocomplete in order to admit the limits or colimits defining the pointwise right or left Kan extensions which we consider.

In Section 2 we treat the case where $\mathcal{T}_0$ contains the cofibration $X \rightarrow Zf$ for any map $f: X \rightarrow EY$ in $\mathcal{T}_1$ (Zf denotes the mapping cylinder of f) and its dual, that is, the case where $\mathcal{T}_0$ contains the mapping path fibration $Y^I \rightarrow X$ for any map $EY \rightarrow X$. In the first case F^E is homotopy invariant and in the second case EF is. Both conditions are clearly satisfied when $\mathcal{T}_0$ contains entire homotopy types, as is the case for categories which are admissible in the sense of [1, 6]. Section 3 contains some categorical facts about Kan extensions, which shall be used in the sequel. Although what is said in this section applies to more general situations, for the sake of shortness and ease of reading, we adhere to the situation of diagram (1.1). In Section 4 we show that if $\mathcal{T}_0$ contains Y^I whenever it contains Y , then F^E is homotopy invariant, and, dually, if it contains $Y \times I$ together with Y , then EF is homotopy invariant. The condition that $\mathcal{T}_0$ contain $Y \times I$ along with Y is rather harmless when $\mathcal{T}_0$ is a category of polyhedra or of CW-complexes as $- \times I$ preserves them and also preserves finiteness, local finiteness and finite dimension, thus left Kan extensions of homotopy invariant functors from any reasonable category of polyhedra or CW-complexes are homotopy invariant. Unfortunately the dual is not adequate for such categories; we shall consider right Kan extensions from categories of polyhedra in another paper.

2. THE CASE WHERE $\mathcal{T}_0$ CONTAINS Zf FOR ALL $f: X \rightarrow EY$.

THEOREM 2.1. *If the category $\mathcal{T}_0$ contains the mapping cylinder Zf for any map $f: X \rightarrow EY$ in $\mathcal{T}_1$, then:*

(i) *F^E exists iff $\tilde{F}^E$ exists.*

(ii) *$\tilde{F}^E H_1 = F^E$, in particular F^E is homotopy invariant.*

PROOF. Let $H_X: (X \downarrow E) \rightarrow (X \downarrow \tilde{E})$ be the functor which takes the object (f, Y) of $(X \downarrow E)$, that is a map $f: X \rightarrow EY$, to its homotopy class $H_1 f$ and a map g of $(X \downarrow E)$ to its homotopy class $H_1 g$. Then the diagram

$$\begin{array}{ccccc} (X \downarrow E) & \xrightarrow{Q} & \mathcal{T}_0 & \xrightarrow{F} & \mathcal{K} \\ H_X \downarrow & & H_0 \downarrow & \nearrow \tilde{F} & \\ (X \downarrow \tilde{E}) & \xrightarrow{Q} & \tilde{\mathcal{T}}_0 & & \end{array}$$

where Q denotes the usual forgetful functor, is commutative. Hence it suffices to show that the functor H_X is initial for all X in $|\mathcal{T}_1|$. Now, every object in $(X \downarrow \tilde{E})$ is clearly of the form $H_X(f, Y)$ and the category $\mathcal{X} = (H_X \downarrow H_X(f, Y))$ is nonempty. In order to show that it is connected we consider two objects

$$(1.2) \quad \begin{array}{ccc} H_X(f_1, Y_1) & & H_X(f_2, Y_2) \\ & \searrow H_1 v_1 \quad \nearrow H_2 v_2 & \\ & H_X(f, Y) & \end{array}$$


in $\mathcal{X}$, i.e., a homotopy commutative diagram

$$\begin{array}{ccccc} EY_1 & \xrightarrow{v_1} & EY & \xleftarrow{v_2} & EY_2 \\ & \searrow f_1 & \uparrow f & \nearrow f_2 & \\ & & X & & \end{array}$$


in $\mathcal{T}_1$. We replace f_1, f_2 by cofibrations c_1, c_2 resp. and obtain a diagram

$$\begin{array}{ccccccc} & & & & EY & & \\ & & & & \uparrow & & \\ EY_1 & \xleftarrow{r_1} & EZf_1 & & EZf_2 & \xrightarrow{r_2} & EY_2 \\ & \searrow f_1 & \nearrow c_1 & & \nearrow c_2 & \searrow f_2 & \\ & & X & & & & \end{array}$$

where r_1 and r_2 are homotopy equivalences and where $v_i r_i c_i \sim f$. By the homotopy extension property there exist maps $u_i: Zf_i \rightarrow Y_i$ with $u_i c_i = f$. Thus we obtain a strictly commutative diagram in $\mathcal{T}_1$



which gives rise to a commutative diagram in $\mathcal{X}$:



that is, to a connection between the objects (1.2).

A dual procedure, replacing the appropriate maps by their mapping path fibrations, gives the proof of:

THEOREM 2.2. *If the category $\mathcal{T}_0$ contains the mapping path fibration for any map $EY \rightarrow X$ in $\mathcal{T}_1$, then:*

- (i) ${}^E F$ exists iff ${}^E \tilde{F}$ exists.
- (ii) ${}^E \tilde{F} H_1 = {}^E F$, in particular ${}^E F$ is homotopy invariant.

When $\mathcal{T}_0$ contains entire homotopy types, then the hypotheses of both theorems are satisfied and we immediately have

COROLLARY 2.3. *If $\mathcal{T}_0$ is closed under homotopy types, then the conclusions of both Theorems, 2.1 and 2.2, hold.*

3. SOME CATEGORICAL FACTS.

As the Kan extension of F along a fully faithful functor E can be chosen so that $F^E E = F$, there is a very explicit form of the limiting cone defining $F^E X$.

PROPOSITION 3.1. *For any object X in $\mathcal{T}_1$ the family*

$$\{ F^E f \mid (f, Y) \in |(X \downarrow E)| \}$$

is a universal cone.

PROOF. Let X be an object in $\mathcal{T}_I$ and let

$$\{ \eta(g, Z): F^E X \rightarrow FQ(g, Z) \},$$

where $Q: (X \downarrow E) \rightarrow \mathcal{T}_I$ is the forgetful functor, be a universal cone. For a map $f: X \rightarrow EY$, the map $F^E f$ renders the diagram

$$\begin{array}{ccc} F^E X & \xrightarrow{F^E f} & F^E EY = FY \\ & \searrow \eta(hf, Z) & \swarrow \lambda(h, Z) \\ & FZ & \end{array}$$

commutative for any (h, Z) in $(EY \downarrow E)$, where λ denotes the universal cone defining $F^E EY$. In particular for $(1, Y)$ we have that $\lambda(1, Y) = 1$, thus $\eta(f, Y) = F^E f$.

The proposition above leads to a criterion for F^E to be homotopy invariant. Thus

PROPOSITION 3.2. *If F^E takes homotopic maps $f, g: X \rightarrow EY$ to the same map, then F^E is homotopy invariant.*

PROOF. Let $h, i: X_1 \rightarrow X_2$ be homotopic maps in $\mathcal{T}_I$. Using the representation given in Proposition 3.1 of the universal cone defining F^E , the map $F^E h$ is the unique one for which $F^E t \cdot F^E h = F^E th$, while $F^E i$ is the unique one for which $F^E t \cdot F^E i = F^E ti$, for all objects (t, Y) in $(X_2 \downarrow E)$. As $th, ti: X_1 \rightarrow EY$ are homotopic, one has, by hypothesis, that

$$F^E th = F^E ti, \text{ hence } F^E h = F^E i$$

by universality.

The next proposition points out a nice feature of quotient functors.

PROPOSITION 3.3. *If $G: \tilde{\mathcal{T}}_I \rightarrow \mathcal{K}$ is any functor, then G is the right and the left pointwise Kan extension of GH_1 along H_1 , with the identity as universal transformation. The same holds, of course, for H_0 .*

PROOF. As H_1 is full and onto objects it induces a bijection

$$\text{Nat}[G, G'] \rightarrow \text{Nat}[GH_1, G'H_1]$$

for all functors $G, G': \tilde{\mathcal{T}}_I \rightarrow \mathcal{K}$. Hence G is not only a pointwise, but even

an absolute Kan extension.

We now proceed to show that precisely when F^E is homotopy invariant, Kan extending at the map or at the homotopy class level give the same. More precisely:

THEOREM 3.4. *In the situation of diagram (1.1) the following are equivalent:*

(i) F^E exists and is homotopy invariant.

(ii) $\tilde{F}^E$ exists and $\tilde{F}^E H_1 = F^E$.

PROOF. (ii) $\Rightarrow$ (i) is trivial. (i) $\Rightarrow$ (ii): As F^E is homotopy invariant there is a unique functor $\bar{F}: \mathcal{T}_1 \rightarrow \mathcal{K}$ with $\bar{F}H_1 = F^E$. Then

$$\begin{aligned}\bar{F} &= (F^E)^{H_1} && \text{by Proposition 3.3,} \\ &= F^{H_1 E} && \text{by Lemma 1.2 (i) of [5],} \\ &= \tilde{F}^E H_0 && \text{as (1.1) is commutative.}\end{aligned}$$

By Proposition 3.3, $\tilde{F} = F^{H_0}$ and by Lemma 1.2 (ii) of [5] there is a unique $\mu: \tilde{F} \tilde{E} \rightarrow \bar{F}$ with $\mu H_0 = 1$, thus $\mu = 1$ as H_0 is onto objects, and $\bar{F} = \tilde{F}^E$.

4. THE CASE WHERE $\mathcal{T}_0$ IS CLOSED UNDER $(-)^I$ OR $-\times I$.

Suppose that for every Y in $\mathcal{T}_0$ also Y^I lies in $\mathcal{T}_0$. Let $f, g: X \rightarrow EY$ be homotopic maps. There is a map $k: X \rightarrow Y^I$ such that

$$d_0 k = f \quad \text{and} \quad d_1 k = g,$$

where d_0 and d_1 are the evaluations at 0 and 1 respectively. As d_0 and d_1 are homotopic and lie in $\mathcal{T}_0$ one has, for any functor $\bar{F}: \mathcal{T}_1 \rightarrow \mathcal{K}$ with $\bar{F}E$ homotopy invariant, that $\bar{F}f = \bar{F}g$. This holds in particular for the right Kan extension F^E of F . Invoking Proposition 3.2 we then have:

THEOREM 4.1. *If $\mathcal{T}_0$ contains X^I whenever it contains X , then F^E is homotopy invariant.*

Dually we have

THEOREM 4.2. *If $\mathcal{T}_0$ contains $X \times I$ whenever it contains X , then ${}^E F$ is homotopy invariant.*

We conclude by observing that, if $\mathcal{J}_0$ is closed under homotopy types, one obtains the conclusion (ii) of the theorems in Section 2 via the theorems in this section and Theorem 3.4 (and its dual). Nevertheless the theorems in Section 2 lead a separate life from those in Section 4 as one sees by taking $\mathcal{J}_0$ to be the category of polyhedra, or CW-complexes or of compact spaces, which are closed under $\times$ - I but not under homotopy types.

REFERENCES.

1. A. DELEANU & P. HILTON, On Kan extensions of cohomology theories and Serre classes of groups, *Fund. Math.* LXXIII (1971), 143-165.
2. A. DELEANU & P. HILTON, Remark on Čech extensions of cohomology functors, *Aarhus Univ. Math. Inst. Series* 13, I (1970), 44-66.
3. A. DELEANU & P. HILTON, Some remarks on general cohomology theories, *Math. Scand.* 22 (1968), 227-240.
4. A. DOLD, *Lectures on algebraic topology* (Appendix), Springer, Grundlehren 200, 1972.
5. A. FREI, On categorical shape theory, *Cahiers Topo. et Géom. Diff.* XVII-3 (1976), 261-294.
6. P. HILTON, On the construction of cohomology theories, *Rendic. di Mat.* 1 (1968), 219-232.
7. R. A. PICCININI, Extension de théories d'homologie, *C.R.A.S. Paris* 274 (1972), 828-829.

Department of Mathematics
 University of British Columbia
 121 - 1984 Mathematics Road
 VANCOUVER, B.C. V6T 1Y4
 CANADA