

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

RONALD BROWN

MAREK GOLASINSKI

A model structure for the homotopy theory of crossed complexes

Cahiers de topologie et géométrie différentielle catégoriques, tome
30, n° 1 (1989), p. 61-82

http://www.numdam.org/item?id=CTGDC_1989__30_1_61_0

© Andrée C. Ehresmann et les auteurs, 1989, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A MODEL STRUCTURE FOR THE HOMOTOPY THEORY OF CROSSED COMPLEXES

by Ronald BROWN and Marek GOLASINSKI

RÉSUMÉ. Les complexes croisés sont analogues à des complexes de chaîne mais avec les propriétés non-abéliennes des modules croisés de dimensions 1 et 2. Ils interviennent dans la théorie homotopique et la cohomologie des groupes. Ici on montre que la catégorie $\mathbf{Czs}$ des complexes croisés a une bonne structure de catégorie avec modèle pour la théorie d'homotopie, en prenant les classes déjà connues pour les équivalence faibles et les fibrations, et une nouvelle notion de cofibration. Les preuves utilisent la structure monoidale fermée sur les complexes croisés développée par Brown et Higgins, laquelle fournit des objets cylindre et cocylindre adéquats pour $\mathbf{Czs}$.

INTRODUCTION.

The definition of *crossed complex* is motivated by the principal example, the *fundamental crossed complex* $\pi\mathbf{X}$ of a *filtered space*

$$\mathbf{X}: X_0 \subset X_1 \subset \dots \subset X_n \subset \dots \subset X_\infty$$

([8], 5). In this crossed complex, $\pi_0\mathbf{X}$ is the set π_0X_0 ; $\pi_1\mathbf{X}$ is the fundamental groupoid $\pi_1(X_1, X_0)$; and for $n \geq 2$, $\pi_n\mathbf{X}$ is the family of relative homotopy groups $\pi_n(X_n, X_{n-1}, p)$, for all p in X_0 . This structure is also equipped with boundary maps from $\pi_n\mathbf{X}$ to $\pi_{n-1}\mathbf{X}$, $n \geq 1$, and operations of $\pi_1\mathbf{X}$ on $\pi_n\mathbf{X}$ for $n \geq 2$, all satisfying appropriate axioms (see Section 1). It is because of the widespread use of crossed complexes (summarised below) that it is necessary to discuss their appropriate homotopy theory, and this is our aim. Crossed complexes with a single vertex and satisfying a freeness condition were used by Whitehead in [26], under the name "homotopy systems", for discussing reali-

sation problems and models of low-dimensional homotopy types. They were also used in his famous paper [27] on simple homotopy types, again for realisation problems, although this application has been neglected up to now. These aspects are taken up in [1], where free, reduced crossed complexes are seen as constituting the first level in a tower of approximations to homotopy theory. The functor

$$\pi : (\text{filtered spaces}) \rightarrow (\text{crossed complexes})$$

satisfies a Generalised Van Kampen Theorem; i.e., it preserves certain colimits [8]. This result includes the usual Van Kampen Theorem, and other basic results in homotopy theory, for example the relative Hurewicz Theorem, and the result of Whitehead that $\pi_2(X \cup \{e_\lambda^2\}, X)$ is a free crossed $\pi_1(X)$ -module. It also implies new results on second homotopy groups [6]. There is a *classifying filtered-space functor*

$$\mathbf{B} : (\text{crossed complexes}) \rightarrow (\text{filtered spaces})$$

such that $\pi\mathbf{B}$ is naturally equivalent to the identity ([8], Section 9. and [11]). For a crossed complex C , the space $(\mathbf{B}C)_\infty$ is written BC and called simply the *classifying space* of the crossed complex C . The main result of [11] is the homotopy classification result, that if $\mathbf{X}$ is the filtered space of skeleta of a CW-complex X , then there is a bijection of homotopy classes

$$[X, BC] \approx [\pi\mathbf{X}, C] \quad (\text{see [5] for a summary}).$$

A crossed complex is of *rank* n if it is zero above dimension n . The crossed complexes of rank 1 are the groupoids, and these are well known to be models of homotopy 1-types. The crossed complexes of rank 2 are the *crossed modules (of groupoids)*. These are models of homotopy 2-types. Thus the Generalised Van Kampen Theorem enables the computation in some cases of the 2-type of a union of spaces.

In homological algebra, it is common to consider a *free resolution* of an algebraic object, for example of a module, and such a resolution is a chain complex of free modules. It is explained in [13] how crossed modules arise in considering identities among relations for a presentation of a group, and the general idea of a *crossed resolution* is explained in the survey article [4]. From this point of view, it is not surprising that

crossed complexes have been used to interpret the cohomology $H^n(G,A)$ of a group G with coefficients in a G -module A (cf. [15,17,20,21]).

It now seems reasonable to regard the replacement of chain complexes by crossed complexes as the first step towards a non-abelian homological algebra. It is these twin relations of crossed complexes to homological algebra and to homotopy theory which make it essential to have a satisfactory homotopy theory for crossed complexes.

The definition of a homotopy of morphisms of crossed complexes is well known and due to Whitehead [26]. It is exploited in [17] for the representation of cohomology of a group and in [9] for the theory of extensions of groups. However the theory of chain complexes has another type of homotopy theory due to Quillen [22], which is important in homological and homotopical algebra, and which involves defining notions of *weak equivalence*, *fibration* and *cofibration*, to obtain the structure of *closed model category*. For crossed complexes, weak equivalences are defined in [7] and fibrations in [16]. We use the methods of [22] to define *cofibrations* of crossed complexes and we prove in Theorem 2.12 that the weak equivalences, fibrations and cofibrations satisfy the axioms for a *closed model category* in the sense of [22]. However, we do not know if one further axiom is satisfied: *is it true that a pushout of a weak equivalence by a cofibration is again a weak equivalence?*

The proof of Theorem 2.12 requires machinery on crossed complexes developed by R. Brown and P.J. Higgins in [10]. They use ω -groupoid methods from [7] to give the category Czs of crossed complexes a symmetric, monoidal closed structure, with internal hom functor $CRS(-,-)$ and tensor product $- \otimes -$, analogous to corresponding functors on chain complexes. If B and C are crossed complexes, then $CRS(B,C)$ is: in dimension 0, the morphisms $B \rightarrow C$; in dimension 1, the homotopies of morphisms; in higher dimensions, the higher homotopies. Thus the closed structure on Czs includes a satisfactory theory of homotopy equivalences. But, in a similar manner to chain complexes, the definitions and applications of fibrations and cofibrations are not so straightforward, and it is useful to take weak equivalences rather than equivalences as basic. It is this theory that we develop.

In §1 we recall some basic definitions of crossed complexes, and weak equivalences. In §2 we follow [16] in defining fibrations, and follow [22] in deriving a definition of cofibration. We then prove that, with respect to these classes of morphisms, $\mathbf{Czs}$ is a closed model category. In §3 we derive a Whitehead type Theorem: *If a morphism of cofibrant objects in $\mathbf{Czs}$ is a weak equivalence, then it is also a homotopy equivalence.* In §4 we point out other homotopical results for crossed complexes which may be obtained using the methods of double categories with connections due to Spencer [23] and Spencer-Wong [24].

A different approach to abstract homotopy theory is given by Kamps-Porter [19] in terms of homotopy and cohomotopy systems. In these, a category is enriched over the category of cubical sets, and certain Kan extension conditions are imposed to allow manipulation of homotopies. This approach is useful for crossed complexes because the equivalence of crossed complexes and ω -groupoids [7], which is used in [10] to obtain the monoidal closed structure on $\mathbf{Czs}$, also allows the category $\mathbf{Czs}$ to be enriched over ω -groupoids. The latter, as cubical sets, satisfy a strong form of the Kan extension condition. The consequences of this will be developed elsewhere by the second author and Kamps.

The second author would like to thank Prof. R. Brown for arranging from the University College of North Wales, Bangor, support for his visit as Academic Visitor in 1986/87 where this work was carried out.

1. PRELIMINARIES.

A *crossed complex* C of groupoids [7] is a sequence

$$\dots \longrightarrow C_n \xrightarrow{\delta} C_{n-1} \longrightarrow \dots \longrightarrow C_2 \xrightarrow{\delta} C_1 \xrightleftharpoons[\delta^1]{\delta^0} C_0$$

satisfying the following conditions:

(i) C_1 is a groupoid with C_0 as its set of vertices and δ^0, δ^1 its initial and final maps. We write $C_1(p, q)$ for the set of arrows from p to q (p, q in C_0) and $C_1(p)$ for the group $C_1(p, p)$.

(ii) For $n \geq 2$, C_n is a family of groups $\{C_n(p)\}_{p \in C}$ and

for $n \geq 3$, the groups $C_n(p)$ are abelian.

(iii) The groupoid C_1 operates on the right on each C_n (for $n \geq 2$) by an action denoted $(x, a) \mapsto x^a$. Here, if $x \in C_n(p)$ and $a \in C_1(p, q)$, then we have $x^a \in C_n(q)$.

We use additive notation for all groups $C_n(p)$ and the groupoid C_1 .

(iv) For $n \geq 2$, $\delta: C_n \rightarrow C_{n-1}$ is a morphism of groupoids over C_0 and preserves the action of C_1 , where C_1 acts on the group $C_1(p)$ by conjugation: $x^a = -a + x + a$.

(v) $\delta\delta = 0: C_n \rightarrow C_{n-2}$ for $n \geq 3$ (and $\delta^0\delta = \delta^1\delta: C_2 \rightarrow C_0$, as follows from (iv)).

(vi) If $c \in C_2$, then δc operates trivially on C_n for $n \geq 3$ and operates on C_2 as conjugation by c , that is

$$x^{\delta c} = -c + x + c \quad (x, c \in C(p)).$$

In any crossed complex C , βc denotes the *base point* of c , that is, if $c \in C_0$ then $\beta c = c$, if $c \in C_1(p, q)$ or $c \in C_n(q)$ for $n \geq 2$, then $\beta c = q$.

A *morphism of crossed complexes* $f: B \rightarrow C$ is a family of morphisms of groupoids $f_n: B_n \rightarrow C_n$ ($n \geq 1$) all inducing the same map of vertices $f_0: B_0 \rightarrow C_0$ and compatible with the *boundary maps* $\delta: B_n \rightarrow B_{n-1}$, $C_n \rightarrow C_{n-1}$ and the actions of B_1 , C_1 on B_n, C_n . We denote by Czs the resulting category of crossed complexes.

The basic example we have in mind is the fundamental crossed complex πX of a filtered space X , as described in the Introduction.

It follows from observations by Brown-Higgins in [7], p.238; that the category Czs is complete and cocomplete. The coproduct in Czs is just disjoint union $\amalg$, while colimits in Czs are constructed in Section 6 of [8]. Moreover the paper [10] defines for any crossed complex B an internal hom functor $CRS(B, -)$ and its left adjoint, a tensor product $- \otimes B$. This gives Czs the structure of a symmetric, monoidal closed category.

Write $\mathbb{C}(n)$ for the crossed complex freely generated by one generator c_n in dimension n . So $\mathbb{C}(0)$ is $*$; $\mathbb{C}(1)$ is the groupoid $\mathcal{D}$; and for $n \geq 2$, $\mathbb{C}(n)$ is in dimensions n and $n-1$ an infinite cyclic group with generators c_n and δc_n respectively, and

is otherwise trivial. Let $\mathbf{S}(n-1)$ be the $(n-1)$ -skeleton of $\mathbf{C}(n)$, with inclusions $\mathbf{S}(n-1) \rightarrow \mathbf{C}(n)$. If $\mathbf{E}^{n-1}$ and $\mathbf{S}^{n-1}$ denote the skeletal filtrations of the standard n -ball

$$\mathbf{E}^n = e^0 \cup e^{n-1} \cup e^n$$

and $(n-1)$ -sphere $\mathbf{S}^{n-1} = e^0 \cup e^{n-1}$, then it is clear that

$$\mathbf{C}(n) \approx \pi \mathbf{E}^n \text{ and } \mathbf{S}^n(n-1) \approx \pi \mathbf{S}^{n-1}.$$

We now define a particular kind of morphism $j: A \rightarrow D$ which we call a *crossed complex morphism of relative free type*. Let A be any crossed complex. A sequence of morphisms $j_n: D^{n-1} \rightarrow D^n$ may be defined inductively as follows. Set $D^0 = A$. Supposing D^{n-1} given, choose any family of morphisms

$$f_n^\lambda: \mathbf{S}(m_\lambda-1) \rightarrow D^{n-1} \text{ for } \lambda \in \Lambda_n$$

and any m_λ , and to construct $j_n: D^{n-1} \rightarrow D^n$ form the pushout:

$$\begin{array}{ccc} \coprod_{\lambda \in \Lambda_n} \mathbf{S}(m_\lambda-1) & \xrightarrow{(f_n^\lambda)} & D^{n-1} \\ \downarrow & & \downarrow \\ \coprod_{\lambda \in \Lambda} \mathbf{C}(m_\lambda) & \longrightarrow & D^n \end{array}$$

Let $D = \text{colim}_n D^n$, and let $j: A \rightarrow D$ be the canonical morphism. We call $j: A \rightarrow D$ a *crossed complex morphism of relative free type*. The images x^{m_λ} of the elements c_{m_λ} in D are called *basis elements* of D relative to A . We can conveniently write:

$$D = A \cup \{x^{m_\lambda}\}_{\lambda \in \Lambda_n, n \geq 0}$$

and may abbreviate this in some cases. For example we may write $D = A \cup x^n \cup x^m$, analogously to standard notations for CW-complexes.

We remark that for $A = \emptyset$ we get by this construction the crossed complexes of *free type* which were considered in [9] under the name "free crossed complexes" and in [11].

If $\{x_\lambda\}_{\lambda \in \Lambda}$ are all the cells of the crossed complex of free type C , then

$$\mathbf{C}(n) \otimes C = (\mathbf{S}(n-1) \otimes C) \cup \{c_n \otimes x_\lambda\}_{\lambda \in \Lambda}.$$

Hence the morphism $\mathbf{S}(n-1) \otimes C \rightarrow C(n) \otimes C$ is also a morphism of relative free type. The functor $-\otimes C$ on $\mathbf{Czs}$ has a right adjoint, and so preserves pushouts. Let $f: A \rightarrow B$ be any morphism. If $j: A \rightarrow D$ is a morphism of relative free type, then so also is the pushout $\bar{j}: B \rightarrow Q$ of j and f . Therefore we get the following

PROPOSITION 1.1. *Let C be a crossed complex of free type. If $A \rightarrow D$ is a morphism of relative free type then $A \otimes C \rightarrow D \otimes C$ is also of relative free type. In particular if D is a crossed complex of free type then the tensor product $D \otimes C$ is also of free type. ■*

We now follow [10] in defining, for $n \geq 0$, the n -fold left homotopies $B \rightarrow C$ from a crossed complex B to a crossed complex C . These homotopies may also be taken to be the elements of $\mathbf{CRS}(B, C)$ in dimension n ([10], Proposition 3.3). A 0-fold left homotopy $B \rightarrow C$ is simply a morphism $B \rightarrow C$. For $n \geq 1$, an n -fold left homotopy $B \rightarrow C$ is to be a pair (H, f) , where $f: B \rightarrow C$ is a morphism of crossed complexes (the *base morphism* of the homotopy) and H is a map of degree n from B to C (i.e., $H: B_k \rightarrow C_{k+n}$ for each $k \geq 0$) satisfying

(i) $\beta H(b) = \beta f(b)$ for all $b \in B$;

(ii) if $b, b' \in B_1$ and $b + b'$ is defined, then

$$H(b + b') = H(b)^{f(b)} + H(b') ;$$

(iii) if $b, b' \in B_n$ ($n \geq 2$) and $b + b'$ is defined, then

$$H(b + b') = H(b) + H(b') ;$$

(iv) if $b \in B_n$ ($n \geq 2$), $b_1 \in B_1$ and b^{b_1} is defined, then

$$H(b^{b_1}) = H(b)^{f(b_1)} .$$

Let $\mathcal{D}$ be the crossed complex which has vertices 0, 1 and is freely generated by an element c_1 from 0 to 1 of dimension 1. Thus $\mathcal{D}$ may be regarded as a groupoid. Put $PC = \mathbf{CRS}(\mathcal{D}, C)$ for any crossed complex C . Then P is a functor on $\mathbf{Czs}$ and there are natural transformations

$$p^0, p^1: P \rightarrow \text{id}_{\mathbf{Crs}} \text{ and } s: \text{id}_{\mathbf{Crs}} \rightarrow P$$

such that

$$p^0 s = p^1 s = \text{id}_{\mathbf{Crs}} .$$

Hence the quadruple $\mathbf{P} = (P; p^0, p^1, s)$ forms a cohomotopy cohomotopy in the Kamps sense [18].

For crossed complexes B, C we have the induced cubical set $Q(B, C)$ such that $Q(B, C)_n = \mathcal{Czs}(B, P^n C)$, for $n \geq 0$. By [8], Corollary 9.6 and [10], Proposition 2.2

$$\mathcal{Czs}(B, P^n C) = \mathcal{Czs}(\otimes^n \mathcal{D}, \mathcal{CRS}(B, C)) = (\lambda \mathcal{CRS}(B, C))_n, \quad n \geq 0,$$

where λ is the functor inducing an equivalence of the category of crossed complexes and ω -groupoids ([7], Theorem 6.2). Therefore $Q(C, D)$, being an ω -groupoid ([8], Corollary 9.6), satisfies the Kan extension condition for all dimensions ([7], Proposition 7.2).

According to [18], the cohomotopy system $\mathbf{P}$ defines in $\mathcal{Czs}$ a notion of homotopy between morphisms. In fact, this is essentially the notion of 1-fold left homotopy given above. This notion of homotopy also leads to the notions of homotopy equivalence, Hurewicz fibration and Hurewicz cofibration, as defined in [18].

Following Kamps [18] and using the analogue of the well-known standard procedure for spaces we get

LEMMA 1.2. *For any crossed complex morphism $f: B \rightarrow C$ there exists the following factorisation*

$$\begin{array}{ccc} B & \xrightarrow{f} & C \\ & \searrow j & \nearrow q \\ & B \times_C P C & \end{array}$$

where q is a Hurewicz fibration and j is a strong deformation retract morphism. hence a homotopy equivalence. ■

Suppose C is a crossed complex and $p \in C_0$. Following Brown-Higgins [7], p. 258 and Howie [16] we define $\pi_0(C)$ to be the set of components of the groupoid C_1 . Define $\pi_1(C, p)$ to be the cokernel of $\delta_2: C_2(p) \rightarrow C_1(p)$ and, for $n \geq 2$, define $\pi_n(C, p)$ to be the subquotient $\text{Ker } \delta_n(p) / \text{Im } \delta_{n+1}(p)$ of $C_n(p)$.

A morphism $f: B \rightarrow C$ in $\mathcal{Czs}$ is said to be a *weak equivalence* if the induced maps

$$\pi_0 B \rightarrow \pi_0 C \text{ and } \pi_n(B, p) \rightarrow \pi_n(C, fp)$$

are isomorphisms, for all $n \geq 1$ and $p \in B_0$. It follows by standard arguments that any homotopy equivalence of crossed complexes is a weak equivalence.

2. CLOSED MODEL CATEGORY STRUCTURE ON Czs .

Recall that a morphism $f: G \rightarrow H$ of groupoids is a fibration [3] if, whenever $p \in G_0$ and $y \in H$ with $\delta^0 y = fp$, there exists $z \in G_1$ such that $fz = y$ and $\delta^0 z = p$.

This notion was extended to morphisms in Czs by Howie in [16] in the following way. A morphism $f: E \rightarrow B$ in Czs is a fibration if each groupoid morphism $f_n: E_n \rightarrow B_n$ ($n \geq 1$) is a fibration of groupoids. Other equivalent descriptions of fibrations in Czs will be given by Brown-Higgins [28] using the notion of what we call here "crossed complex of free type", which is the same notion as that of "free crossed complex" in [9]. A main fact we need is that if $\mathbf{X}$ is the skeletal filtration of a CW-complex, then the fundamental crossed complex $\pi\mathbf{X}$ of $\mathbf{X}$ is of free type.

PROPOSITION 2.1 [28]. *Let $f: E \rightarrow B$ be a morphism of crossed complexes. Then the following conditions are equivalent:*

- (i) *f is a fibration;*
- (ii) *(Covering homotopy property) if C is a crossed complex of free type, $g: C \rightarrow E$ is a morphism, $n \geq 1$, and (H', fg) is an n -fold left homotopy $C \rightarrow B$, then there is an n -fold homotopy $(H, g): C \rightarrow E$ such that $fH = H'$;*
- (ii)' *the covering property holds for $n = 1$;*
- (iii) *if C is a crossed complex of free type, then the induced morphism $f_*: CRS(C, E) \rightarrow CRS(C, B)$ is a fibration. ■*

Note that this Proposition implies that each Hurewicz fibration in Czs is a fibration.

A morphism which is both a fibration and a weak equivalence is said to be a *trivial fibration*.

We will say that a morphism $f: A \rightarrow D$ has the *left lifting property* (LLP) with respect to the class $\mathcal{F}$ of morphisms in Czs if the dotted arrow completion exists in any commutative square

of the form

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & E \\
 f \downarrow & \nearrow \text{dotted arrow} & \downarrow p \\
 D & \xrightarrow{\quad} & B
 \end{array}$$

where p is in the class $\mathcal{F}$. Similarly p has the *right lifting property* (RLP) with respect to $\mathcal{F}$ if the dotted arrow completion exists in any commutative square of the above form, where f is in $\mathcal{F}$.

Following Quillen [22] we define a *cofibration* in $\mathbf{Czs}$ to be a morphism which has the (LLP) with respect to trivial fibrations. *Trivial cofibrations* are morphisms which are cofibrations and weak equivalences. It is easily checked that in a pushout diagram

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & \bar{A} \\
 f \downarrow & & \downarrow \bar{f} \\
 D & \xrightarrow{\quad} & \bar{D}
 \end{array}$$

if f is a cofibration, then so also is $\bar{f}$ ([22], chap.II. §3).

Let $\emptyset$ (resp. $*$) denote the initial (resp. final) object of $\mathbf{Czs}$. An object C is called *cofibrant* if the unique morphism from $\emptyset$ to C is a cofibration. Not all crossed complexes are cofibrant. However for any C the unique morphism $C \rightarrow *$ is a fibration.

The next proposition, which is analogous to Proposition 2.1, is the key to our results on cofibrations. The proof uses explicitly the structure of $\mathbf{CRS}(B, C)$ defined in [10], and which is given above.

PROPOSITION 2.2. *The following are equivalent for a morphism $f: E \rightarrow B$ in $\mathbf{Czs}$:*

- (i) f is a trivial fibration:

(ii) f_0 is surjective; if $p, q \in E$ and $b \in B_1(f_0 p, f_0 q)$, then there is $e \in E_1$ such that $f_1 e = b$; if $n \geq 1$ and $d \in E_n$ satisfies $\delta^0 d = \delta^1 d$ for $n=1$, $\delta d = 0$ for $n \geq 2$, and $b \in B_{n+1}$ satisfies $\delta b = f_n d$, then there is

$$e \in E_{n+1} \text{ such that } f_{n+1} e = b \text{ and } \delta e = d ;$$

(iii) f has the RLP with respect to $\mathbf{S}(n-1) \rightarrow \mathbf{C}(n)$ for all $n \geq 0$;

(iii)' if C is a crossed complex of free type then f has the RLP with respect to $\mathbf{S}(n-1) \otimes C \rightarrow \mathbf{C}(n) \otimes C$ for all $n \geq 0$;

(iv) if C is a crossed complex of free type then the induced morphism $f_*: \text{CRS}(C, E) \rightarrow \text{CRS}(C, B)$ is a trivial fibration.

PROOF. The equivalences (ii) $\Leftrightarrow$ (iii) and (iii)' $\Leftrightarrow$ (iv), and the implication (iv) $\Rightarrow$ (i) are evident.

The implications (i) $\Rightarrow$ (ii) for $n \geq 2$ and (ii) $\Rightarrow$ (i) are straightforward and can be proved by standard procedures in homological algebra. So we prove only the non-Abelian case $n=1$ of (i) $\Rightarrow$ (ii).

Let $p, q \in E_0$ and $b \in B_1(f_0 p, f_0 q)$. By the fibration property there is an element u in $E_1(p, q')$, say, such that $f_1 u = b$. Hence $f_0 q' = f_0 q$ and the isomorphism $\pi_0(E) \rightarrow \pi_0(B)$ determines an element $v \in E_1(q', q)$. The isomorphism $\pi_1(E, q) \rightarrow \pi_1(B, f_0 q)$ shows that there is an element $w \in \pi_1(E, q)$ such that $f_1 w = -f_1 v$. Let $d = u + v + w$. Then $f_1 d = b$.

To prove (i) $\Rightarrow$ (iv) we assume (i) and show that the morphism

$$f_*: \text{CRS}(C, E) \rightarrow \text{CRS}(C, B)$$

satisfies the condition (ii), which can be represented diagrammatically by

$$\begin{array}{ccc} \mathbf{S}(n-1) & \xrightarrow{\hat{d}} & \text{CRS}(C, E) \\ \downarrow & \nearrow \hat{e} & \downarrow f_* \\ \mathbf{C}(n) & \xrightarrow{\hat{b}} & \text{CRS}(C, B) \end{array}$$

for $n \geq 0$, where the morphisms $\hat{b}$, $\hat{d}$ and $\hat{e}$ are defined by their images b , d , e respectively. For $n=0$, we write $\hat{H}$ for $\hat{b}(c_0)$. For $n=1$, we write g^0 , g for $\hat{d}(0)$, $\hat{d}(1)$ respectively and $(\hat{H}, fh)$ for

$\hat{b}(c_1)$, this last being a homotopy from fg^0 to fg . For $n \geq 2$, we write (K, g) for $\hat{d}(\delta c_n)$ and $(\bar{H}, fg)$ for $\hat{b}(c_n)$. Thus if $n = 2$,

$$\delta^0(K, g) = \delta^1(K, g) = g,$$

and for $n \geq 3$, $\delta(K, g) = 0_g$. Also for $n \geq 2$,

$$\delta(\bar{H}, fg) = f_*(K, g) = (fK, fg).$$

Recall that C is of free type. Let X_k be a basis for C_k , $k \geq 1$. We will construct by induction on $k \geq 0$ a family of maps $H_k: C_k \rightarrow E_{n+k}$.

If $n = 0$, then H_0 is to be a morphism $C \rightarrow E$. This H_0 is easily constructed on the basis X using the fact that $f: E \rightarrow B$ is a trivial fibration. Hence H_0 extends over C to give a morphism, also written $H_0: C \rightarrow E$. For $n \geq 1$, we require the explicit formulæ given in [10], Proposition 3.14 for the boundaries of n -fold left homotopies. These formulæ (α_k^n) are as follows:

If (H, g) is a 1-fold left homotopy from g^0 to g , so that $\delta^0(H, g) = g^0$, $\delta^1(H, g) = g$, then

$$\begin{aligned} (\alpha_0^n) \quad g^0(c) &= \delta^0 H_0(c) & \text{if } c \in C_0, \\ (\alpha_1^n) \quad g^0(c) &= H_0(\delta^0 c) + g(c) + \delta(H_0 c) - H_0(\delta^1 c) & \text{if } c \in C_1, \\ (\alpha_k^n) \quad g^0(c) &= [g(c) + H_0(\delta c) + \delta(H_0 c)]^{H_0(\beta c)} & \text{if } c \in C_k \ (k \geq 2). \end{aligned}$$

If $n \geq 2$ and (H, g) is an n -fold left homotopy, then $\delta(H, g) = (K, g)$, where

$$\begin{aligned} (\alpha_0^n) \quad K_0(c) &= \delta H_0(c) & \text{if } c \in C_0, \\ (\alpha_1^n) \quad K_1(c) &= (-1)^{n+1} H_0(\delta^0 c) g(c) + (-1)^n H_0(\delta^1 c) + \delta(H_1 c) & \text{if } c \in C_1, \\ (\alpha_k^n) \quad K_k(c) &= \delta H_k(c) + (-1)^{n+1} H_{k-1}(\delta c) & \text{if } c \in C_k \ (k \geq 2). \end{aligned}$$

The above formulae will be used with H, g, K replaced by $\bar{H}, fg, \bar{K}$ in $\text{CRS}(C, B)$, in order to construct an appropriate element (H, g) in $\text{CRS}(C, E)$. Thus for all $n, k \geq 0$ we require also

$$(\beta_k^n) \quad f_{k+n} H_k = \bar{H}_k.$$

Suppose now that H_i is defined for $0 \leq i \leq k-1$, so that (α_i^n) and (β_i^n) are satisfied for $0 \leq i \leq k-1$. Then H_k is defined using the fact that f is a trivial fibration and C is of free type. With the above information, the details are straightforward and are left to the reader. ■

COROLLARY 2.3. *Let C be a crossed complex of free type. Then the morphism $\mathbf{S}(n-1) \otimes C \rightarrow \mathbf{C}(n) \otimes C$ is a cofibration for all $n \geq 0$. In particular, C is cofibrant.* ■

COROLLARY 2.4. *Let $j: A \rightarrow D$ be a morphism of relative free type. Then j is a cofibration.*

PROOF. By the definition of relative free type, we are given that D is a colimit $\text{colim}_n D^n$, where $D^0 = A$ and each $j_n: D^{n-1} \rightarrow D^n$ is a pushout of a coproduct of inclusions of the form $\mathbf{S}(m_{\lambda}-1) \rightarrow \mathbf{C}(m_{\lambda})$. By Proposition 2.2 (iii), such inclusions are cofibrations. Hence j_n is a cofibration. Hence $j: A \rightarrow D$ is a cofibration. ■

To obtain a description of trivial cofibrations we need

LEMMA 2.5. (i) *Let C be a crossed complex. Then the canonical morphisms $p^0, p^1: PC \rightarrow C$ are trivial fibrations and the induced morphism $(p^0, p^1): PC \rightarrow C \times C$ is a fibration;*

(ii) *for any fibration $f: E \rightarrow B$ the induced morphism $(p^0, Pf): PE \rightarrow E \times_B PB$ is also a fibration.*

PROOF. (i) We prove only that the morphism $(p_0, p_1): PC \rightarrow C \times C$ is a fibration. (Using the same methods one can also show that the canonical morphisms $p^0, p^1: PC \rightarrow C$ are trivial fibrations.) For $n \geq 0$, the elements of $(PC)_n$ are the n -fold left homotopies $(H, f): \mathcal{D} \rightarrow C$. The formulæ for the boundary operators δ^0, δ^1 and δ of the crossed complex PC were given in the proof of Proposition 2.2. Let $n \geq 1$ and let

$$(x, x') \in C_n \times C_n, y \in (PC)_0 = C_1$$

be such that for $n=1$

$$\delta^0(x, x') = (p^0 y, p^1 y)$$

and for $n \geq 2$

$$(x, x') \in C_n(p^0 y) \times C_n(p^1 y).$$

We define a morphism $f: \mathcal{D} \rightarrow C$ and for $k \geq 0$ a family of maps $H: \mathcal{D}_k \rightarrow C_{k+n}$ as follows. For $n=1$ let $f(c_1) = -x+y+x'$ and for $n \geq 2$ let $f(c_1) = y$. We let $H: \mathcal{D}_k \rightarrow C_{k+n}$ be trivial for $k \geq 2$, and be given by $H(0) = x, H(1) = x'$ for $k=1$. Then for $n \geq 1$,

$$(p^0, p^1)(H, f) = (x, x'),$$

and for $n=1$,

$$\delta^0(H, f) = y.$$

(ii) The required property of the morphism

$$(p^0, Pf): PE \rightarrow E \times_B PB$$

is to be that for $n \geq 1$ there is a completion H of the following commutative square

$$\begin{array}{ccc}
 \mathbb{C}(n) \vee \mathcal{D} & \xrightarrow{\quad} & E \\
 \downarrow & \nearrow H & \downarrow f \\
 \mathbb{C}(n) \otimes \mathcal{D} & \xrightarrow{\quad} & B
 \end{array}$$

where $\vee$ means the union in $\mathbf{Czs}$ with base points $0 \in \mathbb{C}(n)$, $0 \in \mathcal{D}$ identified. But this completion exists by the fibration property of the morphism $f: E \rightarrow B$. ■

Now we can follow Quillen's proof ([22], p.3.4) to get:

PROPOSITION 2.6. *The following are equivalent for a morphism $j: A \rightarrow D$ in $\mathbf{Czs}$:*

- (i) *j is a trivial cofibration;*
- (ii) *j has the LLP with respect to the fibrations;*
- (iii) *j is a cofibration and a strong deformation retract morphism.* ■

It follows from Lemma 2.5 that each cofibration $j: A \rightarrow D$ is a Hurewicz cofibration, because j has the LLP with respect to the trivial fibration $p^0: PC \rightarrow C$, for any crossed complex C . Hence j has the homotopy extension property. We do not expect the converse to hold, since for example in chain complexes Hurewicz cofibrations are not necessarily cofibrations in the Quillen sense.

PROPOSITION 2.7. *Any morphism $f: A \rightarrow B$ in $\mathbf{Czs}$ may be factored $f = pj$ where j is of relative free type (and hence a cofibration) and p is a trivial fibration.*

PROOF. The essential fact needed to apply Quillen's *small object argument* ([22], chap. II, p. 3.3 and 3.4) is the characterisation by Proposition 2.2 of trivial fibrations in $\mathbf{Czs}$ by the RLP with respect to the set of morphisms $\{\mathbf{S}(n-1) \rightarrow \mathbb{C}(n)\}_{n \geq 0}$ where each $\mathbf{S}(n-1)$ is "sequentially small" in the sense that $\mathbf{Czs}(\mathbf{S}(n-1), -)$ preserves sequential colimits.

For completeness and the convenience of the reader we give more details. We are given $f: A \rightarrow B$. We construct a diagram

$$\begin{array}{ccccccc}
 A & \xrightarrow{j_0} & E^0 & \xrightarrow{j_1} & E^1 & \xrightarrow{j_2} & \dots \\
 & \searrow f & \searrow p_0 & & \nearrow p_1 & & \\
 & & & & B & &
 \end{array}$$

as follows. Let $E^{-1} = A$ and $p_{-1} = f$. Having obtained E^{n-1} , consider the set Λ of all commutative diagrams λ of the form

$$\begin{array}{ccc}
 \mathbf{S}(m_{\lambda}-1) & \xrightarrow{f_{\lambda}} & E^{n-1} \\
 \downarrow & & \downarrow p_{n-1} \\
 \mathbf{C}(m_{\lambda}) & \xrightarrow{g_{\lambda}} & B
 \end{array}$$

Define $j_n: E^{n-1} \rightarrow E^n$ by the pushout

$$\begin{array}{ccc}
 \coprod_{\lambda \in \Lambda_n} \mathbf{S}(m_{\lambda}-1) & \xrightarrow{(f_n^{\lambda})} & E^{n-1} \\
 \downarrow & & \downarrow j_n \\
 \coprod_{\lambda \in \Lambda} \mathbf{C}(m_{\lambda}) & \xrightarrow{i_n} & E^n
 \end{array}$$

Define $p_n: E^n \rightarrow B$ by

$$p_n j_n = p_{n-1} \text{ and } p_n i_n = (g_{\lambda}).$$

Let $E = \text{colim } E^n$, and let $j: A \rightarrow E$ and $p: E \rightarrow B$ be the canonical morphisms. By the above construction $j: A \rightarrow E$ is a morphism of relative free type. Proposition 2.2 implies that $p: E \rightarrow B$ is a trivial fibration. ■

If B is any crossed complex and $\emptyset \rightarrow B$ is the unique morphism from the initial object then from the above result we get:

COROLLARY 2.8. *Any crossed complex B is weakly equivalent to a crossed complex of free type.* ■

COROLLARY 2.9. *If $f: A \rightarrow D$ is a cofibration then it is a retract*

in the category of maps of Czs of a morphism of relative free type. In particular, each cofibrant object in Czs is a retract of a crossed complex of free type.

PROOF. By Proposition 2.7 $f = pj$ where j is of relative free type and p is a trivial fibration. Hence by the LLP of f with respect to trivial fibrations, there exists a completion of the following commutative square

$$\begin{array}{ccc}
 A & \xrightarrow{j} & E \\
 f \downarrow & \nearrow g & \downarrow p \\
 D & \xrightarrow{id_D} & B
 \end{array}$$

So the following diagram commutes

$$\begin{array}{ccccc}
 A & \xrightarrow{id_A} & A & \xrightarrow{id_A} & A \\
 f \downarrow & & j \downarrow & & f \downarrow \\
 D & \xrightarrow{g} & E & \xrightarrow{p} & D
 \end{array}$$

Hence $pg = id_D$ and f is a retract of j . ■

The next corollary generalises Corollary 2.6.

COROLLARY 2.10. *Let C be cofibrant. If $A \rightarrow D$ is a cofibration then $A \otimes C \rightarrow D \otimes C$ is also a cofibration. In particular, if D is cofibrant then $D \otimes C$ is also cofibrant.* ■

COROLLARY 2.11. *Any morphism $f: B \rightarrow C$ in Czs may be factored $f = pi$ where i is a trivial cofibration and p is a fibration.*

PROOF. By Lemma 1.1 $f = qj$ where q is a fibration and j is a homotopy equivalence. But by Proposition 2.7, $j = pi$ where p is a trivial fibration and i is a cofibration. Also j and p are weak equivalences, so i is a trivial cofibration. Finally,


$$f = qj = (qp)i,$$

where qp is a fibration and i is a trivial cofibration. ■

THEOREM 2.12. *The category Czs of crossed complexes, together with the distinguished classes of weak equivalences, fibrations and cofibrations defined above, satisfies the following axioms:*

CM1: Czs has all finite colimits and limits.


CM2: Suppose given a commutative diagram of the form



in Czs . If any two of f , g or h are weak equivalences, then so is the third.

CM3: The classes of cofibrations, fibrations and weak equivalences are closed under retraction in the category of maps of Czs .

CM4: Suppose given a commutative diagram



in Czs , where p is a fibration and j a cofibration. If either j or p is trivial, then there is a map $h:D \rightarrow E$ such that $ph = g$ and $hj = f$.

CM5: Any crossed complex morphism f may be factored as:

a) $f = pi$, where p is a fibration and i is a trivial cofibration and

b) $f = qj$, where q is a trivial fibration and j is a cofibration.

CM1-CM5 are the closed model axioms (cf. [22]); one says that the category of crossed complexes is a closed model category.

PROOF. CM1 follows, because of the Brown-Higgins result from [8] that $\mathbf{Czs}$ is a complete and cocomplete category.

CM2 and CM3 are completely trivial.

CM4 follows from the definition of cofibrations and Proposition 2.5. The factorisation axiom CM5 was proved by Proposition 2.7 and Corollary 2.11. ■

3. WHITEHEAD THEOREM FOR CROSSED COMPLEXES.

By Proposition 2.5 a cofibration $j: A \rightarrow D$ in $\mathbf{Czs}$ is a trivial cofibration if and only if it is a strong deformation retract. Now we prove a dual fact for fibrations of cofibrant objects in $\mathbf{Czs}$.

PROPOSITION 3.1. *If $p: E \rightarrow B$ is a fibration of cofibrant objects in $\mathbf{Czs}$ then the following are equivalent:*

- (i) p is a trivial fibration;
- (ii) p has the RLP with respect to cofibrations;
- (iii) p is a fibration and a strong deformation coretract.

PROOF. (i) $\Rightarrow$ (ii) follows from the definition of cofibration.

(ii) $\Rightarrow$ (i). In particular, p has the RLP with respect to $\mathbf{S}(n-1) \rightarrow \mathbf{C}(n)$ ($n \geq 0$), hence p is a trivial fibration by Proposition 2.2.

(iii) $\Rightarrow$ (i). This follows from the fact that a strong deformation coretract is a homotopy equivalence, and hence is a weak equivalence.

(i) $\Rightarrow$ (iii). Let $q: E \otimes \mathcal{D} \rightarrow E$ denote the canonical morphism. Thus q is the constant homotopy of the identity morphism on E . The coretract and strong deformation may be constructed by completing in

$$\begin{array}{ccc}
 \emptyset & \xrightarrow{\quad} & E \\
 \downarrow & \nearrow s & \downarrow p \\
 B & \xrightarrow{\text{id}_B} & B
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 E \amalg E & \xrightarrow{(sp, \text{id}_E)} & E \\
 \downarrow (i^0, i^1) & \nearrow h & \downarrow p \\
 E \otimes \mathcal{D} & \xrightarrow{pq} & B
 \end{array}$$

which is possible, since $\emptyset \rightarrow B$, by assumption, and

$$(i^0, i^1): E \amalg E \rightarrow E \otimes \mathcal{D}$$

by Corollary 2.3 for $n=1$ and Corollary 2.10, are cofibrations. ■

THEOREM 3.2 (*The Whitehead Theorem*). *If a morphism $f: C \rightarrow D$ of cofibrant objects in $\mathcal{C}s$ is a weak equivalence then f is also a homotopy equivalence.*

PROOF. By Theorem 2.12 we have the following factorisation of the morphism f :

$$\begin{array}{ccc} C & \xrightarrow{f} & D \\ & \searrow i \quad \nearrow p & \\ & \bar{C} & \end{array}$$

where p is a fibration and i is a trivial cofibration. But f is a weak equivalence and so p is a trivial fibration. Now C is a cofibrant object and $i: C \rightarrow \bar{C}$ is a trivial cofibration. So by lifting in the following diagram

$$\begin{array}{ccccc} \emptyset & \xrightarrow{\quad} & \emptyset & \xrightarrow{\quad} & E \\ \downarrow & & \downarrow & \nearrow \text{dashed} & \downarrow q \\ C & \xrightarrow{i} & \bar{C} & \xrightarrow{\quad} & B \end{array}$$

where q is a trivial fibration, we obtain that $\bar{C}$ is also a cofibrant object. By Proposition 2.5, i is a strong deformation retract and by Proposition 3.1, p is a strong deformation coretract and so, finally, f is a homotopy equivalence. ■

4. DOUBLE CATEGORY METHODS FOR CROSSED COMPLEXES.

Another view of abstract homotopy theory is given by Spencer and Spencer-Wong in [23,24]. Recall first that Gabriel-Zisman [14] derive exact sequences in homotopy theory in the context of a 2-category in which all 2-morphisms are invertible. Spencer shows in [23] that 2-categories are equivalent to *special double categories with connection* or with *thin structure* [24], where the thin squares of the double category derive from the

constant 2-morphisms of the 2-category.

Thus for crossed complexes one obtains a double category with thin structure from the 2-category of crossed complexes, morphisms of crossed complexes, and homotopies of morphisms. The general results of [23,24] now give the following applications to crossed complexes. Recall first that Vogt [25] has shown that for spaces *strong homotopy equivalence* is equivalent to homotopy equivalence. This result is placed in the abstract setting in [23], Proposition 3.1. So we obtain

PROPOSITION 4.1. *A homotopy equivalence of crossed complexes is also a strong homotopy equivalence. ■*

The paper [23] has results on homotopy pullback and homotopy pushout squares - for example, a composite of homotopy pushouts is a homotopy pushout. The paper [24] has results on homotopy commutative cubes and homotopy pushouts and pullbacks. For example, Corollary 4.8 of [24] and its dual apply to give cogluing and gluing theorems for homotopy equivalences of crossed complexes. Roughly speaking, *homotopy pullbacks (pushouts) of homotopy equivalences are homotopy equivalences.*

(4.2) OPEN PROBLEM. *Are homotopy pushouts of weak equivalences also weak equivalences ?*

Note that the paper [2] obtains a type of model structure, there called a cofibration category, for the category of reduced crossed complexes of free type. In this category, all objects are cofibrant, weak equivalences are homotopy equivalences and standard arguments show that homotopy pushouts of homotopy equivalences are homotopy equivalences. However in many cases one wishes to deal with the non-free case, and it is in this context that (4.2) remains open.

REFERENCES.

1. H.J. BAUES, *Algebraic homotopy*, Cambridge University Press, Cambridge, 1989.
2. H.J. BAUES, "The homotopy types of 4-dimensional complexes", Preprint, Bonn 1986.
3. R. BROWN, "Fibrations of groupoids", *J. Algebra* 15 (1970), 103-132.
4. R. BROWN, "Some non-abelian methods in homotopy theory and homological algebra", *Categorical topology, Proc. Conf. on Categorical Topology*, Toledo, Ohio 1983, ed. H.L. Bentley et al, Heldermann-Verlag, Berlin (1984), 108-146.
5. R. BROWN, "Non-Abelian cohomology and the homotopy classification of maps", *Homotopie algébrique et algèbre locale*, Conf. Marseille-Luminy 1982, ed. J.-M.Lemaire et J.-C. Thomas, *Astérisque* 113-114 (1984), 167-172.
6. R. BROWN, "Coproducts of crossed P-modules: applications to second homotopy groups and to the homology of groups", *Topology* 23 (1984), 337-345.
7. R. BROWN & P.J. HIGGINS, "The algebra of cubes", *J. Pure Appl. Algebra* 21 (1981), 233-260.
8. R. BROWN & P.J. HIGGINS, "Colimit theorems for relative homotopy groups", *J. Pure Appl. Algebra* 22 (1981), 11-41.
9. R. BROWN & P.J. HIGGINS, "Crossed complexes and non-Abelian extensions", *Int. Conf. on Category Theory*, Garmersbach (1981), *Lecture Notes in Math.* 962, Springer (1982), 39-50.
10. R. BROWN & P.J. HIGGINS, "Tensor products and homotopies for ω -groupoids and crossed complexes", *J. Pure Appl. Algebra* 47 (1987), 1-33.
11. R. BROWN & P.J. HIGGINS, "The classifying space of a crossed complex" (in preparation).
12. R. BROWN & P.J. HIGGINS, "The relation between crossed complexes and chain complexes with a groupoid of operators" (in preparation).
13. R. BROWN & J. HUEBSCHMANN, "Identities among relations", in *Low-dimensional Topology*, ed. R. Brown & T.L. Thickstun, London Math. Soc. Lecture Notes Series 48, C.U. P. (1982), 153-202.
14. P. GABRIEL & M. ZISMAN, *Calculus of fractions and homotopy theory*, Springer-Verlag, Berlin 1967.
15. D.F. HOLT, "An interpretation of the cohomology groups

- $H^n(G,M)$ ", *J. Alg.* 16 (1970), 307-318.
16. J. HOWIE, "Pullback functors and crossed complexes", *Cahiers de Top. et Géom. Diff.* XX (1979), 281-295.
 17. J. HUEBSCHMANN, "Crossed n -fold extensions and cohomology", *Comm. Math. Helv.* 55 (1980), 302-314.
 18. K.H. KAMPS, "Kan-Bedingungen und abstrakte Homotopietheorie", *Math. Z.* 124 (1972), 215-236.
 19. K.H. KAMPS & T. PORTER, "Abstract homotopy and simple homotopy theory", U.C.N.W. Pure Maths, Preprint 86.9.
 20. A.S.-T. LUE, "Cohomology of groups relative to a variety", *J. Alg.* 69,(1979), 319-320.
 21. S. MAC LANE, "Historical note", *J. Alg.* 69 (1979), 319-320.
 22. D.G. QUILLEN, "Homotopical algebra", *Lecture Notes in Math.* 43, Springer (1967).
 23. C.B. SPENCER, "An abstract setting for homotopy pushouts and pullbacks", *Cahiers de Top. et Géom. Diff.* XVIII (1977), 409-429.
 24. C.B. SPENCER & Y.L. WONG, "Pullback and pushout squares in a special double category with connection", *Cahiers de Top. et Géom. Diff.* XXIV (1983). 161-192.
 25. R.M. VOGT, "A note on homotopy equivalences", *Proc. A.M.S.* 32 (1972), 627-62.
 26. J.H.C. WHITEHEAD, "Combinatorial homotopy II", *Bull. A.M.S.* 55 (1949), 453-469.
 27. J.H.C. WHITEHEAD, "Simple homotopy type", *Amer. J. Math.* 72 (1950), 1-57.
 28. R. BROWN & P.J. HIGGINS, "Non-abelian cohomology with coefficients in a crossed complex" (in preparation).

R. BROWN :
 DEPT. OF PURE MATHEMATICS
 UNIVERSITY COLLEGE OF N.W.
 BANGOR. GWYNEDD LL57 2UW
 U.K

M. GOLASINSKI :
 INSTITUTE OF MATHEMATICS
 UNIVERSITY N. COPERNICUS
 UL. CHOPINA 12/18
 87-100 TORUN. POLAND