

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

F. J. KORKES

T. PORTER

Nilpotent crossed modules and pro- C completions

Cahiers de topologie et géométrie différentielle catégoriques, tome 31, n° 4 (1990), p. 277-282

http://www.numdam.org/item?id=CTGDC_1990__31_4_277_0

© Andrée C. Ehresmann et les auteurs, 1990, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**NILPOTENT CROSSED MODULES AND
Pro- \mathcal{C} COMPLETIONS**

by F.J. KORKES and T. PORTER

RÉSUMÉ. Cet article donne deux résultats sur les pro- \mathcal{C} complétions de groupes simpliciaux, où \mathcal{C} est une sous-catégorie pleine de la catégorie des groupes finis; on étudie leurs relations avec les pro- \mathcal{C} complétions de modules croisés et les actions nilpotentes.

In a earlier paper [2], we have considered the algebraic problem of studying the pro- \mathcal{C} completion of a crossed module and gave a "cofinality condition" that guaranteed that the pro- \mathcal{C} completion of the crossed module could be performed levelwise, i.e., by completing the two groups making up the crossed module individually.

It is a result of Mac Lane and Whitehead that certain equivalence classes of crossed modules correspond via an equivalence of categories with homotopy 2-types. i.e., with homotopy types that have zero homotopy groups above level 2. Any connected homotopy type, X , can be represented by a simplicial group $G(X)$, and any simplicial group G gives rise to a reduced simplicial set $W(G)$, so that these two functors, G and W , give an equivalence on homotopy categories. The work of Bousfield and Kan [1] on p -profinite completions of homotopy types includes the possible use of this adjointness to define such a completion functor algebraically. They also study nilpotent actions and nilpotent fibrations and the interplay of these ideas with their completion processes.

In this paper we investigate the connection between pro- \mathcal{C} completions of crossed modules, levelwise constructions of pro- \mathcal{C} completion of simplicial groups and nilpotent actions.

1. PRELIMINARIES.

In what follows \mathcal{C} will denote a non-trivial full subcategory of the category of finite groups. This category is assumed to be closed under the formation of subgroups, quotients and finite products. For any group G we let $\Omega(G)$ be the directed set

of normal subgroups N of G such that $G/N \in \mathbf{C}$. With this notation the group theoretic pro- \mathbf{C} completion of G is given by

$$\hat{G} = \varprojlim G/N$$

where the inverse limit is taken over the normal subgroups $N \in \Omega(G)$.

A *crossed module* consists of two groups C and G , an action of G on C and a homomorphism $\partial: C \rightarrow G$ satisfying the two conditions:

$$\text{CM1) for all } c \in C \text{ and } g \in G, \\ \partial(g \cdot c) = g \partial(c) g^{-1},$$

$$\text{CM2) for all } c_1, c_2 \in C \\ \partial(c_1) c_2 = c_1 c_2 c_1^{-1}.$$

Morphisms of crossed modules are pairs of homomorphisms preserving the action and giving a commutative square in the obvious way.

A crossed module is said to be a pro- \mathbf{C} crossed module if both groups are pro- \mathbf{C} groups, that is inverse limits of groups in \mathbf{C} , and the action and homomorphism are continuous in the inverse limit topology. Morphisms between two pro- \mathbf{C} crossed modules are continuous morphisms of the algebraic crossed modules underlying them.

There is an obvious forgetful functor from the category of pro- \mathbf{C} crossed modules to that of crossed modules, and in [2] we showed that that functor had a left adjoint which is a pro- \mathbf{C} completion functor. This completion functor is not just the group theoretic pro- \mathbf{C} completion of the two groups. The "bottom" group (G in the above) is sent to its pro- \mathbf{C} completion, but the "top" group on which G acts does not, in general, go to its pro- \mathbf{C} completion. This does happen, however, if the crossed module satisfies the following cofinality condition:

Let $\Omega_G(C)$ be the directed subset of $\Omega(C)$ given by those $N \in \Omega(C)$ which are G -equivariant. We say (C, G, ∂) satisfies the *cofinality condition* if $\Omega_G(C)$ is a cofinal subset of $\Omega(C)$.

We will also need some facts about simplicial groups. We recall that given a simplicial group G , the *Moore complex* (NG, ∂) of G , is the chain complex defined by

$$(NG)_n = \bigcap \{ \text{Ker } d_i^n \mid i \neq 0 \}$$

with $\partial_n: NG_n \rightarrow NG_{n-1}$ given by $\partial_n = d_0^n$ restricted to $(NG)_n$. The

image of ∂_{n+1} is normal in G_n and the homotopy groups of G can be calculated using this complex (NG, ∂) ; in fact,

$$\pi_n(G) \approx \bigcap_{i=0}^n \text{Ker } d_i^n / (d_0^{n+1} (\bigcap_{i=1}^{n+1} \text{Ker } d_i^{n+1})).$$

From G , we can also form a crossed module

$$\partial: (NG_1/d_0NG_2) \rightarrow G_0,$$

in which the "boundary" map ∂ is induced by that of the Moore complex. We will denote this crossed module by $M(G, 1)$ as it represents the 2-type of G . The cokernel of this crossed module is $\pi_0(G)$ and the kernel is $\pi_1(G)$ as is easily seen from the quoted facts about the Moore complex.

Given a simplicial group G , its pro- \mathcal{C} completion will be taken to be the pro- \mathcal{C} simplicial group obtained by applying the group theoretical pro- \mathcal{C} completion in each dimension. (Although this ties in with the definition of Bousfield and Kan, see [1] page 109, the reader should be warned that it does not necessarily coincide with the Artin-Mazur type completion which aims to pro- \mathcal{C} complete the homotopy groups rather than an algebraic model of the homotopy type. The two definitions will coincide in the presence of finiteness conditions.)

Finally we recall (again cf. [1]) the notion of a nilpotent action of a group G on a group C . An action of G on C is said to be *nilpotent* if there is a finite sequence

$$C = C_1 \supset \cdots \supset C_j \supset \cdots \supset C_n = \{e\}$$

of subgroups of C such that for each j

- (i) C_j is closed under the action of G ,
 - (ii) C_{j+1} is normal in C_j and C_j/C_{j+1} is abelian,
- and (iii) the induced G -action on C_j/C_{j+1} is trivial.

We will say that the G -nilpotent length of C in this case is less than or equal to n (denoted $\lambda_G \leq n$).

2. pro- \mathcal{C} COMPLETIONS OF SIMPLICIAL GROUPS AND OF CROSSED MODULES.

Bousfield and Kan proved ([1] page 113): "the homotopy type of $R_\infty X$ in dimensions $\leq k$ " depends only on "the homotopy type of X in dimensions $\leq k$ ". Here R_∞ is a completion functor and we will only be looking at this for the case of a pro- \mathcal{C} completion. For $k=1$, as the 1-type of X is determined by the

fundamental group, this is clear and relatively uninteresting. The next simplest case of this is given by the 2-type and thus, by Mac Lane and Whitehead, by a crossed module. One way of assigning a crossed module to a 2-type is via simplicial groups and the $\hat{M}(-, i)$ construction recalled earlier.

Our situation is not identical with that studied by Bousfield and Kan but we can view their result in a different light representing the 2-type by a crossed module, so it is natural to ask what is the exact relationship between the representing crossed module $M(\hat{G}, 1)$ of the 2-type of the pro- \mathcal{C} completion of a simplicial group G , and $M(G, 1)^\sim$, the crossed module pro- \mathcal{C} completion of $M(G, 1)$ as introduced by us in [2]. The answer is as nice as it could be.

PROPOSITION. *There is a natural isomorphism*

$$M(G, 1)^\sim \approx M(\hat{G}, 1).$$

PROOF. The nerve functor

$$E: \text{CMod} \rightarrow \text{Simp.Groups}$$

is defined as follows: If $M = (C, G, \partial)$ is a crossed module, then $E(M)$ is the simplicial group given as the nerve of the associated cat^1 -group, which is an internal category in the category of groups (see Loday [3]). In dimension 0, $E(M)$ is just G , in dimension 1, it is $C \rtimes G$, and in higher dimensions it is a multiple semi-direct product with many copies of C .

This simplicial group has Moore complex isomorphic to

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \cdots \rightarrow C \rightarrow G,$$

i.e., essentially giving us back M . Now let $T_{\geq 1}$ be the full (reflexive) subcategory of the category Simp.Groups defined by the condition that G is in it if and only if the Moore complex of G has trivial terms in dimensions 2 and above, i.e., $N(G)_i = \{1\}$ for each $i \geq 2$. The reflector $t_{\geq 1}: \text{Simp.Groups} \rightarrow T_{\geq 1}$ is defined by the condition that $N(t_{\geq 1}G)$ is the same as the truncation of $N(G)$ given by:

$$\begin{array}{ll} N(G)_0 & \text{in dimension 0,} \\ N(G)_1 / d_0 N(G)_2 & \text{in dimension 1,} \\ 1 & \text{in dimensions } \geq 2. \end{array}$$

One easily checks that $t_{\geq 1}G$ is isomorphic to $EM(G, 1)$ and that $M(-, 1)$ and E set up an equivalence of categories between CMod and $T_{\geq 1}$. Of course a similar thing happens with pro- \mathcal{C} crossed modules and a reflexive subcategory of pro- \mathcal{C} simplicial groups.

(The notation we will use for the various categories will, it is hoped, be self explanatory.)

Suppose that G , is a (discrete/abstract) simplicial group and M is a pro- \mathcal{C} crossed module, then

$$\text{Simp.Groups}(G, \mathcal{U}(EM)) \approx \text{Simp.pro-}\mathcal{C}(\hat{G}, EM)$$

since EM is a simplicial pro- \mathcal{C} group. This set is itself naturally isomorphic to $T_{\mathbb{1}}^{\mathcal{C}}(t_{\mathbb{1}}^{\mathcal{C}}(\hat{G}), EM)$. Since

$$M(t_{\mathbb{1}}^{\mathcal{C}}(\hat{G}), 1) \approx M(\hat{G}, 1),$$

this gives a natural isomorphism

$$\text{Simp.Groups}(G, \mathcal{U}(EM)) \approx \text{pro-}\mathcal{C}.\text{CMod}(M(\hat{G}, 1), M).$$

The forgetful functor $\mathcal{U}: \text{pro-}\mathcal{C} \rightarrow \text{Groups}$, or more exactly its simplicial and crossed modules extensions, satisfies $\mathcal{U}E \approx E\mathcal{U}$, so one also has

$$\begin{aligned} \text{Simp.Groups}(G, \mathcal{U}(EM)) &\approx \text{Simp.Groups}(G, E(UM)) \\ &\approx \text{CMod}(M(G, 1), UM) \approx \text{pro-}\mathcal{C}.\text{CMod}(M(G, 1)^{\sim}, M). \end{aligned}$$

We thus have that there is a natural isomorphism

$$M(G, 1)^{\sim} \approx M(\hat{G}, 1)$$

as required.

This clarifies and extends Bousfield and Kan's result in the case $k=2$, since for a reduced homotopy type X , the pro- \mathcal{C} completion $W\hat{G}X$ of X has a 2-type represented by $M(\hat{G}X, 1)$ which is isomorphic to the pro- \mathcal{C} completion of the crossed module $M(GX, 1)$, that represents the 2-type of X .

3. NILPOTENT CROSSED MODULES, COFINALITY CONDITIONS AND pro- \mathcal{C} COMPLETIONS.

Crossed modules occur in the work of Loday [3], linked closely to the study of fibrations: if $p: E \rightarrow B$ is a fibration with connected fibre F then the induced map from $\pi_1(F)$ to $\pi_1(E)$ makes $(\pi_1(F), \pi_1(E), p_*)$ into a crossed module. The preservation of certain crossed module structures by termwise pro- \mathcal{C} completion is thus reminiscent of the preservation of nilpotent fibrations by completions as exemplified by the nilpotent fibration lemma of Bousfield and Kan [1], and suggests there should be a link between nilpotent actions and cofinality conditions. The link is the following:

PROPOSITION. *If $M = (C, G, \partial)$ is a crossed module in which the action of G on C is nilpotent, then M satisfies the cofinality*

condition and hence M^\sim is isomorphic to $(\hat{C}, \hat{G}, \hat{\delta})$.

PROOF. The proof is by induction on the G -nilpotent length of C . First we note that if $W \triangleleft C$ is such that C/W is in \mathcal{C} , it is sufficient to prove that $\cap^{\mathcal{E}} W = V$, say, is such that C/V is in \mathcal{C} . If $\lambda_G = 1$, the group C is trivial. If $\lambda_G(C) = 2$, then the group C is abelian with trivial G -action. In neither case is there any difficulty.

Next suppose we have that the conclusion holds provided that $\lambda_G(C) < n$, more precisely we assume that if W is normal in C and $C/W \in \mathcal{C}$, then $V = \cap^{\mathcal{E}} W$ is also such that C/V is in \mathcal{C} . Now if C is such that $\lambda_G(C) = n$, there is a sequence

$$C = C_1 \supset \cdots \supset C_j \supset \cdots \supset C_n = \{e\}$$

as in the definition of Section 1. Taking the normal subgroup C_2 we get a short exact sequence

$$1 \rightarrow C_2 \rightarrow C_1 \xrightarrow{p} C_1/C_2 \rightarrow 1$$

in which $\lambda_G(C_2) < n$ and C_1/C_2 is abelian with trivial G -action.

Now suppose $W \triangleleft C$ is such that $W/C \in \mathcal{C}$. For any $g \in G$, $p(gW) = p(W)$, since the G -action on C_1/C_2 is trivial. Moreover

$$gW \cap C_2 = g(W \cap C_2),$$

since C_2 is closed under the G -action. Thus setting $V = \cap^{\mathcal{E}} W$, we get

$$p(V) = p(W) \text{ and } V \cap C_2 = g(W \cap C_2).$$

As $C_2/C_2 \cap W \in \mathcal{C}$, we apply the induction hypothesis to conclude that $C_2/C_2 \cap V \in \mathcal{C}$. Similarly the quotient of C_1/C_2 by $p(V)$ is in \mathcal{C} as it is the same as that by $p(W)$. The group C/V is thus part of an exact sequence, the other groups of which are in \mathcal{C} , hence it also is in \mathcal{C} as required.

BIBLIOGRAPHY.

1. BOUSFIELD, A. K. & KAN, D. M., Homotopy limits, completions and localizations, *Lecture Notes in Math.* 304, Springer (1972).
2. KORKES, F. J. & PORTER, T., pro- \mathcal{C} completions of crossed modules, *Proc. Edin. Math. Soc.* 33 (1990), 39-51.
3. LODAY, J. L., Spaces with finitely many homotopy groups, *J. Pure & Appl. Algebra* 24 (1982), 179-202.

F. J. KORKES:
DEPARTMENT OF MATHEMATICS
COLLEGE OF SCIENCE
BASRAH UNIVERSITY
BASRAH.

T. PORTER:
SCHOOL OF MATHEMATICS
UNIVERSITY COLLEGE OF
NORTH WALES
BANGOR LL57 1UT. U.K.