

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

J. L. FRITH

The category of uniform frames

Cahiers de topologie et géométrie différentielle catégoriques, tome 31, n° 4 (1990), p. 305-313

http://www.numdam.org/item?id=CTGDC_1990__31_4_305_0

© Andrée C. Ehresmann et les auteurs, 1990, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

THE CATEGORY OF UNIFORM FRAMES

by J. L. FRITH

RÉSUMÉ. Cet article étudie la catégorie des cadres uniformes et ses relations à d'autres catégories. En particulier, les relations à la catégorie des espaces uniformes décrites à l'aide des foncteurs ouvert et spectre sont examinées. Les cadres proximaux de Banaschewski lui sont aussi reliés de façon appropriée.

0. INTRODUCTION.

Frames (and locales, the dual category) are presently structures of considerable interest (see e.g. Johnstone [3]). This paper arises from a desire to consider frames with various uniform-type structures, the topic of the author's thesis. Frame-analogues of the open set, open cover structure of (covering) uniform spaces are considered. These have also been investigated by Pultr [5] where it is established that the uniformizable frames are precisely the completely regular ones.

In this paper, a functor from the category of completely regular frames to uniform frames leaving the underlying frame unchanged is constructed, and it is shown to be the coarsest possible such functor; the result of Pultr is a consequence. Open- and spectrum-type functors are constructed which lift the well-known situation for frames and topologies.

An application includes a demonstration that the proximal structures considered by Banaschewski [1] are just the totally bounded uniform frames, with a particularly simple proof which seems to be new in the spatial setting. In forthcoming papers, non-symmetric analogues of these uniform structures are considered.

The author wishes to thank his supervisor Professor K. Hardie and also Professor B. Banaschewski for considerable help and encouragement.

1. BACKGROUND.

A *frame* is a complete lattice satisfying the (infinite) distributive law:

$$a \wedge \bigvee x_\alpha = \bigvee (a \wedge x_\alpha).$$

Frame maps preserve top and bottom elements (denoted by 1 and 0 resp.), finite meets, arbitrary joins. The resulting category is *FRM*.

The topology of a topological space is a frame, functorially so (but contravariantly, continuous functions are mapped to the associated inverse image lattice map). This functor O (the *open* functor) from *TOP* to *FRM* is adjoint on the right to the (contravariant) *spectrum* functor $\Sigma: FRM \rightarrow TOP$ which may be defined as follows: the points of ΣL are the "points" of L , viz. $\text{hom}(L, 2)$, where 2 is the 2-element chain: the family

$$T_{\Sigma L} = \{\Sigma_a \mid a \in L\}, \quad \text{where } \Sigma_a = \{p \in \Sigma L \mid p(a) = 1\},$$

is the *spectral* topology. This dual adjunction of O and Σ restricts to a dual equivalence on the subcategories of sober spaces (join irreducible closed sets are unique singleton closures) and spatial frames (topologies). (See [3] for the details.)

Two binary relations we shall need are:

(i) $a \overline{\prec} b$ (a is *rather below* b) if there is an element s satisfying

$$a \wedge s = 0, \quad s \vee b = 1.$$

(ii) $a \overline{\prec\prec} b$ (a is *completely below* b) if there is a family of elements, $\{x_\alpha \mid \alpha \in \mathbb{Q} \cap [0, 1]\}$, satisfying

- (i) $x_0 = a, \quad x_1 = b,$
- (ii) $a < \beta \Rightarrow x_\alpha \overline{\prec} x_\beta.$

The completely below relationship interpolates: relabel $x_{1/2}$ as c ; it is clear that $a \overline{\prec\prec} c \overline{\prec\prec} b$. A frame L is *regular* if, for each b in L , $b = \bigvee \{a \mid a \overline{\prec} b\}$; it is *completely regular* if, for each $b \in L$, $b = \bigvee \{a \mid a \overline{\prec\prec} b\}$.

2. COVERS AND UNIFORM FRAMES.

$C \subseteq L$ is a *cover* of L if $\bigvee C = 1$. For C, D covers of L , we say that $C \leq D$ (C *refines* D) if, for each $c \in C$, there is $d \in D$ with $c \leq d$; $C \wedge D$ denotes the cover $\{c \wedge d \mid c \in C, d \in D\}$; for $a \in L$,

$$\text{st}(a, C) = \bigvee \{c \in C \mid c \wedge a \neq 0\}$$

and C^* denotes the cover $\{\text{st}(c, C) \mid c \in C\}$; we say that C *star refines* D if $C^* \leq D$.

DEFINITION 1. (a) Let L be a frame, q a non-empty family of covers of L . (L, q) is a *uniform frame* if:

PROPOSITION 4. *Every compact, completely regular frame has unique compatible uniform structure (the set of **all** covers).*

PROOF. Let L be compact, completely regular; L has at least one compatible uniform structure, q . Suppose C is a cover of L , for $c \in C$ we have

$$c = \bigvee \{t \mid \text{st}(t, D) \leq c \text{ for some } D \in q\}.$$

So

$$\{t \mid \text{st}(t, D) \leq c \text{ for } c \in C, D \in q\}$$

is a cover of L ; select a finite subcover $\{t_i \mid i = 1, 2, \dots, n\}$ of this cover. For each such i , there is $D_i \in q, c_i \in C$ with $\text{st}(t_i, D_i) \leq c_i$. Set $D_0 = \bigwedge D_i$; it is easy to see that D_0 refines C , so C is a uniform cover. ■

Johnstone [3] shows that if $a \overline{\overline{c}} b \in L$, then there is a frame map f from the frame of open sets of the unit interval, $O[0,1]$, to L satisfying $f[0,1] \leq a^*$, $f(0,1] \leq b$. As an immediate consequence, if $T: L \rightarrow (L, q_T(L))$ is functorial from the category of completely regular frames to the category of uniform frames, then every pair $\{a^*, b\}$, where $a \overline{\overline{c}} b$, is indeed a member of $q_T(L)$. This means that $q_C(L) \subset q_T(L)$, so q_C is the coarsest functorial way of constructing compatible uniform structures. It is appropriate then to call $q_C(L)$ the Cech uniformity on L . We turn now to the relationship of *UNIFORM* to other categories.

3. UNIFORM SPACES.

Isbell [2] is a good reference for uniform spaces via covers. Given (X, μ) a uniform space, we denote the canonical topology generated by μ by $T(\mu)$; the $T(\mu)$ -open covers of μ form a base for μ . (We require *no* separation conditions on (X, μ) .)

LEMMA 5. *Let (X, μ) be a uniform space, $U \in T(\mu)$. Then*

$$U = \bigcup \{V \in T(\mu) \mid \text{st}(V, C) \subset U \text{ for some } C \text{ (open) in } \mu\}.$$

PROOF. Suppose $x \in U$; for some $C \in \mu, \text{st}(\{x\}, C) \subset U$. Select a $T(\mu)$ -open member D of μ with D a star-refinement of C . Of course $\text{st}(\{x\}, D)$ is open, but it is also easy to show that we have $\text{st}(\text{st}(\{x\}, D), D) \subset U$, giving the required result. ■

With the help of the above lemma, it is easy to see that the $T(\mu)$ -open subsets of a uniform space (X, μ) together with

THE CATEGORY OF UNIFORM FRAMES

the (T, μ) -open covers give us a uniform frame; denote this uniform frame by $(OX, O\mu)$. In fact, it is almost obvious that we have a functor, O , (contravariant) from the category of uniform spaces and uniformly continuous functions, $UNIF$, to $UNIFORM$. We provide also a spectrum-type functor from $UNIFORM$ to $UNIF$.

DEFINITION 6. Let (L, q) be a uniform frame, ΣL the set of "points" of L . For $C \in q$, set

$$\Sigma C = \{\Sigma_c \mid c \in C\}; \cup \Sigma C = \cup \{\Sigma_c \mid c \in C\} = \Sigma_{\vee C} = \Sigma_1 = \Sigma L$$

so ΣC is indeed a cover of ΣL . Denote by Σq the family of all covers of ΣL refined by a cover of the form ΣC .

PROPOSITION 7. Let (L, q) be a uniform frame; then $(\Sigma L, \Sigma q)$ is a uniform space (possibly empty).

PROOF. We show only that if $C^* \leq D$ ($C, D \in q$), then ΣC star-refines ΣD . To see this, suppose $c \in C$;

$$\begin{aligned} \text{st}(\Sigma_c, \Sigma C) &= \cup \{\Sigma_t \mid t \in C, \Sigma_t \cap \Sigma_c \neq \emptyset\} = \cup \{\Sigma_t \mid t \in C, \Sigma_{t \wedge c} \neq \emptyset\} \\ &\subset \cup \{\Sigma_t \mid t \in C, t \wedge c \neq 0\} = \Sigma_{\text{st}(c, C)} \subset \Sigma_d \end{aligned}$$

for some $d \in D$ since $C^* \leq D$, so ΣC star-refines ΣD . ■

In fact, we can provide the claimed spectrum functor $\Sigma: UNIFORM \rightarrow UNIF$ as follows: suppose $f: (L, q) \rightarrow (M, q')$ is a uniform map. Then define

$$\Sigma f: (\Sigma M, \Sigma q') \rightarrow (\Sigma L, \Sigma q) \text{ by } \Sigma f(p) = p \circ f.$$

It is straightforward to see that Σf is uniformly continuous.

PROPOSITION 8. Let (L, q) be a uniform frame. The topologies $T(\Sigma q)$ and $T_{\Sigma L}$ on ΣL coincide.

PROOF. Suppose $U \in T(\Sigma q)$; then for each $p \in U$, there is $C \in q$ with $\text{st}(p, \Sigma C) \subset U$. But members of ΣC are $T_{\Sigma L}$ -open, so $U \in T_{\Sigma L}$. Conversely, suppose $U \in T_{\Sigma L}$: then $U = \Sigma_a$ for some $a \in L$; but

$$a = \vee \{b \in L \mid \text{st}(b, C) \leq a \text{ for some } C \in q\}$$

and $p \in \Sigma_a$ implies $p(a) = 1$, so for some $b \in L$ with $\text{st}(b, C) \leq a$, we must have $p(b) = 1$ (p is a frame map). Now

$$\text{st}(\{p\}, \Sigma C) = \cup \{\Sigma_c \mid c \in C, p(c) = 1\};$$

but $p(c) = 1$ and

$$p(b) = 1 \Rightarrow p(c \wedge b) = 1 \Rightarrow c \wedge b \neq 0 \Rightarrow c \leq \text{st}(b, C) \leq a \Rightarrow \Sigma_c \subset \Sigma_a .$$

This yields Σ_a as a $T(\Sigma q)$ -open set. (The case where L is a degenerate frame is no problem.) ■

THEOREM 9. *The two contravariant functors O and Σ are adjoint on the right.*

PROOF. One simply verifies that the two (well-known at the "topological" level) maps

$$\eta_L: L \rightarrow O\Sigma L \quad (\eta_L(a) = \Sigma_a),$$

$$\varepsilon_X: X \rightarrow \Sigma OX \quad (\varepsilon_X(x) = p_x \text{ [where } p_x(U) = 1 \Leftrightarrow x \in U])$$

are uniform maps (Proposition 8 says that we are lifting the topological level to the uniform level). Then η_L and ε_X are the required adjunction maps. ■

We naturally call a uniform frame (L, q) *spatial* if it satisfies $(L, q) \approx (OX, O\mu)$ for some uniform space (X, μ) . It is an easy matter to see that the separated uniform spaces and the spatial uniform frames are the fixed objects of the adjunction in Theorem 9; in particular this adjunction induces a dual equivalence on these two subcategories. One may also notice that $(\Sigma O, \Sigma O\mu)$ is the separated reflection of (X, μ) for any uniform space (X, μ) . Note that not all uniform frames are spatial; complete atomless Boolean algebras with the canonical uniform structure are not spatial.

4. PROXIMITY.

Let (L, q) be a uniform frame, as before. For $a, b \in L$, we set $a \ll_q b$ if for some $C \in q$, $st(a, C) \supseteq b$. It is an easy matter to prove:

- P1. $1 \ll_q 1, 0 \ll_q 0$.
- P2. $a \ll_q b \Rightarrow a \leq b$.
- P3. $a \leq b \ll_q c \leq d \Rightarrow a \ll_q d$.
- P4. $a_1, a_2 \ll_q b \Rightarrow a_1 \vee a_2 \ll_q b$; $a \ll_q b_1, b_2 \Rightarrow a \ll_q b_1 \wedge b_2$.
- P5. $a \ll_q b \Rightarrow$ there is $c \in L$ with $a \ll_q c \ll_q b$.
- P6. $a \ll_q b \Rightarrow b^* \ll_q a^*$.
- P7. $a = \bigvee \{ b \mid b \ll_q a \}$ for any $a \in L$.

The reader familiar with proximity spaces (see Naimpally & Warrack [4]) will realize that \ll_q exhibits properties reminiscent of the so-called inclusion relation on a proximity space. We de-

THE CATEGORY OF UNIFORM FRAMES

fine a *proximal frame* to be a pair (L, \ll) where L is a frame and \ll a binary relation on L which satisfies P1 to P7. These are considered by Banaschewski [1]. What we see is that we have constructed a functor from *UNIFORM* to the category of proximal frames and proximal maps, *PROXFRM*. (A map $f: (L, \ll) \rightarrow (L', \ll')$ is proximal if it is a frame map and $a \ll b \Rightarrow f(a) \ll' f(b)$.) We call \ll_q the proximal relation (on L) induced by q .

We consider the converse problem of endowing a proximal frame with a (functorial) compatible uniform structure (in the sense that the induced proximal relation is the original proximal relation).

THEOREM 10. *Let (L, \ll) be a proximal frame, a compatible uniform structure $q_{\ll}(L)$ exist such that $q_{\ll}(L)$ induces \ll .*

PROOF. Suppose $a \ll b \in L$. Set $C_a^b = \{a^*, b\}$; the fact that $a \ll b \Rightarrow a \bar{\ll} b$ ensures that C_a^b is a cover of L ; it is straightforward (mimicking Proposition 3) to see that we may use all such covers to generate a compatible uniform structure $q_{\ll}(L)$ on L . It remains to show that $q_{\ll}(L)$ induces \ll . Suppose $a \ll b$, then $st(a, C_a^b) \leq b$ and so $a \ll_{q_{\ll}(L)} b$. Conversely suppose, for some $C \in q_{\ll}(L)$, that $st(a, C) \leq b$. Our aim is to show that $a \ll b$. We may as well assume that

$$C = C_{a_1}^{b_1} \wedge \dots \wedge C_{a_n}^{b_n} \quad \text{where} \quad a_i \ll b_i \quad (i=1, \dots, n).$$

Using P5 repeatedly, select

$$c_i, d_i \in L \quad (i=1, \dots, n) \quad \text{with} \quad a_i \ll c_i \ll d_i \ll b_i$$

for each i . Note that then $c_i^* \ll a_i^*$. Let t be a typical element of C ; then

$$t = \bigwedge_{i \in I} a_i^* \wedge \bigwedge_{j \in J} b_j \quad \text{where } \{I, J\} \text{ is a partition of } \{1, 2, \dots, n\}.$$

Set $t^r = \bigwedge_{i \in I} c_i^* \wedge \bigwedge_{j \in J} d_j$; note that

$$t^r \in C_{c_1}^{d_1} \wedge \dots \wedge C_{c_n}^{d_n} \quad \text{and} \quad t^r \ll t.$$

Now

$$\begin{aligned} a &\leq \bigwedge \{s^* \mid s \in C, s \wedge a = 0\} = \left(\bigvee \{s \mid s \in C, s \wedge a = 0\} \right)^* \\ &\ll \left(\bigvee \{s^r \mid s \in C, s \wedge a = 0\} \right)^* \leq \bigvee \{s^r \mid s \in C, s \wedge a \neq 0\} \\ &\ll \bigvee \{s \mid s \in C, s \wedge a \neq 0\} = st(a, C), \text{ so } a \ll b. \quad \blacksquare \end{aligned}$$

In fact, if $f: (L, \ll) \rightarrow (L', \ll')$ is proximal, then

$$f: (L, q_{\ll}(L)) \rightarrow (L', q_{\ll'}(L'))$$

is a uniform map, so we have a functor U_{\ll} from *PROXFRM* to *UNIFORM* which is right inverse to the "inducing" functor mentioned at the beginning of §6.

We now establish that the category of totally bounded uniform frames is isomorphic to the category of proximal frames.

PROPOSITION 11. *Suppose (L, \ll) is a proximal frame, q a compatible uniform structure on L which furthermore induces \ll . Then $q_{\ll}(L) \subset q$.*

PROOF. Omitted.

PROPOSITION 12. *Let (L, \ll) be a proximal frame. Suppose q is a totally bounded compatible uniform structure which induces \ll , then $q \subset q_{\ll}(L)$.*

PROOF. Let $C \in q$; select $D \in q$ with $D^* \leq C$. Total boundedness means that there is a finite subset E of D which still covers L (E need not be a member of q). Suppose $E = \{d_1, d_2, \dots, d_n\}$. For each such d_i ($i = 1, \dots, n$), $st(d_i, D) \leq c_i$ (some $c_i \in C$), so in fact $d_i \leq c_i$. Now $C_{d_1}^{c_1} \wedge \dots \wedge C_{d_n}^{c_n}$ refines C (since $d_1^* \wedge \dots \wedge d_n^* = 0!$) showing that $C \in q_{\ll}(L)$. ■

COROLLARY 13. *For a proximal frame (L, \ll) is the unique compatible totally bounded uniform structure which induces \ll .*

The isomorphism of the two categories mentioned above is now clear. Open and spectrum type functors (adjoint) for proximal frames and proximity spaces exist which again lift the well-known adjunction for frames and spaces.

The referee is thanked for his suggestions.

THE CATEGORY OF UNIFORM FRAMES

REFERENCES.

1. BANASCHEWSKI B., Frames and compactifications, *Proc. International Symposium on Extension Theory of topological structures and its applications*, VEB Deutscher Verlag der Wissenschaften, 1969.
2. ISBELL J. R., *Uniform spaces*, A.M.S. Surveys 12, 1964.
3. JOHNSTONE P. T., *Stone spaces*, Cambridge Studies in advanced Math. 3, Cambridge Univ. Press, 1982.
4. NAIMPALLY S.A. & WARRACK B.D., *Proximity spaces*, Cambridge Tracts in Math. and Math. Physics 59, Cambridge Univ. Press, 1970.
5. PULTR A., Pointless uniformities I, *Comment. Math. Univ. Carolinae* 25 (1) (1984), 91-104.

The author acknowledges financial support from the South African Council for Scientific and Industrial Research via the Topology Research group at the University of Cape Town.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CAPE TOWN
PRIVATE BAG
7700 RONDEBOSCH
REPUBLIC OF SOUTH AFRICA