

# GROUPE DE TRAVAIL D'ANALYSE ULTRAMÉTRIQUE

YASUO MORITA

## **A non-archimedean analogue of the discrete series**

*Groupe de travail d'analyse ultramétrique*, tome 9, n° 3 (1981-1982), exp. n° J13, p. J1-J4

[http://www.numdam.org/item?id=GAU\\_1981-1982\\_\\_9\\_3\\_A14\\_0](http://www.numdam.org/item?id=GAU_1981-1982__9_3_A14_0)

© Groupe de travail d'analyse ultramétrique  
(Secrétariat mathématique, Paris), 1981-1982, tous droits réservés.

L'accès aux archives de la collection « Groupe de travail d'analyse ultramétrique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

A NON-ARCHIMEDEAN ANALOGUE OF THE DISCRETE SERIES

By Yasuo MORITA (\*)  
 [Tohoku University]

The purpose of this paper is to define a non-archimedean analogue of the discrete series of  $SL_2(\mathbb{R})$ . We will refer to what is conjectured, what can be proved, and what are the difficulties in studying our representations. For proofs and details, we quote MORITA-MURASE [5].

1. Classical case.

Let  $\mathbb{C}$  and  $\mathbb{R}$  be the complex number field and the real number field, respectively. For any field  $F$ , let  $P^1(F) = F \cup \{\infty\}$  be the one dimensional projective space over  $F$ . Then  $P^1(\mathbb{C}) - P^1(\mathbb{R})$  is the disjoint union of the upper half plane  $H_+ = \{z \in \mathbb{C}; \text{Im}(z) > 0\}$  and the lower half plane  $H_- = \{z \in \mathbb{C}; \text{Im}(z) < 0\}$ . For any integer  $s \leq -2$ , put

$$V_s^+ = \{f : H_+ \rightarrow \mathbb{C}; \text{analytic, } \|f\|_2^2 = \int_{H_+} |f(z)|^2 y^{-s-2} dx dy < \infty\},$$

$$\pi_s^+(g) f(z) = (bz + d)^s f\left(\frac{az + c}{bz + d}\right) \quad (g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R})),$$

where  $f \in V_s^+$  and  $z = x + iy$  ( $x, y \in \mathbb{R}$ ). Then  $V_s^+$  becomes a Hilbert space with the norm  $\|\cdot\|_2$  and  $\pi_s^+(g)$  is a unitary operator on  $V_s^+$ . Hence  $\pi_s^+$  defines a unitary representation of the locally compact group  $SL_2(\mathbb{R})$  on  $V_s^+$ . We can also define  $\pi_{-1}^+$  by modifying the norm suitably. Further, if we use  $H_-$  instead of  $H_+$ , then we obtain another unitary representation  $\pi_s^-$  for any integer  $s \leq -1$ . It is well known :

THEOREM C. - The  $\pi_s^\pm$  are irreducible representations of  $SL_2(\mathbb{R})$ , and no two of them for various  $s$  are equivalent.

2. Definition of  $\pi_s$  in p-adic cases.

Now we are going to construct p-adic analogues of the  $\pi_s^\pm$ . We replace  $\mathbb{R}$  by a finite extension  $L$  of the p-adic number field  $\mathbb{Q}_p$ , and replace  $\mathbb{C}$  by a non-archimedean field  $(k, |\cdot|)$  containing  $L$  such that  $k$  is complete with respect

---

(\*) Yasuo MORITA, Mathematical Institut, Tohoku University, SENDAI 980 (Japan).

to  $k$  and algebraically closed. Hence we consider continuous representations of  $G = \text{SL}_2(L)$  on linear topological  $k$ -vector spaces.

Put  $D = P^1(k) - P^1(L)$ . Since  $L$  is locally compact,  $P^1(L)$  is compact. Hence, for any positive integer  $m$ ,  $P^1(L)$  is covered by  $\{z \in P^1(k) ; |z| > |p^{-m}|\}$  and a finite number of mutually disjoint open balls

$$\{z \in P^1(k) ; |z - a_i| < |p^m|\} \quad (a_i \in L, i = 1, \dots, N).$$

Put

$$D_m = \{z \in P^1(k) ; |z| \leq |p^{-m}|, |z - a_i| \geq |p^m| \quad (i = 1, \dots, N)\}.$$

Then  $\{D_m\}_{m=1}^\infty$  is an increasing sequence of subsets of  $D$  and  $D = \bigcup_m D_m$ . Let  $\mathcal{O}(D_m)$  be the space of  $k$ -valued functions on  $D_m$  of the form

$$f(z) = \sum_{m=0}^\infty c_m z^m + \sum_{i=1}^N \sum_{m=-1}^\infty c_m^{(i)} (z - a_i)^m,$$

where  $c_m$  and  $c_m^{(i)}$  are elements of  $k$ , and we assume that this limit converges (uniformly) on  $D_m$ . It is known that  $\mathcal{O}(D_m)$  becomes a Banach space with the supremum norm  $|f| = \sup_{D_m} |f(z)|$ .

We say that a  $k$ -valued function  $f$  on  $D$  is analytic if, and only if, the restriction of  $f$  to each  $D_m$  belongs to  $\mathcal{O}(D_m)$ . (This is the definition of analytic functions in the theory of rigid analytic spaces (cf. MORITA [4]).) Let  $V$  be the space of all  $k$ -valued functions on  $D$ . Hence  $V$  is the projective limit of the  $\mathcal{O}(D_m)$  with respect to the restriction map  $\mathcal{O}(D_{m+1}) \rightarrow \mathcal{O}(D_m)$ . Therefore  $V$  has a natural Fréchet topology. In particular,  $V$  is a complete linear topological space. We consider this space  $V$  as the analogue of  $V_s^\pm \oplus V_s^\pm$  for any  $s$ .

Let  $s$  be a negative integer. For any  $f \in V$ , put

$$\pi_s(g) f(z) = (bz + d)^s f\left(\frac{az + c}{bz + d}\right) \quad (g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G).$$

Then  $\pi_s$  defines a continuous representation of the locally compact group  $G$  on the Fréchet space  $V$ . This is our analogue of the classical discrete series.

### 3. Conjecture.

Let  $U$  be the space of rational functions of  $z$  (with coefficients in  $k$ ) which have no poles in  $D$ . Then  $U$  is a dense subspace of  $V$ . Further the subspace  $U$  of  $V$  is  $G$ -invariant because  $(bz + d)^s$  belongs to  $U$  for any  $g \in G$ .

Since  $k$  is algebraically closed, any element  $f$  of  $U$  can be expressed as a partial fractional series of the form

$$f(z) = \sum_{m=0}^\infty d_m z^m + \sum_{j=1}^n \sum_{m=-1}^\infty d_m^{(j)} (z - b_j)^m \quad (\text{a finite sum})$$

$(d_m, d_m^{(j)} \in k, n \geq 1, b_j \in L)$ . We define a subspace  $U_s$  of  $U$  by

$$U_s = \{f \in U; d_m^{(j)} = 0 \text{ for any } j \text{ and } m \text{ with } 0 > m > s\}.$$

Then we can prove that  $U_s$  is a closed  $G$ -invariant subspace of  $U$ . Let  $V_s$  be the closure in  $V$  of the subspace  $U_s$ . It is obvious that  $V_s$  is a closed  $G$ -invariant subspace of  $V$ . Hence we obtain representations of  $G$  on  $V_s$  and  $V/V_s$ .

Now we have the following conjecture (cf. Theorem C) :

CONJECTURE V.

- (i)  $V_s$  and  $V/V_s$  are (topologically) irreducible  $G$ -modules.
- (ii) No two of them for various  $s$  are  $G$ -equivalent.

#### 4. The result and remarks.

Though we can not prove the conjecture now, we can prove the corresponding assertion for the dense subspace  $U$  of  $V$ .

Let  $\mathfrak{g} = \{X \in M_2(L); \text{tr}(X) = 0\}$ , and put

$$\exp(X) = \sum_{n=0}^{\infty} \frac{X^n}{n!} \text{ for any } X \in \mathfrak{g}.$$

Then this series converges in  $M_2(L)$  to an element of  $G$  if the eigenvalues  $\pm \lambda$  of  $X$  satisfies  $|\lambda| < |p^{1/(\beta-1)}|$ . It follows that

$$(d\pi_s)(X) f(z) = \lim_{t \rightarrow 0} \frac{1}{t} \{\pi_s(\exp(tX)) f(z) - f(z)\}$$

is well-defined for any  $X \in \mathfrak{g}$  and  $f \in V$ , and  $d\pi_s$  is a representation of the Lie algebra  $\mathfrak{g}$  on the space  $V$ . It is obvious that any closed  $G$ -invariant subspace of  $U$  is  $\mathfrak{g}$ -invariant.

Let  $O$  be the integer ring of  $L$ , and put  $K = SL_2(O)$ . Then  $K$  is a maximal compact subgroup of  $G$ . Since  $K$  is an open subgroup of  $G$ , a closed  $K$ -invariant subspace of  $U$  is  $(\mathfrak{g}, K)$ -invariant.

Now our main result can be stated as :

THEOREM U.

- (i)  $U_s$  and  $U/U_s$  are algebraically irreducible  $(\mathfrak{g}, K)$ -modules (i. e. they have no nontrivial  $(\mathfrak{g}, K)$ -invariant  $k$ -subspaces).
- (ii) No two of them for various  $s$  are isomorphic as  $\mathfrak{g}$ -modules.

Obviously this theorem implies topological irreducibilities of  $U_s$  and  $U/U_s$ , and the non-equivalence of them.

Remark.

- (i). The difficulties in proving the conjecture lie in the fact that Shur's lemma

does not hold in our case. For example, we can prove that the only intertwining operators of  $V$  are the scalar operators, though  $V$  has the closed  $G$ -invariant subspace  $V_{\mathfrak{g}}$ . We prove our theorem by constructing the total space from any non-zero element (cf. MORITA-MURASE [5], 3-3).

(ii). It is remarkable that  $U_{\mathfrak{g}}$  is infinite dimensional and irreducible as a  $K$ -module though  $K$  is a compact group. This phenomenon causes a difficulty in defining the admissible representations.

#### REFERENCES

- [1] Automorphic forms, representations and L-functions, part 1-2. - Providence, American mathematical Society, 1979 (Proceedings of Symposia in pure Mathematics, 33).
- [2] GEL'FAND (I. M.), GRAEV (M. I.) and VILENKIN (N. Ya). - Generalized functions, vol. 5: Integral geometry and representation theory. - New York, London, Academic Press, 1966
- [3] GERRITZEN (L.) and VAN DER PUT (M.). - Schottky groups and Mumford curves. - Berlin, Heidelberg, New York, Springer-Verlag, 1980 (Lecture Notes in Mathematics, 817).
- [4] MORITA (Y.). - Analytic functions on an open subset of  $\mathbb{P}^1(k)$ , J. für reine und angew. Math., t. 311-312, 1979, p. 361-383.
- [5] MORITA (Y.) and MURASE (A.). - Analytic representations of  $SL_2$  over a  $p$ -adic number field, J. of Fac. of Sc., Univ. of Tokyo, Section 1A, t. 28, 1982, p. 891-905.

-----