

J. A. NITSCHÉ

Interior error estimates for semi-discrete Galerkin approximations for parabolic equations

RAIRO. Analyse numérique, tome 15, n° 2 (1981), p. 171-176

http://www.numdam.org/item?id=M2AN_1981__15_2_171_0

© AFCET, 1981, tous droits réservés.

L'accès aux archives de la revue « RAIRO. Analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**INTERIOR ERROR ESTIMATES
FOR SEMI-DISCRETE GALERKIN APPROXIMATIONS
FOR PARABOLIC EQUATIONS (*) (**)**

by J A NITSCHÉ (1)

Abstract — The initial boundary value problem for the heat equation in a domain Ω and the corresponding standard Galerkin method is considered. A certain regularity of the initial data in some sub-domain Ω_1 leads to the same regularity of the solution in Ω_1 and for all times. It is shown that the error between the exact solution and the Galerkin approximation is also of (almost) optimal order in the interior of Ω_1 . Of course certain properties of the underlying approximation spaces are needed, they are typical for finite elements.

Resume — On considere le probleme aux limites avec conditions initiales pour l'equation de la chaleur dans un domaine Ω , ainsi que l'approximation habituelle de Galerkin correspondante. Une certaine regularite des donnees initiales dans un sous-domaine Ω_1 conduit a la même regularite de la solution dans Ω_1 pour tous les temps. On montre que l'erreur entre la solution exacte et l'approximation de Galerkin est aussi d'ordre (presque) optimal dans l'interieur de Ω_1 . Naturellement, certaines proprietes des espaces d'approximations sont utilisees, qui sont caracteristiques des espaces d'elements finis.

1. In order to avoid technical details we restrict ourselves to the model problem

$$\begin{aligned} \dot{u} &= \Delta u & \text{in } \Omega \times (0, T], \\ u &= 0 & \text{on } \partial\Omega \times (0, T], \\ u_{t=0} &= v & \text{in } \Omega \end{aligned} \tag{1}$$

The boundary of $\Omega \subseteq \mathbb{R}^N$ is assumed to be sufficiently smooth. With the help of a finite element space $S_h \subseteq H_1(\Omega)$ the Galerkin approximation $u_h = u_h(t) \in S_h$ is defined by

$$\begin{aligned} (\dot{u}_h, \chi) + D(u_h, \chi) &= 0 \quad \text{for } \chi \in S_h \wedge t > 0, \\ u_h|_{t=0} &= P_h v \end{aligned} \tag{2}$$

(*) Reçu en janvier 1980

(**) Presented at the Conference on Progress in the Theory and Practice of the Finite Element Method, Göteborg, Sweden, August 27-29, 1979

(1) Institut für Angewandte Mathematik, Albert-Ludwigs-Universität, Freiburg, R F A

Here (\cdot, \cdot) resp. $D(\cdot, \cdot)$ is the L_2 -inner product resp. the Dirichlet integral and (for simplicity) P_h is the L_2 -projector onto S_h .

For the corresponding elliptic problem

$$\begin{aligned} -\Delta u &= f & \text{in } \Omega, \\ u &= 0 & \text{on } \partial\Omega \end{aligned} \quad (3)$$

interior estimates of the error $e = u - u_h$ of the Ritz approximation $u_h \in S_h$ defined by

$$D(u_h, \chi) = (f, \chi) \quad \text{for } \chi \in S_h \quad (4)$$

were derived in [1], [5], [6], [7]. They are of the following type (*):

Assume $f \in L_2(\Omega)$ and in addition $f \in H_{k-2}(\Omega_1)$ for some domain $\Omega_1 \subseteq \Omega$ and $k > 2$. Further let Ω_2 be contained properly in Ω_1 . If S_h is of degree r with $r \geq k$ then the error e is of order k in Ω_2 , i.e.

$$\|e\|_{\Omega_2} \leq ch^k \{ \|f\|_{k-2, \Omega_1} + \|f\| \}. \quad (5)$$

The norms of f could be replaced by the appropriate norms of u because of the shift theorem.

Now let us assume the regularity

$$v \in L_2(\Omega) \cap H_k(\Omega_1) \quad (6)$$

and let $\Omega_2 \subset \subset \Omega_1$. Then the solution of (1) has the regularity

$$u \in L_\infty(L_2(\Omega)) \cap L_2(H_1(\Omega)) \cap L_\infty(H_k(\Omega_1)) \quad (7)$$

with the abbreviation $L_p(Z) = L_p(0, T; Z)$ for some $T > 0$ fixed. Corresponding to (5) we would expect in the parabolic case an estimate of the type

$$\|e\|_{L_\infty(L_2(\Omega_2))} \leq ch^k \{ \|v\|_{k, \Omega_1} + \|v\| \}. \quad (8)$$

In the next two sections we prove this error bound for $k < r$ (being the degree of S_h). In case $k = r$ the factor ch^k is to be replaced by $c_\varepsilon h^{r-\varepsilon}$ with $\varepsilon > 0$ arbitrary small.

This problem was already treated in Thomee [9]. There the local error in the $L_\infty(L_2(\Omega_2))$ norm is bounded by the $L_2(L_2(\Omega_1))$ norm besides of a remainder. Although this result does not give the final answer it turns out to be the main step. With respect to the notations as already mentioned as well as to the

(*) We use the notations of [7] resp. [9].

assumptions on S_h we refer to [9]. Since we do not extend our result to difference quotients the uniformity of the subdivisions in defining S_h is not necessary of course.

2. We start with an interior estimate for u_h , see lemma 3.3 in [9] :

LEMMA 1 (Thomée) : Let u_h be the solution of (2). Further assume

$$\hat{\Omega}_2 \subset \subset \Omega_1 \subseteq \Omega$$

and let $q > 0$ be fixed. Then

$$\| u_h(t) \|_{\hat{\Omega}_2}^2 \leq c \left\{ \| P_h v \|_{\Omega_1}^2 + \int_0^t [\| u_h \|_{\Omega_1}^2 + h^q \| \dot{u}_h \|_{\Omega_1}^2] d\tau \right\}. \tag{9}$$

Because of

$$\| \dot{u}_h \|_{\Omega_1}^2 \leq \| \dot{u}_h \|^2 = -\frac{1}{2} \partial_t \| u_h' \|^2 \tag{10}$$

we get

$$\int_0^t \| \dot{u}_h \|_{\Omega_1}^2 d\tau \leq \frac{1}{2} \| u_h'(0) \|^2. \tag{11}$$

Further we have

$$\begin{aligned} \| u_h'(0) \| &\leq ch^{-1} \| u_h(0) \| \\ &\leq ch^{-1} \| v \|. \end{aligned} \tag{12}$$

In this way the last term in (9) may be replaced by $h^{q-2} \| v \|^2$.

Next let us assume there is an additional domain Ω_0 according to

$$\Omega_1 \subset \subset \Omega_0 \subseteq \Omega.$$

Using the result of [6] or of Douglas *et al.* [3] we may estimate

$$\| P_h v \|_{\Omega_1} \leq c \{ \| v \|_{\Omega_0} + h^r \| v \| \}. \tag{13}$$

By replacing Ω_0 by Ω_1 we get with $q = 2r + 2$.

LEMMA 2 : Let $\Omega_2 \subset \subset \Omega_1 \subseteq \Omega$. Then

$$\| u_h(t) \|_{\Omega_2}^2 \leq c \left\{ \| v \|_{\Omega_1}^2 + h^{2r} \| v \|^2 + \int_0^t \| u_h \|_{\Omega_1}^2 d\tau \right\}. \tag{14}$$

Now we use an induction argument. Let Ω_1, Ω_2 with $\Omega_2 \subset \subset \Omega_1$ and $p \in \mathbb{N}$ be given. We can choose domains Ω^ν according to

$$\Omega_2 = \Omega^p \subset \subset \Omega^{p-1} \subset \subset \dots \subset \subset \Omega^0 = \Omega_1. \quad (15)$$

Repeated application of (14) and the interchange of the order of integration leads to

$$\|u_h(t)\|_{\Omega_2}^2 \leq c \left\{ \|v\|_{\Omega_1}^2 + h^{2r} \|v\|^2 + \int_0^t (t-\tau)^{p-1} \|u_h\|_{\Omega_1}^2 d\tau \right\}. \quad (16)$$

Since anyway

$$\|u_h\|_{\Omega_1} \leq \|u_h\| \leq \|v\| \quad (17)$$

we get :

LEMMA 3 : Assume $\Omega_2 \subset \subset \Omega_1 \subseteq \Omega$ and let $p > 0$ be arbitrary. Then

$$\|u_h(t)\|_{\Omega_2} \leq c \{ \|v\|_{\Omega_1} + (t^p + h^r) \|v\| \}. \quad (18)$$

This is the counterpart of the a priori estimate

$$\|u(t)\|_{\Omega_2} \leq c \{ \|v\|_{\Omega_1} + t^p \|v\| \} \quad (19)$$

for the solution of (1) which is easily derived (for instance using the exponential decay of the fundamental solution).

3. Now we are ready to prove (8). Let $\Omega_2 \subset \subset \Omega_1$ be fixed. We choose an Ω' according to

$$\Omega_2 \subset \subset \Omega'_0 \subset \subset \Omega_1. \quad (20)$$

Next let ω be a cut-off function with respect to Ω', Ω_1 , i.e. $\omega \in C^\infty(\Omega)$ with $0 \leq \omega \leq 1$ and

$$\omega = \begin{cases} 1 & \text{in } \Omega' \\ 0 & \text{in } \Omega - \Omega_1 \end{cases}. \quad (21)$$

We will use the splitting

$$v = v^1 + v^2 := \omega v + (1 - \omega) v \quad (22)$$

and denote by u^i, u_h^i the solutions of (1), (2) with the initial data v^i .

We have

$$u = u^1 + u^2, \quad u_h = u_h^1 + u_h^2. \tag{23}$$

The regularity assumption (6) leads to $v^1 \in H_k(\Omega)$ and

$$\|v^1\|_k \leq c \|v\|_{k,\Omega_1} \tag{24}$$

Therefore (see e.g. Bramble *et al.* [2])

$$\|u^1 - u_h^1\| \leq ch^k \|v\|_{k,\Omega_1} \tag{25}$$

is guaranteed. On the other hand v^2 is in $L_2(\Omega)$ with $\|v^2\| \leq \|v\|$ and v^2 vanishes in Ω' . The estimates (18), (19) for u^2 and u_h^2 with Ω_1 replaced by Ω' give

$$\|u^2 - u_h^2\|_{\Omega_2} \leq c(t^p + h^r) \|v\|. \tag{26}$$

In this way

$$\|e(t)\|_{\Omega_2} = \|u - u_h\|_{\Omega_2} \leq c \{ h^k \|v\|_{k,\Omega_1} + (t^p + h^r) \|v\| \} \tag{27}$$

is shown. In order to eliminate the time dependence we make use of the time dependent error estimate

$$\|e(t)\| \leq ch^r t^{-r/2} \|v\| \tag{28}$$

due to Helfrich [4], Thomee [8] which in connection with (27) gives

$$\|e(t)\|_{\Omega_2} \leq c \{ h^k \|v\|_{k,\Omega_1} + [h^r + \text{Min}(t^p, h^r t^{-r/2})] \|v\| \}. \tag{29}$$

The minimum is maximal for $t = c_p h^\alpha$ with $\alpha = 2r/(r + 2p)$ leading to

$$\text{Min}(t^p, h^r t^{-r/2}) = ch^\beta \tag{30}$$

with

$$\beta = \frac{2pr}{2p+r} = r - \frac{r^2}{2p+r}. \tag{31}$$

For any $k < r$ we may choose p such that $\beta \geq k$.

Then (29) gives (8). In case of $k = r$ and $\varepsilon > 0$ fixed we can choose p such that $\beta \geq r - \varepsilon$. Since the choice of p influences the number of iterations the constant in (8) then depends on p resp. ε .

Remark : In the previous estimates we did not control the constants. The-

refore we did not take into account a contribution of $(p - 1)^{-1}$ in the denominator of the integral in (16) By a careful analysis the "garbage" term $h^{-\varepsilon}$ in (8) in case of $k = r$ could be replaced by a logarithmic one

REFERENCES

- 1 J H BRAMBLE, J NITSCHÉ and A SCHATZ, *Maximum-norm interior estimates for Ritz-Galerkin methods* Math Comp 29, 677-688 (1975)
- 2 J H BRAMBLE, A SCHATZ, V THOMÉE and L B WAHLBIN, *Some convergence estimates for semidiscrete Galerkin type approximations for parabolic equations* SIAM J Numer Anal 14, 218-241 (1977)
- 3 J JR DOUGLAS, T DUPONT and L B WAHLBIN, *The stability in L^q of the L^2 -projection into finite element function spaces* Numer Math 23, 193-197 (1975)
- 4 H-P HELFRICH, *Fehlerabschätzungen für das Galerkinverfahren zur Lösung von Evolutionsgleichungen* Manuscripta Math 13, 219-235 (1974)
- 5 J NITSCHÉ, *Interior error estimates of projection methods* Proceedings Equadiff 3, Czechoslovak Conference on Differential Equations and their Applications, 235-239, Brno (1972)
- 6 J NITSCHÉ and A SCHATZ, *On local approximation properties of L_2 -projection on spline-subspaces* Appl Anal 2, 161-168 (1972)
- 7 J NITSCHÉ and A SCHATZ, *Interior estimates for Ritz-Galerkin methods* Math Comp 28, 937-958 (1974)
- 8 V THOMÉE, *Some convergence results for Galerkin methods for parabolic boundary value problems* Proceedings of a Symposium on Mathematical Aspects of Finite Elements in Partial Differential Equations, Madison, Wisc, Apr 1-3, 1974, C de Boor ed, Academic Press, 55-88 (1974)
- 9 V THOMÉE, *Some interior estimates for semidiscrete Galerkin approximations for parabolic equations* Math Comp 33, 37-62 (1979)