

DAVID RUELLE

Integral representation of measures associated with a foliation

Publications mathématiques de l'I.H.É.S., tome 48 (1978), p. 127-132

http://www.numdam.org/item?id=PMIHES_1978__48__127_0

© Publications mathématiques de l'I.H.É.S., 1978, tous droits réservés.

L'accès aux archives de la revue « Publications mathématiques de l'I.H.É.S. » (<http://www.ihes.fr/IHES/Publications/Publications.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

INTEGRAL REPRESENTATION OF MEASURES ASSOCIATED WITH A FOLIATION

by DAVID RUELLE

To Jean Leray

Let M be a compact differentiable manifold, \mathcal{F} a foliation of codimension k , and \mathcal{S} the set of open submanifolds of dimension k transversal to \mathcal{F} . A *transverse measure* ρ for \mathcal{F} is a collection of real measures ρ_Σ on the $\Sigma \in \mathcal{S}$, such that these measures correspond to each other by the *canonical isomorphisms* defined by \mathcal{F} . For a discussion of these notions, and applications, see Plante [8], Ruelle and Sullivan [10], Schwartzmann [11], Edwards, Millett and Sullivan [6], Sullivan [14], Garnett [7]. We note that we can, as in [10], assume that \mathcal{F} is only a partial foliation of M , and that the orientation assumptions of [10] are unnecessary here.

We generalize the notion of transverse measure by introducing measures associated with a cocycle. We call *cocycle* a family (f_τ) indexed by the canonical isomorphisms, such that:

(a) If τ maps $\Sigma \in \mathcal{S}$ onto $\Sigma' \in \mathcal{S}$, then f_τ is a continuous function on Σ' with strictly positive real values.

(b) If τ' maps Σ' onto Σ'' , then:

$$f_{\tau' \circ \tau} = f_{\tau'} \cdot (f_\tau \circ \tau'^{-1}).$$

We say that a collection $\rho = (\rho_\Sigma)$ of real measures on the $\Sigma \in \mathcal{S}$ is a *measure associated with the cocycle* (f_τ) , or is a (f_τ) -*measure*, if the image of the measure ρ_Σ by $\tau : \Sigma \rightarrow \Sigma'$ is $f_\tau \cdot \rho_{\Sigma'}$. Otherwise stated:

$$f_\tau = \frac{d(\tau\rho_\Sigma)}{d\rho_{\Sigma'}} \quad \text{a.e.}$$

for each local isomorphism $\tau : \Sigma \rightarrow \Sigma'$. The transverse measures are those associated with the trivial cocycle (1_τ) . The notion of (f_τ) -measure occurs naturally in the work of Connes [5]; see also Bowen [1].

The (f_τ) -measures form a real vector space \mathcal{J} . We call *vague topology* the topology defined on \mathcal{J} by the semi-norms:

$$\rho \mapsto |\rho_\Sigma(\varphi)|$$

where φ is a real continuous function with compact support in $\Sigma \in \mathcal{S}$. We write $\rho \geq 0$ if $\rho_\Sigma \geq 0$ for all $\Sigma \in \mathcal{S}$. With these definitions \mathcal{J} is an ordered topological vector space.

Choose $\Sigma \in \mathcal{S}$ and a compact set $K \subset \Sigma$. There is a map α_K of \mathcal{J} in the space $\mathcal{C}(K)^*$ of measures on K , such that $\alpha_K \rho$ is the restriction of ρ_Σ to K . The map α_K is linear and order-preserving.

Lemma 1. — *Let Σ, K be such that each leaf of \mathcal{F} intersects the interior of K in Σ . Then α_K is an isomorphism of the ordered vector space \mathcal{J} onto a subspace of $\mathcal{C}(K)^*$ closed for the vague topology.*

Remember that the vague topology is the w^* -topology of $\mathcal{C}(K)^*$ as dual of the space $\mathcal{C}(K)$ of real continuous functions on K . Note that α_K need not be continuous for the vague topologies.

To prove the lemma we remark that if K' is compact in $\Sigma' \in \mathcal{S}$, there are finitely many open L_i in Σ' covering K' , and canonical isomorphisms $\tau_i : L_i$ into Σ such that the closure of $\tau_i L_i$ lies in the interior of K . Therefore, using a partition of unity, and the fact that ρ is associated with the cocycle (f_τ) , we obtain an order preserving map π from the continuous functions on Σ' with support in K' to the continuous functions on Σ with support in K , such that $\rho_{\Sigma'}(\varphi) = \rho_\Sigma(\pi\varphi) = (\alpha_K \rho)(\pi\varphi)$. Thus α_K is injective, and $\rho \geq 0$ if and only if $\alpha_K \rho \geq 0$. Furthermore, if $\alpha_K \rho$ tends to a limit vaguely, $\alpha_{K'} \rho$ also converges vaguely, hence ρ converges vaguely, and the limit is obviously associated with the cocycle (f_τ) .

Lemma 2. — *Let \mathcal{G} be a linear subspace of the space $\mathcal{C}(K)^*$ of real measures on the compact set K . If $\rho \in \mathcal{G}$ implies $|\rho| \in \mathcal{G}$, then the cone \mathcal{G}_+ of positive measures in \mathcal{G} is simplicial. If ρ, ρ' belong to distinct extremal generatrices of the cone \mathcal{G}_+ , they are disjoint measures.*

Remember that a cone C in a real vector space is simplicial if the order that it defines on itself is a lattice (any two points have a min and a max). The easy proof of Lemma 2 is left to the reader.

Theorem. — *The cone C of positive elements of \mathcal{J} is simplicial. If ρ, ρ' belong to distinct extremal generatrices of C , then the measures $\rho_\Sigma, \rho'_\Sigma$ are disjoint for all $\Sigma \in \mathcal{S}$.*

In view of Lemma 1, the theorem immediately follows from Lemma 2 applied to $\mathcal{G} = \alpha_K \mathcal{J}$.

The cone C is closed and has a basis B which is convex, compact, and metrizable. For instance, if Σ, K are as in Lemma 1, let φ have compact support in Σ , $\varphi \geq 0$, and $\varphi(x) = 1$ if $x \in K$; one can take:

$$B = \{ \rho \in \mathcal{J} : \rho \geq 0 \text{ and } \rho(\varphi) = 1 \}.$$

According to Choquet's theory [4], the theorem implies that each $\rho \geq 0$ has a unique integral representation in terms of extremal elements of B :

$$\rho = \int_B \sigma m_\rho(d\sigma)$$

where m_ρ is carried by the set of extremal points of B . The arbitrariness in the choice of B corresponds to the fact that there is no natural normalization of positive (f_τ) -measures, but all choices of B give equivalent decompositions. If ρ is an extremal point of some B (i.e. if $\rho \neq 0$ and ρ belongs to an extremal generatrix of C) we say that ρ is a *pure (f_τ) -measure* (respectively a *pure transverse measure* in the case of the trivial cocycle). The theorem gives thus a unique decomposition of (f_τ) -measures into pure (f_τ) -measures, and states that two pure (f_τ) -measures are either proportional or disjoint ⁽¹⁾.

Given a positive (f_τ) -measure ρ , we let \mathcal{A}_ρ be the algebra of classes of bounded real functions on M which are constant on leaves of \mathcal{F} , and such that their restriction to each $\Sigma \in \mathcal{S}$ is ρ_Σ -measurable. Two functions are in the same class if their restrictions to each $\Sigma \in \mathcal{S}$ are equal ρ_Σ -almost everywhere.

Proposition. — *A positive (f_τ) -measure ρ is pure if and only if \mathcal{A}_ρ is trivial (consisting of the constant functions).*

If ρ is not pure, let $\rho = \rho^1 + \rho^2$ with non proportional (f_τ) -measures $\rho^1, \rho^2 \geq 0$. Choose Σ, K as in Lemma 1, and let $\sigma^i = \rho_\Sigma^i - \inf(\rho_\Sigma^1, \rho_\Sigma^2)$. There are ρ_Σ measurable functions $\psi_1, \psi_2 \geq 0$ such that $\sigma^i = \psi^i \rho_\Sigma$. We have $\psi^1 + \psi^2 \neq 0$ (because

$$\sigma^1 + \sigma^2 = \sup(\rho_\Sigma^1, \rho_\Sigma^2) \neq 0)$$

and $\psi_1 \cdot \psi_2 = 0$ a.e. (because σ^1, σ^2 are disjoint). Choosing some Riemann metric d on the leaves of \mathcal{F} , let:

$$\Psi^i(x) = \lim_{n \rightarrow \infty} \min\{\psi_i(y), y \in K, d(x, y) \leq n\}.$$

Clearly Ψ^1, Ψ^2 belong to \mathcal{A}_ρ and are not proportional, so that \mathcal{A}_ρ is non trivial. Conversely, if \mathcal{A}_ρ is non trivial, it is immediate that ρ is not pure.

Interpretation of the decomposition. — Let h be a diffeomorphism of a compact manifold B , and \mathcal{F} be the foliation by the orbits of the suspension of h . We identify B with a submanifold of codimension 1 of M , transverse to \mathcal{F} . The transverse measures of \mathcal{F} correspond then to the h -invariant measures on B . The pure transverse measures correspond to the h -ergodic measures, and the decomposition into pure transverse measures corresponds to the ergodic decomposition. The integral representation of positive (f_τ) -measures appears thus as an extension of ergodic theory. A different, deeper, relation is with the theory of Gibbs states in statistical mechanics, as discussed in the following example.

Example. — Let $A \in \text{SL}_n(\mathbf{Z})$ be hyperbolic, i.e. the spectrum of A is disjoint from $\{z : |z| = 1\}$. Let V^s (respectively V^u) be the subspace of \mathbf{R}^n associated with the eigenvalues less than 1 (respectively larger than 1) in absolute value. We call \hat{A} the map

⁽¹⁾ For cases where there is only one pure (f_τ) -measure, see Bowen and Marcus [2], and also the Example below.

induced by A on $\mathbf{T}^n = \mathbf{R}^n / \mathbf{Z}^n$, and W^s, W^u the images of V^s, V^u in \mathbf{T}^n . It is readily seen that $G = W^s \cap W^u$ is a n -generator subgroup of \mathbf{T}^n , G is dense in \mathbf{T}^n because W^s, W^u are dense.

Choose $a_1, \dots, a_n \in \mathbf{R}^n$ such that their images in \mathbf{T}^n are generators of G . Write $a_i = (a_{i1}, \dots, a_{in})$, take $x = (x_1, \dots, x_n) \in \mathbf{R}^n$, and define:

$$\Gamma_x = \left\{ \left(x_1 + \sum_i t_i a_{i1}, \dots, x_n + \sum_i t_i a_{in}, t_1, \dots, t_n \right) \in \mathbf{R}^{2n} : t_1, \dots, t_n \in \mathbf{R} \right\}.$$

The images F_x of the Γ_x in $M = \mathbf{T}^n \times \mathbf{T}^n$ constitute a codimension n foliation of M , with holonomy group G with respect to the Section $\mathbf{T}^n = \mathbf{T}^n \times \{0\}$. We shall define functions $f_\tau : \mathbf{T}^n \rightarrow \mathbf{R}$ when $\tau \in G$, i.e. for the canonical isomorphisms of the section \mathbf{T}^n . It is easy to extend this definition to that of a cocycle for \mathcal{F} .

Let φ be a real Hölder continuous function on \mathbf{T}^n . We let:

$$f_\tau(x) = \exp \sum_{k=-\infty}^{\infty} (\varphi(\hat{A}^k \tau^{-1} x) - \varphi(\hat{A}^k x)).$$

There is one and only one measure ρ associated with this cocycle. In fact:

$$\rho_{\mathbf{T}^n} = \lim_{m \rightarrow +\infty} \frac{1}{N_m} \left(\exp \sum_{k=-m}^m \varphi(\hat{A}^k x) \right) dx$$

where dx is Haar measure on \mathbf{T}^n , and N_m a normalizing factor. These statements have their origin in a relation between statistical mechanics and differentiable dynamical systems introduced by Sinai: $\rho_{\mathbf{T}^n}$ is a *Gibbs state* for the function φ (see Sinai [13], Capocaccia [3], Ruelle [9], Chapter 7). We notice that if $\varphi = 0$ then $\rho_{\mathbf{T}^n} = dx$, and uniqueness follows from the fact that G is a dense subgroup of \mathbf{T}^n . For the general case the reader is referred to the papers quoted above.

In view of the frequent non-uniqueness of Gibbs states we conjecture that, for the foliation discussed here, there exist cocycles with several non proportional associated measures.

Invariance under a diffeomorphism. — Let g be a diffeomorphism of M preserving \mathcal{F} (i.e. permuting the leaves). Suppose that (f_τ) is a cocycle *compatible* with g , i.e. such that:

$$f_{g \circ \tau \circ g^{-1}} = f_\tau \circ g^{-1}.$$

This condition is for instance always satisfied by the trivial cocycle (I_τ) .

If $\rho = (\rho_\Sigma)$ is a (f_τ) -measure, then $g\rho = (g\rho_{g^{-1}\Sigma})$ is again a (f_τ) -measure. This is because:

$$\begin{aligned} \tau(g\rho_{g^{-1}\Sigma}) &= g(g^{-1} \circ \tau \circ g) \rho_{g^{-1}\Sigma} = g(f_{g^{-1}\tau g} \rho_{g^{-1}\Sigma}) \\ &= g((f_\tau \circ g) \cdot \rho_{g^{-1}\Sigma}) = f_\tau \cdot (g\rho_{g^{-1}\Sigma}). \end{aligned}$$

Thus $g\mathcal{J} = \mathcal{J}$, and in fact $g\mathbf{C} = \mathbf{C}$, where \mathbf{C} is the cone of positive measures in \mathcal{J} . Suppose $\mathcal{J} \neq 0$, and let B be a compact basis of \mathbf{C} . We have $B = \mathbf{C} \cap \{\rho : \lambda(\rho) = 1\}$

for some continuous linear functional ρ on \mathcal{F} . The map $\rho \mapsto g\rho/\lambda(g\rho)$ has a fixed point $\rho_0 \in B$. Therefore $g\rho_0 = \lambda_0\rho_0$, where $\lambda_0 = \lambda(g\rho_0) > 0$, and λ_0 is in general different from 1.

Consider now the case of the trivial cocycle, *i.e.* of transverse measures. Under suitable conditions, discussed in [10], [14], λ_0 is an eigenvalue of the action of g on cohomology, and the corresponding class is associated with a geometric current determined by ρ_0 . If the class is nonzero, λ_0 is thus an algebraic number (in fact, a unit in the ring of algebraic integers).

Question: under what conditions do the numbers λ_0 associated with the transverse measures of a foliation form a finite set of algebraic numbers? A. Connes has pointed out to me that this is not always the case.

Diffeomorphisms which expand leaves. — Let the foliation \mathcal{F} contain a leaf with polynomial growth (*i.e.* the Riemann volume of a ball $B(x, r) \subset L$ increases polynomially with its radius r) then Plante [8] has shown that \mathcal{F} has a transverse measure $\rho \neq 0$ with support in the closure of L .

If the diffeomorphism g preserves \mathcal{F} and expands the leaves (*i.e.* multiplies sufficiently small distances on leaves, with respect to some Riemann metric, by a factor $> C > 1$), then the leaves have polynomial growth. This was proved by Sullivan and Williams [15]; see also Shub [12]. In particular \mathcal{F} has a transverse measure $\rho \neq 0$, and by the preceding Section we may assume that $g\rho_0 = \lambda_0\rho_0$. We recover thus a result stated in another context by Sullivan (see [14], III, 13): *if the diffeomorphism g preserves \mathcal{F} and expands the leaves, there is a transverse measure $\rho_0 \neq 0$ such that $g\rho_0 = \lambda_0\rho_0$.*

Acknowledgements. — My thanks are due to Dennis Sullivan who convinced me that measures associated with a foliation are interesting, and to Alain Connes who explained to me his beautiful recent results.

REFERENCES

- [1] R. BOWEN, *Anosov foliations are hyperfinite*. Preprint.
- [2] R. BOWEN and B. MARCUS, *Unique ergodicity for horocycle foliation*. Preprint.
- [3] D. CAPOCACCIA, A definition of Gibbs state for a compact set with Z^v action, *Commun. Math. Phys.*, **48** (1976), 85-88.
- [4] G. CHOQUET et P.-A. MEYER, Existence et unicité des représentations intégrales dans les convexes compacts quelconques, *Ann. Inst. Fourier*, **13** (1963), 139-154.
- [5] A. CONNES. Unpublished.
- [6] R. EDWARDS, K. MILLETT and D. SULLIVAN, Foliations with all leaves compact, *Topology*, **16** (1977), 13-32.
- [7] L. GARNETT, *An ergodic theory for foliations*. Preprint.
- [8] J. PLANTE, Foliations with measure preserving holonomy, *Ann. Math.*, **102** (1975), 327-362.
- [9] D. RUELLE, *Thermodynamic formalism*, Addison-Wesley, Reading, Mass., 1978.

- [10] D. RUELLE and D. SULLIVAN, Currents, flows and diffeomorphisms, *Topology*, **14** (1975), 319-327.
- [11] S. SCHWARTZMANN, Asymptotic cycles, *Ann. Math.*, **66** (1957), 270-284.
- [12] M. SHUB, Endomorphisms of compact differentiable manifolds, *Amer. J. Math.*, **91** (1969), 175-199.
- [13] Ia. G. SINAI, Gibbsian measures in ergodic theory, *Uspehi Mat. Nauk*, **27**, n° 4 (1972), 21-64. English translation, *Russian Math. Surveys*, **27**, n° 4 (1972), 21-69.
- [14] D. SULLIVAN, Cycles for the dynamical study of foliated manifolds and complex manifolds, *Inventiones math.*, **36** (1976), 225-255.
- [15] D. SULLIVAN and R. F. WILLIAMS, On the homology of attractors, *Topology*, **15** (1976), 259-262.

IHES/PM/77/181,
mai 1977.