RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

L. CARLITZ

Another Saaschützian theorem for double series

Rendiconti del Seminario Matematico della Università di Padova, tome 34 (1964), p. 200-203

<http://www.numdam.org/item?id=RSMUP_1964__34__200_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1964, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (http://rendiconti.math.unipd.it/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

ANOTHER SAASCHÜTZIAN THEOREM FOR DOUBLE SERIES

Nota *) *di* L. CARLITZ (*a Durham*)

Put

$$S(m, n) = \sum_{r=0}^{m} \sum_{s=0}^{n} \frac{(-m)_r(-n)_s(\alpha)_{r+s}(\beta)_r(\beta')_s}{r!\, s!\, (\gamma)_{r+s}(\delta)_r(\delta')_s}.$$

The writer [1] has proved that if

$$\text{(1)} \qquad \begin{cases} \gamma = \beta + \beta', \\ \gamma + \delta = \alpha + \beta - m + 1, \\ \gamma + \delta' = \alpha + \beta' - n + 1, \end{cases}$$

then

$$\text{(2)} \qquad S = \frac{(\beta + \beta' - \alpha)_{m+n}(\beta')_m(\beta)_n}{(\beta + \beta')_{m+n}(\beta' - \alpha)_m(\beta - \alpha)_n}.$$

Moreover (2) is equaivalent to the transformation formula

$$\text{(3)} \qquad F_1(\gamma - \alpha;\ \beta', \beta;\ \gamma;\ x, y) = (1 - x)^{\beta' - \alpha}(1 - y)^{\beta - \alpha} F_1(\alpha;\ \beta, \beta';\ \gamma;\ x, y),$$

where $\gamma = \beta + \beta'$ and

$$F_1(\alpha;\ \beta, \beta';\ \gamma;\ x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n}(\beta)_m(\beta')_n}{m!\, n!\, (\gamma)_{m+n}} x^m y_n.$$

---

*) Pervenuto in Redazione il 31 ottobre 1963.
Indirizzo dell'A.: Depart. of Mathematics - Duke University - *Durham*, North Carolina, U.S.A.

It may be of interest to note another result somewhat similar to (2). Put

$$S(n) = \sum_{r+s \geqslant n} \frac{(-n)_{r+s}(\alpha)_r(\alpha')_s(\beta)_r(\beta')_s}{r!\, s!\,(\gamma)_{r+s}(\delta)_r(\delta')_s}.$$

Then

$$S(n) = \sum_{r=0}^{n} \frac{(-n)(\alpha)(\beta)_r}{r!\,(\gamma)_r(\delta)_r} \sum_{s=0}^{n-r} \frac{(-n+r)(\alpha')(\beta')_s}{s!\,(\gamma+r)_s(\delta')_s}.$$

If we assume that

$$\gamma + \delta' = \alpha' + \beta' - n + 1, \tag{4}$$

then inner sum is Saaschutzian and therefore equal to

$$\frac{(\gamma-\alpha'+r)_{n-r}(\gamma-\beta'+r)_{n-r}}{(\gamma+r)_{n-r}(\gamma-\alpha'-\beta'+r)_{n-r}} =$$

$$= \frac{(\gamma-\alpha')_n(\gamma-\beta')_n}{(\gamma)_n(\gamma-\alpha'-\beta')_n} \frac{(\gamma)_r(\gamma-\alpha'-\beta')_r}{(\gamma-\alpha')_r(\gamma-\beta')_r}.$$

It follows that

$$S_n = \frac{(\gamma-\alpha')_n(\gamma-\beta')_n}{(\gamma)_n(\gamma-\alpha'-\beta')_n} \sum_{r=0}^{n} \frac{(-n)_r(\alpha)_r(\beta)_r(\gamma-\alpha'-\beta')_r}{r!\,(\gamma-\alpha')_r(\gamma-\beta')_r(\delta)_r}.$$

If we assume that

$$\gamma = \beta + \beta', \tag{5}$$

the last sum becomes

$$\sum_{r=0}^{n} \frac{(-n)_r(\alpha)_r(\beta-\alpha')_r}{r!\,(\gamma-\alpha')_r(\delta)_r}.$$

If in addition we take

$$\gamma + \delta = \alpha + \beta - n + 1, \tag{6}$$

the sum reduces to

$$\frac{(\gamma-\alpha-\alpha')_n(\gamma-\beta)_n}{(\gamma-\alpha')_n(\gamma-\alpha-\beta)_n},$$

so that finally

$$S(n) = \frac{(\beta)_n(\beta')_n(\gamma-\alpha-\alpha')_n}{(\gamma)_n(\beta-\alpha')_n(\beta'-\alpha)_n}, \tag{7}$$

provided

$$\begin{cases} \gamma = \beta + \beta' \\ \gamma + \delta = \alpha + \beta - n + 1, \\ \gamma + \delta' = \alpha' + \beta' - n + 1 . \end{cases} \tag{8}$$

If $\gamma = \beta + \beta'$ and λ is arbitrary, it follows from (7) that

$$\sum_{n=0}^{\infty} \frac{(\beta)_n(\beta')_n(\gamma-\alpha-\alpha')_n}{n!\,(\gamma)_n(\lambda)_n} x^n = \sum_{n=0}^{\infty} \frac{(\beta-\alpha')_n(\beta'-\alpha)_n}{n!\,(\lambda)_n} x^n,$$

$$\sum_{r+s\geqslant n} \frac{(-n)_{r+s}(\alpha)_r(\alpha')_s(\beta)_r(\beta')_s}{r!\,s!\,(\gamma)_{r+s}(\alpha-\beta'-n+1)_r(\alpha'-\beta-n+1)_s} .$$

Since

$$(\alpha-\beta'-n+1)_r = (-1)^r \frac{(\beta'-\alpha)_n}{(\beta'-\alpha)_{n-r}},$$

the right member reduces to

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{x^n} \sum_{r+s\leqslant n} (-1)^{r+s} \frac{(\alpha)_r(\alpha')_s(\beta)_r(\beta')_s}{(n-r-s)!\,r!\,s!\,(\gamma)_{r+s}} \cdot$$

$$\cdot (\beta'-\alpha)_{n-r}(\beta-\alpha')_{n-s} =$$

$$= \sum_{r,s=0}^{\infty} \frac{(\alpha)_r(\alpha')_s(\beta)_r(\beta')_s(\beta-\alpha')_r(\beta'-\alpha)_s}{r!\,s!\,(\gamma)_{r+s}(\lambda)_{r+s}} x^{r+s} \cdot$$

$$\cdot \sum_{n=0}^{\infty} \frac{(\beta-\alpha'+r)_n(\beta'-\alpha+s)_n}{n!\,(\lambda+r+s)_n} x^n .$$

We have therefore

$${}_3F_2\left[\begin{matrix}\beta, \beta', \gamma-\alpha-\alpha'; \\ \gamma, \lambda\end{matrix}\; x\right] = \tag{9}$$

$$= \sum_{r,s=0}^{\infty} \frac{(\alpha)_r(\alpha')_s(\beta)_r(\beta')_s(\beta-\alpha')_r(\beta'-\alpha)_s}{r!\,s!\,(\gamma)_{r+s}(\lambda)_{r+s}} x^{r+s} \cdot$$

$$\cdot F[\beta-\alpha'+r, \beta'-\alpha+s; \lambda+r+s; x],$$

where $\gamma = \beta + \beta'$ and λ is arbitrary. Alternatively this can be written in the form

$$(10) \qquad {}_3F_2\left[\frac{\beta, \beta', \gamma - \alpha - \alpha' ; x}{\gamma, \lambda}\right] =$$

$$= \sum_{n=0}^{\infty} \frac{(\beta - \alpha')_n (\beta' - \alpha)_n}{n\,!\,(\lambda)_n} \cdot$$

$$\cdot \sum_{r,s=0}^{\infty} \frac{(\alpha)_r (\alpha')_s (\beta)_r (\beta')_s (\beta - \alpha' + n)_r (\beta' - \alpha + n)_s}{r\,!\,s\,!\,(\gamma)_{r+s} (\lambda + n)_{r+s}} x^{r+s} .$$

REFERENCE

[1] CARLITZ L.: *A Saalschützian theorem for double series.* Journal of the London Mathematical Society, Vol. 38 (1963), 415-418.