

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

PAUL C. EKLOF

SAHARON SHELAH

**A calculation of injective dimension over
valuation domains**

Rendiconti del Seminario Matematico della Università di Padova,
tome 78 (1987), p. 279-284

http://www.numdam.org/item?id=RSMUP_1987__78__279_0

© Rendiconti del Seminario Matematico della Università di Padova, 1987, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A Calculation of Injective Dimension over Valuation Domains.

PAUL C. EKLOF - SAHARON SHELAH (*)

This paper takes up a problem which was posed in a paper by S. Bazzoni [B], about the injective dimension of certain direct sums of divisible modules over a valuation domain. We refer the reader to that paper for the motivation for the problem. We shall make use of the same notation as in [B], which we now proceed to review.

Let R be a valuation domain of global dimension $n + 1$, where $n \geq 2$. Let $\{L_\alpha: \alpha \in A\}$ be a family of archimedean ideals of R , where A is a set of cardinality $\geq \aleph_{n-2}$. For each α let I_α be the injective envelope of R/L_α . Let $I = \prod_{\alpha \in A} I_\alpha$, and for each $1 \leq k \leq n$, let D_{n-k} be the submodule of I consisting of those elements having support of cardinality $< \aleph_{n-k}$, i.e. for all $y \in I$, y belongs to D_{n-k} if and only if the cardinality of

$$\{\alpha \in A: y(\alpha) \neq 0\}$$

is strictly less than \aleph_{n-k} .

Bazzoni proves in [B] that the injective dimension of D_{n-k} is at most k . She also shows that the injective dimension of D_{n-1} is exactly 1 and that it is consistent with ZFC that the injective dimension of

(*) Indirizzo degli AA.: P. C. EKLOF: University of California, Department of Math., Irvine; S. SHELAH: Hebrew University and Rutgers University, New Brunswick, New Jersey, 08903 - U.S.A.

Partially supported by N.S.F. Grant DMS-8600451.

D_{n-2} is exactly 2. It is the main purpose of this paper to prove this latter result in ZFC. In fact we prove:

THEOREM. *The injective dimension of D_{n-k} is ≥ 2 if $2 \leq k \leq n$.*

Before proving the theorem we prove some lemmas. The first of these is a combinatorial fact. (Compare [Sh; § 6].)

LEMMA 1. *Let κ be a regular cardinal. There exists a family $\{w_\nu^\alpha : \alpha < \kappa^+, \nu < \kappa\}$ of subsets of κ^+ satisfying for all $\alpha < \kappa^+$:*

- (1) $\alpha = \bigcup_{\nu < \kappa} w_\nu^\alpha$;
- (2) for all $\nu < \mu < \kappa$, $w_\nu^\alpha \subseteq w_\mu^\alpha$;
- (3) for all $\nu < \kappa$ and all $\beta < \alpha$, $\beta \in w_\nu^\alpha \Rightarrow w_\nu^\beta = w_\nu^\alpha \cap B$;
- (4) for all $\nu < \kappa$, the cardinality of w_ν^α is $< \kappa$.

PROOF. We shall define the w_ν^α for all ν by induction on α . Let $w_\nu^0 = \emptyset$ for all ν . Now suppose that w_ν^β has been defined for all $\beta < \alpha$. If α is a successor ordinal, say $\alpha = \gamma + 1$, then let $w_\nu^\alpha = w_\nu^\gamma \cup \{\gamma\}$ for all ν . It is easy to see that (1)-(4) hold for α if they hold for γ .

If α is a limit ordinal, let $\lambda =$ the cofinality of α , and let $\eta: \lambda \rightarrow \alpha$ be a strictly increasing function such that the supremum of its range is α . Define a function $f: \lambda \rightarrow \kappa$ by the rule:

$$f(\mu) = \text{the least } \nu < \kappa \text{ such that}$$

$$\text{for all } \tau < \sigma \leq \mu, \eta(\tau) \in w_\nu^{\eta(\sigma)}.$$

It is easy to see that f is well-defined because of (1) and (2) and because κ is regular and $\geq \lambda > |\mu|$. Now for each $\nu < \kappa$ let

$$w_\nu^\alpha = \bigcup \{w_\tau^{\eta(\mu)} : \mu < \nu \text{ and } f(\mu) \leq \nu\}.$$

Conditions (2) and (4) are easily verified. To see that (1) holds, suppose $\gamma < \alpha$ and choose μ such that $\eta(\mu) > \gamma$. Then $\gamma \in w_\tau^{\eta(\mu)}$ for some τ , so if $\nu > \max\{\tau, \mu, f(\mu)\}$, then $\gamma \in w_\nu^\alpha$. To prove (3), let us fix α and ν and let $Y = \{\mu < \nu : f(\mu) \leq \nu\}$. Thus

$$w_\nu^\alpha = \bigcup_{\mu \in Y} w_\nu^{\eta(\mu)}.$$

Notice first that if $\tau < \mu$ and $\mu \in Y$, then $\eta(\tau) \in w_v^{\eta(\mu)}$; so by induction $w_v^{\eta(\tau)} = w_v^{\eta(\mu)} \cap \eta(\tau)$. Now if $\beta \in w_v^\alpha$, then $\beta \in w_v^{\eta(\mu)}$ for some $\mu \in Y$; in this case it is easy to see, using the previous observation, that $\beta \in w_v^{\eta(\tau)}$ for any $\tau \in Y$ such that $\beta < \eta(\tau)$. Clearly

$$w_v^\beta = w_v^{\eta(\mu)} \cap \beta \subseteq w_v^\alpha \cap B,$$

so we are left with proving the opposite inclusion. Suppose $\gamma \in w_v^\alpha \cap \beta$; then $\gamma \in w_v^{\eta(\tau)}$ for some $\tau \in Y$. As above, $\gamma \in w_v^{\eta(\sigma)}$ for any $\sigma \in Y$ such that $\gamma < \eta(\sigma)$, so without loss of generality $\beta < \eta(\tau)$. But then $\gamma \in w_v^{\eta(\tau)} \cap \beta = w_v^\beta$, since $\beta \in w_v^{\eta(\tau)}$.

The second lemma will be used to show that for certain submodules $K' \supseteq K$ of I_α , the quotient K'/K has sufficiently large cardinality. (K and K' will have the form $\{u \in I_\alpha: ru = 0\}$ for an appropriate r .) Here $\mathfrak{F}(\gamma)$ is the set of all subsets of γ .

LEMMA 2. *Let $\{r_\nu: \nu < \gamma\}$ be a sequence of elements of R , and let N be a pure-injective module such that for all $\mu < \gamma$ there exists an element $a_\mu \in N$ such that $r_\mu a_\mu = 0$ and $r_{\mu+1} a_\mu \neq 0$. Then for each $S \in \mathfrak{F}(\gamma)$ there exist an element x_S of N such that*

$$(*) \quad \text{for all } \beta < \gamma \text{ and all } S, T \in \mathfrak{F}(\gamma), \text{ if } S \cap \beta = T \cap \beta, \text{ then } r_{\beta+1}(x_S - x_T) = 0 \text{ if and only if } S \cap (\beta + 1) = T \cap (\beta + 1).$$

PROOF. The idea of the construction is that x_S should « be » $\sum_{\mu \in S} a_\mu$.

The actual construction is by induction on γ . If γ is finite and $S \subseteq \gamma$, let $x_S = \sum_{\mu \in S} a_\mu$. (We let $x_\emptyset = 0$.) Now suppose that for all $\delta < \gamma$ and all $S \subseteq \delta$ we have defined x_S so that $(*)$ holds. We consider two cases.

Case 1: $\gamma = \delta + 1$ for some δ . We let $x_S = x_{S \cap \delta}$ if $\delta \notin S$, and we let $x_S = x_{S \cap \delta} + a_\delta$ if $\delta \in S$. It is easy to check, using the inductive hypothesis, that $(*)$ holds.

Case 2: $\gamma = \lambda$, a limit ordinal. Here we use the fact that since N is pure-injective it is algebraically compact: see, for example, [FS; p. 215]. For any $S \subseteq \lambda$ we let x_S be a solution of the set of equations

$$\{r_{\beta+1}(x - x_{S \cap (\beta+1)}) = 0: \beta < \lambda\}$$

in the single unknown x . (The elements $x_{S \cap (\beta+1)}$ of N have been defined by induction.) This system of equations is finitely solvable

in N : indeed, for any finite subset F of λ , if $\delta > \sup(F)$, then $x_{S \cap \delta}$ is a solution of

$$\{_{\beta+1}(x - x_{S \cap (\beta+1)}) = 0 : \beta \in F\}.$$

Hence by algebraic compactness there is a global solution, x_S . It remains to check that $(*)$ is satisfied. So suppose that S and T are subsets of λ , and $\beta < \lambda$ such that $S \cap \beta = T \cap \beta$. We have:

$$x_S - x_T = (x_S - x_{S \cap (\beta+1)}) + (x_{S \cap (\beta+1)} - x_{T \cap (\beta+1)}) + (x_{T \cap (\beta+1)} - x_T)$$

so $r_{\beta+1}(x_S - x_T) = 0 + r_{\beta+1}(x_{S \cap (\beta+1)} - x_{T \cap (\beta+1)}) + 0$; hence we are done by induction. \square

The third lemma will guarantee us the existence of the elements a_μ in Lemma 2 provided that $r_{\mu+1} \notin r_\mu R$. (Of course, over a valuation domain, injective = pure-injective + divisible.)

LEMMA 3. *Suppose L is an archimedean ideal and N is a divisible module containing R/L . Suppose also that r, s, t are elements of R such that t is a non-unit and $r = st$. Then there exists $a \in N$ such that $ra = 0$ and $sa \neq 0$.*

PROOF. We shall let \bar{b} denote the coset, $b + L$ of $b \in R$ in $R/L \subseteq N$. Since L is archimedean there is an element $b \in L \setminus tL$. If $bt^{-1} \in R$, let $a \in N$ such that $sa = bt^{-1} + L$. Then $ra = \bar{b} = 0$, but $sa \neq 0$ since $bt^{-1} \notin L$ (because $b \notin tL$). If $tb^{-1} \in R$, let $a \in N$ such that $s(tb^{-1})a = \bar{1}$. Then $ra = \bar{b} = 0$, but $sa \neq 0$ since $tb^{-1}(sa) = \bar{1}$. \square

We are now ready to give the:

PROOF OF THE THEOREM. Let $D = D_{n-k}$. As Bazzoni observes, we can assume that $|A| = \aleph_{n-k}$ since we can replace D by the direct summand of D consisting of elements whose support lies in a fixed subset of A of size \aleph_{n-k} . It suffices to prove that $\text{Ext}^1(J, D) \neq 0$ for some ideal J of R , for then $\text{Ext}^2(R/J, D) \neq 0$ (cf. [FS; VI.5.2]). For this it suffices to prove that the canonical map: $\text{Hom}(J, I) \rightarrow \text{Hom}(J, I/D)$ is not surjective. In fact we shall show that this map is not surjective whenever J is an ideal of R which is not generated by a set of size \aleph_{n-k} but is generated by a set of size \aleph_{n-k+1} ; there is such an ideal because $\text{gl. dim } R > n - k + 2$ (cf. [0] or [FS; IV.2.3].)

Let $\{j_{\alpha+1} : \alpha < \aleph_{n-k+1}\}$ be a set of generators of J ordered so that for all $\beta < \alpha$, $j_{\beta+1} \in Rj_{\alpha+1}$ and $j_{\alpha+1} \notin Rj_{\beta+1}$. Thus for every pair of

ordinals $\beta < \alpha$ we have a non-unit r_β^α of R such that $r_\beta^\alpha j_{\alpha+1} = j_{\beta+1}$. Moreover, for all $\beta < \gamma < \alpha$ we have $r_\beta^\alpha = r_\beta^\gamma r_\gamma^\alpha$.

Let $\kappa = \aleph_{n-k}$. We may as well suppose that $\Lambda = \kappa$. So defining $f: J \rightarrow I/D$ amounts to choosing, for each $\nu < \kappa$, elements $x_\nu^\alpha \in I_\nu$ ($\alpha < \kappa^+ = \aleph_{n-k+1}$) so that for all $\beta < \alpha$, $|\{\nu < \kappa: r_\beta^\alpha x_\nu^\alpha \neq x_\nu^\beta\}| < \kappa$; for then we can define $f(j_{\alpha+1}) = x^\alpha + D$, where $x^\alpha = \langle x_\nu^\alpha: \nu < \kappa \rangle \in I$. We are going to use the sets w_ν^α ($\alpha < \kappa^+$, $\nu < \kappa$) constructed in Lemma 1 in order to define the x_ν^α 's; in fact, we shall construct them so that $r_\beta^\alpha x_\nu^\alpha = x_\nu^\beta$ if $\beta \in w_\nu^\alpha$. Then f will be defined because, by (1) of Lemma 1, for any $\beta < \alpha$ there exists $\mu < \kappa$ so that $\beta \in w_\mu^\alpha$, and hence by (2), the set of ν such that $r_\beta^\alpha x_\nu^\alpha \neq x_\nu^\beta$ is contained in μ , and thus has cardinality less than κ .

In order to make f not liftable to a homomorphism into I we shall also require that the x_ν^α be chosen so that if $\text{sup}(w_\nu^\alpha) + \kappa < \beta < \alpha$, then $r_\beta^\alpha x_\nu^\alpha \neq x_\nu^\beta$. (The sum is ordinal addition.) Indeed, if there were a $g: J \rightarrow I$ which lifted f , then we would have $g(j_\alpha) = y^\alpha$ for some $y^\alpha \in I$ such that $y^\alpha = x^\alpha + d^\alpha$ for some $d^\alpha \in D$, for all $\alpha < \kappa^+$. For each $\mu < \kappa$, let

$$Y_\mu \stackrel{\text{def}}{=} \{\alpha < \kappa^+: \mu \notin \text{supp}(d^\alpha)\};$$

then for some $\nu < \kappa$, Y_ν is a stationary subset of κ^+ since $\bigcup Y_\mu = \kappa^+$ (cf. [J; Lemma 7.4]). Now by (4), $\text{sup}(w_\nu^\alpha) < \alpha$ if $\text{cf}(\alpha) = \kappa$, so by Fodor's Lemma ([J; p. 59]) there is a stationary subset Y' of Y_ν and an ordinal γ such that for all $\alpha \in Y'$ $\text{sup}(w_\nu^\alpha) = \gamma$. Hence there are elements β, α of Y' such that $\gamma + \kappa < \beta < \alpha$. But then $y^\alpha(\nu) = x_\nu^\alpha$ and $y^\beta(\nu) = x_\nu^\beta$, and by construction $r_\beta^\alpha x_\nu^\alpha \neq x_\nu^\beta$, which means that g is not a homomorphism.

Thus it remains only to construct for each ν the elements x_ν^α of I_ν so that for all $\beta < \alpha < \kappa^+$:

- (i) $r_\beta^\alpha x_\nu^\alpha = x_\nu^\beta$ if $\beta \in w_\nu^\alpha$;
- (ii) $r_\beta^\alpha x_\nu^\alpha \neq x_\nu^\beta$ if $\beta > \text{sup}(w_\nu^\alpha) + \kappa$.

We shall do this for each fixed ν by induction on α . Let $x_\nu^0 = \bar{1}$. Suppose now that x_ν^β has been defined for all $\beta < \alpha$ so that (i) and (ii) hold where defined. In order to satisfy (i) it is enough to choose x_ν^α to be a solution, z , of the system of equations

$$(\dagger) \quad \{r_\beta^\alpha z = x_\nu^\beta: \beta \in w_\nu^\alpha\}.$$

Since I is pure-injective, it suffices to show that this system is finitely solvable in I_ν . If F is a finite subset of w_ν^α and $\sigma = \max(F)$, we claim that any z such that $r_\sigma^\alpha z = x_\nu^\sigma$ will be a solution of

$$\{r_\beta^\alpha z = x_\nu^\beta : \beta \in F\}.$$

In fact, if $\beta \in F$ and $\beta < \sigma$, then since $\sigma, \beta \in w_\nu^\alpha$, (3) implies that $\beta \in w_\nu^\sigma$, so $r_\beta^\sigma x_\nu^\sigma = x_\nu^\beta$ and hence $r_\beta^\alpha z = r_\beta^\sigma r_\sigma^\alpha z = r_\beta^\sigma x_\nu^\sigma = x_\nu^\beta$.

Now consider (ii). Let $\delta = \sup(w_\nu^\alpha)$. Let z be a fixed solution of (\dagger) . Then (i) will hold if x_ν^α is of the form $z + u$ where $r_\delta^\alpha u = 0$. Let $\beta = \delta + \kappa + 1$. It suffices to choose u so that $r_\delta^\alpha u = 0$ and for each γ such that $\beta \leq \gamma < \alpha$, $r_\beta^\alpha u \neq r_\beta^\gamma x_\nu^\gamma - r_\beta^\alpha z$. (We let $r_\beta^\beta = 1$.) For then, since $r_\beta^\alpha = r_\beta^\gamma r_\gamma^\alpha$, we have that $r_\gamma^\alpha(z + u) \neq x_\nu^\gamma$. But Lemma 2 (with $r_\nu = r_{\delta+\nu}^\alpha$, for $\nu < \kappa$) in conjunction with Lemma 3 implies that the quotient group

$$\{u \in I_\nu : r_\delta^\alpha u = 0\} / \{u \in I_\nu : r_\beta^\alpha u = 0\}$$

has cardinality $\geq 2^\kappa$. Thus there certainly is a u with the desired properties. This completes the inductive step of the construction, and hence completes the proof of the theorem. \square

COROLLARY. *If gI , $\dim(R) \geq 3$, and for each $n \in \omega$, I_n is an injective nodule containing R/L_n for some archimedean ideal L_n of R , then the injective dimension of $\bigoplus_{n \in \omega} I_n$ is ≥ 2 . \square*

REFERENCES

- [B] S. BAZZONI, *Injective dimension of some divisible modules over a valuation domain*, to appear in Proc. Amer. Math. Soc.
- [FS] L. FUCHS - L. SALCE, *Modules over Valuation Domains*, Marcel Dekker, 1985.
- [J] T. JECH, *Set Theory*, Academic Press, 1978.
- [O] B. OSOFSKY, *Global dimension of valuation rings*, Trans. Amer. Math. Soc., **127** (1967), pp. 136-149.
- [Sh] S. SHELAH, *Uncountable constructions for B.A. e.c. groups and Banach spaces*, Israel. J. Math., **51** (1985), pp. 273-297.

Manoscritto pervenuto in redazione il 14 marzo 1987.