

RENDICONTI  
*del*  
SEMINARIO MATEMATICO  
*della*  
UNIVERSITÀ DI PADOVA

JAVIER OTAL

JUAN MANUEL PEÑA

**Groups with minimal conditions related to finiteness  
properties on conjugacy classes**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 81 (1989), p. 79-84

[http://www.numdam.org/item?id=RSMUP\\_1989\\_\\_81\\_\\_79\\_0](http://www.numdam.org/item?id=RSMUP_1989__81__79_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1989, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## **Groups with Minimal Conditions Related to Finiteness Properties on Conjugacy Classes.**

JAVIER OTAL - JUAN MANUEL PEÑA (\*)

### **1. Introduction.**

In a series of papers Belyaev [2], Belyaev-Sesekin [3] and Bruno-Phillips [4] (see also [11; section 8]) have obtained results on minimal non  $FC$ -groups and classified them when they have a non-trivial finite factor. Furthermore, in [5], Bruno-Phillips have generalized these results studying minimal non finite-by-nilpotent groups; this generalization is based on the well-known fact that a finite-by-abelian group, that is, a group with finite derived group, is a particular type of an  $FC$ -group (see [10; 4.35]). In any case these minimal are special cases of Cernikov groups (see [5; section 7] and [11; 8.11 and 8.13]).

Lately, in [8] and [9], the authors have extended the above problems in a direction of research that, roughly speaking, consists in replacing the term « finite group » by « Cernikov group » and so «  $FC$ -group » by «  $CC$ -group », although it should be remarked that the results we have found have a rather different nature than those of Bruno-Phillips because our theorems give subgroup characterizations of the conditions under consideration.

The aim of the present paper is to combine these ideas considering stronger conditions on the conjugacy classes of a group, namely the

(\*) Indirizzo degli AA.: J. OTAL: Departamento de Matemáticas, Facultad de Ciencias, Universidad de Zaragoza, 50009 Zaragoza, Spain; J. M. PEÑA: Departamento de Matemática Aplicada, EUITI, Universidad de Zaragoza, 50009 Zaragoza, Spain.

condition of being an *FL*-group or a *CL*-group (the terminology will be explain below). We shall first prove the Cernikov case and then deduce the finite case from it in the following way.

**THEOREM 1.** *Let  $\mathfrak{X}$  be one of the following classes of groups (i) the theoretical union of the class of *CL*-groups and that of Cernikov-by-abelian groups; (ii) the class of *CL*-groups. Then a locally graded group  $G$  is an  $\mathfrak{X}$ -group if and only if every proper subgroup of  $G$  is an  $\mathfrak{X}$ -group.*

**THEOREM 2.** *Let  $\mathfrak{Y}$  be one of the following classes of groups: (i) the theoretical union of the class of *FL*-groups and that of finite-by-abelian groups; (ii) the class of *FL*-groups. Then*

(1) *A locally graded minimal non  $\mathfrak{Y}$ -group is a non-perfect minimal non *FC*-group.*

(2) *A locally graded group  $G$  satisfies the minimal condition for non  $\mathfrak{Y}$ -subgroups if and only if  $G$  is either a  $\mathfrak{Y}$ -group or a Cernikov group.*

We recall some facts about a locally graded minimal non *FC*-group  $G$  (see [11]); such  $G$  is countable and locally finite. If  $G$  is perfect, then it is  $\mathcal{F}$ -perfect by [11; 8.9]; otherwise  $G$  is a Cernikov group and its structure is given by the results quoted above. Thus the structure of the groups in (1) of Theorem 2 is known.

Finally a comment on minimals; if  $\mathcal{C}$  is a class of groups, by a *minimal non  $\mathcal{C}$ -group* we shall mean a group  $G$  which is not a  $\mathcal{C}$ -group but in which each proper subgroup is a  $\mathcal{C}$ -group. Attempts to classify these groups, for  $\mathcal{C}$  adequate, are certain to fail because of the existence of the Tarski groups, infinite nonabelian groups in which each proper subgroup is finite ([7]); indeed, these monsters show that our results are false without some sort of restriction on the groups under consideration. This difficulty has been avoided by the imposition of the local gradedness condition: a group  $G$  is said to be *locally graded* if every non-trivial finitely generated subgroup of  $G$  contains a proper subgroup of finite index.

## 2. Notation.

Throughout our group-theoretic notation is standard and is taken from [10]. For the reader's convenience we quickly review the definitions and some elementary facts to be used in the sequel. A group  $G$  is said to be an *FC*-group (a *CC*-group) if  $G/C_G(x^G)$  is a finite group

(a Cernikov group) for all  $x$  in  $G$ ; Polovickii's theorems of characterization of these groups assure that  $G$  is an  $FC$ -group if and only if  $x^a$  is finite-by-cyclic for every  $x$  in  $G$  ([10; Corollary 3 to 4.32]) and that if  $G$  is a  $CC$ -group, then  $[G, x]$  is Cernikov and so  $x^a$  is Cernikov-by-cyclic for every  $x$  in  $G$  ([10; 4.36]). More properties of  $FC$ -groups can be found in [11] and of  $CC$ -groups in [1] and [8].

Let  $G$  be a group and let  $m$  be a positive integer or infinity; then the subgroup  $G_m$  generated by all elements of  $G$  with order  $m$  is called the  $m$ -layer of  $G$ . A group each of whose layers is a Cernikov group is called a  $CL$ -group and a group whose layers are all finite is called an  $FL$ -group. The classification of  $CL$ - and  $FL$ -groups, again due to Polovickii, is collected in [10; 4.42 and 4.43]. Using the well-known Sunkov-Kegel-Wehrfritz theorem [6; 5.8], the more convenient way of dealing with these groups is as follows: *A group  $G$  is a  $CL$ -group (an  $FL$ -group) if and only if  $G$  is a periodic  $CC$ -group (a periodic  $FC$ -group) in which every Sylow subgroup is a Cernikov group.* Clearly both classes are closed under subgroups and taking homomorphic images. They are not closed under extensions but satisfy a certain form of weak extensibility, which can be stated as follows:

(\*) *Let  $G$  be a periodic  $CC$ -group (a periodic  $FC$ -group) with a normal subgroup  $N$  such that  $G/N$  is Cernikov (finite). Then  $G$  is a  $CL$ -group (an  $FL$ -group) if and only if  $N$  is a  $CL$ -group (an  $FL$ -group).*

$FL$ -groups or finite-by-abelian groups are  $FC$ -groups and clearly  $CL$ -groups are  $CC$ -groups. In general Cernikov-by-abelian groups need not to be  $CC$ -groups; for example, the semidirect product of a Prüfer  $p$ -group  $P$  by the group generated by an automorphism  $\alpha$  of infinite order of  $P$  given by  $x^\alpha = x^r$ ,  $x$  in  $P$ , where  $(p, r) = 1$ . However a periodic Cernikov-by-abelian group, that is, a periodic group  $G$  with Cernikov derived group, is a  $CC$ -group by [10; 4.36], which will be frequently used in what follows. Finally note that each notion defined in the finite case trivially implies its corresponding in the Cernikov case.

### 3. Proof of the Theorems.

PROOF OF THEOREM 1. Suppose that the case (i) has been proved but the case (ii) is false. Let  $G$  be a counterexample, that is, a locally graded minimal non  $CL$ -group. By (i)  $G'$  is Cernikov and, since  $G$  is clearly periodic, it follows that  $G$  is a  $CC$ -group. Then there exists

a Sylow subgroup  $P$  of  $G$  that does not satisfy Min so that, by minimality,  $G = P$  and every proper subgroup of  $G$  satisfies Min. Therefore  $G$  satisfies Min, a contradiction which shows (ii).

Thus it remains to show (i). As above we suppose that the result is false and take  $G$  to be a counterexample. We claim that  $G$  must be periodic. If not, we note that whenever  $G/L$  is a non-trivial periodic image of  $G$ , then  $L$  is a Cernikov group because  $L$  cannot be periodic. If  $G$  is finitely generated, then  $G$  has a proper normal subgroup  $N$  of finite index so that  $N$  is Cernikov. Clearly  $G/N$  satisfies Max and it follows immediately that every proper subgroup of  $G/N$  is finite-by-abelian. By [9; Theorem 1] (or [5; 4.1])  $G/N$  is finite-by-abelian, a contradiction. Therefore every finitely generated subgroup  $H$  of  $G$  is proper and, in any case,  $H$  is Cernikov. Thus  $G$  is locally finite. If  $T$  is the periodic part of  $G$ , then  $G' \leq T$  and  $G/T$  is a non-trivial torsion-free abelian group. Choose a proper subgroup  $M$  of  $G$  containing  $T$  such that  $G/M$  is Cernikov; the election of a such  $M$  is clear if  $G/T$  is not  $\mathcal{F}$ -perfect and it is a consequence of the well-known structure of divisible torsion-free abelian groups otherwise. As we remarked above  $M'$  must be Cernikov and so  $T'$  is Cernikov. Therefore every  $CL$ -subgroup of  $G$ , being contained in  $T$ , has Cernikov derived group. Hence every proper subgroup of  $G$  is Cernikov-by-abelian and so is  $G$  by [9; Theorem 1], a contradiction which shows that  $G$  must be periodic as claimed. As a consequence every proper subgroup of  $G$  is a  $CC$ -group.

If  $G$  itself is a  $CC$ -group, then  $G$  cannot be  $\mathcal{F}$ -perfect because a periodic  $\mathcal{F}$ -perfect  $CC$ -group is abelian by [8; 1.2]. Hence  $G$  has a proper normal subgroup  $K$  of finite index and, since  $K$  cannot be a  $CL$ -group by the property (\*), it follows that  $K'$  is Cernikov. On the other hand  $G/K'$  is an abelian-by finite  $CC$ -group and so it is Cernikov-by-abelian by [8; 1.1] and [10; 4.23]. Therefore  $G'$  is Cernikov, a contradiction.

Thus  $G$  is a minimal non  $CC$ -group. By the result of [8],  $G$  is countable, locally finite, perfect,  $\mathcal{F}$ -perfect and every non-trivial image of  $G$  is again a minimal non  $CC$ -group having the same properties mentioned above.  $G/\zeta(G)$  has trivial center by Grün's Lemma ([10; part 1, p. 48]) so that replacing  $G$  by  $G/\zeta(G)$  we may assume that  $\zeta(G) = 1$ . Since  $G$  is perfect and  $\mathcal{F}$ -perfect, any Cernikov normal subgroup of  $G$  is central in  $G$  by [10; 3.29] and so trivial. Therefore proper and non-trivial normal subgroups of  $G$  cannot have Cernikov layers and so they all are abelian.

Let  $J$  be a non-trivial simple image of  $G$ . By the result of [6; 1.D], for every prime  $p$ ,  $J$  has a Sylow  $p$ -subgroup that contains a conjugate of every finite  $p$ -subgroup of  $J$ . Clearly a proper  $p$ -subgroup of  $J$  is soluble-by-finite so that it has finite derived length. Hence  $J$  cannot be enormous (terminology of [6; p. 122]) and this contradicts [8; 4.2]. Therefore  $G$  has no non-trivial simple images so that  $G$  can be expressed as the union of a chain of proper normal subgroups. As we showed in the previous paragraph every proper normal subgroup of  $G$  is abelian. Therefore  $G$  is abelian, a contradiction which shows our theorem.

The proof of the second part of the next Theorem needs an auxiliary result.

**LEMMA.** *Let  $G$  be a periodic CC-group in which every proper subgroup is either an FC-group or a Cernikov group. Then  $G$  is either an FC-group or a Cernikov group.*

**PROOF.** Again we assume that the result is false and take  $G$  to be a counterexample. Pick  $x$  in  $G$  such that  $|G:C_G(x)|$  is infinite. Since  $x^\sigma$  is Cernikov,  $G/x^\sigma$  cannot be Cernikov and so there exists a proper subgroup  $H$  of  $G$  containing  $x^\sigma$  such that  $H/x^\sigma$  is not Cernikov. Thus  $H$  is an FC-group and then  $x^\sigma$  is an abelian-by-finite FC-group. Therefore  $x^\sigma$  is central-by-finite and then it is known that  $(x^\sigma)'$  is finite (see [10], for example). Hence  $x^\sigma$  is a finite-by-abelian Cernikov group generated by elements of order  $|x|$ . This implies  $x^\sigma$  finite, a contradiction.

**PROOF OF THEOREM 2.** (1) Let  $G$  be a locally graded minimal non  $\mathfrak{Y}$ -group; clearly every proper subgroup of  $G$  is an FC-group and, in any case,  $G$  is either a CL-group or a Cernikov-by-abelian group by Theorem 1. We claim that it suffices to show that  $G$  is not an FC-group, which is clear if  $G$  is a CL-group. For,  $G$  will be then a minimal non FC-group. If  $G$  were perfect, then it would be  $\mathcal{F}$ -perfect and all this would give  $G$  abelian, reasoning directly in case of  $G$  being Cernikov-by-abelian or using [8; 1.2] in case of  $G$  being a CL-group, so that this assumption cannot occur and our claim follows.

Therefore it remains to show that  $G$  is not an FC-group in case (i) provided  $G$  is a Cernikov-by-abelian group. Assume that  $G$  is an FC-group. Since  $G$  is not abelian, it has a proper normal subgroup  $N$  of finite index. Obviously  $N$  cannot be an FL-group by the property (\*) so that  $N'$  is finite. However  $G/N'$  is finite-by-abelian and therefore  $G$  is finite-by-abelian, a contradiction.

(2) We assume that there exists a locally graded group  $G$  satisfying the minimal condition for non  $\mathcal{Y}$ -subgroups which is neither a  $\mathcal{Y}$ -group nor a Cernikov group. Then  $G$  has a subgroup  $H$  which is minimal with respect to being neither a  $\mathcal{Y}$ -group nor a Cernikov group. In case (ii)  $H$  is a  $CL$ -group by Theorem 1 and so an  $FC$ -group by the Lemma, which is a contradiction. In case (i), as above,  $H$  cannot be a  $CL$ -group so that  $H'$  is Cernikov. As we did in the proof of Theorem 1, arguing with the periodic part of  $H$ , it is easy to show that  $H$  is periodic and so a  $CC$ -group. Then the Lemma again gives that  $H$  is in fact an  $FC$ -group and, as in the last part of the proof of the above (1), this yields to a contradiction, and the proof of Theorem 2 is now complete.

## REFERENCES

- [1] J. ALCAZAR - J. OTAL, *Sylow subgroups of groups with Cernikov conjugacy classes*, J. Algebra, **110** (1987), pp. 507-513.
- [2] V. V. BELYAEV, *Groups of Miller-Moreno type*, Siberian Math. J., **19** (1978), pp. 350-360.
- [3] V. V. BELYAEV - N. F. SESEKIN, *Infinite groups of Miller-Moreno type*, Acta Math. Acad. Hungar., **26** (1975), pp. 369-376.
- [4] B. BRUNO - R. E. PHILLIPS, *Minimal non FC-groups*, Abstracts Amer. Math. Soc., **2** (1980), p. 565.
- [5] B. BRUNO - R. E. PHILLIPS, *On minimal conditions related to Miller-Moreno type groups*, Rend. Sem. Mat. Univ. Padova, **69** (1983), pp. 153-168.
- [6] O. H. KEGEL - B. A. F. WEHRFRUTZ, *Locally finite groups*, North Holland, Amsterdam (1973).
- [7] A. YU. OL'SANSKII, *Infinite groups with cyclic subgroups*, Soviet Math. Dokl., **20** (1979), pp. 343-346.
- [8] J. OTAL - J. M. PEÑA, *Minimal non CC-groups*, Comm. Algebra, **16** (1988), pp. 1231-1242.
- [9] J. OTAL - J. M. PEÑA, *Groups in which every proper subgroup is Cernikov-by-nilpotent or nilpotent-by-Cernikov*, Arch. Math., **51** (1988), pp. 193-197.
- [10] D. J. S. ROBINSON, *Finiteness conditions and generalized soluble groups*, Springer-Verlag, Berlin (1972).
- [11] M. J. TOMKINSON, *FC-groups*, Pitman, London (1984).

Manoscritto pervenuto in redazione il 24 agosto 1987 e in forma revisionata il 3 febbraio 1988.