

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

JOHN C. LENNOX

JAMES WIEGOLD

On a question of Deaconescu about automorphisms

Rendiconti del Seminario Matematico della Università di Padova,
tome 89 (1993), p. 83-86

http://www.numdam.org/item?id=RSMUP_1993__89__83_0

© Rendiconti del Seminario Matematico della Università di Padova, 1993, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On a Question of Deaconescu About Automorphisms.

JOHN C. LENNOX - JAMES WIEGOLD (*)

1. Introduction.

At a recent meeting in Barnaul, Siberia, Marian Deaconescu posed the following problem.

1.1. *Do there exist infinite groups G such that $\text{Aut } H \cong \cong N_G(H)/C_G(H)$ for all subgroups H of G ?*

Let us call (finite or infinite) groups with this property MD-groups. We note at the outset that the infinite dihedral group D has the MD-property, and this is fairly easy to prove. Theorem 2.1 characterizes the infinite metabelian MD-groups; there are just 2^{\aleph_0} of them, and they all have very simple structure. A corollary is that D is the only finitely generated infinite metabelian MD-group.

It is quite probable that the groups covered in Theorem 2.1 are the only infinite MD-groups, because automorphism groups tend to be insoluble and big. For example, every free group F_m of infinite rank m has F_{2^m} in its automorphism group; thus, by the splitting property of free groups, any MD-group containing a subgroup whose automorphism group contain a non-abelian free subgroup would have to be of order bigger than all the cardinals obtained from \aleph_0 by related exponentiation ω times; that is, bigger than $\aleph_0, 2^{\aleph_0}, 2^{2^{\aleph_0}}, \dots$.

On the other hand, there are very few finite MD-groups:

1.2. *The only non-trivial finite MD-groups are Z_2 and S_3 .*

PROOF. Let G be a finite non-trivial MD-group. We claim first that every non-trivial Sylow subgroup is of prime order. If not, G has a p -subgroup P of order bigger than p , and the celebrated Gaschütz theo-

(*) Indirizzo degli AA.: School of Mathematics, University of Wales College of Cardiff, Senghennydd Road, Cardiff, Wales.

rem [2] shows that $\text{Out } P$ has an element of order p . Let $X/C_G(P)$ be a Sylow p -subgroup of $N_G(P)/C_G(P)$. Then $X/C_G(P)$ is of order more than $P/Z(P)$, so that X contains a p -subgroup strictly bigger than P . This process continues and we can construct p -subgroups of arbitrarily large order. This is impossible since G is finite.

So G is metacyclic, and we can choose a Carter subgroup H of G , that is, a self-normalizing nilpotent subgroup. Then $\text{Aut } H \cong N_G(H)/C_G(H) = H/Z(H) = 1$, so again by Gaschütz's theorem, H must be cyclic of order 2, with generator h , say. But then $C_G(h) = H$, and h has $|G|/2$ conjugates, so that $|G'| = |G|/2$. Let K be any subgroup of G' ; then $C_G(K) \geq G'$ and so $\text{Aut } K$ has order at most 2. This means that G' is of order 1, 2, 4 or 6, and a routine calculation completes the proof.

The only general property of MD-groups we shall require in the sequel is this:

1.3. *For every MD-group, G/G' is of exponent dividing 2.*

PROOF. For every x in G , $N_G(\langle x \rangle)/C_G(\langle x \rangle) \cong \text{Aut } \langle x \rangle$ so there must exist $g \in G$ such that $x^g = x^{-1}$. Thus $[x, g] = x^{-2}$, so every square is a commutator.

2. Infinite metabelian ND-groups.

Here is the characterization promised in the introduction.

THEOREM 2.1. *An infinite metabelian group G is an MD-group if and only if G can be generated by a non-trivial torsion-free locally cyclic group A and an element x of order 2 such that*

- (i) $a^x = a^{-1}$ for all a in A ,
- (ii) *the type of A is finite at every prime.*

COROLLARY 2.2. *There are precisely 2^{\aleph_0} infinite metabelian MD-groups.*

The corollary follows since $G' = A^2$ and there are 2^{\aleph_0} different structures for A ; for any locally cyclic group A of the type under consideration, $A \cong A^2$.

Before embarking on the proof of Theorem 2.1, we establish the following simple lemma, which is probably known:

LEMMA 2.3. *Every subgroup H of a torsion-free abelian group A*

has automorphism group of exponent at most 2 if and only if A is locally cyclic with type that is finite at each prime. In fact, $\text{Aut } H$ is of order at most 2 in all such cases.

PROOF. Let A be a torsion-free abelian group such that $\text{Aut } H$ is of exponent at most 2 for all subgroups H . Then A is certainly locally cyclic since $Z \times Z$ has non-abelian automorphism group. If A is p -divisible for some prime p , the map $a \rightarrow a^p$ is an automorphism of infinite order; so the type of A is finite at all primes. Thus A is isomorphic to the subgroup A^* of the additive group Q of rationals generated by reciprocals $p_n^{-k_n}$, where p_n is the n -th prime and k_n is a non-negative integer for each n (see Fuchs [1]). Let α be an automorphism of A^* , and write $1\alpha = rs^{-1}$, where r and s are relatively prime integers. We claim that $s = 1$. If not, then s is divisible by p_m for some m , so that $k_m > 0$; but, since A^* is torsion-free, this means that $p_m^{-k_m}\alpha = rs^{-1}p_m^{-k_m}$, whereas this number has denominator divisible by $p_m^{k_m+1}$, and so is not in A^* . Thus $1\alpha = r$, and a similar argument on α^{-1} shows that $r = \pm 1$, so that $\text{Aut } Z = Z_2$, as required. The same holds for all subgroups H of A , of course.

The converse is clear, and the proof of the lemma is complete.

We turn now to the proof of Theorem 2.1. Let G be a group having a subgroup A and an element x with the properties mentioned in the statement, and let H be a non-trivial subgroup of G . If $H \leq A$, then $H \triangleleft G$ and $C_G(H) = A$, so that $N_G(A)/C_G(A) \cong Z_2$, and so $N_G(H)/C_G(H) \cong \text{Aut } H$, by Lemma 2.3. If $H \not\leq A$, then $H = \langle K, bx \rangle$ for some $K \leq A$ and some element b of A . If $K = 1$, all is well. If not, $C_G(K) = A$ and $C_G(bx) = \langle bx \rangle$, so $C_G(H) = 1$. A very quick calculation shows that $N_G(H) = \langle \sqrt{K}, bx \rangle$, where $\sqrt{K} = \{a: a \in A \text{ and } a^2 \in K\}$. However, we have $\sqrt{K} \cong K$ since K/K^2 is of order precisely 2 (this is true for any nontrivial subgroup of A), and the isomorphism is the squaring map. This means that $N_G(H) \cong H$ and we simply need to show that $\text{Aut } H \cong H$ in this case. Without loss of generality, we may suppose that $b = 1$, so we are considering a group of the form $H = \langle K, x \rangle$, where K is a non-trivial subgroup of A .

Now K is characteristic in H , since every element of $H-K$ is of order 2. Thus, for every automorphism α of H , $\alpha|_K$ is the identity or the inverting map, and $\alpha: x \rightarrow bx$ for some $b \in K$. Conversely, every map $\phi_{\varepsilon, l}$ of the form

$$k \rightarrow k^\varepsilon \quad \text{for } k \in K, \quad x \rightarrow lx,$$

where $\varepsilon = \pm 1$ and $l \in L$, yields an automorphism of K .

Those with $\varepsilon = 1$ form a subgroup isomorphic with K ; the automorphism $\xi = \phi_{-1,1}$ is of order 2, and

$$\xi^{-1} \phi_{1,l} \xi = \phi_{1,l^{-1}},$$

so that $\text{Aut } H \cong H$, as required.

Conversely, let G be an infinite metabelian MD-group, and A a maximal abelian subgroup containing G' . For every subgroup H of A , $C_G(H) \geq A \geq G'$, so $N_G(H)/C_G(H)$ is of exponent at most 2, by 1.3. Thus the periodic part of A is of order at most 6 and so $A = S \times P$ for some finite group P and torsion-free group S . Since $C_G(A) = A$ by maximality we have $S \neq 1$ since G is infinite. If $P \neq 1$, then A contains a subgroup isomorphic to $Z \times Z_n$, $n \neq 1$, which has non-abelian automorphism group. So A is torsion-free and thus it is locally cyclic since $Z \times Z$ has non-abelian automorphism group. By 1.3 and 2.3, A must be a locally cyclic group, and its type is finite for each prime, so $\text{Aut } A$ is of order 2. Thus $Z_2 \cong N_G(A)/C_G(A) = G/A$, so $G = \langle x, A \rangle$ for some x such that $x^2 \in A$. But x^2 is central and this means that $x^2 = 1$ else $N_G(\langle x^2 \rangle)/C_G(\langle x^2 \rangle)$ is of order 1, not 2 as it is for infinite cyclic groups. This completes the proof of the theorem.

Finally, the proof of our last corollary is obvious:

COROLLARY 2.4. *This only infinite finitely generated metabelian MD-group is D .*

REFERENCES

- [1] L. FUCHS, *Abelian Groups*, Hungarian Academy of Sciences, Budapest (1958).
- [2] W. GASCHÜTZ, *Nichtabelsche p -Gruppen besitzen äussere p -Automorphismen*, *J. Algebra*, 4 (1966), pp. 1-2.

Manoscritto pervenuto in redazione il 17 dicembre 1991.