

# RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

PAZ JIMÉNEZ SERAL

## **A lattice of homomorphs**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 90 (1993), p. 39-43

[<http://www.numdam.org/item?id=RSMUP\\_1993\\_\\_90\\_\\_39\\_0>](http://www.numdam.org/item?id=RSMUP_1993__90__39_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1993, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques*  
<http://www.numdam.org/>

## A Lattice of Homomorphs.

PAZ JIMÉNEZ SERAL (\*)

### Preliminary notes.

In this paper all groups are finite and soluble. The homomorph  $h(\mathcal{B})$  for a boundary  $\mathcal{B}$  consists of all « $\mathcal{B}$ -perfect groups», namely all those groups that have no  $\mathcal{B}$ -groups among their epimorphic images. The boundary  $b(\mathcal{H})$  for a homomorph  $\mathcal{H}$  consists of all groups  $G$  such that  $G \notin \mathcal{H}$  and if  $1 \neq N \leq G$ , then  $G/N \in \mathcal{H}$ . The maps  $h$  and  $b$  are mutually inverse bijections between the set of non-empty homomorphs and the set of boundaries. Let  $\mathcal{H}$  be a homomorph. We recall from [4] that the class  $D\mathcal{H}$  of  $\mathcal{H}$  comprises all groups  $G$  such that  $\text{Cov}_{\mathcal{H}}(G) \neq \emptyset$  namely all those groups that have  $\mathcal{H}$ -covering subgroups.  $D\mathcal{H}$  is also a homomorph. We study in [6] the set

$$\mathbb{H}(\mathcal{U}) = \{\mathcal{H} \mid D\mathcal{H} = \mathcal{U}\}, \text{ where } \mathcal{U} \text{ is a homomorph.}$$

Those homomorphs  $\mathcal{H}$  such that  $D\mathcal{H} = \mathcal{U}$  behave with regard to  $\mathcal{U}$  in a somewhat similar way to the Schunck classes with regard to the whole universe of soluble groups. The class  $\mathcal{A}(\mathcal{U})$  (see (2.1) of [6]) is introduced in order to characterize the homomorphs  $\mathcal{H}$  of  $\mathbb{H}(\mathcal{U})$ , when  $\mathbb{H}(\mathcal{U}) = \emptyset$  or  $|\mathbb{H}(\mathcal{U})| = 1$  and to study the relation of usual containment in  $\mathbb{H}(\mathcal{U})$ . The class  $\mathcal{A}(\mathcal{U})$  consists of those primitive groups  $G$  in  $\mathcal{U}$  that satisfy:

If  $M < X$  and  $X/\text{core}_X M \cong G$ , we have  $M \in \mathcal{U}$  if and only if  $X \in \mathcal{U}$ .

Let  $\mathcal{P}$  denote the class of finite soluble primitive groups.

If  $\mathbb{H}(\mathcal{U}) \neq \emptyset$ , the minimum in  $\mathbb{H}(\mathcal{U})$  with regard to the relation of containment is  $\mathcal{M} = h((b(\mathcal{U}) - \mathcal{P}) \cup \mathcal{A}(\mathcal{U}))$  (see [6], (3.3)).

(\*) Indirizzo dell'A.: Departamento de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, España.

In this paper we study the relation of strong containment in  $\mathbb{H}(\mathcal{U})$  given by

1 DEFINITION. Let  $\mathcal{U}$  be a homomorph. Let  $\mathcal{X}, \mathcal{Y} \in \mathbb{H}(\mathcal{U})$ . We say that  $\mathcal{X}$  is strongly contained in  $\mathcal{Y}$ , and write  $\mathcal{X} \ll \mathcal{Y}$  if, for each  $G \in \mathcal{U}$  an  $\mathcal{X}$ -covering subgroup of  $G$  is contained in some  $\mathcal{Y}$ -covering subgroup of  $G$ .

For a homomorph  $\mathcal{H}$ , we denote  $\tilde{\mathcal{H}} := h(b(\mathcal{H}) \cap \mathcal{P})$ . For every group  $G \in \mathbf{D}\mathcal{H}$  we have:  $\text{Cov}_{\mathcal{H}}(G) = \text{Cov}_{\tilde{\mathcal{H}}}(G)$  (see [6], (1.8)).

2 LEMMA. Let  $\mathcal{H}$  be a homomorph. We denote  $a(\mathcal{H}) := \{G \in \mathbf{D}\mathcal{H} \mid \text{if } H \in \text{Cov}_{\mathcal{H}}(G), H \cap \text{Soc } G = 1\}$ .

We have:

- a)  $a(\mathcal{H}) = a(\tilde{\mathcal{H}}) \cap \mathbf{D}\mathcal{H}$ .
- b)  $\tilde{\mathcal{H}} = h(a(\mathcal{H}))$ .

PROOF. a) It is evident by the definition.

b) Since  $b(\tilde{\mathcal{H}}) = b(\mathcal{H}) \cap \mathcal{P}$ , we have  $b(\tilde{\mathcal{H}}) \subseteq a(\tilde{\mathcal{H}}) \cap \mathbf{D}\mathcal{H} = a(\mathcal{H})$  and therefore  $h(a(\mathcal{H})) \subseteq h(b(\tilde{\mathcal{H}})) = \tilde{\mathcal{H}}$ . Since  $\tilde{\mathcal{H}} = h(a(\tilde{\mathcal{H}}))$  (see [2], VI (1.4)) and  $a(\mathcal{H}) \subseteq a(\tilde{\mathcal{H}})$ , we have  $\tilde{\mathcal{H}} = h(a(\tilde{\mathcal{H}})) \subseteq h(a(\mathcal{H}))$ .

Let us recall now the following

3 DEFINITION ([5] and [3] (8.2)). Let  $\mathcal{B} \subset \mathcal{P}$ . We define  $\mathcal{B}_0 = \mathcal{B}$ , and if  $\mathcal{B}_i$  has already been defined, let

$$\mathcal{B}_{i+1} = \left\{ (X/C_X(V))[V] \mid H \leq X \leq K < G = KF(G) \in \mathcal{B}_i, \right. \\ \left. H \in \text{Cov}_{h(\mathcal{B}_i)}(K), \text{ is } X\text{-composition factor of } F(G) \right\}.$$

We denote by  $\mathcal{B}^\infty$  the union of all class  $\mathcal{B}_i$  previously defined.

In a similar way to (8.3) from [3] we have

4 PROPOSITION. Let  $\mathcal{H}$  be a homomorph and  $\mathcal{B} \subset \mathcal{P}$  such that  $\mathcal{B} \subseteq a(\mathcal{H})$ . We have that  $\mathcal{B}^\infty \subseteq a(\mathcal{H})$  (in particular  $a(\mathcal{H})^\infty = a(\mathcal{H})$ ).

PROOF. Let us prove that  $\mathcal{B}_i \subseteq a(\mathcal{H})$  for every  $i \in \mathbb{N}$ . We proceed by induction on  $i$ . We have that  $\mathcal{B} = \mathcal{B}_0 \subseteq a(\mathcal{H})$ . Suppose  $\mathcal{B}_i \subseteq a(\mathcal{H})$ . Let  $B \in \mathcal{B}_{i+1}$ . There exists  $G \in \mathcal{B}_i \subseteq a(\mathcal{H})$ ,  $Y \leq X \leq K$ ,  $K$  complement of  $F(G)$ ,  $H \in \text{Cov}_{h(\mathcal{B}_i)}(K)$ ,  $V, W, X$ -subgroups of  $F(G)$ ,  $V/W, X$ -composition

of  $F(G)$  such that  $B = X/C_X(V/W)[(V/W)]$ . Since  $\mathcal{B}_i \subseteq a(\mathcal{H}) \subseteq a(\tilde{\mathcal{H}})$ , by [1] (2.2), we have  $\tilde{\mathcal{H}} \ll h(\mathcal{B}_i)$ , hence there exists  $H \in \text{Cov}_{\tilde{\mathcal{H}}}(K)$  such that  $H \leq Y$ . As  $G \in a(\mathcal{H}) \subseteq \mathbf{D}\mathcal{H}$ , we have  $H \in \text{Cov}_{\mathcal{H}}(K) \subseteq \text{Cov}_{\mathcal{H}}(G)$ . Besides, it can be confirmed that

$$B = X/C_X(V/W)[(V/W)] \cong XV/C_X(V/W)W.$$

By the properties of covering subgroups  $H \in \text{Cov}_{\mathcal{H}}(XV)$  and

$$HC_X(V/W)W/C_X(V/W)W \in \text{Cov}_{\mathcal{H}}(XV/C_X(V/W)W),$$

therefore  $B \in \mathbf{D}\mathcal{H}$ . We know from [3] (8.3), that  $B \in a(\tilde{\mathcal{H}})$ , so we can deduce that  $B \in a(\tilde{\mathcal{H}}) \cap \mathbf{D}\mathcal{H} = a(\mathcal{H})$ .

Below we study the relation « $\ll$ » in  $\mathbb{H}(\mathcal{U})$ .

**5 PROPOSITION.** Let  $\mathcal{X}, \mathcal{Y} \in \mathbb{H}(\mathcal{U})$ . We have  $\mathcal{X} \ll \mathcal{Y}$  if and only if  $\tilde{\mathcal{X}} \ll \tilde{\mathcal{Y}}$ .

**PROOF.**  $\Leftarrow$ ) It is evident from that comment before Lemma 2.

$\Rightarrow$ ) We have  $b(\tilde{\mathcal{Y}}) = b(\mathcal{Y}) \cap \mathcal{P}$ . By definition of  $\ll$  and  $a(\mathcal{X})$ , we have that  $b(\mathcal{Y}) \cap \mathcal{P} = b(\mathcal{Y}) \cap \mathbf{D}\mathcal{Y} \subseteq a(\mathcal{X})$ . Moreover,  $a(\mathcal{X}) \subseteq a(\tilde{\mathcal{X}})$ , hence  $b(\tilde{\mathcal{Y}}) \subseteq b(\tilde{\mathcal{X}})$  and by [1] (2.2),  $\tilde{\mathcal{X}} \ll \tilde{\mathcal{Y}}$ .

Since the mapping  $\mathcal{H} \rightarrow \tilde{\mathcal{H}}$  from  $\mathbb{H}(\mathcal{U})$  to the set of Schunck classes is injective (see [6], 3.1),  $\mathbb{H}(\mathcal{U})$  can be considered a subset of the Schunck classes ordered by  $\ll$ .

In the examples described in [6] (1.9), (3.8), (3.9),  $(\mathbb{H}(\mathcal{U}), \ll)$  has a lattice structure. In these examples we have  $\mathcal{A}(\mathcal{U}) = a(\mathcal{H})$ . In this respect, we can say:

**6 PROPOSITION.** Let  $\mathcal{U}$  be a homomorph and  $\mathcal{H}$  the minimum for  $\subseteq$  in  $\mathbb{H}(\mathcal{U})$ . The following statements are equivalent:

- a)  $\mathcal{A}(\mathcal{U}) = a(\mathcal{H})$ ;
- b)  $\mathcal{A}(\mathcal{U})^\infty = \mathcal{A}(\mathcal{U})$ .

**PROOF.**  $a) \Rightarrow b)$  It follows immediately from Proposition 4.

$b) \Rightarrow a)$  By  $b)$  we obviously have  $\mathcal{A}(\mathcal{U})^\infty \cap h(\mathcal{A}(\mathcal{U})) = \emptyset$ . By [3] (8.4), we have  $\mathcal{A}(\mathcal{U}) \subseteq a(h(\mathcal{A}(\mathcal{U})))$ . By [6] (3.3),  $h(\mathcal{A}(\mathcal{U})) = \tilde{\mathcal{H}}$  and therefore  $\mathcal{A}(\mathcal{U}) \subseteq a(\tilde{\mathcal{H}})$ .

Besides,  $\mathfrak{A}(\mathfrak{U}) \subseteq \mathfrak{U} = \mathbf{D}\mathfrak{M}$  implies  $\mathfrak{A}(\mathfrak{U}) \subseteq a(\tilde{\mathfrak{M}}) \cap \mathbf{D}\mathfrak{M} = a(\mathfrak{M})$ . By [6] (1.7), we have  $a(\mathfrak{M}) \subseteq \mathfrak{A}(\mathfrak{U})$  and therefore the equality.

7 THEOREM. Let  $\mathfrak{U}$  be a homomorph such that  $b(\mathfrak{U}) \cap \mathcal{P} = \emptyset$ . (These homomorphs are known as totally unsaturated).

$(\mathbb{H}(\mathfrak{U}), \ll)$  is a lattice if and only if  $\mathfrak{A}(\mathfrak{U})^\infty = \mathfrak{A}(\mathfrak{U})$ .

PROOF.  $\Rightarrow$  By the proposition above and [6] (1.7), it suffices to prove that  $\mathfrak{A}(\mathfrak{U}) \subseteq a(\mathfrak{M})$ . Let  $G \in \mathfrak{A}(\mathfrak{U})$ . Let  $\mathcal{X} = h(b(\mathfrak{U}) \cup \{G\})$ . By [6] (2.3),  $\mathcal{X} \in \mathbb{H}(\mathfrak{U})$ . Since  $\ll$  implies  $\subseteq$ , the infimum of  $\{\mathcal{X}, \mathfrak{M}\}$  must be  $\mathfrak{M}$ . Thus  $\mathfrak{M} \ll \mathcal{X}$ , therefore  $\tilde{\mathfrak{M}} \ll \tilde{\mathcal{X}}$  and consequently  $b(\tilde{\mathcal{X}}) \subseteq a(\tilde{\mathfrak{M}})$ . As  $\{G\} = b(\tilde{\mathcal{X}})$ , we have that

$$G \in a(\tilde{\mathfrak{M}}) \cap \mathfrak{U} = a(\tilde{\mathfrak{M}}) \cap \mathbf{D}\mathfrak{M} = a(\mathfrak{M}).$$

$\Leftarrow$ ) Let  $\mathcal{X}, \mathcal{Y} \in \mathbb{H}(\mathfrak{U})$ . Recall from [5] Theorem A that

$$\tilde{\mathcal{X}} \wedge \tilde{\mathcal{Y}} = h((b(\tilde{\mathcal{X}}) \cup b(\tilde{\mathcal{Y}}))^\infty).$$

By Proposition 6 we have  $\mathfrak{A}(\mathfrak{U})^\infty = \mathfrak{A}(\mathfrak{U}) = a(\mathfrak{M})$ . Since  $b(\tilde{\mathcal{X}}) \cup b(\tilde{\mathcal{Y}}) \subseteq a(\mathfrak{M})$ , by Proposition 4, we have that  $(b(\tilde{\mathcal{X}}) \cup b(\tilde{\mathcal{Y}}))^\infty \subseteq a(\mathfrak{M})$  and therefore  $b(\tilde{\mathcal{X}} \wedge \tilde{\mathcal{Y}}) \subseteq \mathfrak{A}(\mathfrak{U})$ . By [6] (2.3), we have that  $\mathcal{X} = h(b(\mathfrak{U}) \cup b(\tilde{\mathcal{X}} \wedge \tilde{\mathcal{Y}})) \in \mathbb{H}(\mathfrak{U})$  and it can easily be confirmed that  $\mathcal{X} = \mathcal{X} \wedge \mathcal{Y}$ .

Now let,  $\mathcal{Y} = h(a(\mathcal{X}) \cap a(\mathcal{Y}))$ . Again by the characterization in [6] (2.3) and (3.1), of the homomorphs in  $\mathbb{H}(\mathfrak{U})$  we have that  $\mathcal{Z} = \mathcal{Y} \cap \mathfrak{U} \in \mathbb{H}(\mathfrak{U})$ , and  $\mathcal{Y} = \tilde{\mathcal{Z}}$ . It can be confirmed that  $\mathcal{Z} = \mathcal{X} \vee \mathcal{Y}$ .

8 PROPOSITION. Let  $\mathfrak{U}$  be a totally unsaturated homomorph such that  $(\mathbb{H}(\mathfrak{U}), \ll)$  is a lattice. For every  $\mathcal{X}, \mathcal{Y} \in \mathbb{H}(\mathfrak{U})$  we have:

a)  $\widetilde{\mathcal{X} \wedge \mathcal{Y}} = \tilde{\mathcal{X}} \wedge \tilde{\mathcal{Y}}$ .

b)  $\mathcal{X} \ll \mathcal{Z} \neq \mathfrak{U}$  implies  $\mathcal{X} = \mathcal{Z}$  if and only if  $|b(\mathcal{X}) \cap \mathcal{P}| = 1$ .

PROOF. a) It is clear from the previous proof that

$$b(\mathcal{X} \wedge \mathcal{Y}) \cap \mathcal{P} = b(\tilde{\mathcal{X}} \wedge \tilde{\mathcal{Y}}).$$

b)  $\Rightarrow$ ) If  $|b(\mathcal{X}) \cap \mathcal{P}| \neq 1$ , we can have  $\emptyset \neq \mathcal{B} \subset b(\mathcal{X}) \cap \mathcal{P} \subseteq \mathfrak{A}(\mathfrak{U})$ . Now  $\mathcal{Z} = h(b(\mathfrak{U}) \cup \mathcal{B}) \in \mathbb{H}(\mathfrak{U})$ ,  $\mathcal{Z} \neq \mathcal{X}$  and  $\mathcal{X} \ll \mathcal{Z} \neq \mathfrak{U}$  in contradiction with the hypothesis.

$\Leftarrow$ ) As  $\tilde{\mathcal{X}} = h(b(\mathcal{X}) \cap \mathcal{P})$ ,  $\tilde{\mathcal{X}}$  is maximal, hence  $\tilde{\mathcal{X}} \ll \tilde{\mathcal{Z}} \neq \mathfrak{U}$  implies  $\tilde{\mathcal{X}} = \tilde{\mathcal{Z}}$  and by Proposition 5 we have the thesis.

## REFERENCES

- [1] K. DOERK, *Über Homomorph endlicher auflösbarer Gruppen*, J. Algebra, **30** (1974), pp. 12-30.
- [2] K. DOERK - T. O. HAWKES, *Finite Soluble Groups*, Walter de Gruyter, Berlin, New York (1992).
- [3] P. FÖRSTER, *Charakterisierungen einiger Schunckklassen endlicher auflösbarer Gruppen III*, J. Algebra, **62** (1980), pp. 124-153.
- [4] P. FÖRSTER, *Über die iterierten Definitionsbereiche von Homomorphen endlicher auflösbarer Gruppen*, Arch. Math., **35** (1980), pp. 27-41.
- [5] T. HAWKES, *The family of Schunck classes as a lattice*, J. Algebra, **39** (1976), pp. 527-550.
- [6] P. JIMÉNEZ-SERAL, *Homomorphs with relative Schunck classes*, Communication in Algebra, **19**, no. 10 (1991), pp. 2687-2707.

Manoscritto pervenuto in redazione il 6 febbraio 1992.