

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

ERIC A. CARLEN

DANIEL W. STROOCK

An application of the Bakry-Émery criterion to infinite dimensional diffusions

Séminaire de probabilités (Strasbourg), tome 20 (1986), p. 341-348

http://www.numdam.org/item?id=SPS_1986__20__341_0

© Springer-Verlag, Berlin Heidelberg New York, 1986, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

An Application of the Bakry-Emerly Criterion to Infinite Dimensional Diffusions

by

Eric A. Carlen

and

Daniel W. Stroock*

The note [1] by Bakry and Emery contains an important criterion with which to check whether a diffusion semigroup is hypercontractive. Although Bakry and Emery's interest in their criterion stems from its remarkable ability to give best constants in certain finite dimensional examples, what will concern us here is its equally remarkable ability to handle some infinite dimensional situations.

We begin by recalling their criterion in the setting with which we will be dealing. Let M be a connected, compact, N -dimensional smooth manifold with Riemannian metric g . Let Φ be a smooth function on M and define the differential operator L by

$$Lf = 1/2 \exp(\Phi) \operatorname{div}(\exp(-\Phi) \operatorname{grad}(f)), \quad f \in C^\infty(M),$$

and the probability measure m by

$$m(dx) = \exp(-\Phi(x)) \lambda(dx) / \int \exp(-\Phi(y)) \lambda(dy),$$

where λ denotes the Riemann measure on M associated with the metric g . Next, use $\{P_t : t > 0\}$ to denote the diffusion semigroup determined by L . The following facts about $\{P_t : t > 0\}$ are easy to check:

i) $\{P_t : t > 0\}$ on $C(M)$ is a strongly continuous, conservative Markov semigroup under which $C^\infty(M)$ is invariant.

ii) $\{P_t : t > 0\}$ is m -reversible (i.e. P_t is symmetric in $L^2(m)$ for each $t > 0$) and $\|P_t f - \int f dm\|_{C(M)} \rightarrow 0$ as $t \rightarrow \infty$ for each $f \in C(M)$. In particular, for all $t > 0$ and $p \in [1, \infty)$,

$$\|P_t\|_{L^p(m) \rightarrow L^p(m)} = 1$$

and there is a unique strongly continuous semigroup $\{\bar{P}_t : t > 0\}$ on $L^2(m)$ such that $\bar{P}_t f = P_t f$ for all $t > 0$ and $f \in C(M)$.

As a consequence, note that, for each $f \in L^2(M)$, $t \rightarrow (f - \bar{P}_t f, f)_{L^2(M)} / t$ is a non-negative, non-increasing function and that, therefore, the Dirichlet form given by

$$\mathfrak{E}(f, f) = \lim_{t \downarrow 0} (f - \bar{P}_t f, f)_{L^2(M)} / t$$

exists (as an element of $[0, \infty]$) for each $f \in L^2(M)$.

Theorem (Bakry and Emery): Denote by $H_{\mathfrak{g}}$ the (covariant) Hessian tensor of \mathfrak{g} (i.e. $H_{\mathfrak{g}}(X, Y) = X \cdot Y \mathfrak{g} - \nabla_X Y \mathfrak{g}$ for $X, Y \in \Gamma(T(M))$) and let Ric be the Ricci curvature on (M, g) . If, as quadratic forms, $\text{Ric} + H_{\mathfrak{g}} \geq a g$ for some $a > 0$, then the logarithmic Sobolev inequality :

$$(L.S.) \quad \int f^2 \log f^2 dm \leq 4/a \mathfrak{E}(f, f) + \|f\|_{L^2(M)}^2 \log \|f\|_{L^2(M)}^2, \quad f \in L^2(M)$$

and, therefore, the hypercontractive estimate :

$$(H.C.) \quad \begin{aligned} & \|P_t\|_{L^p(M) \rightarrow L^q(M)} = 1, \\ & 1 < p \leq q < \infty \text{ and } t > 0 \text{ with } e^{at} \geq (q-1)/(p-1) \end{aligned}$$

hold.

Remark : Actually, Bakry and Emery's result is somewhat more refined than the one just stated. However, the refinement seems to become less and less significant as N becomes large. Since we are interested here in what happens as $N \rightarrow \infty$, the stated result will suffice.

Remark : Several authors (e.g. O. Raths [5]) have observed that a logarithmic Sobolev inequality implies a gap in the spectrum of L . To be precise, (L.S.) implies that:

$$(S.G.) \quad \|f - \int f dm\|_{L^2(M)}^2 \leq 2/a \mathfrak{E}(f, f), \quad f \in L^2(M),$$

or, equivalently,

$$(S.G.') \quad \|\bar{P}_t f - \int f dm\|_{L^2(M)} \leq \exp(-\alpha t/2) \|f\|_{L^2(M)}, \quad f \in L^2(M).$$

We now turn to the application of the Bakry-Emery result to infinite

dimensional diffusions. For the sake of definiteness, let $d \geq 2$ and $\nu \geq 1$ be given, and, for $n \geq 1$, set

$$M_n = (S^d)^{\wedge n},$$

where $\wedge_n = \{k \in Z^\nu : \|k\| = \max_{1 \leq i \leq \nu} |k_i| \leq n\}$, and give M_n the product Riemannian structure which it inherits from the standard structure on S^d . Let π_k be the natural projection map from M_n onto the k^{th} sphere S^d , and, for $X \in \Gamma(T(M_n))$, set $X^{(k)} = (\pi_k)_* X$. Noting, as was done in [1], that on S^d the Ricci curvature is equal to $(d - 1)$ times the metric, we see that the Ricci curvature Ric_n and the metric g_n on M_n satisfy the same relationship. Finally, let $\Phi_n \in C^\infty(M_n)$ be given and define the operator L_n , the measure m_n , the semigroup $\{P_t^n : t > 0\}$, and the Dirichlet form \mathcal{E}_n accordingly. As an essentially immediate consequence of the Bakry-Emery theorem, we have the following.

Theorem : Assume that for all $X \in \Gamma(T(M_n))$:

$$|\mathbb{H}_{\Phi_n}(X, X)| \leq \sum_{k, \lambda \in \wedge_n} \gamma(k - \lambda) \|X^{(k)}\| \|X^{(\lambda)}\|$$

where $\gamma : Z^\nu \rightarrow [0, \infty)$ satisfies

$$\sum_{k \in Z^\nu} \gamma(k) \leq (1 - \epsilon)(d - 1)$$

for some $0 < \epsilon < 1$. Set $\alpha = \epsilon(d - 1)$. Then:

$$(L.S.)_n \quad \int f^2 \log f^2 dm_n \leq (4/\alpha) \mathcal{E}_n(f, f) + \|f\|_{L^2(m_n)}^2 \log \|f\|_{L^2(m_n)}^2,$$

for $f \in L^2(m_n)$ and

$$(H.C.)_n \quad \begin{aligned} &\|P_t^n\|_{L^p(m_n) \rightarrow L^q(m_n)} = 1, \\ &1 < p \leq q < \infty \text{ and } t > 0 \text{ with } \exp(\alpha t) \geq (q - 1)/(p - 1). \end{aligned}$$

In particular,

$$(S.G.)_n \quad \|f - \int f dm_n\|_{L^2(m_n)}^2 \leq (2/\alpha) \mathcal{E}_n(f, f), \quad f \in L^2(m_n),$$

and

$$(S.G.')_n \quad \|P_t^n f - \int f dm_n\|_{L^2(m_n)} \leq \exp(-\alpha t/2) \|f\|_{L^2(m_n)}, \quad f \in L^2(m_n).$$

Proof : Simply observe that, by Young's inequality, the bound

on H_{Φ_n} (as a quadratic form) in terms of g can be dominated by $\|\gamma\|_{\mathcal{Q}^1(Z^{\nu})}$.

To complete our program, set $M_{\infty} = (S^d)^{Z^{\nu}}$, $\mathcal{F} = \{F \subseteq Z^{\nu} : \text{card}(F) < \infty\}$, and, for $F \in \mathcal{F}$, denote by π_F the natural projection of M_{∞} onto $(S^d)^F$. (Thus, in the notation used before, $\pi_k = \pi_{\{k\}}$ and $M_n = (S^d)^{\wedge n}$.) Next, set $\mathcal{D}_F = \{f \circ \pi_F : f \in C^{\infty}((S^d)^F)\}$, $\mathcal{D} = \bigcup \{\mathcal{D}_F : F \in \mathcal{F}\}$, and let $\Gamma(T(M_{\infty}))$ be the set of derivations from \mathcal{D} into itself. We now suppose that we are given a potential $\mathcal{J} = \{J_F : F \in \mathcal{F}\}$, where:

i) for each $F \in \mathcal{F}$, $J_F \in \mathcal{D}_F$, and for each $k \in Z^{\nu}$ there are only a finite number of $F \ni k$ for which J_F is not identically zero,

ii) there is a constant $B < \infty$ such that

$$\sum_{F \ni k} |X^{(k)} J_F| \leq B \|X^{(k)}\|, \quad k \in Z^{\nu} \text{ and } X \in \Gamma(T(M_{\infty})),$$

iii) there is a $\gamma : Z^{\nu} \rightarrow [0, \infty)$ such that

$$\sum_{k \in Z^{\nu}} \gamma(k) < \infty$$

and

$$\sum_{F \supseteq \{k, \ell\}} |H_{J_F}(X^{(k)}, X^{(\ell)})| \leq \gamma(k - \ell) \|X^{(k)}\| \|X^{(\ell)}\|$$

for all $k, \ell \in Z^{\nu}$ and $X \in \Gamma(T(M_{\infty}))$.

Set $H_k = \sum_{F \ni k} J_F$ and define L_{∞} on \mathcal{D} by

$$L_{\infty} f = 1/2 \sum_{k \in Z^{\nu}} \exp(H_k) \text{div}_k (\exp(-H_k) \text{grad}_k f)$$

where "div_k" and "grad_k" refer to the corresponding operations in the directions of the k^{th} sphere.

In order to describe the measure m_{∞} , we will need to introduce the concept of a Gibbs state and this, in turn, requires us to develop a little more notation. For $n \geq 1$ and $x, y \in M_{\infty}$, define $Q_n(x|y) \in M_{\infty}$ by

$$Q_n(x|y)_k = \begin{cases} x_k & \text{if } k \in \Lambda_n \\ y_k & \text{if } k \notin \Lambda_n. \end{cases}$$

(It will be convenient, and should cause no confusion, for us to sometimes consider $x \rightarrow Q_n(x|y)$, for fixed $y \in M_\infty$, as a function on M_n and $y \rightarrow Q_n(x|y)$, for fixed $x \in M_\infty$, as a function on $(S^d)^{\Lambda_n^c}$.) Define

$$\Phi_n(x|y) = \sum_{F \wedge \Lambda_n \neq \emptyset} J_F \circ Q_n(x|y)$$

and let $m_n(\cdot|y)$ denote the probability measure on M_n associated with $\Phi_n(\cdot|y)$.

We will say that a probability measure m_∞ on M_∞ is a Gibbs state with potential \mathcal{J} and will write $m_\infty \in \mathcal{G}(\mathcal{J})$ if, for each $n \geq 1$, $y \rightarrow m_n(\cdot|y)$ is a regular conditional probability distribution of m_∞ given $\sigma(x_k: k \in \Lambda_n^c)$.

The following lemma summarizes some of the reasonably familiar facts about the sort of situation described above (cf. [2] and [3]).

Lemma : There is exactly one conservative Markov semigroup $\{P_t^\infty: t > 0\}$ on $C(M_\infty)$ such that

$$P_T^\infty f - f = \int_0^T P_t^\infty L_\infty f dt, \quad f \in \mathcal{D}.$$

Moreover, if, for each $n \geq 1$, $\Phi_n \in C^\infty(M_n)$ and the associated operator L_n are given, and if $[L_n(f \circ Q_n(\cdot|y))](x) \rightarrow L_\infty f(x)$ uniformly in $x, y \in M_\infty$ for every $f \in \mathcal{D}$, then the associated semigroups $\{P_t^n: t > 0\}$ have the property that

$$[P_t^n f \circ Q_n(\cdot|y)](x) \rightarrow P_t^\infty f(x)$$

uniformly in $(t, x, y) \in [0, T] \times M_\infty \times M_\infty$ for every $T > 0$ and $f \in C(M_\infty)$. Finally:

$\mathcal{G}(\mathcal{J})$ is a non-empty, compact, convex set; $m_\infty \in \mathcal{G}(\mathcal{J})$ if and only if it is a $\{P_t^\infty: t > 0\}$ -reversible measure; and for each extreme element m_∞ of $\mathcal{G}(\mathcal{J})$ there is a $y \in M_\infty$ such that $m_n(\cdot|y) \rightarrow m_\infty$.

Theorem : Referring to the situation described above, assume that

$$\sum_{k \in \mathbb{Z}^v} \gamma(k) \leq (1 - \epsilon)(d - 1)$$

for some $0 < \epsilon < 1$ and that m_∞ is an extreme element of $\mathfrak{M}(\mathfrak{T})$ (cf. the remark below). Denote by ξ_∞ the Dirichlet form determined on $L^2(m_\infty)$ by $\{P_t^\infty : t > 0\}$.

Then, for $f \in L^2(m_\infty)$:

$$(L.S.)_\infty \quad \int f^2 \log f^2 dm_\infty \leq (4/\alpha) \xi_\infty(f, f) + \|f\|_{L^2(m_\infty)}^2 \log \|f\|_{L^2(m_\infty)}^2,$$

where $\alpha = \epsilon(d - 1)$. In particular,

$$(H.C.)_\infty \quad \begin{aligned} & \|P_t^\infty\|_{L^p(m_\infty) \rightarrow L^q(m_\infty)} = 1, \\ & 1 < p \leq q < \infty \text{ and } t > 0 \text{ with } \exp(\alpha t) \geq (q - 1)/(p - 1), \end{aligned}$$

$$(S.G.)_\infty \quad \|f - \int f dm_\infty\|_{L^2(m_\infty)}^2 \leq 2/\alpha \xi_\infty(f, f), \quad f \in L^2(m_\infty),$$

and

$$(S.G.')_\infty \quad \|P_t^\infty f - \int f dm_\infty\|_{L^2(m_\infty)} \leq \exp(-\alpha t/2) \|f\|_{L^2(m_\infty)}, \quad f \in L^2(m_\infty),$$

where $\{P_t^\infty : t > 0\}$ is the contraction semigroup on $L^2(m_\infty)$ determined by $\{P_t^\infty : t > 0\}$.

Proof : Choose $y \in M_\infty$ so that $m_n = m_n(\cdot | y) \rightarrow m_\infty$. Set $\Phi_n = \Phi_n(\cdot | y)$ and define L_n and $\{P_t^n : t > 0\}$ accordingly. It is easy to check that the hypotheses of the previous theorem are satisfied for each n . In particular, $(H.C.)_n$ holds for all $n \geq 1$. Moreover, the preceding lemma allows us to conclude that

$$\begin{aligned} & \|P_t^\infty\|_{L^p(m_\infty) \rightarrow L^q(m_\infty)} \\ & \leq \liminf_{n \rightarrow \infty} \|P_t^n\|_{L^p(m_n) \rightarrow L^q(m_n)} \end{aligned}$$

for all $1 \leq p \leq q < \infty$ and $t > 0$. Hence, we now know that $(H.C.)_\infty$ holds.

Since $(L.S.)_\infty$, $(S.G.)_\infty$, and $(S.G.')_\infty$ all follow from $(H.C.)_\infty$, the proof is complete.

Remark: It turns out that the hypotheses in the preceding theorem allow one to conclude that $\mathfrak{D}(\mathfrak{V})$ contains precisely one element. In addition, when the potential \mathfrak{V} is shift invariant (ie. $J_{F+k} = J_F \circ S^k$ for all k and F , where S is the natural shift operation on $C(M_\infty)$) and has finite range (ie. there is a cube Λ such that $J_F = 0$ for all $F \ni 0$ for which $F \not\subset \Lambda$), one can show that for each shift invariant probability measure μ on M_∞ and all $f \in \mathfrak{D}$ there is an $A(f) \in (0, \infty)$ (not depending on μ) with the property that

$$|\int P_t^\infty f d\mu - \int f d\mu_\infty| \leq A(f) \exp(-at/2).$$

These and related results will be the topic of a forthcoming article by the second of the present authors and R. Holley [4].

REFERENCES

- [1] Bakry, D. and Emery, M., "Hypercontractivite de semi-groupes de diffusion", C.R. Acad. Sc. Paris, t. 299, Serie I no.15 (1984).
- [2] Holley, R. and Stroock, D., " L^2 theory for the stochastic Ising model", Z. Wahr. 35, pp. 87-101 (1976).
- [3] Holley, R. and Stroock, D., "Diffusions on an infinite dimensional Torus", J. Fnal. Anal., vol. 42 no.1, pp. 29-63 (1981).
- [4] Holley, R. and Stroock, D., "Logarithmic Sobolev inequalities and stochastic Ising models", to appear in the issue of J. Statistical Physics dedicated to the memory of M. Kac.
- [5] Rothaus, O., "Logarithmic Sobolev inequalities and the spectrum of Schroedinger operators", J. Fnal. Anal., vol. 42 no.1, pp. 110-120 (1981).

*During the period of this reseach, this author was partially supported by NSF DMS-8415211 and ARO DAAG29-84-K-0005. The address of both authors is Dept. of Mathematics, M.I.T., Cambridge, MA 02140, U.S.A.