



Partial differential equations

On the stability of the state 1 in the non-local Fisher–KPP equation in bounded domains

Sur la stabilité de l'état 1 dans l'équation de Fisher–KPP non locale en domaine borné

Camille Pouchol^{a,b}

^a Sorbonne Université, UPMC Université Paris-6, CNRS UMR 7598, Laboratoire Jacques-Louis-Lions, 75005 Paris, France

^b INRIA Team Mamba, INRIA Paris, 2, rue Simone- Iff, CS 42112, 75589 Paris, France

ARTICLE INFO

Article history:

Received 17 January 2018

Accepted after revision 12 April 2018

Available online 25 April 2018

Presented by Haïm Brézis

ABSTRACT

We consider the non-local Fisher–KPP equation on a bounded domain with Neumann boundary conditions. Thanks to a Lyapunov function, we prove that, under a general hypothesis on the kernel involved in the non-local term, the homogenous steady state 1 is globally asymptotically stable. This assumption happens to be linked to some conditions given in the literature, which ensure that travelling waves link 0 to 1.

© 2018 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

RÉSUMÉ

Nous considérons l'équation de Fisher–KPP non locale en domaine borné, avec conditions de Neumann au bord. À l'aide d'une fonction de Lyapunov, nous montrons que, sous une hypothèse générale sur le noyau présent dans le terme non local, l'état stationnaire 1 est globalement asymptotiquement stable. Cette hypothèse se trouve être reliée à certaines conditions données dans la littérature, qui assurent que les fronts de propagation relient 0 et 1.

© 2018 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

E-mail address: pouchol@jll.math.upmc.fr.

<https://doi.org/10.1016/j.crma.2018.04.016>

1631-073X/© 2018 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

We consider the so-called non-local Fisher–KPP equation endowed with Neumann boundary conditions

$$\begin{aligned} \frac{\partial u}{\partial t}(t, x) &= \mu \left(1 - \int_{\Omega} K(x, y) u(t, y) dy \right) u(t, x) + \Delta u(t, x), \quad x \in \Omega, \quad t > 0, \\ \frac{\partial u}{\partial n}(t, x) &= 0, \quad x \in \partial\Omega, \quad t > 0, \\ u(0, x) &= u^0(x) \geq 0 \quad x \in \Omega, \end{aligned} \tag{1}$$

where Ω is a regular bounded domain of \mathbb{R}^d and $K > 0$ a kernel modelling an additional death rate due to non-local interactions.

We will sometimes write in short $K[u] = \int_{\Omega} K(x, y) u(y) dy$ for a generic function u .

Assuming

$$\forall x \in \Omega, \int_{\Omega} K(x, y) dy = 1, \tag{2}$$

and in the limit $K(x, y) \rightarrow \delta_{x-y}$, we recover the classical Fisher KPP equation

$$\frac{\partial u}{\partial t} = \mu(1 - u)u + \Delta u. \tag{3}$$

The assumption (2) ensures that 1 remains a homogeneous stationary solution to (1).

The classical Fisher–KPP equation (3) is often analysed on the whole space for the investigation of travelling waves, which are known to exist since the pioneering works of Fisher, Kolmogorov, Petrovsky, and Piskunov [7] for any speed above $2\sqrt{\mu}$. Furthermore, any non-zero initial condition eventually converges locally uniformly to 1, which is therefore a globally asymptotically stable for non-zero initial conditions.

When one adds a non-local term, it does not remain true that travelling waves exist and when they do, whether they link 0 to 1 or to another non-homogeneous steady state of the equation. 1 can indeed become unstable: Turing patterns appear [8,9].

A natural question is thus to understand under which conditions the status of 1 is changed due to the non-local term. When $K(x, y)$ is given by a convolution $\phi(x - y)$, several results have already been obtained in the full space, in dimension $d = 1$. If the Fourier transform is everywhere positive or if μ is small enough, it is known that travelling waves necessarily connect 0 to 1 [2]. See also [1,5].

In this note, we provide a general result on the global asymptotic stability on 1 on a bounded domain, based on a Lyapunov functional. The result holds provided that the following general assumption on the kernel K is satisfied:

$$\forall f \in L^2(\Omega), \int_{\Omega \times \Omega} K(x, y) f(x) f(y) dx dy \geq 0. \tag{4}$$

K is then referred to as being a positive kernel, and (4) can be thought of as a strong competition assumption. These types of Lyapunov functionals have been used successfully in selection equations in [6,10,11], and are inspired by Lyapunov functions for Lotka–Volterra ODEs [4].

It remains an open question to know whether this condition leads to the same conclusion on the whole space. As such, our Lyapunov function requires integrability for $u(t) - 1 - \ln(u(t))$, which is too much to ask in \mathbb{R}^d . We still believe that the condition (4) is highly relevant. Indeed, when $\Omega = \mathbb{R}^d$, and if K is a convolution $K(x, y) = \phi(x - y)$, then condition (4) becomes

$$\forall f \in L^2(\mathbb{R}^d), \int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(x - y) f(x) f(y) dx dy \geq 0. \tag{5}$$

It is easy to check that, if ϕ has a non-negative Fourier transform, then condition is satisfied, see [6]. The converse is almost true, as evidenced by Bochner’s Theorem [12]: if ϕ is bounded and continuous, then (4) holds if and only if ϕ is the Fourier transform of a finite bounded measure on \mathbb{R}^d .

Consequently, condition (4) shows that the condition on the Fourier transform of ϕ used in dimension 1 in the literature can be appropriate in any dimension, and may not only be a sufficient but also a necessary condition when it comes to the stability of the state 1.

2. The Lyapunov function approach

We make the following regularity assumption on the kernel K :

$$K \in C^{0,1}(\overline{\Omega} \times \overline{\Omega}), \tag{6}$$

where $C^{0,1}(\overline{\Omega} \times \overline{\Omega})$ denotes the set of Lipschitz continuous functions on $\overline{\Omega} \times \overline{\Omega}$.

Under the previous assumption (6), for $u^0 \in L^1(\Omega)$, we know from [3] that there exists a unique non-negative classical solution in $C([0, +\infty), L^1(\Omega)) \cap C^1((0, +\infty), C^{2,\alpha}(\Omega))$, which we denote $t \mapsto S_t u^0$.

It will also be convenient to introduce the set $Z := \{u \in C^{2,\alpha}(\Omega), u \geq 0\}$. Finally, we define the non-negative function $H(w) := w - 1 - \ln(w)$ for $w > 0$, and for u in Z

$$V(u) := \int_{\Omega} (u(x) - 1 - \ln(u(x))) \, dx, \tag{7}$$

the last integral possibly being equal to $+\infty$.

Our result is then the following.

Theorem 1. *Assume (4), (6), (2). Then, for any initial datum u^0 in $L^1(\Omega)$, $u^0 \geq 0$, $u^0 \neq 0$, the solution to (1) satisfies*

$$u(t, \cdot) \longrightarrow 1$$

uniformly in Ω .

Proof. *First step: computation of the Lyapunov functional.*

First, let us remark that by the parabolic strong maximum principle, $u(t, x) > 0$ for all $t > 0, x \in \Omega$. Now, let us check that this holds true also for $x \in \partial\Omega$, from which we will infer that $V(u(t))$ is well defined for all $t > 0$. By the parabolic strong maximum principle at the boundary, we have the following alternative for $x \in \partial\Omega$: either $u(t, x) > 0$ or $u(t, x) = 0$ and $\frac{\partial u}{\partial n}(t, x) < 0$. Only $u(t, x) > 0$ can hold due to the Neumann boundary conditions.

We now consider $g(t) := V(u(t))$ for $t > 0$, where $\{u(t)\}_{t \geq 0}$ is the trajectory emanating from u_0 . Let us prove that this is a Lyapunov functional, by computing for $t > 0$

$$\begin{aligned} g'(t) &= \int_{\Omega} \frac{\partial u}{\partial t}(t) \left(1 - \frac{1}{u(t)}\right) \\ &= \int_{\Omega} \Delta u(t) \left(1 - \frac{1}{u(t)}\right) - \mu \int_{\Omega} (1 - K[u(t)])(1 - u(t)) \\ &= - \int_{\Omega} \frac{|\nabla(u(t, x))|^2}{u^2(t, x)} \, dx - \mu \int_{\Omega^2} K(x, y) (1 - u(t, x))(1 - u(t, y)) \, dx \, dy, \end{aligned}$$

after integration by part for the first term. For the second one, we used $1 - K[u] = K[1 - u]$, owing to (2).

Thanks to (4), this yields $g'(t) \leq 0$, that is, g is non-increasing over \mathbb{R}_+ . Since $g \geq 0$, we infer the convergence of $g(t)$ as t tends to $+\infty$, and we denote by l its limit.

Second step: compactness of trajectories.

Since $C^{2,\alpha}(\Omega)$ is compactly embedded into $C(\overline{\Omega})$, the trajectory $\{S_t u^0\}_{t \geq \delta}$ (for some fixed $\delta > 0$) is relatively compact in $C(\overline{\Omega})$, meaning that one can find $\bar{u} \geq 0$ in $C(\overline{\Omega})$ and a sequence (t_k) tending to $+\infty$ in k , such that $u(t_k)$ converges to \bar{u} as k goes to $+\infty$, in $C(\overline{\Omega})$. Note that the limit cannot be identically 0, since otherwise $g(t)$ would go to $+\infty$, in contradiction with its convergence to l .

Our aim is to prove that $\bar{u} = 1$, which will mean that the whole trajectory converges to \bar{u} , hence the expected result.

Third step: identifying the limit.

Let us now consider the trajectory starting from the initial datum \bar{u} , namely $\{S_t \bar{u}\}_{t \geq 0}$, which we also denote by $\{\tilde{u}(t)\}_{t \geq 0}$. Because $\bar{u} \geq 0, \bar{u} \neq 0$, we again have $\tilde{u}(t, x) > 0$ for all $t > 0, x \in \overline{\Omega}$. Let us prove that V is constant along the trajectory $\{S_t \bar{u}\}_{t \geq 0}$ for $t > 0$.

For this, we write $V(\tilde{u}(t)) = V(S_t \bar{u}) = V(S_t \lim_{k \rightarrow +\infty} S_{t_k} u^0) = V(\lim_{k \rightarrow +\infty} S_{t+t_k} u^0)$. It is also easy to see that, for any u in $C(\overline{\Omega})$ which is furthermore positive on $\overline{\Omega}$, V (seen as acting on $C(\overline{\Omega})$) is continuous at u , and this implies $V(\tilde{u}(t)) = \lim_{k \rightarrow +\infty} V(S_{t+t_k} u^0) = l$. As claimed, the function $t \mapsto V(S_t \bar{u})$ is constant (equal to l) for $t > 0$.

Hence its derivative must be zero for $t > 0$: from the computations made in the first step, it must hold that both $\int_{\Omega} \frac{|\nabla(\tilde{u}(t))|^2}{\tilde{u}^2(t)}$ and $\int_{\Omega^2} K(x, y) (\tilde{u}(t, x) - 1)(\tilde{u}(t, y) - 1) \, dx \, dy$ vanish identically for $t > 0$. Let us now fix $t > 0$, and from the

first term, we know that $\tilde{u}(t)$ is a constant. From the second term and owing to $K > 0$, this constant must be equal to 1. By continuity of the trajectory, this also holds true at $t = 0$, i.e. $\tilde{u} = 1$, which ends the proof. \square

References

- [1] M. Alfaro, J. Coville, Rapid traveling waves in the nonlocal Fisher equation connect two unstable states, *Appl. Math. Lett.* 25 (12) (2012) 2095–2099.
- [2] H. Berestycki, G. Nadin, B. Perthame, L. Ryzhik, The non-local Fisher–KPP equation: travelling waves and steady states, *Nonlinearity* 22 (12) (2009) 2813.
- [3] J. Coville, Convergence to equilibrium for positive solutions of some mutation–selection model, Preprint arXiv:1308.6471, 2013.
- [4] B.S. Goh, Global stability in many-species systems, *Am. Nat.* 111 (1977) 135–143.
- [5] F. Hamel, L. Ryzhik, On the nonlocal Fisher–KPP equation: steady states, spreading speed and global bounds, *Nonlinearity* 27 (11) (2014) 2735.
- [6] P.-E. Jabin, G. Raoul, On selection dynamics for competitive interactions, *J. Math. Biol.* 63 (3) (2011) 493–517.
- [7] A.N. Kolmogorov, I.G. Petrovsky, N. Piskunov, Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Bull. Univ. État Moscou Sér. Int. A 1* (1937) 1–26.
- [8] G. Nadin, B. Perthame, M. Tang, Can a traveling wave connect two unstable states? The case of the nonlocal Fisher equation, *C. R. Acad. Sci. Paris, Ser. I* 349 (9–10) (2011) 553–557.
- [9] B. Perthame, *Parabolic Equations in Biology*, Springer, 2015.
- [10] C. Pouchol, J. Clairambault, A. Lorz, E. Trélat, Asymptotic analysis and optimal control of an integro-differential system modelling healthy and cancer cells exposed to chemotherapy, *J. Math. Pures Appl.* (9) (2018), <https://doi.org/10.1016/j.matpur.2017.10.007>.
- [11] C. Pouchol, E. Trélat, Global stability with selection in integro-differential Lotka–Volterra systems modelling trait-structured populations, Preprint arXiv:1702.06187, 2017.
- [12] M. Reed, B. Simon, *Methods of Modern Mathematical Physics, vol. II*, Academic Press, 1975.