

MÉMOIRES DE LA S. M. F.

PRZEMYSŁAW WOJTASZCZYK

Approximation properties and universal Banach spaces

Mémoires de la S. M. F., tome 31-32 (1972), p. 395-398

<http://www.numdam.org/item?id=MSMF_1972__31-32__395_0>

© Mémoires de la S. M. F., 1972, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

APPROXIMATION PROPERTIES AND UNIVERSAL BANACH SPACES

by

Przemysław WOJTASZCZYK

1. - In this talk we are dealing with two kinds of concepts.

The first one, approximation properties, has its origin in the concepts of Schauder basis [10] and metric approximation property of Grothendieck [2]. By "approximation property" we mean the answer to the question in what a way we can approximate the identity operator on a Banach space X by finite dimensional operators. The second one, universality, has its origin in the classical Banach-Mazur theorem on the universality of the space of continuous functions on the Cantor discontinuum [1]. The question is, to find for a given class of Banach spaces a space which contains (in a nice way) any element of the given class.

2. - Definitions.

DEFINITION 1. - A Banach space X has the bounded approximation property, shortly BAP (resp. unconditional bounded approximation property, shortly UBAP) iff there exists a sequence of finite dimensional operators $A_n : X \rightarrow X$, $n = 1, 2, \dots$, such that for each $x \in X$ $x = \sum_{n=1}^{\infty} A_n(x)$ (and the series is unconditionally convergent).

Those concepts are modifications of the metric approximation property of Grothendieck [2].

DEFINITION 2. - A Banach space X has a (unconditional) basis of finite dimensional subspaces iff there exists a sequence $(X_n)_{n=1}^{\infty}$ of finite dimensional subspaces of X such that for each $x \in X$ we have a unique decomposition $x = \sum_{n=1}^{\infty} x_n$ where $x_n \in X_n$ (and the series is unconditionally convergent). If moreover $\dim X_n = 1$ for $n = 1, 2, \dots$ the Banach space X has the Schauder basis (resp. unconditional Schauder basis).

It is easily seen that those concepts are in fact "approximation properties".

DEFINITION 3. - Let us have a class \mathcal{B} of Banach spaces. A space X is complementably universal for the class \mathcal{B} iff for each $Y \in \mathcal{B}$ there exists in X a complemented subspace Y_1 which is isomorphic to Y .

3. - About this concepts the following results are proved.

THEOREM 1. - [9] A Banach space X has UBAP iff X is isomorphic to a complemented subspace of a Banach space with the unconditional basis of finite dimensional subspaces. Moreover there exists one space U_{fd} with the unconditional basis of finite dimensional subspaces such that any X having UBAP is isomorphic to a complemented subspace of U_{fd} . The space U_{fd} is unique up to isomorphism among spaces with UBAP.

Remark 1. - Some constructions used in the proof of theorem 1 has applications in simultaneous extensions of continuous functions (cf. [9]).

THEOREM 2. - A Banach space X has BAP iff X is isomorphic to a complemented subspace of a Banach space with the Schauder basis. Moreover there exists one space B with the Schauder basis which is complementably universal for the class of all Banach spaces with BAP. The space B is unique up to isomorphism among spaces with BAP.

This theorem was proved independently in [4] and [8] using results of [9] and [7].

Remark 2. - In [3] among others is proved the following fact.

There exists a family of separable Banach spaces C_p , $1 \leq p \leq \infty$ such that for any Banach space X such that X^* has BAP and any p the space $X \oplus C_p$ has the Schauder basis.

Remark 3. - The space B was constructed by different ways in [7], [9] and [5]. The proof that spaces constructed in this papers are exactly the same follows from [8]. The construction of Kadec [5] has interesting applications to the theory of preduals of L_1 (cf. [9] and [12]).

Remark 4. - Many interesting results concerning various approximation properties are proved in [4].

4. - To finish this talk we are going to state some unsolved problems.

Problem 1. - Is any Banach space with UBAP isomorphic to a complemented subspace of a Banach space with an unconditional basis ?

Problem 2. - Find an example of a separable Banach space with UBAP not having an unconditional basis.

It is probably that such an example can be found among spaces constructed in [6].

Problem 3. - Prove that in a reflexive Banach space X with BAP there exists a sequence of finite dimensional projections P_n such that $P_n(X) \subset P_{n+1}(X)$ for $n = 1, 2, \dots$, and $P_n(x) \rightarrow x$ for any $x \in X$.

Problem 4. - Does any reflexive Banach space with a basis of finite dimensional subspaces have a Schauder basis?

The positive solution of problems 3 and 4 together with results of [2] and [4] would imply that a separable, reflexive space with approximation property of Grothendieck [2] has a Schauder basis.

Problem 5. - [7]. Does there exist a separable Banach space complementably universal for the class of separable Banach spaces?

The solution of this problem would have important consequences connected with "basis problem" and "approximation problem" (cf. [11] p. 386).

BIBLIOGRAPHIE

- [1] BANACH (S.) and MAZUR (S.). - Zur theorie der linearen Dimension, Studia Math. 4, 1933, p. 100-112.
- [2] GROTHENDIECK (A.). - Produits tensoriels topologiques et espaces nucléaires Mem. Amer. Math. Soc. n° 16, 1955.
- [3] JOHNSON (W.B.). - Factoring compact operators, Israël J. Math. 9, 1971, p. 337-45
- [4] JOHNSON (W.B.), ROSENTHAL (H.), ZIPPIN (M.). - On bases, finite dimensional decompositions and weaker structures in Banach spaces. Israel J. Math. 9, 1971, p. 488-506.
- [5] KADEC (M.I.). - On complementably universal spaces. Studia Math. 40, 1971, p. 85-89.
- [6] KWAPIEN (S.) and PELCZYNSKI (A.). - The main triangle projection in matrix spaces and its applications, Studia Math. 34, 1970, p. 43-68.
- [7] PELCZYNSKI (A.). - Universal bases, Studia Math. 32, 1969, p. 247-68.
- [8] PELCZYNSKI (A.). - Any separable Banach space with the bounded approximation property is a complemented subspace of a Banach space with a basis, Studia Math. 40, 1971, p. 239-43.
- [9] PELCZYNSKI (A.) and WOJTASZCZYK (P.). - Banach spaces with finite dimensional expansions of identity and universal bases of finite dimensional subspace Studia Math. 40, 1971, p. 91-108.

- [10] SCHAUDER (J.). - Zur theorie stetiger Abbildungen in funktionenräumen, Math. Zeitsch. 26, 1927, p. 47-65.
- [11] SINGER (I.). - Bases in Banach spaces I, Springer-Verlag Berlin-Heidelberg-New-York, 1970.
- [12] WOJTASZCZYK (P.). - Some remarks on the Gurarij space, (to appear in Studia Math. 41.)

Institute of Mathematics of the Polish
Academy of Sciences
Sniadeckich 8
VARSOVIE 1
(Pologne)
