

MÉMOIRES DE LA S. M. F.

RICARDO BAEZA

On the classification of quadratic forms over semi local rings

Mémoires de la S. M. F., tome 59 (1979), p. 7-10

<http://www.numdam.org/item?id=MSMF_1979__59__7_0>

© Mémoires de la S. M. F., 1979, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE CLASSIFICATION OF QUADRATIC FORMS OVER SEMI LOCAL RINGS

by

Ricardo BAEZA

1. Notations and definitions

Let A be a semi local ring. In this note we shall only consider non singular quadratic spaces over A . Let $W_q(A)$ be the Witt-group of quadratic spaces over A and $W(A)$ be the Witt-ring of bilinear spaces over A (see [1], [2] for the definitions). It is well-known, that $W_q(A)$ is a $W(A)$ -algebra. We shall use the notation $q \sim 0$ to express the fact that the class $[q]$ is 0 in $W_q(A)$. More generally, if q_1, q_2 are two quadratic spaces with $[q_1] = [q_2]$ in $W_q(A)$, we shall write $q_1 \sim q_2$. Let $\Delta(A)$ be the group of isomorphism classes of quadratic separable algebras over A , i.e. of algebras $A(p^{-1}(b)) = A \oplus Az$ with $z^2 = z + b$, $1 + 4b \in A^* =$ groups of unities of A . The product in $\Delta(A)$ is defined by

$$A(p^{-1}(a)) \circ A(p^{-1}(b)) = A(p^{-1}(a + b + 4ab))$$

For example if $z \in A^*$, then $\Delta(A) \cong A^*/A^{*2}$, and if $4 = 0$, then $\Delta(A) \cong A/p(A)$, where $p(A) = \{a^2 - a \mid a \in A\}$.

Let $Br(A)$ be the Brauer group of A . Then we have the following usual invariants for quadratic forms (see [1], [2])

$$\begin{aligned} d : W_q(A) &\rightarrow \mathbb{Z}/2\mathbb{Z} && (\text{dimension}) \\ a : W_q(A) &\rightarrow \Delta(A) && (\text{Arf-invariant}) \\ w : W_q(A) &\rightarrow Br(A) && (\text{Witt-invariant}) \end{aligned}$$

The Arf- and Witt-invariants of a quadratic space (E, q) are defined as follows : let $C(E)$ be the Clifford algebra of (E, q) and $D(E)$ be the centralizer of the sub-algebra $C(E)^+$ of elements of degree 0 in $C(E)$. It is easy to see that $D(E)$ is a quadratic separable algebra over A . Thus we define $a(q) = [D(E)] \in \Delta(A)$. a is a group homomorphism on the subgroup $W_q(A)_o$ of $W_q(A)$, which consist of the elements of even dimension, i.e. $W_q(A)_o = \text{Ker}(d)$. If $\dim E$ is even, then $C(E)$ is an Azumaya algebra over A , and we define in this case $w(q) = [C(E)] \in Br(A)$. If $\dim E$ is odd, then $C(E)^+$ is an Azumaya algebra over A , and we set $w(q) = [C(E)^+]$. For example let us consider the quadratic space $\langle d \rangle \otimes [1, b]$ with $d \in A^*$, $1 - 4b \in A^*$. Here $\langle d \rangle$ is the one dimensional bilinear space defined by d

and $[1, b]$ is the quadratic space $(Ae \oplus Af, q)$ with $q(e) = 1$, $q(f) = b$, $b_q(e, f) = 1$. Then we have

$$a(\langle d \rangle \otimes [1, b]) = [A(p^{-1}(-b))]$$

$$w(\langle d \rangle \otimes [1, b]) = [(-d, -b)],$$

where $(-d, -b]$ is the quaternion algebra $A \oplus Az \oplus Ae \oplus Aze$ with $z^2 = z-b$, $e^2 = -d$, $ze + ez = e$.

Let I_A be the maximal ideal of $W(A)$ of bilinear spaces of even dimension. If $2 \in A^*$ we may identify I_A with $W_q(A)_0$, but if $2 \notin A^*$ we have $W_q(A)_0 = W_q(A)$. Then it is easy to show that

$$I_A W_q(A)_0 = \text{Ker}(a|_{W_q(A)_0})$$

$$I_A^2 W_q(A)_0 \subseteq \text{Ker}(w|_{I_A W_q(A)_0})$$

A long standing question of Pfister is whether the equality $I_A^2 W_q(A)_0 = \text{Ker}(w|_{I_A W_q(A)_0})$ is true (if A is a field of characteristic 2 the answer is yes (see [5])). In the next section we shall prove a weak version of the equality above for semi local rings.

Now we introduce another type of invariants, namely the signatures of quadratic forms. A signature of the ring A is a ring homomorphism $\sigma : W(A) \rightarrow \mathbb{Z}$ (= ring of integers). Let $\text{Sig}(A)$ be the set of all signatures of A . The canonical homomorphism $\beta : W_q(A) \rightarrow W(A)$, which assigns to every quadratic form q its associated bilinear form b_q , induces a ring homomorphism $\bar{\sigma} = \sigma \circ \beta : W_q(A) \rightarrow \mathbb{Z}$ for every $\sigma \in \text{Sig}(A)$, such that $\text{Ker}(\bar{\sigma})$ is a $W(A)$ -submodule of $W_q(A)$. Let us denote the set of such ring homomorphisms $\bar{\sigma} : W_q(A) \rightarrow \mathbb{Z}$ by $\overline{\text{Sig}}(A)$. The correspondence $\sigma \mapsto \bar{\sigma} = \sigma \circ \beta$ defines a bijection $\text{Sig}(A) \rightarrow \overline{\text{Sig}}(A)$. Then it can be shown that

$$W(A)_t = \bigcap_{\sigma \in \text{Sig}(A)} \text{Ker}(\sigma)$$

$$W_q(A)_t = \bigcap_{\bar{\sigma} \in \overline{\text{Sig}}(A)} \text{Ker}(\bar{\sigma})$$

(see [2]). Now we define the total signature map

$$s : W_q(A) \rightarrow \prod_{\bar{\sigma} \in \overline{\text{Sig}}(A)} \mathbb{Z}_{\bar{\sigma}}$$

$$(\mathbb{Z}_{\bar{\sigma}} = \mathbb{Z}) \quad \text{by} \quad s(q) = (\bar{\sigma}(q)) \in \prod_{\bar{\sigma} \in \overline{\text{Sig}}(A)} \mathbb{Z}_{\bar{\sigma}}$$

2. A classification theorem for quadratic forms

Combining the maps of section 1 we can define

$$\begin{aligned} \Phi : W_q(A) &\rightarrow \mathbb{Z}/2\mathbb{Z} \times \Delta(A) \times \text{Br}(A) \times \prod_{\sigma \in \text{Sig}(A)} \frac{\mathbb{Z}}{\sigma} \\ \Phi &= (d, a, w, s) \end{aligned}$$

Then the following theorem is a generalization of some results of Elman and Lam (see [3]).

Theorem. Assume $|A/m| \geq 3$ for all maximal ideals of A . Then Φ is injective if and only if $I_A^2 W_q(A)_0$ is torsion free.

This result means, that if $I_A^2 W_q(A)_0$ is torsion free, quadratic forms over A are classified by its dimension, Arf-invariant, Witt-invariant and total signature. The main step in the proof of the theorem is the following.

Lemma. Assume $|A/m| \geq 3$ for all $m \in \max(A)$. Let $\Sigma(A)$ be the set of elements of A of the form $b = d + d^2 + \sum_1 c_i^2$ with $1 + 4b \in A^*$. Define $B = A(p^{-1}(b))$ for some $b \in \Sigma(A)$. Then if $I_A^2 W_q(A)_0$ is torsion free, it follows that $I_B^2 W_q(B)_0$ is torsion free, too.

Let us now use this lemma to prove the theorem.

Let q be an anisotropic quadratic form over A with $\Phi(q) = 0$. We want to show $q = 0$. Thus let us assume $q \neq 0$. Since $s(q) = 0$, it follows that $q \in W_q(A)_t$, and since $a(q) = 0$, we have $q \in (I_A W_q(A)_0)_t$. Using (7.13), V in [1] or (8.10), V in [2], we obtain

$$q \sim \bigoplus_{i=1}^r \langle a_i \rangle \otimes [1, -b_i]$$

with $b_i \in \Sigma(A)$. If $r \leq 2$, then comparing invariants on both sides, we conclude $q \sim 0$, which is a contradiction. Assume now $r > 2$. Taking $B = A(p^{-1}(b))$ with $b = b_1$, it follows that

$$q \otimes B \sim \bigoplus_{i=2}^r \langle a_i \rangle \otimes [1, -b_i]$$

On the other hand $I_B^2 W_q(B)_0$ is still torsion free (see the lemma) and $\Phi(q \otimes B) = 0$, thus we obtain $q \otimes B \sim 0$ by induction on r . Now q was assumed to be anisotropic, thus we get from this last relation

$$q \cong \varphi \otimes [1, -b]$$

for a suitable $\varphi = \langle c_1, \dots, c_r \rangle$, $c_i \in A^*$ (see (4.9), V in [1] or (4.10), V in [B]). Hence

$$q \sim \langle c_1 \rangle \otimes \langle 1, d_1 c_2 \rangle \otimes \langle 1, c_1 c_3 \rangle \otimes [1, -b] + \varphi_1 \otimes [1, -b]$$

with $\varphi_1 = \langle d_1, \dots, d_{r-2} \rangle$ for some $d_i \in A^*$. But

$\langle 1, c_1 c_2 \rangle \otimes \langle 1, c_1 c_3 \rangle \otimes [1, -b] \in (I_A^2 W_q(A)_0)_t = 0$, thus

$$q \sim \varphi_1 \otimes [1, -b].$$

Now we apply again induction to the right side and obtain $q \sim 0$. This is a contradiction. Hence $q = 0$, proving the theorem.

Corollary. If A is a semi local ring with $I_A^2 W_q(A)_0 = 0$, then

$$w : I_A W_q(A)_0 \rightarrow B_r(A)$$

is a monomorphism.

This follows from the fact, that $I_A^2 W_q(A)_0$ implies $\text{Sig}(A) = \emptyset$, and from the theorem above. This corollary was proved by Mandelberg in [4].

Remark. Let $u(A)$ be the u -invariant of A , i.e. the maximal dimension of anisotropic quadratic forms over A , which are torsion elements in $W_q(A)$. Then if $u(A) < 8$, it follows that $I_A^2 W_q(A)_0$ is torsion free. This fact can be seen as follows. Take $[q] \in (I_A^2 W_q(A)_0)_t$ and assume that q is anisotropic. Since $u(A) < 8$ implies $u(A) \leq 6$ (see Appendix B in [2]), we have $\dim q \leq 6$. But $[q] \in I_A^2 W_q(A)_0$ implies $\dim q \equiv 0(2)$, $a(q) = 1$, $w(q) = 1$, so that we can apply (4.13), V in [1] or (4.14), V in [2], to conclude that $q \sim 0$.

REFERENCES.

- [1] R. BAEZA, Quadratische Formen über semi lokalen Ringer. Habilitations schrift, Saarbrücken, 1975.
- [2] R. BAEZA, Quadratic forms over semi local rings. Lecture Notes in Math. n° 655.
- [3] R. ELMAN, T.Y. LAM, Classification theorems for quadratic forms over fields. Com. Math. Helv. 49, 373-381 (1974).
- [4] K.I. MANDELBERG, On the classification of quadratic forms over semi local rings. J. of algebra, 33, 463-471, (1975).
- [5] C-H, SAH, Symmetric bilinear forms and quadratic forms. J. of algebra, 20, 144-160 (1972).

Ricardo BAEZA

Mathematisches Institut
der Universität des Saarlandes
D - 66 - SAARBRÜCKEN