

# MÉMOIRES DE LA S. M. F.

W. DALE BROWNAWELL

## **On the orders of zero of certain functions**

*Mémoires de la S. M. F. 2<sup>e</sup> série*, tome 2 (1980), p. 5-20

[<http://www.numdam.org/item?id=MSMF\\_1980\\_2\\_2\\_5\\_0>](http://www.numdam.org/item?id=MSMF_1980_2_2_5_0)

© Mémoires de la S. M. F., 1980, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

ON THE ORDERS OF ZERO OF CERTAIN FUNCTIONS

by

W. Dale BROWNAWELL♦

I. INTRODUCTION

This is a report on joint work with D. W. Masser into the possible order of vanishing of a certain class of analytic functions. A complete exposition of our most general result and its relation to the previous work of Ju. Nesterenko [4] is given in [1]. That paper was written expressly for the use of fellow practitioners of transcendence theory.

For this conference it seemed appropriate to present a variant of the proof of the main theorem of [1], this time assuming familiarity with commutative algebra from the outset. The major change is the use here of the Hilbert characteristic function  $H_a(\alpha, t)$  for inhomogeneous ideals  $\alpha$ . In this way we avoid the technicalities involved in keeping track of the order of vanishing while homogenizing and dehomogenizing ideals. (These technicalities seem indispensable however for handling denominators in some of Masser's most recent work). Since we have not found a reference for the properties of  $H_a(\alpha, t)$  in the literature (see [2, p. 157] however), we discuss them in a short appendix following the body of the proof.

We are concerned here with solutions of a fixed system of differential equations

---

♦ Research supported in part by the National Science Foundation.

$$(1) \quad f'_i = F_i(f_1, \dots, f_n) \quad (1 \leq i \leq n)$$

where  $F_1, \dots, F_n$  are non-zero polynomials of total degree at most  $d$ . For a given set of initial values  $\theta = (\theta_1, \dots, \theta_n)$ , we denote by  $\bar{f}(z; \theta)$  the corresponding solution of (1) analytic at the origin, i.e. the coordinates  $f_1(z; \theta), \dots, f_n(z; \theta)$  satisfy (1) and  $\bar{f}(0, \theta) = \theta$ . In this paper we will deal with  $M+1$  fixed such initial conditions given by  $\theta_0, \theta_1, \dots, \theta_M$ .

Let  $\alpha$  be an ideal of  $R = \mathbb{C}[X_1, \dots, X_n]$ . For  $0 \leq m \leq M$  we define

$$\text{ord}_m \alpha = \min_{\substack{P \in \alpha \\ P \neq 0}} \text{ord } P(\bar{f}(z; \theta_m)),$$

where  $\text{ord}$  on the right hand side denotes the order of zero at the origin with the usual proviso that  $\text{ord } 0 = \infty$ . Say  $\alpha$  is generated by  $P_1, \dots, P_s$ . Then clearly

$$\text{ord}_m \alpha = \min_i \text{ord } P_i(\bar{f}(z; \theta_m)).$$

If  $\alpha$  has rank  $r$  (we avoid the term "height" because of other associations in transcendence theory), we can suppose that the indices are chosen in such a way that  $P_1, \dots, P_r$  have total degrees at most  $D_1 \geq \dots \geq D_r$ , respectively, whereas  $P_{r+1}, \dots, P_s$  have total degrees at most  $D_r$ . Let  $T = D_1 \dots D_r$ , and recall that  $d \leq \max_i \deg F_i$ . Then we can state our main result.

Theorem : Assume that  $\text{ord}_m \alpha$  is finite for  $0 \leq m \leq M$ . Then if  $r < n$  and  $d > 1$ , we have

$$\sum_{m=0}^M \text{ord}_m \alpha \leq (dT)^{2^{n-r}} + M(dT)^{2^{n-r-1}},$$

while if  $r < n$  and  $d = 1$ , we have

$$\sum_{m=0}^M \text{ord}_m \alpha \leq (n-r+1)D_1^{n-r}T + (n-r)MTD_1^{n-r-1}.$$

Finally if  $r = n$ , then

$$\sum_{m=0}^M \text{ord}_m \alpha \leq T.$$

## II. UNMIXING

Because we can estimate the degree of an ideal of the principal class (rank = minimal number of generators) through Proposition 4A, we would be very happy if  $r = s$ . Since that is not always the case, we show that for our purposes, it is always possible to replace  $\alpha$  by an ideal  $\alpha_r$  of rank  $r$  and generated by  $r$  polynomials of degrees at most  $D_1, \dots, D_r$ , respectively, such that

$$\text{ord}_m \alpha_r = \text{ord}_m \alpha \quad (0 \leq m \leq M).$$

For that the following general remark is useful.

Lemma 1 : Let  $P_1, \dots, P_s$  be polynomials and let  $\mathcal{I}_1, \dots, \mathcal{I}_k$  be ideals such that for each  $1 \leq \kappa \leq k$ , not all of  $P_1, \dots, P_s$  lie in  $\mathcal{I}_\kappa$ . Then for some integer,  $1 \leq \lambda \leq ks$ ,

$$P_1 + \lambda P_2 + \dots + \lambda^{s-1} P_s \notin \bigcup_{\kappa=1}^k \mathcal{I}_\kappa.$$

Proof : If each of the  $Q_\lambda = P_1 + \lambda P_2 + \dots + \lambda^{s-1} P_s$  ( $1 \leq \lambda \leq ks$ ) lies in some  $\mathcal{I}$ , then by the Box Principle, at least  $s$  of the  $Q_\lambda$  lie in the same  $\mathcal{I}$ . Inverting the corresponding Vandermonde determinant shows then that  $P_1, \dots, P_s$  all lie in that same  $\mathcal{I}$  as well. This contradiction establishes the lemma.

For a vector  $\theta = (\theta_1, \dots, \theta_n)$  of  $\mathbb{C}^n$  we denote by  $\mathcal{M}(\theta)$  the corresponding maximal ideal  $(x_1 - \theta_1, \dots, x_n - \theta_n)$ . For brevity we write  $\mathcal{M}_0, \dots, \mathcal{M}_M$  instead of  $\mathcal{M}(\theta_0), \dots, \mathcal{M}(\theta_M)$ , respectively. For  $0 \leq m \leq M$  we write  $\alpha^{(m)}$  for the contracted extension

$$\alpha^{(m)} = \alpha_m \cap R,$$

where  $\alpha_m$  denotes the extension of  $\alpha$  to the localization of  $R = \mathbb{C}[x_1, \dots, x_n]$  at  $\mathcal{M}_m$ . Further we write  $\alpha^*$  for the contracted extension with respect to the multiplicative set  $S = R \setminus \bigcup_{m=0}^M \mathcal{M}_m$ , i.e.  $\alpha^* = (\alpha \otimes_{R_S} R_S) \cap R$ . We see that  $\alpha^*$  is also obtained by deleting from a primary decomposition of  $\alpha$  components not lying in any  $\mathcal{M}_m$  ( $0 \leq m \leq M$ ), just as  $\alpha^{(m)}$  is obtained on deleting components not in  $\mathcal{M}_m$ . Therefore

$$(2) \quad \alpha^* = \bigcap_{m=0}^M \alpha^{(m)}.$$

Since every element of  $\alpha^{(m)}$  is the quotient of an element of  $\alpha$  by an element outside  $\mathcal{H}_m$ , we see that  $\text{ord}_m \alpha^{(m)} = \text{ord}_m \alpha$ . Therefore from  $\alpha \subseteq \alpha^* \subseteq \alpha^{(m)}$  we deduce that

$$(3) \quad \text{ord}_m \alpha = \text{ord}_m \alpha^* = \text{ord}_m \alpha^{(m)} \quad (0 < m < M)$$

Proposition 2 : If in addition to the hypotheses of the Main Theorem we have

$$0 < \text{ord}_m \alpha \quad (0 < m < M),$$

then there are polynomials  $Q_1, \dots, Q_r$  in  $\alpha$  with  $\deg Q_1 < D_1, \dots, \deg Q_r < D_r$  such that the ideal  $\alpha_r = (Q_1, \dots, Q_r)$  satisfies

$$\text{rank } \alpha_r = r,$$

$$\text{ord}_m \alpha_r = \text{ord}_m \alpha \quad (0 < m < M),$$

and

$$\deg \alpha_r^* < D_1 \dots D_r = T.$$

Proof : Since  $\text{ord}_m \alpha$  is assumed to be finite for each  $m$ , the polynomials  $P$  such that  $\text{ord}_m P > \text{ord}_m \alpha$  form an ideal which we denote  $\mathcal{I}_m$ . By Lemma 1 we can select  $\lambda \in \mathbb{Z}$  such that

$$Q_1 = P_1 + \lambda P_2 + \dots + \lambda^{s-1} P_s \notin \bigcup_{m=0}^M \mathcal{I}_m.$$

Set  $\alpha_1 = (Q_1)$ . Then  $\text{rank } \alpha_1 = 1$  and  $\text{ord}_m \alpha_1 = \text{ord}_m \alpha$ ,  $0 < m < M$ . Now  $\alpha_1^*$  is principal with generator obtained from  $Q_1$  by deleting all factors not in any  $\mathcal{H}_m$ ,  $0 < m < M$ . Since  $\text{ord}_m \alpha_1^* = \text{ord}_m \alpha_1 > 0$ ,  $Q_1$  is not constant and  $\text{rank } \alpha_1^* = 1$ . We deduce from Proposition 4A of the appendix that

$$\deg \alpha_1^* < D_1,$$

which is what was claimed for  $r = 1$ .

# ORDERS OF ZERO OF FUNCTIONS

For  $r > 1$  we choose polynomials  $Q_2, \dots, Q_r$  inductively of the form

$$(4) \quad Q_i = P_i + \lambda_1 P_{i+1} + \dots + \lambda_1^{s-i} P_s$$

such that the ideals  $\alpha_i = (Q_1, \dots, Q_i)$  satisfy

$$\text{rank } \alpha_i = i, \deg \alpha_i^* < D_1 \dots D_i.$$

Moreover the form of selection given in (4) guarantees that for  $1 < i < r$

$$\text{ord}_m \alpha_i = \text{ord}_m \alpha \quad (0 < m < M)$$

The selection has already been made for  $i = 1$ , and now we show how to find  $Q_{i+1}$  with the desired properties once  $\alpha_i$  has been obtained ( $i < r$ ).

Fix a prime component  $\mathcal{P}$  of  $\alpha_i$  of rank  $i$  (necessarily so by the Cohen-Macauley Theorem [5-II, p.310]). If  $P_{i+1}, \dots, P_s$  all lay in  $\mathcal{P}$ , Then we see inductively from (4) that so do  $P_i, P_{i-1}, \dots, P_1$ . Thus  $\alpha$  would be contained in  $\mathcal{P}$  and  $\text{rank } \alpha < i$ , contrary to our assumption that  $\text{rank } \alpha = r$ . Thus at least one of  $P_{i+1}, \dots, P_s$  does not lie in  $\mathcal{P}$ . So by Lemma 1, there is a  $Q_{i+1}$  of the form (4) not in any prime component of  $\alpha_i$ , i.e.  $\alpha_i : Q_{i+1} = \alpha_i$ . Therefore by Proposition 4A of the appendix, for  $\alpha_{i+1} = (\alpha_i, Q_{i+1})$  we have  $\text{rank } \alpha_{i+1} = i+1$  and  $\deg \alpha_{i+1} < (\deg \alpha_i) D_{i+1}$ , which establishes Proposition 2.

## III. THE CASE $n = r$ .

To deal with this case we require a fundamental result concerning exponents of ideals.

Lemma 3 : If  $\mathcal{Q}$  is a primary ideal of length  $l$  and exponent  $e$ , then  $e < l$ .

Proof : Say that  $\mathcal{Q}$  is  $\mathcal{P}$ -primary. Then  $\mathcal{P}^e \subseteq \mathcal{Q}$ , and  $e$  is the least positive power of  $\mathcal{P}$  lying in  $\mathcal{Q}$ . If  $e = 1$ , then  $\mathcal{P} = \mathcal{Q}$  and there is nothing to show. If  $e > 2$ , the ideals  $\mathcal{Q}_i = \mathcal{Q} : \mathcal{P}^i$  ( $0 < i < e$ ) are  $\mathcal{P}$ -primary [5-I, p.154]. Since  $\mathcal{P}^{e-1} \mathcal{P} = \mathcal{P}^e \subseteq \mathcal{Q}$ ,  $\mathcal{P} \subseteq \mathcal{Q}_{e-1}$ , and so  $\mathcal{P} = \mathcal{Q}_{e-1}$ . Thus we obtain  $e$  primary ideals

$$\mathcal{Q} = \mathcal{Q}_0 \subseteq \mathcal{Q}_1 \subseteq \dots \subseteq \mathcal{Q}_{e-1} = \mathcal{P}$$

If we can show that these inclusions are strict, then the lemma will follow. But  $p\mathfrak{q}_{k+1} \subseteq \mathfrak{q}_k$  ( $0 \leq k < e-1$ ). So if some  $\mathfrak{q}_k = \mathfrak{q}_{k+1}$ , then

$p\mathfrak{q}_{k+2} \subseteq \mathfrak{q}_{k+1} = \mathfrak{q}_k$ . Therefore  $p^{k+1}\mathfrak{q}_{k+2} \subseteq p^k\mathfrak{q}_k \subseteq \mathfrak{q}$  and  $\mathfrak{q}_{k+2} \subseteq \mathfrak{q}_{k+1}$ . Consequently  $\mathfrak{q}_k = \mathfrak{q}_{k+1} = \mathfrak{q}_{k+2}$ . Repeating the argument shows that

$$\mathfrak{q}_k = \mathfrak{q}_{k+1} = \dots = \mathfrak{q}_{e-1}.$$

Thus  $p = \mathfrak{q} : p^k$  and  $p^{k+1} \subseteq \mathfrak{q}$ . By the definition of  $e$ ,  $e \leq k+1$ . But  $k+1 \leq e-1$  by assumption. This contradiction shows that the inclusions are strict and establishes the lemma.

Proposition 4 : Let  $\mathfrak{q}$  be primary of rank  $n$  such that  $\text{ord}_m \mathfrak{q}$  is positive but finite for some  $0 < m \leq M$ . Then  $\mathcal{M}_m$  is the associated prime ideal of  $\mathfrak{q}$  and

$$\text{ord}_m \mathfrak{q} < \deg \mathfrak{q}.$$

Proof : Since  $\text{ord}_m \mathfrak{q} > 0$ , we have  $\mathfrak{q} \subseteq \mathcal{M}_m$ . Because  $\text{rank } \mathfrak{q} = n$ ,  $\mathcal{M}_m$  is the associated prime. Let  $e$  be the exponent of  $\mathfrak{q}$ . Then

$$\mathcal{M}_m^e \subseteq \mathfrak{q} \subseteq \mathcal{M}_m.$$

Since for any ideals  $k, \mathcal{L}$

$$\text{ord}_m (k \cap \mathcal{L}) \leq \text{ord}_m (k\mathcal{L}) = \text{ord}_m k + \text{ord}_m \mathcal{L},$$

we see that

$$\text{ord}_m \mathfrak{q} \leq e \text{ord}_m \mathcal{M}_m.$$

From Proposition 2A of the appendix and Lemma 4 we see that

$$e < \text{length } \mathfrak{q} < \deg \mathfrak{q}.$$

If  $\mathcal{O}_m = (\theta_1, \dots, \theta_n)$ , then  $\text{ord}_m \mathcal{M}_m = \min_i \text{ord}_m (f_i(z; \mathcal{O}_m) - \theta_i)$ , and so if  $\text{ord}_m \mathcal{M}_m > 1$ , then each  $f_i'(z; \mathcal{O}_m)$  vanishes at the origin. Now differentiation of (1) leads to the relations

$$f_i''(z; \theta_m) = \sum_{j=1}^n \frac{\partial^2 f_i}{\partial x_j^2} \bigg|_{\tilde{f}(z; \theta_m)} f_j'(z; \theta_m) \quad (1 \leq i \leq n).$$

Thus

$$\text{ord}_{z=0} f_i''(z; \theta_m) > \min_j \text{ord}_{z=0} f_j'(z; \theta_m) > 0 \quad (1 \leq i \leq n).$$

This implies that  $\text{ord}_{z=0} f_i'(z; \theta_m) = \infty$ ,  $1 \leq i \leq n$ . Then  $\text{ord}_m \mathcal{M}_m = \infty$ , a contradiction which completes the proof of the proposition.

Proof of the case  $n = r$ .

We may clearly assume that  $\text{ord}_m \alpha > 0$  ( $0 \leq m \leq M$ ) simply by renumbering and taking  $M$  smaller if necessary. For if the analogous bound holds on the sum of the remaining  $\text{ord}_m \alpha$ , then the desired bound will hold on the full sum. We apply Proposition 2 to  $\alpha$  to obtain  $\alpha_n^*$ . If  $\alpha_n^* = \mathcal{G}_0 \cap \dots \cap \mathcal{G}_N$  is the primary decomposition of  $\alpha_n^*$ , then by the first half of Proposition 4,  $N = M$ , and we may take  $\mathcal{G}_m$  to be  $\mathcal{G}_m$ -primary ( $0 \leq m \leq M$ ). By the second half of Propositions 3A and 4A of the appendix and by Proposition 4,

$$\begin{aligned} T > \deg \alpha_n^* &= \sum_{m=0}^M \deg \mathcal{G}_m \\ &> \sum_{m=0}^M \text{ord}_m \mathcal{G}_m = \sum_{m=0}^M \text{ord}_m \alpha, \end{aligned}$$

since  $\text{ord}_m \alpha = \text{ord}_m \alpha_n = \text{ord}_m \alpha_n^{(m)} = \text{ord}_m \mathcal{G}_m$  ( $0 \leq m \leq M$ ). This proves the assertion of the theorem for  $r = n$ .

#### IV. INCREASING THE LOCAL RANK

In this section we develop a procedure which allows us to cope with ideals of rank less than  $n$ . We inductively produce polynomials  $Q_{r+1}, \dots, Q_n$  of predictably bounded degrees  $D_{r+1}, \dots, D_n$ , respectively, through differentiation of the generators of  $\alpha_r$  such that for  $r < i \leq n$  the ideals

$$(7) \quad \mathcal{L}_i = (\alpha_r, Q_{r+1}, \dots, Q_i)$$

satisfy

$$(8) \quad \text{rank } \mathcal{G}_i^{(m)} = i \quad (0 < m < M),$$

$$(9) \quad \deg \mathcal{G}_i^* < D_1 \dots D_i$$

and

$$(10) \quad \text{ord}_m \mathcal{A}_{i-r-1} < \text{ord}_m \mathcal{G}_i \neq \infty \quad (0 < m < M)$$

where

$$W_k = \begin{cases} \sum_{j=0}^k d^{2^{j-1}} T^{2^j}, & \text{if } d > 1 \\ T \sum_{j=0}^k D_1^j, & \text{if } d = 1. \end{cases}$$

(Recall that  $d$  is our upper bound on the degree of the polynomials  $F_1, \dots, F_n$  in (1) and  $D_1$  bounds the degree of all the generators of  $\mathcal{A}$ ). As we shall see,  $W_{i-r-1}$  is a bound on the number of derivatives we may have to take to obtain  $Q_{r+1}, \dots, Q_i$ . Because of its importance for the Main Theorem, we state the result for  $i = n$  as a proposition. Set

$$D_i = \begin{cases} (dT)^{2^{i-r-1}}, & \text{if } d > 1 \\ D_1, & \text{if } d = 1 \end{cases}$$

Proposition 5 : Given  $\mathcal{A}_r$  of Proposition 2 with

$$W_{n-r-1} < \text{ord}_m \mathcal{A}_r \neq \infty \quad (0 < m < M),$$

there are then polynomials  $Q_{r+1}, \dots, Q_n$  with  $\deg Q_i < D_i$  ( $r < i < n$ ) such that the following holds for  $\mathcal{G}_n = (\mathcal{A}_r, Q_{r+1}, \dots, Q_n)$  :

$$\text{rank } \mathcal{G}_n^{(m)} = n, \text{ord}_m \mathcal{A}_r - W_{n-r-1} < \text{ord}_m \mathcal{G}_n \quad (0 < m < M)$$

and

$$\deg \mathcal{G}_n^* < D_1 \dots D_n.$$

Proof : As mentioned above, the construction of the  $\mathcal{L}_{r+1}, \dots, \mathcal{L}_n$  with the properties (8), (9), (10) is inductive. If we set  $W_{-1} = 0$ , we may consider  $\mathcal{L}_r = \mathcal{Q}_r$  to start it off. For the inductive step assume that (8), (9), (10) hold for some  $\mathcal{L}_i$  of the form (7),  $r \leq i < n$ . Consider the derivation

$$\Delta = \sum_{j=1}^n F_j \frac{\partial}{\partial x_j}$$

defined on  $R = \mathbb{C}[x_1, \dots, x_n]$ . Clearly if  $P \in R$  has total degree  $G$ , then  $\Delta P$  has total degree at most  $G + d - 1$ . Also if  $\bar{f} = (f_1, \dots, f_n)$  is an analytic solution of (1) then for any polynomial  $P \in R$ ,

$$\Delta P(\bar{f}) = \frac{d}{dz} P(\bar{f}).$$

So if  $\text{ord}_m P > 1$ , then  $\text{ord}_m \Delta P = \text{ord}_m P - 1$  ( $C \leq m \leq M$ ). Since the ideals  $\mathcal{L}_i^{(m)}$  have rank  $i$  and  $\mathcal{L}_i$  is generated by  $i$  elements,  $\mathcal{L}_i^{(m)}$  is unmixed by the Cohen-Macaulay Theorem [5-II, p.310]. Therefore in particular all primary components of  $\mathcal{L}_i^*$  have rank  $i$ . For the next few paragraphs we consider one fixed such component  $\mathcal{Q}$  and its associated prime  $\mathcal{P}$ .

From Lemma 3, Propositions 2A and 3A of the appendix, and our induction assumption, we deduce that

$$(11) \quad e < \deg \alpha_i^* < D_1 \dots D_i$$

for the exponent  $e$  of  $\mathcal{Q}$ . We claim that

$$(12) \quad \Delta^e \mathcal{L}_i^* \not\subseteq \mathcal{P}.$$

Since  $\mathcal{L}_i^* \subseteq \mathcal{P} \subseteq \mathcal{M}_m$  for some  $0 < m \leq M$ ,  $\text{ord}_m \mathcal{P}$  is positive but bounded by  $\text{ord}_m \mathcal{L}_i^*$ . Choose a polynomial  $P \in \mathcal{P}$  with  $\text{ord}_m P = \text{ord}_m \mathcal{P}$ . Since

$$\text{ord}_m \Delta P = \text{ord}_m P - 1 < \text{ord}_m \mathcal{P},$$

$\Delta P$  does not lie in  $\mathcal{P}$ . Let  $Q$  be a polynomial lying in every primary component  $\mathcal{Q}' \neq \mathcal{Q}$  of  $\mathcal{L}_i^*$  and not lying in  $\mathcal{P}$  — for example the product of elements from each  $\mathcal{Q}' \setminus \mathcal{P}$ . From the definition of exponent we know that  $P^e$  lies in  $\mathcal{Q}$ . So  $P^e Q$  lies in  $\mathcal{L}_i^*$ . Moreover since  $P$  lies in  $\mathcal{P}$ ,

$$\Delta^e (P^e Q) \equiv e! (\Delta P)^e Q \pmod{\mathcal{P}}.$$

Since neither  $Q$  nor  $\Delta P$  is in  $\mathcal{P}$ , this establishes (12), a crucial step in our proof.

Now let  $t$  be the least positive integer such that  $\Delta^t \mathcal{L}_1^* \notin \mathcal{P}$ . From (11) and (12) we see that

$$t \leq e \leq D_1 \dots D_i.$$

We claim that

$$(13) \quad \Delta^t \mathcal{L}_i \notin \mathcal{P}.$$

For there is a  $B$  in  $\mathcal{L}_i^*$  such that  $\Delta^t B$  is not in  $\mathcal{P}$ . From the definition of  $\mathcal{L}_i^*$ , there is an  $A$  outside  $\bigcup_{m=0}^M \mathcal{M}_m$  such that  $C = AB$  lies in  $\mathcal{L}_i$ . From the minimality of  $t$  we see that each  $\Delta^\tau B$  lies in  $\mathcal{P}$  for  $0 \leq \tau < t$  and thus that

$$\Delta^t C \equiv A \Delta^t B \pmod{\mathcal{P}}.$$

Since  $A$  is not in any  $\mathcal{M}_m$ , it is not in  $\mathcal{P}$  either, for, as we noted above,  $\mathcal{P} \subseteq \mathcal{M}_m$  for some  $0 \leq m \leq M$ . We deduce that  $\Delta^t C$  is not in  $\mathcal{P}$ , which verifies (13).

Since for  $0 \leq \tau < t$

$$\Delta^\tau \mathcal{L}_i \subseteq \Delta^\tau \mathcal{L}_i^* \subseteq \mathcal{P},$$

the integer  $t$  is also minimal with respect to the property (13). Write

$$C = A_1 Q_1 + \dots + A_i Q_i$$

with  $A_1, \dots, A_i$  in  $R$ . Then by the minimality of  $t$ ,

$$\Delta^t C = A_1 \Delta^t Q_1 + \dots + A_i \Delta^t Q_i \pmod{\mathcal{P}}.$$

Hence  $\Delta^t Q_j$  is not in  $\mathcal{P}$  for some  $1 \leq j \leq i$ . By (11), since  $t \leq e$ ,  $\Delta^t Q_j$  has total degree at most

$$\max(D_1, D_i) + (d-1)D_1 \dots D_i \leq D_{i+1}.$$

Moreover

$$\begin{aligned}
 \text{ord}_m \Delta^t Q_j &> \text{ord}_m \mathcal{B}_i - t \\
 &> \text{ord}_m \alpha_r - w_{i-r-1} - t \\
 &> \text{ord}_m \alpha_r - w_{i-r} .
 \end{aligned}$$

When we carry out this construction for each of the prime components  $\mathcal{P}_1, \dots, \mathcal{P}_k$  of each  $\mathcal{B}_i^{(m)}$  ( $0 \leq m \leq M$ ), we obtain polynomials  $L_1, \dots, L_k$  of degrees at most  $D_{i+1}$  such that

$$\text{ord}_m \alpha_r - w_{i-r} < \text{ord}_m L_j \neq \infty \quad (1 \leq j \leq k, 0 \leq m \leq M).$$

Through Lemma 1 we can choose integers  $\lambda_1, \dots, \lambda_k$  such that

$$Q_{i+1} = \lambda_1 L_1 + \dots + \lambda_k L_k$$

does not lie in any  $\mathcal{P}_1, \dots, \mathcal{P}_k$ , and moreover  $\deg Q_{i+1} \leq D_{i+1}$ ,

$$\text{ord}_m \alpha_r - w_{i-r} < \text{ord}_m Q_{i+1} \neq \infty \quad (0 \leq m \leq M).$$

Thus  $\mathcal{B}_{i+1} = (Q_1, \dots, Q_{i+1})$  satisfies (10) with  $i$  replaced by  $i+1$ . We must now check (8) and (9). The construction of  $Q_{i+1}$  guarantees that

$$(14) \quad \mathcal{B}_i^{(m)} : Q_{i+1} = \mathcal{B}_i^{(m)} \quad (0 \leq m \leq M).$$

So we conclude from (8) and Krull's Principal Ideal Theorem [5-I, p.238] that

$$\text{rank } \mathcal{B}_{i+1}^{(m)} = i+1,$$

which verifies (8) for  $i+1$ . By (14) we also have  $\mathcal{B}_i^* : Q_{i+1} = \mathcal{B}_i^*$ . So by Proposition 4A of the appendix we have

$$\deg \mathcal{B}_{i+1}^* < \deg (\mathcal{B}_i^*, Q_{i+1}) < (D_1 \dots D_i) D_{i+1},$$

which verifies (9) for  $i+1$ . Therefore Proposition 5 follows.

V. PROOF FOR  $r < n$ .

Without affecting the validity of the assertion, we may, as in the proof of the case  $n = r$ , reindex  $\theta_0, \dots, \theta_M$  and decrease  $M$  if necessary to insure that each

$$\text{ord}_m Q > \begin{cases} (dT)^{2^{n-r-1}} & \text{if } d > 1 \\ (n-r)D_1^{n-r-1}T & \text{if } d = 1, \end{cases} \quad (0 < m < M).$$

Consider the primary decomposition of the ideal  $\mathcal{B}_n^*$  produced by Propositions 3 and 5 :

$$\mathcal{B}_n^* = \mathcal{G}_0 \cap \dots \cap \mathcal{G}_M.$$

Since  $\text{rank } \mathcal{B}_n^* = n$  and  $\text{ord}_m \mathcal{B} > 0$  for each  $0 < m < M$ , it follows from Proposition 4 that  $N = M$  and, after renumbering if necessary, the associated primes can be taken to be precisely  $\mathcal{M}_0, \dots, \mathcal{M}_M$ . By (3),

$$\text{ord}_m \mathcal{B}_n = \text{ord}_m \mathcal{B}_n^* = \text{ord}_m \mathcal{G}_m \quad (0 < m < M).$$

By Proposition 4 we know that

$$\text{ord}_m \mathcal{G}_m < \deg \mathcal{G}_m.$$

From Proposition 3A of the appendix we see that

$$\sum_{m=0}^M \deg \mathcal{G}_m = \deg \mathcal{B}_n^*.$$

Finally Proposition 5 furnishes the inequalities

$$\deg \mathcal{B}_n^* < D_1 \dots D_n$$

and

$$\text{ord}_m Q_r < \text{ord}_m \mathcal{B}_n + w_{n-r-1}.$$

Putting this all together with Proposition 3 gives

# ORDERS OF ZERO OF FUNCTIONS

$$\begin{aligned}
 \sum_{m=0}^M \text{ord}_m \alpha &= \sum_{m=0}^M \text{ord}_m \alpha_r < \sum_{m=0}^M \text{ord}_m \mathcal{G}_n + (M+1)W_{n-r-1} \\
 &< \sum_{m=0}^M \text{ord}_m \mathcal{G}_m + (M+1)W_{n-r-1} \\
 &< \sum_{m=0}^M \deg \mathcal{G}_m + (M+1)W_{n-r-1} \\
 &< \deg \mathcal{G}_n^* + (M+1)W_{n-r-1} \\
 &< D_1 \dots D_n + (M+1)W_{n-r-1}.
 \end{aligned}$$

This establishes the theorem after some straightforward computations.

We remark that the theorem applies to formal power series solutions of (1) over any field of characteristic zero.

## APPENDIX.

### Inhomogeneous Hilbert Functions

Recall that for a homogeneous ideal  $\mathcal{h}$  in  $K[x_0, \dots, x_n]$ , where  $K$  is an infinite field, the associated volume function  $V(t, \mathcal{h})$  counts the number of  $K$ -linearly independent forms of degree  $t$  in  $\mathcal{h}$  [2, pp.154-162]. The Hilbert characteristic function of  $\mathcal{h}$  is given by

$$H(t, \mathcal{h}) = \binom{n+t}{n} - V(t, \mathcal{h}).$$

$H(t, \mathcal{h})$  counts the number of  $K$ -linearly independent forms of degree  $t$  modulo  $\mathcal{h}$ .

Similarly for an arbitrary ideal  $\mathcal{A}$  of  $K[x_1, \dots, x_n]$ , the associated affine volume function  $V_a(t, \mathcal{A})$  counts the number of  $K$ -linearly independent polynomials in  $\mathcal{A}$  of degree at most  $t$  [2, p.157]. The affine characteristic function of  $\mathcal{A}$ ,  $H_a(t, \mathcal{A})$ , counts the number of  $K$ -linearly independent polynomials modulo  $\mathcal{A}$  of degree at most  $t$  in  $K[x_1, \dots, x_n]$ , and so

$$H_a(t, \mathcal{A}) = \binom{n+t}{n} - V_a(t, \mathcal{A}).$$

The parallel with the usual characteristic function is obviously strong, but still one cannot carry over the standard proofs to the affine case in a straightforward way.

The following basic property of  $V(t, \mathfrak{f})$  comes from the fact that the dimensions of the sum and intersection of a pair of vector subspaces add up to the sum of the dimensions of the original subspaces : for homogeneous ideals  $\mathfrak{f}, \mathfrak{k}$

$$V(t, \mathfrak{f}) + V(t, \mathfrak{k}) = V(t, \mathfrak{f} + \mathfrak{k}) + V(t, \mathfrak{f} \cap \mathfrak{k}).$$

However for affine ideals  $\mathcal{A}, \mathcal{B}$  one has merely

$$V_a(t, \mathcal{A}) + V_a(t, \mathcal{B}) \leq V_a(t, \mathcal{A} + \mathcal{B}) + V_a(t, \mathcal{A} \cap \mathcal{B}).$$

For example, when we set  $\mathcal{A} = (x^2 + 1)$ ,  $\mathcal{B} = (x^2)$  in  $K[x]$ , then  $\mathcal{A} + \mathcal{B} = K[x]$  and  $\mathcal{A} \cap \mathcal{B} = (x^4 + x^2)$ . So for  $t = 2$ ,  $V_a(2, \mathcal{A}) = V_a(2, \mathcal{B}) = 1$ ,  $V_a(2, \mathcal{A} + \mathcal{B}) = 3$ , and  $V_a(2, \mathcal{A} \cap \mathcal{B}) = 0$ .

In spite of this divergence of behavior, one can still deduce from many of the properties of  $H(t, \mathfrak{f})$  the corresponding ones for  $H_a(t, \mathcal{A})$ . Now the map

$$f(x_1, \dots, x_n) \longmapsto x_0^t f(x_1/x_0, \dots, x_n/x_0)$$

is a  $K$ -vector space isomorphism between the subspace of polynomials of degree at most  $t$  lying in an ideal  $\mathcal{A}$  and the subspace of the corresponding graded ideal  $^{gr}\mathcal{A}$  consisting of forms of degree  $t$ . Since  $^{gr}\mathcal{A}$  is made up of all forms of the corresponding homogeneous ideal  $^h\mathcal{A}$ , we see that

$$(15) \quad V_a(t, \mathcal{A}) = V(t, ^h\mathcal{A}), \quad H_a(t, \mathcal{A}) = H(t, ^h\mathcal{A}).$$

In particular we have the following result :

**THEOREM 1A :** The characteristic function of an ideal  $\mathcal{A}$  in  $K[x_1, \dots, x_n]$  of rank  $r$  has the following representation for large enough values of  $t$  in terms of binomial coefficients :

$$H_a(t, \mathcal{A}) = h_0 \binom{t}{n-r} + h_1 \binom{t}{n-r-1} + \dots + h_{n-r}$$

with  $h_0, \dots, h_r$  in  $\mathbb{Z}$  and  $h_0 > 0$ .

We call  $h_0$  the degree of  $\mathcal{Q}$  and write also  $h_0(\mathcal{Q})$ . The theorem follows from (15) and the corresponding representation for  $H(t, {}^h\mathcal{Q})$  [2, p.161] .

Recall that if  $\mathcal{Q} = \mathcal{Q}_1 \cap \dots \cap \mathcal{Q}_N$  is an irredundant primary decomposition with associated prime ideals  $\mathcal{P}_1, \dots, \mathcal{P}_N$ , respectively, then  ${}^h\mathcal{Q} = {}^h\mathcal{Q}_1 \cap \dots \cap {}^h\mathcal{Q}_N$  is also an irredundant primary decomposition with associated primes  ${}^h\mathcal{Q}_1, \dots, {}^h\mathcal{Q}_N$  [5-II, p.181] . Note also that the homogeneous ideals which are equivalent to their dehomogenized ideals are those whose prime components do not contain  $x_0$  [5-11, p.184] .

Proposition 2A : The degree of a  $p$ -primary ideal  $\mathcal{Q}$  of length  $\ell$  satisfies

$$h_0(\mathcal{Q}) = h_0(\mathcal{P}).$$

Proof : Since the map  $\mathcal{Q} \rightarrow {}^h\mathcal{Q}$  is injective [5-II, p.182] , the homogenization of a maximal chain of  $p$ -primary ideals

$$\mathcal{Q} = \mathcal{Q}_1 \subsetneq \dots \subsetneq \mathcal{Q}_\ell = \mathcal{P}$$

is a maximal chain of  ${}^hp$ -primary ideals lying above  ${}^h\mathcal{Q}$ . Now the claim follows from the corresponding property for homogeneous ideals [2, p.166] .

The following result is obtained similarly :

Proposition 3A : Let  $\mathcal{Q} = \mathcal{Q}_1 \cap \dots \cap \mathcal{Q}_s \cap \mathcal{Q}_{s+1} \cap \dots \cap \mathcal{Q}_t$  be an irredundant primary decomposition where  $\text{rank } \mathcal{Q} = \text{rank } \mathcal{Q}_1 = \dots = \text{rank } \mathcal{Q}_s$ , but  $\text{rank } \mathcal{Q}_{s+1}, \dots, \text{rank } \mathcal{Q}_t$  are larger. Then

$$h_0(\mathcal{Q}) = h_0(\mathcal{Q}_1) + \dots + h_0(\mathcal{Q}_s)$$

Proposition 4A : Let  $\mathcal{Q}$  be unmixed,  $\text{rank } \mathcal{Q} \leq n-1$  and let  $f$  be a polynomial of degree  $D \geq 1$  such that  $\mathcal{Q} : (f) = \mathcal{Q}$  and  $\mathcal{Q} + (f) \neq K[x_1, \dots, x_n]$ . Then

$$\text{rank } \mathcal{Q} + (f) = 1 + \text{rank } \mathcal{Q},$$

$$h_0(f) = D,$$

and

$$h_0(\mathcal{Q} + (f)) \leq Dh_0(\mathcal{Q})$$

Proof : The first assertion follows from Krull's Principal Ideal Theorem [3,p.37], [5-I, p.238] . The second assertion holds for any non-zero polynomial and follows from the corresponding property for homogeneous polynomials [2,p.167] .

For the final assertion note that  $\mathcal{Q} : (f) = \mathcal{Q}$  is equivalent to saying that  $f$  does not lie in any prime ideals associated with  $\mathcal{Q}$  . This property remains true under homogenization. Since  ${}^h\mathcal{Q} + ({}^hf) \subseteq {}^h(\mathcal{Q} + (f))$ , we find

$$H_a(t, \mathcal{Q} + (f)) \leq H(t, {}^h\mathcal{Q} + ({}^hf)).$$

So when we apply the Bezout theorem for homogeneous ideals [3, p.64] or [2,p.167] , we obtain the desired inequality. Let us finally note that every time we use Proposition 4.A, the positivity of the orders of  $\mathcal{Q}$  and  $f$  enables us to verify the condition that  $\mathcal{Q} + (f) \neq \mathbb{C}[x_1, \dots, x_n]$ .

---

#### BIBLIOGRAPHY

- [1] W. D. Brownawell and D. W. Masser, Multiplicity estimates for analytic functions II.
- [2] W. Gröbner, Moderne Algebraische Geometrie, die idealtheoretischen Grundlagen, Springer Verlag, Wien und Innsbruck, 1949.
- [3] W. Krull, Idealtheorie, 2<sup>nd</sup> expanded printing, Springer Verlag, Berlin, 1968.
- [4] Ju. V. Nesterenko, Estimates for the orders of zero of some functions of a certain class and their application in the theory of transcendental numbers, Izv. Akad. Nauk SSR, Ser. Mat. 41 (1977), 253-284, Math. USSR Izv. 11 (1977), 239-279, (see also these Proceedings).
- [5] O. Zariski and P. Samuel, Commutative Algebra, Volumes I and II, Springer Verlag, New York, 1968.

The Pennsylvania State University  
 Department of Mathematics  
 215, Mc Allistair Building  
 University Park, Penn. 16802  
 U. S. A.