

ANNALES DE LA FACULTÉ DES SCIENCES DE TOULOUSE Mathématiques

D. LE PEUTREC

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Tome XIX, n° 3-4 (2010), p. 735-809.

http://afst.cedram.org/item?id=AFST_2010_6_19_3-4_735_0

© Université Paul Sabatier, Toulouse, 2010, tous droits réservés.

L'accès aux articles de la revue « Annales de la faculté des sciences de Toulouse Mathématiques » (<http://afst.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://afst.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

D. LE PEUTREC⁽¹⁾

ABSTRACT. — This article follows the previous works [HeKlNi, HeNi] by Helffer-Klein-Nier and Helffer-Nier about the metastability in reversible diffusion processes via a Witten complex approach. Again, exponentially small eigenvalues of some self-adjoint realization of $\Delta_{f,h}^{(0)} = -h^2\Delta + |\nabla f(x)|^2 - h\Delta f(x)$ are considered as the small parameter $h > 0$ tends to 0. The function f is assumed to be a Morse function on some bounded domain Ω with boundary $\partial\Omega$. Neumann type boundary conditions are considered. With these boundary conditions, some possible simplifications in the Dirichlet problem studied in [HeNi] are no more possible. A finer treatment of the three geometries involved in the boundary problem (boundary, metric, Morse function) is here carried out.

RÉSUMÉ. — Cet article est dans la continuation des travaux [HeKlNi, HeNi] de Helffer-Klein-Nier et Helffer-Nier sur l'étude de la métastabilité dans des processus de diffusions réversibles via une approche de Witten. Nous considérons encore ici les valeurs propres exponentiellement petites d'une réalisation auto-adjointe de $\Delta_{f,h}^{(0)} = -h^2\Delta + |\nabla f(x)|^2 - h\Delta f(x)$ lorsque le paramètre $h > 0$ tend vers 0. La fonction f est une fonction de Morse sur un domaine borné Ω de bord $\partial\Omega$. Des conditions au bord de type Neumann sont considérées ici. Avec ces conditions, certaines simplifications utilisées pour l'étude du problème de Dirichlet dans [HeNi] ne sont plus possibles. Un traitement plus fin des trois géométries intervenant dans le problème à bord (bord, métrique, fonction de Morse) est donc nécessaire.

(*) Reçu le ??/??/20??, accepté le 30/03/2010

(1) Laboratoire de Mathématiques, UMR-CNRS 8628, Université Paris-Sud 11, Bâtiment 425, 91405 Orsay, France.
dorian.lepeutrec@math.u-psud.fr

Contents

1	Introduction and result	735
2	Witten Laplacien with Neumann boundary condition	742
2.1	Introductions and notations	742
2.2	Stokes and Green formulas	744
2.3	Normal Neumann realization	746
3	First localization of the spectrum	751
3.1	Introduction and result	751
3.2	A few preliminary lemmas	752
	3.2.1 Variational results for the Witten Laplacian on \mathbb{R}^k	752
	3.2.2 The model half-space problem	754
	3.2.3 Small eigenvalues for the model half-space problem	757
	3.3 Proof of Theorem 3.1.5	765
4	Accurate WKB analysis near the boundary for $\Delta_{f,h}^{(1)}$	770
4.1	Introduction	770
4.2	Local WKB construction	771
4.3	Another local Neumann realization of $\Delta_{f,h}^{(1)}$	772
4.4	Exponential decay of eigenvectors of $\Delta_{f,h}^{N,D,(1)}$	775
4.5	Small eigenvalues are exponentially small	781
4.6	Accurate comparison with the WKB solution	781
5	Labelling of local minima and construction of the quasimodes	788
5.1	Preliminaries	788
5.2	Generalized critical points and local structure of the level sets of a Morse function	788
5.3	Labelling of local minima and first consequence	791
5.4	Construction of the quasimodes	794
5.5	Quasimodal estimates	797
6	Final proof	804
6.1	Main result	804
6.2	Finite dimensional reduction and final proof	805

1. Introduction and result

In this work, we focus on the exponentially small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian $\Delta_{f,h}^{(0)}$ (acting on 0-forms) on a connected compact Riemannian manifold with regular boundary.

Our purpose is to derive with the same accuracy as in [HeKlNi] and in [HeNi] asymptotic formulas for the smallest non zero eigenvalues of the Neumann realization of $\Delta_{f,h}^{(0)}$

A similar problem was considered by many authors via a probabilistic approach in [FrWe], [HoKuSt], [Mic], and [Kol]. More recently, in the case of \mathbb{R}^n , accurate asymptotic forms of the exponentially small eigenvalues were obtained in [BoEcGaKl] and [BoGaKl].

These results were improved and extended to the cases of boundaryless compact manifolds in [HeKlNi] and of compact manifolds with boundaries for the Dirichlet realization of the Witten Laplacian in [HeNi].

We want here to extend these last results to the case of compact manifolds with boundaries for the Neumann realization of the Witten Laplacian, that is with coherently deformed Neumann boundary conditions.

Let us also make mention of an other recent work, [KoPrSh], about semiclassical asymptotics for the eigenvalues of the Witten Laplacian on compact manifolds with boundary, and where some techniques developed in [HeNi] are used. In [KoPrSh], the purpose is nevertheless quite different and does not consist in giving full semiclassical asymptotic expansions like it is done in [HeNi] and in the present paper.

The function f is assumed to be a Morse function on $\bar{\Omega} = \Omega \cup \partial\Omega$ with no critical points at the boundary. Furthermore, its restriction to the boundary $f|_{\partial\Omega}$ is also assumed to be a Morse function.

From [ChLi], which completed results yet obtained in the boundaryless case (see [Sim][Wit][CyFrKiSi][Hen][HeSj4][Hel3]), the number m_p of eigenvalues of the Neumann realization of the Witten Laplacian $\Delta_{f,h}^{(p)}$ (acting on p -forms) in some interval $[0, Ch^{\frac{3}{2}}]$ (for $h > 0$ small enough) relies closely on the number of critical points of f with index p .

In the boundaryless case, these numbers are exactly the numbers of critical points of f with index p in Ω . Like in [HeNi], this definition has to be *generalized* in the case with boundary, taking into account the structure

of the function f at the boundary, $f|_{\partial\Omega}$. Note furthermore that m_0 is here the number of local minima of f in $\overline{\Omega}$.

Moreover, the first eigenvalue in our case is 0 and the other small eigenvalues are actually exponentially small as $h \rightarrow 0$, i.e. of order $e^{-\frac{C}{h}}$, where C is a positive number independent of the small parameter $h > 0$.

The point of view of [HeKlNi] and [HeNi] intensively uses, together with the techniques of [HeSj4], the two facts that the Witten Laplacian is associated with a cohomology complex and that the function $x \mapsto \exp -\frac{f(x)}{h}$ is a distributional solution in the kernel of the Witten Laplacian on 0-forms, consequently allowing to construct very efficiently quasimodes.

Recall that the Witten Laplacian is defined as

$$\Delta_{f,h} = d_{f,h}d_{f,h}^* + d_{f,h}^*d_{f,h} , \tag{1.0.1}$$

where $d_{f,h}$ is the distorted exterior differential

$$d_{f,h} := e^{-f(x)/h} (hd) e^{f(x)/h} = hd + df(x) \wedge , \tag{1.0.2}$$

and where $d_{f,h}^*$ is its adjoint for the L^2 -scalar product canonically associated with the Riemannian structure (see for example [GaHuLa][Gol][Sch]). The restriction of $d_{f,h}$ to p -forms is denoted by $d_{f,h}^{(p)}$. With these notations, the Witten Laplacian on functions is

$$\Delta_{f,h}^{(0)} = d_{f,h}^{(0)*}d_{f,h}^{(0)} . \tag{1.0.3}$$

In the Witten complex spirit and due to the relation

$$d_{f,h}^{(0)}\Delta_{f,h}^{(0)} = \Delta_{f,h}^{(1)}d_{f,h}^{(0)} , \tag{1.0.4}$$

it is more convenient to consider the singular values of the restricted differential $d_{f,h}^{(0)} : F^{(0)} \rightarrow F^{(1)}$. The space $F^{(\ell)}$ is the m_ℓ -dimensional spectral subspace of $\Delta_{f,h}^{(\ell)}$, $\ell \in \{0, 1\}$,

$$F^{(\ell)} = \text{Ran } 1_{I(h)}(\Delta_{f,h}^{(\ell)}) , \tag{1.0.5}$$

with $I(h) = [0, Ch^{\frac{3}{2}}]$ and the property¹

$$1_{I(h)}(\Delta_{f,h}^{(1)})d_{f,h}^{(0)} = d_{f,h}^{(0)}1_{I(h)}(\Delta_{f,h}^{(0)}) . \tag{1.0.6}$$

⁽¹⁾ The right end $a(h) = Ch^{\frac{3}{2}}$ of the interval $I(h) = [0, a(h)]$ is suitable for technical reasons. What is important is that $a(h) = o(h)$. The value of $C > 0$ does not play any role provided h is small enough.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

The restriction $d_{f,h}|_{F^{(\ell)}}$ will be more shortly denoted by $\beta_{f,h}^{(\ell)}$,

$$\beta_{f,h}^{(\ell)} := (d_{f,h}^{(\ell)})_{/F^{(\ell)}}. \quad (1.0.7)$$

We will mainly focus on the case $\ell = 0$.

It appears that working with singular values of $\beta_{f,h}^{(0)}$ is efficient in order to compute the eigenvalues of $\Delta_{f,h}^{(0)}$ (which are their squares). Those quantities agree better with the underlying Witten complex structure.

Note that in our case, 0 is the smallest eigenvalue of the (deformed) Neumann realization of the Witten Laplacian on 0-forms due to the fact that $x \mapsto \exp -\frac{f(x)}{h}$ belongs to the domain of this operator (see Proposition 2.7 for the exact definition).

Let us now state the main result. Let $\mathcal{U}^{(0)}$ and $\mathcal{U}^{(1)}$ denote respectively the set of local minima and the set of *generalized* critical points with index 1, or *generalized* saddle points, of the Morse function f on $\bar{\Omega}$ (see Definition 5.1 for the exact meaning of “generalized”). The analysis is easier under a generic assumption which ensures that the exponentially small eigenvalues are simple with different logarithmic equivalent as $h \rightarrow 0$. Although it is possible to consider more general cases like in [HeKINi] and in [HeNi], we will follow the point of view presented in [Nie] and work directly in a generic case which avoids some technical and unnecessary considerations.

ASSUMPTION 1.1. — *The critical values of f and $f|_{\partial\Omega}$ are all distinct and the quantities $f(U^{(1)}) - f(U^{(0)})$, with $U^{(1)} \in \mathcal{U}^{(1)}$ and $U^{(0)} \in \mathcal{U}^{(0)}$ are distinct.*

Following this assumption, a one to one mapping j can be defined from $\mathcal{U}^{(0)} \setminus \{U_1^{(0)}\}$ where $U_1^{(0)}$ is the global minimum, into the set $\mathcal{U}^{(1)}$. The local minima are denoted by $U_k^{(0)}$, $k \in \{1, \dots, m_0\}$, and the generalized saddle points by $U_j^{(1)}$, $j \in \{1, \dots, m_1\}$. The ordering of the local minima as well as the one to one mapping j will be specified in Subsection 5.3.

The following fundamental quantities will enter in the expression of the final result.

DEFINITION 1.2. — *For $k \in \{2, \dots, m_0\}$, we define:*

$$\gamma_k(h) = \begin{cases} \frac{|\det \text{Hess } f(U_k^{(0)})|^{\frac{1}{4}}}{(\pi h)^{\frac{n}{4}}} & \text{if } U_k^{(0)} \in \Omega \\ \left(\frac{-2\partial_n f(U_k^{(0)})}{h}\right)^{\frac{1}{2}} \frac{|\det \text{Hess } f|_{\partial\Omega}(U_k^{(0)})|^{\frac{1}{4}}}{(\pi h)^{\frac{n-1}{4}}} & \text{if } U_k^{(0)} \in \partial\Omega, \end{cases}$$

$$\delta_{j(k)}(h) = \begin{cases} \frac{|\det \text{Hess } f(U_{j(k)}^{(1)})|^{\frac{1}{4}}}{(\pi h)^{\frac{n}{4}}} & \text{if } U_{j(k)}^{(1)} \in \Omega \\ \left(\frac{-2\partial_n f(U_{j(k)}^{(1)})}{h}\right)^{\frac{1}{2}} \frac{|\det \text{Hess } f|_{\partial\Omega}(U_{j(k)}^{(1)})|^{\frac{1}{4}}}{(\pi h)^{\frac{n-1}{4}}} & \text{if } U_{j(k)}^{(1)} \in \partial\Omega, \end{cases}$$

and,

$$\theta_{j(k)}(h) = \begin{cases} \frac{h^{\frac{1}{2}}}{\pi^{\frac{1}{2}}} \frac{(\pi h)^{\frac{n}{2}} |\widehat{\lambda}_1^\Omega|^{\frac{1}{2}}}{|\det \text{Hess } f(U_{j(k)}^{(1)})|^{\frac{1}{2}}} & \text{if } U_{j(k)}^{(1)} \in \Omega \\ \frac{h^2}{-2\partial_n f(U_{j(k)}^{(1)})} \frac{(\pi h)^{\frac{n-2}{2}} |\widehat{\lambda}_1^{\partial\Omega}|^{\frac{1}{2}}}{|\det \text{Hess } f|_{\partial\Omega}(U_{j(k)}^{(1)})|^{\frac{1}{2}}} & \text{if } U_{j(k)}^{(1)} \in \partial\Omega, \end{cases}$$

where $\widehat{\lambda}_1^W$ is the negative eigenvalue of $\text{Hess } f|_W(U_{j(k)}^{(1)})$ for $W = \Omega$ or $W = \partial\Omega$.

THEOREM 1.3. — *Under Assumption 1.1 and after the ordering specified in Subsection 5.3, there exists h_0 such that, for $h \in (0, h_0]$, the spectrum in $[0, h^{\frac{3}{2}})$ of the Neumann realization of $\Delta_{f,h}^{(0)}$ in $\overline{\Omega}$ consists of m_0 eigenvalues $0 = \lambda_1(h) < \dots < \lambda_{m_0}(h)$ of multiplicity 1.*

Moreover, the above $m_0 - 1$ non zero eigenvalues are exponentially small and admit the following asymptotic expansions:

$$\lambda_k(h) = \gamma_k^2(h) \delta_{j(k)}^2(h) \theta_{j(k)}^2(h) e^{-2\frac{f(U_{j(k)}^{(1)}) - f(U_k^{(0)})}{h}} (1 + hc_k^1(h))$$

where $\gamma_k(h)$, $\delta_{j(k)}(h)$, and $\theta_{j(k)}(h)$ are defined in the above definition and $c_k^1(h)$ admits a complete expansion: $c_k^1(h) \sim \sum_{m=0}^{\infty} h^m c_{k,m}$.

This theorem is an extension to the case with Neumann boundary conditions of the previous results of [BoGaKl] and its improvements in [HeKlNi] and [HeNi] (see also non-rigorous formal computations of [KoMa], who look also at cases with symmetry and the books [FrWe] and [Kol] and references therein).

To prove this theorem, we will follow the same strategy as in [HeKlNi] and in [HeNi] and some intermediary results will be reused without demonstration (what will be indicated in the article). Moreover, some proofs will be

improved (see for example the final proof reduced now to a simple Gaussian elimination, explained in [Lep1]).

Finally, let us underline that the geometry of the Neumann case is different from the geometry of the above references. This leads to different results (compare Theorem 1.3 and the main theorem of [HeNi]) and some proofs have to be entirely reconsidered. In fact, the study of the Dirichlet realization of the Witten Laplacian done in [HeNi] agreed better with the local geometry near the boundary, which led to simpler computations (see the local WKB construction in Section 4 for example).

The article is organized as follows.

In the second section, we analyze in detail the boundary complex adapted to our analysis in order to keep the commutation relation (1.0.4). A part of the answer already exists in the literature (see [Sch], [Duf], [DuSp], [Gue], and [ChLi]) in connection with the analysis of the relative or absolute cohomology as defined in [Gil].

The third section is devoted to the proof of rough estimates (to get a first localization of the spectrum of the Laplacian) replacing the harmonic oscillator approximation in the case without boundary.

These two sections bring no additional difficulties in comparison with what was done in [HeNi].

In the fourth section, we give the WKB construction for an eigenform of the Witten Laplacian on 1-forms localized near a critical point of the boundary, according to the analysis done in [Lep2]. In the Dirichlet case, it was possible to use only one single coordinate system for the WKB construction but in the present case different coordinate systems arise naturally. Lemma 3.18 will play a crucial role to juggle with these different coordinate systems.

In the first part of the fifth section we label the local minima and we construct the above injective map j under Assumption 1.1.

In the second part of this fifth section we build quasimodes adapted to our analysis and we make some scalar estimates thanks to the Laplace method. This leads directly to the proof of the final result in Section 6 using a result of [Lep1]. Again, we cannot use a single one coordinate system like in [HeNi] and we must again call on Lemma 3.18 to be able to use the Laplace method. This is due to the local geometry near a *generalized* critical point with index 1 which is rather more complicated than in [HeNi].

2. Witten Laplacian with Neumann boundary condition

2.1. Introduction and notations

This section is analogous to the second section of [HeNi] and we will use the same notations that we recall here.

Let $\overline{\Omega}$ be a C^∞ connected compact oriented Riemannian n -dimensional manifold. We will denote by g_0 the given Riemannian metric on $\overline{\Omega}$; Ω and $\partial\Omega$ will denote respectively its interior and its boundary.

The cotangent (resp. tangent) bundle on Ω is denoted by $T^*\Omega$ (resp. $T\Omega$) and the exterior fiber bundle by $\Lambda T^*\Omega = \bigoplus_{p=0}^n \Lambda^p T^*\Omega$ (resp. $\Lambda T\Omega = \bigoplus_{p=0}^n \Lambda^p T\Omega$).

The fiber bundles $\Lambda T\partial\Omega = \bigoplus_{p=0}^{n-1} \Lambda^p T\partial\Omega$ and $\Lambda T^*\partial\Omega = \bigoplus_{p=0}^{n-1} \Lambda^p T^*\partial\Omega$ are defined similarly.

The space of C^∞ , C_0^∞ , L^2 , H^s , etc. sections in any of these fiber bundles, E , on $O = \Omega$ or $O = \partial\Omega$, will be denoted respectively by $C^\infty(O; E)$, $C_0^\infty(O; E)$, $L^2(O; E)$, $H^s(O; E)$, etc..

When no confusion is possible we will simply use the short notations $\Lambda^p C^\infty$, $\Lambda^p C_0^\infty$, $\Lambda^p L^2$ and $\Lambda^p H^s$ for $E = \Lambda^p T^*\Omega$ or $E = \Lambda^p T^*\partial\Omega$.

Note that the L^2 spaces are those associated with the unit volume form for the Riemannian structure on Ω or $\partial\Omega$ (Ω and $\partial\Omega$ are oriented).

The notation $C^\infty(\overline{\Omega}; E)$ is used for the set of C^∞ sections up to the boundary.

Finally since $\partial\Omega$ is C^∞ , $C^\infty(\overline{\Omega}; E)$ is dense in $H^s(\Omega; E)$ for $s \geq 0$ and the trace operator $\omega \rightarrow \omega|_{\partial\Omega}$ extends to a surjective operator from $H^s(\Omega; E)$ onto $H^{s-1/2}(\partial\Omega; E)$ as soon as $s > 1/2$.

Let d be the exterior differential on $C_0^\infty(\Omega; \Lambda T^*\Omega)$

$$\left(d^{(p)} : C_0^\infty(\Omega; \Lambda^p T^*\Omega) \rightarrow C_0^\infty(\Omega; \Lambda^{p+1} T^*\Omega) \right)$$

and d^* its formal adjoint with respect to the L^2 -scalar product inherited from the Riemannian structure

$$\left(d^{(p)*} : C_0^\infty(\Omega; \Lambda^{p+1} T^*\Omega) \rightarrow C_0^\infty(\Omega; \Lambda^p T^*\Omega) \right).$$

Remark 2.1. — Note that d and d^* are both well defined on $C^\infty(\overline{\Omega}; \Lambda T^*\Omega)$.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

For a function $f \in \mathcal{C}^\infty(\overline{\Omega}; \mathbb{R})$ and $h > 0$, the distorted operators are defined on $\mathcal{C}^\infty(\overline{\Omega}; \Lambda T^* \Omega)$ by:

$$d_{f,h} = e^{-f(x)/h} (hd) e^{f(x)/h} \quad \text{and} \quad d_{f,h}^* = e^{f(x)/h} (hd^*) e^{-f(x)/h}.$$

The Witten Laplacian is the differential operator defined on $\mathcal{C}^\infty(\overline{\Omega}; \Lambda T^* \Omega)$ by:

$$\Delta_{f,h} = d_{f,h}^* d_{f,h} + d_{f,h} d_{f,h}^* = (d_{f,h} + d_{f,h}^*)^2. \quad (2.1.1)$$

Remark 2.2. — The last equality follows from the property $dd = d^*d^* = 0$ which implies:

$$d_{f,h} d_{f,h} = d_{f,h}^* d_{f,h}^* = 0. \quad (2.1.2)$$

It means, by restriction to the p -forms in $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$:

$$\Delta_{f,h}^{(p)} = d_{f,h}^{(p),*} d_{f,h}^{(p)} + d_{f,h}^{(p-1)} d_{f,h}^{(p-1),*}.$$

Note that (2.2) implies for all u in $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$ that:

$$\Delta_{f,h}^{(p+1)} d_{f,h}^{(p)} u = d_{f,h}^{(p)} \Delta_{f,h}^{(p)} u \quad (2.1.3)$$

and

$$\Delta_{f,h}^{(p-1)} d_{f,h}^{(p-1),*} u = d_{f,h}^{(p-1),*} \Delta_{f,h}^{(p)} u. \quad (2.1.4)$$

We end up this section by a few relations with exterior and interior products (respectively denoted by \wedge and \mathbf{i}), gradients (denoted by ∇) and Lie derivatives (denoted by \mathcal{L}) which will be very useful:

$$(df \wedge)^* = \mathbf{i}_{\nabla f} \quad (\text{in } L^2(\overline{\Omega}; \Lambda^p T^* \Omega)), \quad (2.1.5)$$

$$d_{f,h} = hd + df \wedge, \quad (2.1.6)$$

$$d_{f,h}^* = hd^* + \mathbf{i}_{\nabla f}, \quad (2.1.7)$$

$$d \circ \mathbf{i}_X + \mathbf{i}_X \circ d = \mathcal{L}_X, \quad (2.1.8)$$

$$\Delta_{f,h} = h^2(d + d^*)^2 + |\nabla f|^2 + h(\mathcal{L}_{\nabla f} + \mathcal{L}_{\nabla f}^*), \quad (2.1.9)$$

where X denotes a vector field on Ω or $\overline{\Omega}$.

Remark 2.3. — We work here on a Riemannian manifold and the operators introduced depend on the Riemannian metric g_0 . Nevertheless, we have omitted here this dependence for conciseness.

2.2. Stokes and Green formulas

In order to define suitably the self-adjoint Neumann realization of the Witten Laplacian $\Delta_{f,h}$, we need variants from the Stokes and the Green formulas.

For that, we use some notations and properties which are very convenient for boundary problems and which are introduced for example in [Sch] and recalled in [HeNi].

DEFINITION 2.4. — *We denote by \vec{n}_σ the outgoing normal at $\sigma \in \partial\Omega$ and by \vec{n}_σ^* the 1-form dual to \vec{n}_σ for the Riemannian scalar product.*

For any $\omega \in \mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$, the form $\mathbf{t}\omega$ is the element of $\mathcal{C}^\infty(\partial\Omega; \Lambda^p T^* \Omega)$ defined by:

$$(\mathbf{t}\omega)_\sigma(X_1, \dots, X_p) = \omega_\sigma(X_1^T, \dots, X_p^T), \quad \forall \sigma \in \partial\Omega,$$

with the decomposition into the tangential and normal components to $\partial\Omega$ at σ : $X_i = X_i^T \oplus x_i^\perp \vec{n}_\sigma$.

Moreover,

$$(\mathbf{t}\omega)_\sigma = \mathbf{i}_{\vec{n}_\sigma}(\vec{n}_\sigma^* \wedge \omega_\sigma).$$

The projected form $\mathbf{t}\omega$, which depends on the choice of \vec{n}_σ (i.e. on g_0), can be compared with the canonical pull-back $j^*\omega$ associated with the imbedding $j : \partial\Omega \rightarrow \Omega$. Actually the exact relationship is $j^*\omega = j^*(\mathbf{t}\omega)$. With an abuse of notation, the form $j^*(\mathbf{t}\omega)$ will be simply written $\mathbf{t}\omega$ for example in Stokes formula without any possible confusion.

The normal part of ω on $\partial\Omega$ is defined by:

$$\mathbf{n}\omega = \omega|_{\partial\Omega} - \mathbf{t}\omega \quad \in \mathcal{C}^\infty(\partial\Omega; \Lambda^p T^* \Omega).$$

If necessary $\mathbf{t}\omega$ and $\mathbf{n}\omega$ can be considered as elements of $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$ by a variant of the collar theorem (see [HeNi] or [Sch] for details).

The Hodge operator \star is locally defined in a pointwise orthonormal frame (E_1, \dots, E_n) by:

$$(\star\omega_x)(E_{\sigma(p+1)}, \dots, E_{\sigma(n)}) = \varepsilon(\sigma) \omega_x(E_{\sigma(1)}, \dots, E_{\sigma(p)}),$$

for $\omega_x \in \Lambda^p T_x^* \Omega$ and with any permutation $\sigma \in \Sigma(n)$ of $\{1, \dots, n\}$ preserving $\{1, \dots, p\}$ ($\varepsilon(\sigma)$ denotes the signature of σ).

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

We recall the formulas:

$$\star(\star\omega_x) = (-1)^{p(n-p)}\omega_x, \quad \forall \omega_x \in \Lambda^p T_x^* \Omega, \quad (2.2.1)$$

$$\langle \omega_1 | \omega_2 \rangle_{\Lambda^p L^2} = \int_{\Omega} \omega_1 \wedge \star \overline{\omega_2}, \quad \forall \omega_1, \omega_2 \in \Lambda^p L^2, \quad (2.2.2)$$

and:

$$\star d^{*,(p-1)} = (-1)^p d^{(n-p)} \star, \quad \star d^{(p)} = (-1)^{p+1} d^{*,(n-p-1)} \star, \quad (2.2.3)$$

$$\star \mathbf{n} = \mathbf{t} \star, \quad \star \mathbf{t} = \mathbf{n} \star, \quad (2.2.4)$$

$$\mathbf{t} d = d \mathbf{t}, \quad \mathbf{n} d^* = d^* \mathbf{n}. \quad (2.2.5)$$

With the previous convention $j^*(\mathbf{t}\omega) = \mathbf{t}\omega$, the Stokes formula writes:

$$\forall \omega \in C^\infty(\overline{\Omega}; \Lambda^p T^* \Omega), \quad \int_{\Omega} d\omega = \int_{\partial\Omega} j^* \omega = \int_{\partial\Omega} \mathbf{t}\omega, \quad (2.2.6)$$

and a first deformed Green formula given in [HeNi] states that

$$\begin{aligned} & \langle d_{f,h}\omega | d_{f,h}\eta \rangle_{\Lambda^{p+1} L^2} + \langle d_{f,h}^* \omega | d_{f,h}^* \eta \rangle_{\Lambda^{p-1} L^2} \\ &= \langle \Delta_{f,h}\omega | \eta \rangle_{\Lambda^p L^2} + h \int_{\partial\Omega} (\mathbf{t}\overline{\eta}) \wedge (\star \mathbf{n} d_{f,h}\omega) - h \int_{\partial\Omega} (\mathbf{t} d_{f,h}^* \omega) \wedge (\star \mathbf{n} \overline{\eta}) \end{aligned} \quad (2.2.7)$$

holds for all $\omega \in \Lambda^p H^2$ and $\eta \in \Lambda^p H^1$. This formulation of (2.2.7) does not depend on the choice of an orientation. If μ and $\mu_{\partial\Omega}$ denote the volume forms in Ω and $\partial\Omega$, the orientation is chosen such that $(\mu_{\partial\Omega})_\sigma(X_1, \dots, X_{n-1}) = \mu_\sigma(\vec{n}_\sigma, X_1, \dots, X_{n-1})$. A simple computation in normal frames (see [Sch], prop. 1.2.6) leads to:

$$\mathbf{t}\omega_1 \wedge \star \mathbf{n} \overline{\omega_2} = \langle \omega_1 | \mathbf{i}_{\vec{n}_\sigma} \omega_2 \rangle_{\Lambda^p T_\sigma^* \Omega} d\mu_{\partial\Omega}, \quad (2.2.8)$$

for $\omega_1 \in C^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$ and $\omega_2 \in C^\infty(\overline{\Omega}; \Lambda^{p+1} T^* \Omega)$.

DEFINITION 2.5. — We denote by $\frac{\partial f}{\partial n}(\sigma)$ or $\partial_n f(\sigma)$ the normal derivative of f at σ :

$$\frac{\partial f}{\partial n}(\sigma) = \partial_n f(\sigma) := \langle \nabla f(\sigma) | \vec{n}_\sigma \rangle.$$

As a consequence of (2.2.8) we get the following useful decomposition formula.

LEMMA 2.6 (*Normal Green Formula*). — *The identity*

$$\begin{aligned} \|d_{f,h}\omega\|_{\Lambda^{p+1}L^2}^2 + \|d_{f,h}^*\omega\|_{\Lambda^{p-1}L^2}^2 &= h^2 \|d\omega\|_{\Lambda^{p+1}L^2}^2 + h^2 \|d^*\omega\|_{\Lambda^{p-1}L^2}^2 \\ &+ \|\ |\nabla f|\omega\|_{\Lambda^p L^2}^2 + h\langle(\mathcal{L}_{\nabla f} + \mathcal{L}_{\nabla f}^*)\omega|\omega\rangle_{\Lambda^p L^2} \\ &+ h \int_{\partial\Omega} \langle\omega|\omega\rangle_{\Lambda^p T_\sigma^* \Omega} \left(\frac{\partial f}{\partial n}\right)(\sigma) d\mu_{\partial\Omega} \end{aligned}$$

holds for any $\omega \in \Lambda^p H^1$ such that $\mathbf{n}\omega = 0$.

Proof. — Since $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$ is dense in $\Lambda^p H^1$, while both terms of the identity are continuous on $\Lambda^p H^1$, the form ω can be assumed to be in $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^* \Omega)$.

We use the relation (2.2.7) with both $f = 0$ ($d_{0,h} = hd$ and $d_{0,h}^* = hd^*$) and a general $f \in \mathcal{C}^\infty(\overline{\Omega}; \mathbb{R})$. We obtain:

$$\begin{aligned} \|d_{f,h}\omega\|_{\Lambda^{p+1}L^2}^2 + \|d_{f,h}^*\omega\|_{\Lambda^{p-1}L^2}^2 - h^2 \|d\omega\|_{\Lambda^{p+1}L^2}^2 - h^2 \|d^*\omega\|_{\Lambda^{p-1}L^2}^2 &= \\ \langle(\Delta_{f,h} - \Delta_{0,h})\omega|\omega\rangle_{\Lambda^p L^2} + h \int_{\partial\Omega} (\mathbf{t}\overline{\omega}) \wedge \star \mathbf{n}(df \wedge \omega) - h \int_{\partial\Omega} (\mathbf{t}\mathbf{i}_{\nabla f}\omega) \wedge (\star \mathbf{n}\overline{\omega}) &= \\ = \langle(\Delta_{f,h} - \Delta_{0,h})\omega|\omega\rangle_{\Lambda^p L^2} + h \int_{\partial\Omega} \langle\omega|\mathbf{i}_{\overline{\mathbf{n}}_\sigma}(df \wedge \omega)\rangle_{\Lambda T_\sigma^* \Omega} d\mu_{\partial\Omega} . \end{aligned}$$

By (2.1.9):

$$\langle(\Delta_{f,h} - \Delta_{0,h})\omega|\omega\rangle_{\Lambda^p L^2} = \|\ |\nabla f|\omega\|_{\Lambda^p L^2}^2 + h\langle(\mathcal{L}_{\nabla f} + \mathcal{L}_{\nabla f}^*)\omega|\omega\rangle_{\Lambda^p L^2} .$$

For the integral term, we write:

$$\begin{aligned} \mathbf{i}_{\overline{\mathbf{n}}_\sigma}(df \wedge \omega)(X_1, \dots, X_p) &= (df \wedge \omega)(\overline{\mathbf{n}}_\sigma, X_1, \dots, X_p) \\ &= df(\overline{\mathbf{n}}_\sigma) \cdot \omega(X_1, \dots, X_p) \quad \text{because } \mathbf{n}\omega = 0 \\ &= \langle \nabla f(\sigma) | \overline{\mathbf{n}}_\sigma \rangle \cdot \omega(X_1, \dots, X_p) \\ &= \left(\frac{\partial f}{\partial n}(\sigma)\right) \omega(X_1, \dots, X_p) , \end{aligned}$$

which proves the lemma. \square

2.3. Normal Neumann realization

In this subsection, we specify the self-adjoint realization of $\Delta_{f,h}^{(0)}$ in which we are interested. Like in [HeNi], we want this self-adjoint realization (denoted by $\Delta_{f,h}^N$) to coincide with the Neumann realization on 0-forms and to

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

preserve the complex structure:

$$\begin{aligned} (1 + \Delta_{f,h}^{N,(p+1)})^{-1} d_{f,h}^{(p)} &= d_{f,h}^{(p)} (1 + \Delta_{f,h}^{N,(p)})^{-1} \\ \text{and} \\ (1 + \Delta_{f,h}^{N,(p-1)})^{-1} d_{f,h}^{(p-1),*} &= d_{f,h}^{(p-1),*} (1 + \Delta_{f,h}^{N,(p)})^{-1} \end{aligned}$$

on the form domain of $\Delta_{f,h}^{N,(p)}$.

Having in mind the works [Sch] and [ChLi] about cohomology complexes and boundary problems, we introduce the space:

$$\Lambda^p H_{0,\mathbf{n}}^1 = H_{0,\mathbf{n}}^1(\Omega; \Lambda^p T^* \Omega) = \{ \omega \in H^1(\Omega; \Lambda^p T^* \Omega); \mathbf{n}\omega = 0 \}. \quad (2.3.1)$$

In the case $p = 0$, it coincides with the space $H^1(\Omega)$, while for $p \geq 1$ the condition says only that the form vanishes on $\partial\Omega$ when applied to non tangential p -vectors. Since the boundary $\partial\Omega$ is assumed to be regular, the space

$$\Lambda^p \mathcal{C}_{0,\mathbf{n}}^\infty = \mathcal{C}_{0,\mathbf{n}}^\infty(\Omega; \Lambda^p T^* \Omega) = \{ \omega \in \mathcal{C}^\infty(\overline{\Omega}, \Lambda^p T^* \Omega); \mathbf{n}\omega = 0 \}$$

is dense in $\Lambda^p H_{0,\mathbf{n}}^1$. The following construction is a variant of known results in the case $f = 0$ (see [Sch]). We will use the notations:

$$\mathcal{D}_{f,h}(\omega, \eta) = \langle d_{f,h}\omega \mid d_{f,h}\eta \rangle_{\Lambda^{p+1}L^2} + \langle d_{f,h}^*\omega \mid d_{f,h}^*\eta \rangle_{\Lambda^{p-1}L^2}$$

and

$$\mathcal{D}_{f,h}(\omega) = \mathcal{D}_{f,h}(\omega, \omega) = \|d_{f,h}\omega\|_{\Lambda^{p+1}L^2}^2 + \|d_{f,h}^*\omega\|_{\Lambda^{p-1}L^2}^2.$$

PROPOSITION 2.7. — *The non negative quadratic form $\omega \rightarrow \mathcal{D}_{f,h}(\omega)$ is closed on $\Lambda^p H_{0,\mathbf{n}}^1$. The associated (self-adjoint) Friedrichs extension is denoted by $\Delta_{f,h}^{N,(p)}$. Its domain is:*

$$D(\Delta_{f,h}^{N,(p)}) = \{ u \in \Lambda^p H^2; \mathbf{n}\omega = 0 \text{ and } \mathbf{n}d_{f,h}\omega = 0 \},$$

and we have:

$$\forall \omega \in D(\Delta_{f,h}^{N,(p)}), \quad \Delta_{f,h}^{N,(p)}\omega = \Delta_{f,h}^{(p)}\omega \quad \text{in } \Omega.$$

Proof. — By the same argument as in the proof of Proposition 2.4 of [HeNi], the space $\Lambda^p H_{0,\mathbf{n}}^1$ is isomorphic to the direct sum:

$$\Lambda^p H_0^1 \oplus \mathfrak{t} \Lambda^p H^{1/2}(\partial\Omega; \Lambda^p T^* \Omega)$$

with continuous embedding. Hence, since $\partial\Omega$ is a regular boundaryless manifold, its dual is the direct sum of $\Lambda^p H^{-1}$ and $\mathfrak{t}\Lambda^p H^{-1/2}(\partial\Omega; \Lambda^p T^*\Omega)$:

$$(\Lambda^p H_{0,\mathbf{n}}^1)' = \Lambda^p H^{-1} \oplus \mathfrak{t}\Lambda^p H^{-1/2}(\partial\Omega; \Lambda^p T^*\Omega) .$$

We have to check that $\omega \mapsto \mathcal{D}_{f,h}^{(p)}(\omega) + C \|\omega\|_{\Lambda^p L^2}^2$ is equivalent to the square of the $\Lambda^p H^1$ norm on $\Lambda^p H_{0,\mathbf{n}}^1$. By (2.1.6)-(2.1.9) this is equivalent to the same result for $f = 0$ and $h = 1$. This last case is known as Gaffney's inequality which is a consequence of the Weitzenböck formula (see [Sch], Theorem 2.1.7).

Hence the quadratic form $\omega \rightarrow \mathcal{D}_{f,h}(\omega)$ is closed on $\Lambda^p H_{0,\mathbf{n}}^1$ and the identity

$$\forall \eta \in \Lambda^p H_{0,\mathbf{n}}^1, \quad \mathcal{D}_{f,h}^{(p)}(\omega, \eta) = \langle A^{(p)}\omega, \eta \rangle$$

defines an isomorphism $A^{(p)} : \Lambda^p H_{0,\mathbf{n}}^1 \rightarrow (\Lambda^p H_{0,\mathbf{n}}^1)'$.

The self-adjoint Friedrichs extension $\Delta_{f,h}^{N,(p)}$ is then defined as the operator:

$$D(\Delta_{f,h}^{N,(p)}) = \left\{ \omega \in \Lambda^p H_{0,\mathbf{n}}^1, A^{(p)}\omega \in \Lambda^p L^2 \right\}, \quad \Delta_{f,h}^{N,(p)}\omega = A^{(p)}\omega .$$

It remains to identify this domain and the explicit action of $A^{(p)}$.

If ω belongs to $D(\Delta_{f,h}^{N,(p)})$, by the first Green formula (2.2.7) we get:

$$\forall \eta \in \Lambda^p \mathcal{C}_0^\infty, \quad \langle \omega | A^{(p)}\eta \rangle = \mathcal{D}_{f,h}^{(p)}(\omega, \eta) = \langle \omega | \Delta_{f,h}^{(p)}\eta \rangle .$$

The inequality:

$$|\mathcal{D}_{f,h}^{(p)}(\omega, \eta)| \leq C \|\omega\|_{\Lambda^p H^1} \|\eta\|_{\Lambda^p H^1} ,$$

together with the density of $\Lambda^p \mathcal{C}_0^\infty$ in $\Lambda^p H_0^1$ implies that the current $\Delta_{f,h}^{(p)}\omega \in \mathcal{D}'(\Omega; \Lambda^p T^*\Omega)$ is indeed the $\Lambda^p H^{-1}$ component of $A^{(p)}\omega$.

Assume that ω belongs to $\Lambda^p H_{0,\mathbf{n}}^1 \cap \Lambda^p H^2$; then the Green formula (2.2.7) gives:

$$h \int_{\partial\Omega} (\mathfrak{t}\bar{\eta}) \wedge (\star \mathbf{nd}_{f,h}\omega) = \mathcal{D}_{f,h}^{(p)}(\omega, \eta) - \langle \Delta_{f,h}^{(p)}\omega | \eta \rangle_{\Lambda^p L^2}, \quad \forall \eta \in \Lambda^p H_{0,\mathbf{n}}^1 .$$

By density, one can define, for any ω in $\Lambda^p H_{0,\mathbf{n}}^1$ such that $\Delta_{f,h}^{(p)}\omega \in \Lambda^p L^2$, a trace of $\mathbf{nd}_{f,h}\omega$ by the previous identity, observing that the r.h.s. defines an

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

antilinear continuous form with respect to η . With this generalized definition of $\mathbf{n}d_{f,h}^{(p)}\omega$ we claim that:

$$D(\Delta_{f,h}^{N,(p)}) = \left\{ \omega \in \Lambda^p H_{0,\mathbf{n}}^1, \Delta_{f,h}^{(p)}\omega \in \Lambda^p L^2 \text{ and } \mathbf{n}d_{f,h}^{(p)}\omega = 0 \right\}.$$

The last point consists in observing that the boundary value problem

$$\Delta_{f,h}^{(p)}u = g, \quad \mathbf{n}u = g_1, \quad \mathbf{n}d_{f,h}^{(p)}u = g_2 \quad (2.3.2)$$

satisfies the Lopatinski-Shapiro conditions. At the principal symbol level ($h > 0$ fixed), these conditions are indeed the same as for

$$(dd^* + d^*d)^{(p)}u = g, \quad \mathbf{n}u = g_1, \quad \mathbf{n}d^{(p)}u = g_2.$$

This is checked in [Sch]. Hence any solution to (2.3.2) with $g \in \Lambda^p L^2$, $g_1 = g_2 = 0$ belongs to $\Lambda^p H^2$. \square

PROPOSITION 2.8. — *For any $p \in \{0, \dots, n\}$, the self-adjoint unbounded operator $\Delta_{f,h}^{N,(p)}$ introduced in Proposition 2.7 has a compact resolvent.*

Moreover, if $z \in \mathbb{C} \setminus \mathbb{R}_+$, the commutation relations

$$(z - \Delta_{f,h}^{N,(p+1)})^{-1}d_{f,h}^{(p)}v = d_{f,h}^{(p)}(z - \Delta_{f,h}^{N,(p)})^{-1}v,$$

and

$$(z - \Delta_{f,h}^{N,(p-1)})^{-1}d_{f,h}^{(p-1),*}v = d_{f,h}^{(p-1),*}(z - \Delta_{f,h}^{N,(p)})^{-1}v,$$

hold for any $v \in \Lambda^p H_{0,\mathbf{n}}^1$.

Proof. — The domain of the operator is contained in $\Lambda^p H^2$, which is compactly embedded in $\Lambda^p L^2$, by Sobolev injections. This yields the first statement.

Since $\Lambda^p \mathcal{C}_{0,\mathbf{n}}^\infty$ is dense in $\Lambda^p H_{0,\mathbf{n}}^1$, it is sufficient to consider the case when $v \in \Lambda^p \mathcal{C}_{0,\mathbf{n}}^\infty$. For such a v and for $z \in \mathbb{C} \setminus \mathbb{R}_+$, we set:

$$u = (z - \Delta_{f,h}^{N,(p)})^{-1}v.$$

Due to the ellipticity of the associated boundary problem (the Lopatinski-Shapiro conditions are verified) u belongs to $\mathcal{C}^\infty(\overline{\Omega}; \Lambda^p T^*\Omega)$. The commutation relations (2.1.3) and (2.1.4) can be applied since here $f \in \mathcal{C}^\infty(\overline{\Omega}; \mathbb{R})$:

$$(z - \Delta_{f,h}^{(p+1)})d_{f,h}^{(p)}u = d_{f,h}^{(p)}(z - \Delta_{f,h}^{(p)})u = d_{f,h}^{(p)}v \quad (2.3.3)$$

and

$$(z - \Delta_{f,h}^{(p-1)})d_{f,h}^{(p-1),*}u = d_{f,h}^{(p-1),*}(z - \Delta_{f,h}^{(p)})u = d_{f,h}^{(p-1),*}v. \quad (2.3.4)$$

Since $u \in D(\Delta_{f,h}^{N,(p)})$, we have $\mathbf{n}u = 0$ and $\mathbf{n}d_{f,h}^{(p)}u = 0$.

Then, $\mathbf{n}d_{f,h}u = 0$ and $\mathbf{n}d_{f,h}d_{f,h}u = 0$ imply $d_{f,h}u \in D(\Delta_{f,h}^{N,(p+1)})$. So by (2.3.3) we have:

$$d_{f,h}^{(p)}(z - \Delta_{f,h}^{N,(p)})^{-1}v = d_{f,h}u = (z - \Delta_{f,h}^{N,(p+1)})^{-1}d_{f,h}^{(p)}v .$$

In order to show the second commutation relation, we first use the relation (2.2.5) which implies:

$$\mathbf{n}d_{f,h}^*u = hd^*nu + \mathbf{n}(i_{\nabla}fu) = 0 .$$

For the normal trace of the differential, we write $(\Delta_{f,h}u = zu - v)$:

$$\mathbf{n}d_{f,h}(d_{f,h}^*u) = z\mathbf{n}u - \mathbf{n}v - \mathbf{n}d_{f,h}^*d_{f,h}u = -d_{f,h}^*\mathbf{n}d_{f,h}u = 0 .$$

Hence $d_{f,h}^{(p-1),*}u$ belongs to $D(\Delta_{f,h}^{N,(p-1)})$ and the identity (2.3.4) yields the last commutation relation to show. \square

DEFINITION 2.9. — *For any Borel subset $E \subset \mathbb{R}$ and $p \in \{0, \dots, n\}$, we will denote by $1_E(\Delta_{f,h}^{N,(p)})$ the spectral projection of $\Delta_{f,h}^{N,(p)}$ on E .*

From Proposition 2.8 and Stone's Formula we deduce:

COROLLARY 2.10. — *For any Borel subset $E \subset \mathbb{R}$, the identities*

$$1_E(\Delta_{f,h}^{N,(p+1)})d_{f,h}^{(p)}v = d_{f,h}^{(p)}1_E(\Delta_{f,h}^{N,(p)})v$$

and

$$1_E(\Delta_{f,h}^{N,(p-1)})d_{f,h}^{(p-1),*}v = d_{f,h}^{(p-1),*}1_E(\Delta_{f,h}^{N,(p)})v$$

hold for all $v \in \Lambda^p H_{0,\mathbf{n}}^1$.

In the particular case when v is an eigenvector of $\Delta_{f,h}^{N,(p)}$ corresponding to the eigenvalue λ , then $d_{f,h}^{(p)}v$ (resp. $d_{f,h}^{(p-1),}v$) belongs to the spectral subspace $\text{Ran } 1_{\{\lambda\}}(\Delta_{f,h}^{N,(p+1)})$ (resp. $\text{Ran } 1_{\{\lambda\}}(\Delta_{f,h}^{N,(p-1)})$).*

Proposition 2.8 and Corollary 2.10 were stated for p -forms $v \in \Lambda^p H_{0,\mathbf{n}}^1(\Omega)$, belonging to the form domain of $\Delta_{f,h}^{N,(p)}$. It is convenient to work in this framework because the multiplication by any cut-off function preserves the form domain $\Lambda H_{0,\mathbf{n}}^1(\Omega)$:

$$(\omega \in \Lambda H_{0,\mathbf{n}}^1(\Omega), \quad \chi \in C^\infty(\overline{\Omega})) \Rightarrow (\chi\omega \in \Lambda H_{0,\mathbf{n}}^1(\Omega)) ,$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

while this property is no more true for $D(\Delta_{f,h}^N)$. In this spirit, we will often refer to the following easy consequence of the spectral theorem.

LEMMA 2.11. — *Let A be a non negative self-adjoint operator on a Hilbert space \mathcal{H} with associated quadratic form $q_A(x) = (x | Ax)$ and with form domain $Q(A)$. Then for any $a, b \in (0, +\infty)$, the implication*

$$(q_A(u) \leq a) \Rightarrow \left(\|1_{[b, +\infty)}(A)u\|^2 \leq \frac{a}{b} \right)$$

holds for any $u \in Q(A)$.

3. First localization of the spectrum

3.1. Introduction and result

Let us first recall that we are working with the fixed Riemannian metric g_0 on $\overline{\Omega}$. Like in the third section of [HeNi] for their tangential Dirichlet realization of the Witten Laplacian, we check here that the number of eigenvalues of $\Delta_{f,h}^{N,(p)}$ smaller than $h^{3/2}$ equals a Morse index which involves in its definition the boundary conditions. To this end, we will adapt [HeNi] which uses techniques yet presented in [Sim], [CyFrKiSi], [ChLi], [Bis], [Bur], and in [Hel1]. These techniques are also used in the same spirit in [KoPrSh].

In order to make the connection between the normal Neumann realization of the Witten Laplacian $\Delta_{f,h}^N$ and the Morse theory, we assume additional properties for the function f up to the boundary $\partial\Omega$.

ASSUMPTION 3.1. — *The real-valued function $f \in C^\infty(\overline{\Omega})$ is a Morse function on Ω with no critical points in $\partial\Omega$. In addition its restriction $f|_{\partial\Omega}$ is a Morse function on $\partial\Omega$.*

Remark 3.2. — With this assumption, the function f has a finite number of critical points with index p in Ω . Note furthermore that the assumption ensures that there is no critical point on $\partial\Omega$, which implies that the outgoing normal derivative $\frac{\partial f}{\partial n}(U)$ is not 0 when U is a critical point of $f|_{\partial\Omega}$.

DEFINITION 3.3. — *For $\ell \in \{0, \dots, n\}$, the integer $m_{\ell,-}^{\partial\Omega}$ is the number of critical points U of $f|_{\partial\Omega}$ with index ℓ such that $\frac{\partial f}{\partial n}(U) < 0$ (with the additional convention $m_{n,-}^{\partial\Omega} = 0$).*

For $p \in \{0, \dots, n\}$, let

$$m_p^{\overline{\Omega}} = m_p^\Omega + m_{p,-}^{\partial\Omega} .$$

Remark 3.4. — In [HeNi], the authors worked with the tangential Dirichlet conditions ($\mathbf{t}\omega = 0$ and $\mathbf{t}d_{f,h}^*\omega = 0$) and the corresponding definition was similar with $m_{\ell,-}^{\partial\Omega}$ and $\frac{\partial f}{\partial n}(U) < 0$ replaced respectively by $m_{\ell-1,-}^{\partial\Omega}$ and $\frac{\partial f}{\partial n}(U) > 0$.

The aim of this section is to prove the following theorem:

THEOREM 3.5. — *Under Assumption 3.1, there exists $h_0 > 0$, such that the normal Neumann realization of the Witten Laplacian $\Delta_{f,h}^N$ introduced in Subsection 2.3 has, for $h \in (0, h_0]$, the following property:*

For any $p \in \{0, \dots, n\}$, the spectral subspace $F^{(p)} = \text{Ran}1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(p)})$ has rank: $\dim F^{(p)} = m_p^{\bar{\Omega}}$.

To prove this theorem, we will adapt for the normal Neumann realization of the Witten Laplacian the proof given in [HeNi] for the tangential Dirichlet realization. Many points of this demonstration do not require any modification, so we will only recall these results without any demonstration.

The theorem will be proved in the Subsection 3.3.

3.2. A few preliminary lemmas

In this subsection, we recall some results of [HeNi] needed to prove Theorem 3.5.

3.2.1 Variational results for the Witten Laplacian on \mathbb{R}^k

Let g be a C^∞ metric on \mathbb{R}^k which equals the Euclidean metric outside a compact set K .

ASSUMPTION 3.6 (g). — *The function f is a Morse C^∞ real-valued function and there exist $C_1 > 0$ and a compact K such that, for the metric g :*

$$\forall x \in \mathbb{R}^k \setminus K, \quad |\nabla f(x)| \geq C_1^{-1} \quad \text{and} \quad |\text{Hess } f(x)| \leq C_1 |\nabla f(x)|^2. \quad (3.2.1)$$

Note that the above assumption ensures that f has a finite number of critical points and m_p will denote the number of critical points with index p .

Let us recall the Propositions 3.6 and 3.7 of [HeNi]. They gather consequences of Simader's Theorem in [Sima] about the essential self-adjointness of non negative Schrödinger operators, of Persson's Lemma in [Per] about

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

the localization of the essential spectrum and of the semiclassical analysis à la Witten in [Wit] leading to Morse inequalities. We also refer the reader to [CyFrKiSi][Hen] [Hel3] or [Zha] for the Witten approach to Morse inequalities in the boundaryless case and to [Mil1] and [Lau] for a topological presentation of Morse theory.

PROPOSITION 3.7 — *Under Assumption 3.6, there exist $h_0 > 0$, $c_0 > 0$ and $c_1 > 0$ such that the following properties are satisfied for any $h \in (0, h_0]$:*

i) *The Witten Laplacian $\Delta_{f,h}$ considered as an unbounded operator on $L^2(\mathbb{R}^k; \Lambda T^* \mathbb{R}^k)$ is essentially self-adjoint on $\mathcal{C}_0^\infty(\mathbb{R}^k; \Lambda T^* \mathbb{R}^k)$.*

ii) *For any Borel subset E in \mathbb{R} , the identities*

$$\begin{aligned} 1_E(\Delta_{f,h}^{(p+1)})d_{f,h}^{(p)}u &= d_{f,h}^{(p)}1_E(\Delta_{f,h}^{(p)})u \\ \text{and} \\ 1_E(\Delta_{f,h}^{(p-1)})d_{f,h}^{(p-1),*}u &= d_{f,h}^{(p-1),*}1_E(\Delta_{f,h}^{(p)})u \end{aligned} \quad (3.2.2)$$

hold for any u belonging to the form domain of $\Delta_{f,h}^{(p)}$.

In particular, if v is an eigenvector of $\Delta_{f,h}^{(p)}$ associated with the eigenvalue λ , then $d_{f,h}^{(p)}v$ (resp. $d_{f,h}^{(p-1),}v$) belongs to the spectral subspace $\text{Ran } 1_{\{\lambda\}}(\Delta_{f,h}^{(p+1)})$ (resp. $\text{Ran } 1_{\{\lambda\}}(\Delta_{f,h}^{(p-1)})$).*

iii) *The essential spectrum $\sigma_{\text{ess}}(\Delta_{f,h}^{(p)})$ is contained in $[c_1, +\infty)$.*

iv) *The range of $1_{[0, c_0 h]}(\Delta_{f,h}^{(p)})$ has dimension m_p , for all $h \in (0, h_0]$.*

PROPOSITION 3.8. — *If the Morse function f satisfies Assumption 3.6 and admits a unique critical point at $x = 0$ with index p_0 , so $m_p = \delta_{p, p_0}$, then there exist $h_0 > 0$ and $c_0 > 0$, such that the following properties hold for $h \in (0, h_0]$:*

i) *For $p \neq p_0$, $\Delta_{f,h}^{(p)} \geq c_0 h \text{Id}$.*

ii) *If $\psi_{p_0}^h$ is a normalized eigenvector of the one dimensional spectral subspace $\text{Ran } 1_{[0, c_0 h]}(\Delta_{f,h}^{(p_0)})$, it satisfies*

$$d_{f,h}\psi_{p_0}^h = 0, \quad d_{f,h}^{(p_0-1),*}\psi_{p_0}^h = 0 \quad \text{and} \quad \Delta_{f,h}^{(p_0)}\psi_{p_0}^h = 0,$$

so that $\text{Ran } 1_{[0, c_0 h]}(\Delta_{f,h}^{(p_0)}) = \text{Ker } \Delta_{f,h}^{(p_0)}$. Moreover

$$\sigma(\Delta_{f,h}^{(p_0)}) \setminus \{0\} \subset [c_0 h, \infty).$$

iii) If $\chi \in C_0^\infty(\mathbb{R}^k)$ satisfies $\chi = 1$ in a neighborhood of 0, then there exists $C_\chi \geq 1$, such that, for all $h \in (0, h_0/C_\chi)$, the inequality,

$$(1 - \chi)\Delta_{f,h}^{(p)}(1 - \chi) \geq C_\chi^{-1} [1 - \chi]^2 ,$$

holds in the sense of quadratic form on $\Lambda^p H^1(\mathbb{R}^k)$.

3.2.2 The model half-space problem

We work here on $\mathbb{R}_-^n = \mathbb{R}^{n-1} \times (-\infty, 0)$ with a Riemannian metric \tilde{g}_0 . Assume furthermore that there are coordinates $x = (x', x_n)$ such that $\tilde{g}_0 = \sum_{i,j=1}^n \tilde{g}_{ij}^0(x) dx_i dx_j$ satisfies

$$\tilde{g}_{i,n}^0 = \tilde{g}_{n,i}^0 = 0 \quad \text{for } i < n \tag{3.2.3}$$

and

$$\forall x \in \overline{\mathbb{R}_-^n} \setminus K_1, \quad \partial_x \tilde{g}_{ij}^0(x) = 0 , \tag{3.2.4}$$

for some compact set $K_1 \subset \overline{\mathbb{R}_-^n}$.

In this paragraph, the coordinates (x', x_n) are fixed while different metrics on $\overline{\mathbb{R}_-^n}$ are considered. The notation $G(\cdot)$ will be used for the matrix valued map $x \mapsto G(x) = {}^t G(x) = (g_{ij}(x))_{i,j} \in GL^n(\mathbb{R})$, which is assumed to be a C^∞ function. According to the standard notation, the coefficients of $G(x)^{-1}$ are written $g^{ij}(x)$.

Consider also a function f which has a specific form in the same coordinates (x', x_n) .

ASSUMPTION 3.9. — *The function $f \in C^\infty(\overline{\mathbb{R}_-^n})$ satisfies:*

- i) *The estimates $|\nabla f(x)| \geq C^{-1}$ and $|\partial_x^\alpha f(x)| \leq C_\alpha$ hold, for all $x \in \overline{\mathbb{R}_-^n}$ and all $\alpha \in \mathbb{N}^n$, $\alpha \neq 0$.*
- ii) *The function f is the sum $f(x', x_n) = -\frac{1}{2}f_+(x_n) + \frac{1}{2}f_-(x')$. Moreover, there exists $C_1 > 0$ such that*

$$\forall x_n \in (-\infty, 0) , \quad C_1^{-1} \leq |\partial_{x_n} f_+(x_n)| \leq C_1 ,$$

and f_- is a Morse function on \mathbb{R}^{n-1} which satisfies Assumption 3.6 for the metric $\sum_{i,j=1}^{n-1} \tilde{g}_{ij}^0(x', 0) dx_i dx_j$ and admits a unique critical point at $x' = 0$ with index p_0 .

The boundedness of $|\partial_x^\alpha f|$, $1 \leq |\alpha| \leq 2$, avoids any subtle questions about the domains.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

PROPOSITION 3.10. — *Under Assumption 3.9-i), the unbounded operator $\Delta_{f,h}^N$ on $L^2(\mathbb{R}_-^n; \Lambda T^* \mathbb{R}_-^n)$, with domain*

$$D(\Delta_{f,h}^N) = \{ \omega \in \Lambda H^2(\mathbb{R}_-^n), \quad \mathbf{n}\omega = 0, \quad \mathbf{n}d_{f,h}\omega = 0 \},$$

is self-adjoint.

If E is any Borel subset of \mathbb{R} , the relations

$$\begin{aligned} 1_E(\Delta_{f,h}^{N,(p+1)}) d_{f,h}^{(p)} u &= d_{f,h}^{(p)} 1_E(\Delta_{f,h}^{N,(p)}) u, \\ \text{and} \\ 1_E(\Delta_{f,h}^{N,(p-1)}) d_{f,h}^{(p-1),*} u &= d_{f,h}^{(p-1),*} 1_E(\Delta_{f,h}^{N,(p)}) u, \end{aligned} \quad (3.2.5)$$

hold for any $u \in \Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}_-^n)$.

Proof. — The uniform estimate on ∇f allows the same proof as for Proposition 2.8 and Corollary 2.10 (here $\mathcal{C}_{0,\mathbf{n}}^\infty$ denotes the space of \mathcal{C}^∞ compactly supported functions in $\overline{\mathbb{R}_-^n}$ with a vanishing normal component on $\{x_n = 0\}$). \square

We are looking for a result similar to Proposition 3.7 and Proposition 3.8 for the case with normal boundary condition on \mathbb{R}_-^n (this result will be stated in Subsection 3.2). One difficulty here comes from the metric which, although diagonal in the coordinates (x', x_n) , is not constant. The general case can be reduced to a simpler situation where $g_{ij}(x) = g_{ij}(x')$ with $g_{nn} = 1$ after several steps.

We need some notations.

DEFINITION 3.11. — *For a metric g which satisfies (3.2.4), the corresponding H^s -norm on the space $\Lambda^p H^s(\overline{\mathbb{R}_-^n})$ is denoted by $\| \cdot \|_{\Lambda^p H^s, g}$ and the notation $\| \cdot \|_{\Lambda^p H^s}$ is kept for the Euclidean metric $g_e = \sum_{i=1}^n dx_i^2$. Similarly, the quadratic form associated with $\Delta_{f,h}^{N,(p)}$ is written*

$$\mathcal{D}_{g,f,h}(\omega) = \|d_{g,f,h}^* \omega\|_{\Lambda^{p-1} L^2, g}^2 + \|d_{f,h} \omega\|_{\Lambda^{p+1} L^2, g}^2, \quad \forall \omega \in \Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}_-^n),$$

where the codifferential $d_{g,f,h}^$ also depends on g .*

Remark 3.12. — For the considered metrics satisfying (3.2.3) and (3.2.4), the different (L^2, g) -norms are equivalent.

The required accuracy while comparing the quadratic forms $\mathcal{D}_{g,f,h}$ needs some care.

We will work further with partitions of unity and the following proposition, similar to the standard IMS localization formula (see [CyFrKiSi]), but in the case with boundary, will be useful.

PROPOSITION 3.13 (*IMS Localization Formula*). — For $W = \Omega$ or $W = \mathbb{R}_-^n$, consider $\{\chi_k\}_{1,\dots,N}$ a partition of unity of \overline{W} (i.e. satisfying $\sum_{k=1}^N \chi_k^2 = 1$ on \overline{W}).

Let g and f be respectively a Riemannian metric and a C^∞ function (satisfying Assumption 3.9-i) in the case \mathbb{R}_-^n) on \overline{W} .

The following IMS localization formula is then valid:

$$\forall \omega \in \Lambda H_{0,\mathbf{n}}^1, \quad \mathcal{D}_{g,f,h}(\omega) = \sum_{k=1}^N \mathcal{D}_{g,f,h}(\chi_k \omega) - h^2 \|\|\nabla \chi_k | \omega\|_{\Lambda L^2, g}^2. \quad (3.2.6)$$

Proof. — For clarity, we omit the dependence on g in the proof.

Recall, from $\sum_{k=1}^N \chi_k^2 = 1$, than for any $\eta \in \Lambda H^1$:

$$\sum_{k=1}^N \chi_k d\chi_k \wedge \eta = 0, \text{ and by duality (2.1.5), } \sum_{k=1}^N \chi_k \mathbf{i}_{\nabla \chi_k} \eta = 0. \quad (3.2.7)$$

Now, for any $\omega \in \Lambda H_{0,\mathbf{n}}^1$ and $k \in \{1, \dots, N\}$,

$$\mathcal{D}_{f,h}(\chi_k \omega) = \|d_{f,h}(\chi_k \omega)\|^2 + \|d_{f,h}^*(\chi_k \omega)\|^2.$$

From (2.1.6) and (2.1.7),

$$d_{f,h}(\chi_k \omega) = h d\chi_k \wedge \omega + \chi_k d_{f,h} \omega \quad \text{and} \quad d_{f,h}^*(\chi_k \omega) = h \mathbf{i}_{\nabla \chi_k} \omega + \chi_k d_{f,h}^* \omega.$$

Hence, from $\sum_{k=1}^N \chi_k^2 = 1$, (2.1.6), and (2.1.7), for any $\omega \in \Lambda H_{0,\mathbf{n}}^1$,

$$\begin{aligned} \sum_{k=1}^N \mathcal{D}_{f,h}(\chi_k \omega) &= \mathcal{D}_{f,h}(\omega) + \sum_{k=1}^N h^2 (\langle d\chi_k \wedge \omega | d\chi_k \wedge \omega \rangle + \langle \mathbf{i}_{\nabla \chi_k} \omega | \mathbf{i}_{\nabla \chi_k} \omega \rangle) \\ &+ \sum_{k=1}^N 2 \operatorname{Re} (\langle h d\chi_k \wedge \omega | h \chi_k d\omega + \chi_k df \wedge \omega \rangle + \langle h \mathbf{i}_{\nabla \chi_k} \omega | h \chi_k d^* \omega + \chi_k \mathbf{i}_{\nabla f} \omega \rangle). \end{aligned}$$

Using (3.2.7),

$$\sum_{k=1}^N \mathcal{D}_{f,h}(\chi_k \omega) = \mathcal{D}_{f,h}(\omega) + h^2 \sum_{k=1}^N (\langle d\chi_k \wedge \omega | d\chi_k \wedge \omega \rangle + \langle \mathbf{i}_{\nabla \chi_k} \omega | \mathbf{i}_{\nabla \chi_k} \omega \rangle).$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

At least, the identity

$$\mathbf{i}_X(\alpha \wedge \beta) = (\mathbf{i}_X\alpha) \wedge \beta + (-1)^{\deg \alpha} \alpha \wedge (\mathbf{i}_X\beta)$$

implies

$$\begin{aligned} \langle d\chi_k \wedge \omega \mid d\chi_k \wedge \omega \rangle + \langle \mathbf{i}_{\nabla\chi_k}\omega \mid \mathbf{i}_{\nabla\chi_k}\omega \rangle &= \langle \mathbf{i}_{\nabla\chi_k}(d\chi_k \wedge \omega) + d\chi_k \wedge (\mathbf{i}_{\nabla\chi_k}\omega) \mid \omega \rangle \\ &= \langle (\mathbf{i}_{\nabla\chi_k}d\chi_k)\omega \mid \omega \rangle = \langle |\nabla\chi_k|^2 \omega \mid \omega \rangle, \end{aligned}$$

which proves the proposition. \square

Let us give now two lemmas whose proofs are the same than the proofs of Lemmas 3.11 and 3.12 of [HeNi].

The first lemma provides a reduction to the case $\partial_{x_n}G = 0$ and the second allows us to consider again a simpler metric with $g_{nn} = 1$.

LEMMA 3.14. — *Let g_1 and g_2 be two metrics which satisfy (3.2.4) and coincide on $\{x_n = 0\}$. Let f be a function satisfying Assumption 3.9. There exist constants $C_{12} \geq 1$ and $h_0 > 0$ such that the inequality,*

$$\mathcal{D}_{g_2, f, h}(\omega) \geq (1 - C_{12}h^{2/5})\mathcal{D}_{g_1, f, h}(\omega) - C_{12}h^{7/5} \|\omega\|_{\Lambda^p L^2, g_1}^2, \quad (3.2.8)$$

holds for $\omega \in \Lambda^p H_{0, \mathbf{n}}^1(\mathbb{R}_-^n)$, with $p \in \{0, \dots, n\}$ and $h \in (0, h_0)$, as soon as $\text{supp } \omega \subset \{x_n \geq -C_0 h^{2/5}\}$.

LEMMA 3.15. — *Let g_1 and g_2 be two conformal metrics (which satisfy (3.2.4)) in the sense:*

$$g_2 = e^{\varphi(x)} g_1.$$

Let f be a function satisfying Assumption 3.9. Then there exist constants $C_{12} \geq 1$ and $h_0 > 0$, such that the inequality,

$$\forall \omega \in \Lambda^p H_{0, \mathbf{n}}^1(\mathbb{R}_-^n), \quad \mathcal{D}_{g_2, f, h}(\omega) \geq C_{12}^{-1} \mathcal{D}_{g_1, f, h}(\omega) - C_{12}h^2 \|\omega\|_{\Lambda^p L^2, g_1}^2, \quad (3.2.9)$$

holds, for all $p \in \{0, \dots, n\}$ and all $h \in (0, h_0)$.

3.2.3 Small eigenvalues for the model half-space problem

Before giving the proof of Theorem 3.5, we state the main result for the model half-space problem which is similar to Proposition 3.7 and Proposition 3.8.

PROPOSITION 3.16. — *Assume that the metric \tilde{g}_0 satisfies (3.2.3) and (3.2.4) and let f be a Morse function satisfying Assumption 3.9 for some*

$p_0 \in \{0, \dots, n\}$. Then there exist constants $h_0 > 0$, $c_0 > 0$ and $c_1 > 0$, such that the self-adjoint operator $\Delta_{f,h}^N$ satisfies the following properties for $h \in (0, h_0]$:

i) For $p \in \{0, \dots, n\}$, the essential spectrum $\sigma_{\text{ess}}(\Delta_{f,h}^{N,(p)})$ is contained in $[c_1, +\infty)$.

ii) For $p \in \{0, \dots, n\}$, the range of $1_{[0,c_0h)}(\Delta_{f,h}^{N,(p)})$ has dimension

$$\begin{cases} \delta_{p,p_0} & \text{if } \partial_{x_n} f(0) = -\frac{1}{2}\partial_{x_n} f_+(0) < 0, \\ 0 & \text{if } \partial_{x_n} f(0) = -\frac{1}{2}\partial_{x_n} f_+(0) > 0. \end{cases}$$

iii) In the first case,

$$\text{Ran } 1_{[0,c_0h)}(\Delta_{f,h}^{N,(p_0)}) = \text{Ker } \Delta_{f,h}^{N,(p_0)} = \mathbb{C}\varphi^h,$$

where

$$\|\varphi^h - (e^{f_+(x_n)/2h})\psi_{p_0}^h\|_{\Lambda^p L^2} = \mathcal{O}(h^{1/10}),$$

and $\psi_{p_0}^h$ belongs to the kernel of a $(n-1)$ -dimensional Witten Laplacian $\Delta_{g',f_-/2,h}^{(p_0)}$ in a metric g' , which is conformal to $\tilde{g}_0 = \sum_{i,j=1}^{n-1} \tilde{g}_{ij}^0(x', 0)dx_i dx_j$ on \mathbb{R}^{n-1} .

iv) For any $\chi \in C_0^\infty(\overline{\mathbb{R}^n_-})$ such that $\chi = 1$ in a neighborhood of 0, there exists $C_\chi > 0$ such that the lower bounds

$$(1 - \chi)\Delta_{f,h}^{N,(p)}(1 - \chi) \geq C_\chi^{-1} [1 - \chi]^2, \quad 0 \leq p \leq n,$$

hold, for any $h \in (0, h_0/C_\chi)$, in the sense of quadratic forms on $\Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}^n_-)$.

Remark 3.17. — This proposition is an adaptation of Proposition 3.13 of [HeNi] in the case with normal boundary conditions: we have mainly replaced $f_+(x_n)$ by $-f_+(x_n)$ and $p_0 + 1$ by p_0 and the proof is similar.

We also emphasize the fact that, as in [HeNi], there is no need of any quadratic approximation to get these spectral results. The different approximations done are obtained by combination of the two previous lemmas with different suitable metrics.

Proof. — The clue of this result is an accurate lower bound for the quadratic form $\mathcal{D}_{\tilde{g}_0,f,h}(\eta)$, when evaluated for η such that $\text{supp } \eta \subset \{x_n \geq -C_0 h^{2/5}\}$. By Lemmas 3.14 and 3.15, one can find a metric g , which satisfies

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

(3.2.3) and (3.2.4), with $G(x) = G(x')$ independent of the x_n -coordinate, $g_{nn} = 1$ and a constant $C > 1$ such that

$$\mathcal{D}_{\tilde{g}_0, f, h}(\eta) \geq C^{-1} \mathcal{D}_{g, f, h}(\eta) - Ch^{7/5} \|\eta\|_{\Lambda L^2, g}^2. \quad (3.2.10)$$

Take two cut-off functions $\tilde{\chi}_i \in C^\infty(\mathbb{R})$, such that $\tilde{\chi}_1 \in C_0^\infty(\mathbb{R})$, $\tilde{\chi}_1 = 1$ in a neighborhood of 0 such that $\tilde{\chi}_1^2 + \tilde{\chi}_2^2 = 1$.

By the IMS localization formula (3.2.6), for any $\omega \in \Lambda H_{0, \mathbf{n}}^1(\mathbb{R}_-^n)$,

$$\begin{aligned} \mathcal{D}_{\tilde{g}_0, f, h}(\omega) &\geq \mathcal{D}_{\tilde{g}_0, f, h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega) + \mathcal{D}_{\tilde{g}_0, f, h}(\tilde{\chi}_2(h^{-2/5}x_n)\omega) \\ &\quad - Ch^{6/5} \|\omega\|_{\Lambda L^2, \tilde{g}_0}^2. \end{aligned}$$

By (2.6), since $|\nabla f(x)|^2 \geq C^{-1}$ on \mathbb{R}_-^n , the second term of the r.h.s. is bounded from below by a constant times $\|\tilde{\chi}_2(h^{-2/5}x_n)\omega\|_{\Lambda L^2, \tilde{g}_0}^2$ and we get:

$$\begin{aligned} \mathcal{D}_{\tilde{g}_0, f, h}(\omega) &\geq \mathcal{D}_{\tilde{g}_0, f, h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega) - Ch^{6/5} \|\tilde{\chi}_1(h^{-2/5}x_n)\omega\|_{\Lambda L^2, \tilde{g}_0}^2 \\ &\quad + \frac{C^{-1}}{2} \|\tilde{\chi}_2(h^{-2/5}x_n)\omega\|_{\Lambda L^2, \tilde{g}_0}^2. \end{aligned}$$

Finally after changing the constant $C \geq 1$, the inequality (3.2.10) yields

$$\begin{aligned} \mathcal{D}_{\tilde{g}_0, f, h}(\omega) &\geq C^{-1} \mathcal{D}_{g, f, h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega) - Ch^{6/5} \|\tilde{\chi}_1(h^{-2/5}x_n)\omega\|^2 \\ &\quad + C^{-1} \|\tilde{\chi}_2(h^{-2/5}x_n)\omega\|^2, \end{aligned} \quad (3.2.11)$$

where the L^2 -norms in the r.h.s. can be computed with the metric g or \tilde{g}_0 while possibly adapting the constant C , owing to Remark 3.12. Here and in the sequel, we omit the subscript $(\Lambda L^2, g)$ for L^2 -norms.

Now the problem is reduced to the analysis of $\mathcal{D}_{g, f, h}$ with the metric g . The product structure of the metric g allows an explicit analysis of the spectrum.

(a) The case $n = 1$.

We have $x = x_n \in \mathbb{R}_-$, $f(x) = -\frac{1}{2}f_+(x_n)$. Here the metric is $g = dx_n^2$. We keep the reference to the index n for the later application.

The spaces $\Lambda^0 H_{0, \mathbf{n}}^1(\mathbb{R}_-)$ and $\Lambda^1 H_{0, \mathbf{n}}^1(\mathbb{R}_-)$ are respectively $H^1(\mathbb{R}_-)$ and

$$\{\beta(x_n) dx_n, \beta \in H_0^1(\mathbb{R}_-)\}.$$

By identity (2.6), for any 1-form βdx_n with $\beta \in H_0^1(\mathbb{R}_-)$:

$$\mathcal{D}_{g,-f_+/2,h}(\beta dx_n) = h^2 \|\partial_{x_n} \beta\|^2 + \frac{1}{4} \|\partial_{x_n} f_+ \beta\|^2 - \frac{h}{2} \langle \partial_{x_n}^2 f_+(x_n) \beta \mid \beta \rangle . \quad (3.2.12)$$

From (3.2.12), we get:

$$\mathcal{D}_{g,-f_+/2,h}(\beta dx_n) \geq (C^{-2} - hC) \|\beta\|^2 ,$$

and deduce that there exist $c_1(\partial_{x_n} f_+, \partial_{x_n}^2 f_+) = c_1 > 0$ and $h_0 > 0$ such that, for all $h \in (0, h_0]$,

$$\Delta_{g,-f_+/2,h}^{N,(1)} \geq c_1 \text{Id} . \quad (3.2.13)$$

Again by identity (2.6), we have for any 0-form $\alpha \in H^1(\mathbb{R}_-)$:

$$\begin{aligned} \mathcal{D}_{g,-f_+/2,h}(\alpha) &= h^2 \|\partial_{x_n} \alpha\|^2 + \frac{1}{4} \|\partial_{x_n} f_+ \alpha\|^2 + \frac{h}{2} \langle \partial_{x_n}^2 f_+(x_n) \alpha \mid \alpha \rangle \\ &\quad - \frac{h}{2} \partial_{x_n} f_+(0) |\alpha(0)|^2 , \end{aligned} \quad (3.2.14)$$

and there are two subcases:

(a1) Subcase $\partial_{x_n} f_+(0) < 0$:

In this case, identity (3.2.14) implies:

$$\forall \alpha \in \Lambda^0 H_{0,\mathbf{n}}^1, \quad \mathcal{D}_{g,-f_+/2,h}(\alpha) \geq (C^{-2} - hC) \|\alpha\|^2 ,$$

which provides the existence of $c_1(\partial_{x_n} f_+, \partial_{x_n}^2 f_+) = c_1 > 0$ and $h_0 > 0$ such that:

$$\Delta_{-f_+/2,h}^{N,(0)} \geq c_1 \text{Id} , \quad \forall h \in (0, h_0] . \quad (3.2.15)$$

(a2) Subcase $\partial_{x_n} f_+(0) > 0$:

If $\Delta_{-f_+/2,h}^{N,(0)}(\alpha) = \lambda_h \alpha$, with $\lambda_h < c_1$, we have by Proposition 3.10:

$$\Delta_{-f_+/2,h}^{N,(1)}(d_{-f_+/2,h} \alpha) = \lambda_h d_{-f_+/2,h} \alpha .$$

Now suppose for a while that $d_{-f_+/2,h} \alpha$ is not 0. Then it is an 1-eigenform of $\Delta_{-f_+/2,h}^{N,(1)}$ for $\lambda_h < c_1$, but this is in contradiction with (3.2.13). As a consequence it is zero and we get by (2.1.6):

$$d_{-f_+/2,h} \alpha = h \partial_{x_n} \alpha - \frac{1}{2} (\partial_{x_n} f_+) \alpha = 0 .$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Hence:

$$\alpha(x_n) = C e^{f_+(x_n)/2h}.$$

Since the 0-form $e^{f_+(x_n)/2h}$ belongs to $\text{Ker}(\Delta_{-f_+/2,h}^{N,(0)})$, $\lambda_h = 0$ is the only eigenvalue (with multiplicity one) of $\Delta_{-f_+/2,h}^{N,(0)}$ smaller than c_1 .

(b) The case $n > 1$.

First note that any $\omega \in \Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}_-^n)$ is a sum

$$\omega = \sum_{\#I=p-1} \alpha_I(x) dx'^I \wedge dx_n + \sum_{\#J=p} \beta_J(x) dx'^J =: \alpha \wedge dx_n + \beta,$$

with $\alpha_I, \beta_J \in H^1(\mathbb{R}_-^n)$, $\alpha_I(x', 0) = 0$, while $dx'^I = dx'_{i_1} \wedge \dots \wedge dx'_{i_{\#I}}$, $I = \{i_1 < \dots < i_{\#I}\} \subset \{1, \dots, n-1\}$ and $J = \{j_1 < \dots < j_{\#J}\} \subset \{1, \dots, n-1\}$.

If in addition $\omega \in \Lambda^p H^2(\mathbb{R}_-^n)$, the condition $\mathbf{nd}\omega = 0$ reads, with the metric g , $\partial_{x_n} \beta_J(x', 0) = 0$.

Secondly, owing to the Weitzenböck formula with the product metric g , the Riemannian connection, the Riemann tensor, $R_{(4)}$ (see [CyFrKiSi] pp. 266-267 for the definition of this operator) and therefore the Hodge Laplacian split like direct sums:

$$\begin{aligned} \nabla_X Y &= \nabla_{X_n}^n Y_n + \nabla_{X'}^n Y', \\ \text{Riem}(x, y, z, t) &= \text{Riem}^n(x_n, y_n, z_n, t_n) + \text{Riem}'(x', y', z', t'), \\ R_{(4)} &= \sum_{ijkl} \text{Riem}_{ijkl}(dx_i \wedge) \circ \mathbf{i}_{\nabla x_j} \circ (dx_k \wedge) \circ \mathbf{i}_{\nabla x_\ell} = R_{(4)}^n + R_{(4)}', \\ (d + d^*)^2 &= (d_{x_n} + d_{x_n}^*)^2 + (d_{x'} + d_{x'}^*)^2. \end{aligned}$$

We also refer the reader to [GaHuLa] (p. 110 and p. 70) for details and more general statements.

Thirdly, the decomposition $f(x) = -\frac{1}{2}f_+(x_n) + \frac{1}{2}f_-(x')$ with the product metric g gives

$$\begin{aligned} |\nabla f|^2 &= |\nabla_{x_n} f|^2 + |\nabla_{x'} f|^2 \\ \mathcal{L}_{\nabla f} + \mathcal{L}_{\nabla f}^* &= -\frac{1}{2} \left(\mathcal{L}_{\nabla f_+} + \mathcal{L}_{\nabla f_+}^* \right) + \frac{1}{2} \left(\mathcal{L}_{\nabla f_-} + \mathcal{L}_{\nabla f_-}^* \right). \end{aligned}$$

For $\omega = \alpha \wedge dx_n + \beta \in D(\Delta_{f,h}^N)$ (with the product metric g), we have

$$\mathcal{D}_{g,f,h}(\omega) = \langle \omega | \Delta_{f,h} \omega \rangle = \left\langle \omega | \Delta_{-f_+/2,h}^n \omega \right\rangle + \left\langle \omega | \Delta'_{-f_-/2,h} \omega \right\rangle.$$

Since the two operators $\Delta_{-f_+/2,h}^n$ (acting only in the variable x_n) and $\Delta'_{f_-/2,h}$ (acting only in the variable x') preserve the partial degree in dx_n , we get

$$\begin{aligned} \mathcal{D}_{g,f,h}(\omega) &= \left\langle \alpha \wedge dx_n \mid \Delta_{-f_+/2,h}^n(\alpha \wedge dx_n) \right\rangle + \left\langle \beta \mid \Delta_{-f_+/2,h}^n \beta \right\rangle \\ &\quad + \left\langle \alpha \wedge dx_n \mid \Delta'_{f_-/2,h}(\alpha \wedge dx_n) \right\rangle + \left\langle \beta \mid \Delta'_{f_-/2,h} \beta \right\rangle \end{aligned} \tag{3.2.16}$$

Hence the variables (x', x_n) can be separated. The equivalence between the norms $\left\| \sum_J \gamma_J(x') dx'^J \right\|$ and $\sum_J \|\gamma_J(x')\|$ on $\Lambda^p T^* \mathbb{R}_-^n$, where $J = \{j_1 < \dots < j_{\#J}\} \subset \{1, \dots, n-1\}$, leads to ²⁾:

$$\begin{aligned} \mathcal{D}_{g,f,h}(\omega) &\geq \\ \frac{1}{c} \int_{\mathbb{R}^{n-1}} &\left[\sum_{\#I=p-1} \mathcal{D}_{-f_+/2,h}^n(\alpha_I(x', \cdot) dx_n) + \sum_{\#J=p} \mathcal{D}_{-f_+/2,h}^n(\beta_J(x', \cdot)) \right] d\lambda(x') \\ &\quad + \int_{-\infty}^0 \mathcal{D}'_{f_-/2,h}(\alpha(\cdot, x_n)) + \mathcal{D}'_{f_-/2,h}(\beta(\cdot, x_n)) dx_n, \end{aligned} \tag{3.2.17}$$

where we used the notations $\mathcal{D}'_{f_-/2,h}$ for the quadratic form of the Witten Laplacian on \mathbb{R}^{n-1} and $\mathcal{D}_{-f_+/2,h}^n$ for the quadratic form of the 1-dimensional Witten Laplacian on \mathbb{R}_- with boundary conditions. The measure $d\lambda(x')$ simply equals $(\det G(x'))^{1/2} dx'$. The absence of $\alpha - \beta$ cross product term is due to (3.2.16).

Again there are two subcases.

(b1) Subcase $\partial_{x_n} f_+(0) < 0$:

The analysis of the one dimensional problem implies the existence of $c_1 > 0$ independent of x' such that:

$$\begin{aligned} \mathcal{D}_{-f_+/2,h}^n(\alpha_I(x', \cdot) dx_n) &\geq c_1 \|\alpha_I(x', \cdot)\|^2 \\ \text{and} \\ \mathcal{D}_{-f_+/2,h}^n(\beta_J(x', \cdot)) &\geq c_1 \|\beta_J(x', \cdot)\|^2. \end{aligned}$$

Hence there exists $c_2 > 0$ such that:

$$\forall \omega \in \Lambda^p H_{0,\mathbf{n}}^1, \quad \mathcal{D}_{g,f,h}(\omega) \geq c_2 \|\omega\|^2$$

(²⁾ In [HeNi], at this level of the proof, one should read “ $\mathcal{D}_{\bar{g},f,h}(\omega) \geq \frac{1}{c} \int_{\mathbb{R}^{n-1}} \dots$ ” instead of “ $\mathcal{D}_{\bar{g},f,h}(\omega)$ equals $\int_{\mathbb{R}^{n-1}} \dots$ ” accordingly to 2.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

and

$$\Delta_{f,h}^{N,(p)} \geq c_2 \text{Id} , \quad \forall p \in \{0, \dots, n\} .$$

(b2) Subcase $\partial_{x_n} f_+(0) > 0$:

Then there exists $c_1 > 0$ such that

$$\begin{aligned} \mathcal{D}_{g,f,h}(\omega) &\geq \frac{1}{c} \int_{\mathbb{R}^{n-1}} \sum_{\#J=p} \mathcal{D}_{-f_+/2,h}^n(\beta_J(x', \cdot)) d\lambda(x') \\ &\quad + \int_{-\infty}^0 \mathcal{D}'_{f_-/2,h}(\beta(\cdot, x_n)) + c_1 \|\alpha\|^2 . \end{aligned} \tag{3.2.18}$$

If ω is a p -form with $p \neq p_0$ ($\deg \beta = \deg \omega$), the lower bound

$$\mathcal{D}'_{f_-/2,h}(\beta) \geq C_1^{-1} h \|\beta\|^2 ,$$

which was given in Proposition 3.8, yields:

$$\mathcal{D}_{g,f,h}(\omega) \geq C^{-1} h \|\omega\|^2 ,$$

while the equality $\mathcal{D}_{g,f,h}(\omega) = 0$ implies that $p = p_0$ and that $\omega = c (e^{f_+(x_n)/2h}) \psi_{p_0}^h$, where ψ_{p_0} belongs to the kernel of the $(n-1)$ -dimensional Witten Laplacian associated with the metric

$$g' = \sum_{i,j=1}^{n-1} g_{i,j}(x', 0) dx_i dx_j .$$

We have now all the ingredients to check every statement for the metric \tilde{g}_0 .

We focus on the subcase $\partial_{x_n} f_+(0) > 0$, which covers all possibilities.

Statements i) and iv)

Statement i) is a consequence of iv) together with Persson's Lemma in [Per]. It is sufficient to check that, for all $R > 0$, there exists $c_R > 0$, such that, for all $\omega \in \Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}_-^n)$ supported in $\{\min(|x'|, |x_n|) > R\}$, one has

$$\mathcal{D}_{\tilde{g}_0,f,h}(\omega) \geq c_R \|\omega\|^2 .$$

The inequalities (3.2.11) and (3.2.18), together with the estimate

$$\mathcal{D}'_{f_-/2,h}(\beta(\cdot, x_n)) \geq c'_R \|\beta(\cdot, x_n)\|^2 \quad \text{if } \text{supp } \omega \subset \{|x'| > R\} ,$$

provided by Proposition 3.8-iii), yield the result.

Statements ii) and iii)

If $p \neq p_0$ the inequalities (3.2.11), (3.2.18) and the inequality

$$\mathcal{D}'_{f_{-/2},h}(\beta(\cdot, x_n)) \geq C^{-1}h \|\beta(\cdot, x_n)\|^2 ,$$

imply

$$\mathcal{D}_{\tilde{g}_0,f,h}(\omega) \geq c_0h \|\omega\|^2 ,$$

and

$$\Delta_{f,h}^{N,(p)} \geq c_0h \text{Id} . \tag{3.2.19}$$

If $p = p_0$, by Proposition 3.10, the only possibility for $\lambda_h \in [0, c_0h)$ to be an eigenvalue of $\Delta_{f,h}^{N,(p_0)}$ is $\lambda_h = 0$.

Assume indeed $\Delta_{f,h}^{N,(p_0)}u_h = \lambda_h u_h$ with $\lambda_h \in [0, c_0h)$ and $\|u_h\| = 1$.

By Proposition 3.10 and (3.2.19), $d_{f,h}^{(p_0)}u_h = d_{f,h}^{(p_0-1),*}u_h = 0$. Thus:

$$\lambda_h = \left\langle \Delta_{f,h}^{N,(p_0)}u_h \mid u_h \right\rangle = \mathcal{D}_{\tilde{g}_0,f,h}(u_h) = 0 .$$

When the metric is g , the corresponding spectral subspace is one dimensional and equals $\mathbb{C} \left(e^{f_+(x_n)/2h} \right) \psi_{p_0}^h$.

For the metric \tilde{g}_0 , equation $\Delta_{\tilde{g}_0,f,h}^{N,(p_0)}\omega = 0$ with $\|\omega\| = 1$ (which implies $\mathcal{D}_{\tilde{g}_0,f,h}(\omega) = 0$) and inequality (3.2.11) lead to:

$$C^2h^{6/5} \left\| \tilde{\chi}_1(h^{-2/5}x_n)\omega \right\|^2 \geq \mathcal{D}_{g,f,h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega) + \left\| \tilde{\chi}_2(h^{-2/5}x_n)\omega \right\|^2 .$$

Without the last term, Lemma 2.11 implies:

$$\text{dist}_{L^2}(\tilde{\chi}_1(h^{-2/5}x_n)\omega, \mathbb{C} \left(e^{f_+(x_n)/2h} \right) \psi_{p_0}^h) \leq Ch^{1/10} .$$

The upper bound of the last term,

$$\left\| \tilde{\chi}_2(h^{-2/5}x_n)\omega \right\|^2 \leq C^2h^{6/5} ,$$

implies:

$$\text{dist}_{L^2}(\omega, \mathbb{C} \left(e^{f_+(x_n)/2h} \right) \psi_{p_0}^h) = \mathcal{O}(h^{1/10}) .$$

It remains to check that $\text{Ker } \Delta_{f,h}^{N,(p_0)}$ is not reduced to $\{0\}$. The statements of Lemma 3.14 and Lemma 3.15 are symmetric with respect to the choice

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

of the metric. Hence the reverse inequality of (3.2.11) (with exchange of g and \tilde{g}_0),

$$\begin{aligned} \mathcal{D}_{g,f,h}(\omega) \geq C^{-1} \mathcal{D}_{\tilde{g}_0,f,h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega) - Ch^{6/5} \left\| \tilde{\chi}_1(h^{-2/5}x_n)\omega \right\|^2 \\ + C^{-1} \left\| \tilde{\chi}_2(h^{-2/5}x_n)\omega \right\|^2, \quad (3.2.20) \end{aligned}$$

also holds for any $\omega \in \Lambda H_{0,\mathbf{n}}^1(\mathbb{R}^n)$. We apply it with $\omega = (e^{f_+(x_n)/2h}) \psi_{p_0}^h$ and this leads to:

$$\mathcal{D}_{\tilde{g}_0,f,h}(\tilde{\chi}_1(h^{-2/5}x_n)\omega^h) \leq Ch^{6/5} \left\| \tilde{\chi}_1(h^{-2/5}x_n)\omega \right\|^2.$$

The Min-Max principle then says that $\Delta_{f,h}^{N,(p_0)}$ admits an eigenvalue smaller than $Ch^{6/5}$. It has to be 0 due to the above argument. \square

3.3. Proof of Theorem 3.5

We end here the proof of Theorem 3.5 by introducing, after a partition of unity, convenient coordinates which allow the comparison with the model half-space problem.

That proof is almost the same as the proof of the corresponding theorem in [HeNi], but we recall it for completeness.

Proof of Theorem 3.5. — Let $\{U_k, 1 \leq k \leq K\}$ denote the critical points of f and $f|_\Omega$. Consider a partition of unity of $\bar{\Omega}$, $\sum_{k=1}^N \chi_k^2 = 1$, such that the $C_0^\infty(\bar{\Omega})$ function χ_k identically equals 1 in a neighborhood of U_k when $1 \leq k \leq K$. A refinement of this partition of unity will be specified later by the local construction of adapted coordinates.

We recall that the operator $\Delta_{f,h}^N$ is the Friedrichs extension associated with the quadratic form:

$$\mathcal{D}_{g_0,f,h}(\omega) = \|d_{f,h}\omega\|_{\Lambda L^2,g_0}^2 + \left\| d_{f,h}^{*,g_0}\omega \right\|_{\Lambda L^2,g_0}^2,$$

on $\Lambda H_{0,\mathbf{n}}^1(\Omega)$. The IMS localization formula 3.2.6 gives, for any $\omega \in \Lambda H_{0,\mathbf{n}}^1$,

$$\mathcal{D}_{g_0,f,h}(\omega) = \sum_{k=1}^N \mathcal{D}_{g_0,f,h}(\chi_k\omega) - h^2 \left\| |\nabla \chi_k| \omega \right\|_{\Lambda L^2,g_0}^2.$$

If $\text{supp } \chi_k$ **does not meet the boundary**, the term $\mathcal{D}_{g_0,f,h}(\chi_k\omega)$ behaves like in the boundaryless case (see [HeKlNi] for details):

- If $k > K$, then we have

$$\forall \omega \in \Lambda H^1, \quad \mathcal{D}_{g_0, f, h}(\chi_k \omega) \geq C^{-1} \|\chi_k \omega\|_{\Lambda L^2, g_0}^2.$$

- If $k \leq K$ and U_k is a critical point of f with index $p_k \neq p$, then

$$\forall \omega \in \Lambda H^1, \quad \mathcal{D}_{g_0, f, h}(\chi_k \omega) \geq C^{-1} h \|\chi_k \omega\|_{\Lambda L^2, g_0}^2.$$

- If $k \leq K$ and U_k is a critical point of f with index $p_k = p$, then there exists a fixed 1-dimensional space $F_k^{(p)}$ (determined by Hess $f(U_k)$) such that,

$$\forall \omega \in \Lambda H^1, \quad \mathcal{D}_{g_0, f, h}(\chi_k \omega) \leq C^{-1} h^{6/5} \|\chi_k \omega\|_{\Lambda^p L^2, g_0}^2$$

implies

$$\forall \omega \in \Lambda H^1, \quad \text{dist}(\chi_k \omega, F_k^{(p)}) \leq C h^{1/10} \|\omega\|_{\Lambda^p L^2, g_0}.$$

Again like in the proof of Proposition 3.16-iii), this last statement refers to Lemma 2.11 at the level of quadratic forms.

Consider now the case when $\text{supp } \chi_k \cap \partial\Omega \neq \emptyset$, with the support of χ_k centered around a point $U_0 \in \partial\Omega$. There are two cases: U_0 is a critical point of $f|_{\partial\Omega}$ with $\frac{\partial f}{\partial n}(U_0) < 0$ which is equivalent to $-\frac{\partial f}{\partial n}(U_0) = |\nabla f(U_0)|$ or U_0 is not a critical points of $f|_{\partial\Omega}$ with $\frac{\partial f}{\partial n} < 0$ which is equivalent to $(-\frac{\partial f}{\partial n})(U_0) < |\nabla f(U_0)|$. Indeed, U_0 is either a critical point of $f|_{\partial\Omega}$ with $\frac{\partial f}{\partial n}(U_0) > 0$, i.e. $\frac{\partial f}{\partial n}(U_0) = |\nabla f(U_0)|$ or U_0 is not a critical point of $f|_{\partial\Omega}$, i.e. $\left| \frac{\partial f}{\partial n}(U_0) \right| < |\nabla f(U_0)|$.

Case 1) $(-\frac{\partial f}{\partial n})(U_0) < |\nabla f(U_0)|$.

Then the cut-off χ_k is chosen so that, in a neighborhood \mathcal{V} of $\text{supp } \chi_k$,

$$\forall x \in \mathcal{V} \cap \partial\Omega, \quad \left(-\frac{\partial f}{\partial n}\right)(x) < (1 - \delta) |\nabla f(x)|,$$

for some $\delta > 0$. Locally it is possible to construct a function \hat{f} such that $-\partial_n \hat{f} = |\nabla \hat{f}|$ in $\mathcal{V} \cap \partial\Omega$ and $|\nabla \hat{f}| = |\nabla f|$ in \mathcal{V} . By setting $\tilde{\omega} = \chi_k \omega$ for $\omega \in \Lambda H_{0, \mathbf{n}}^1$, the Green formula (2.6) and the inequality $\mathcal{D}_{g_0, \hat{f}, h}(\tilde{\omega}) \geq 0$ imply ($\mathcal{L}_{\nabla f} + \mathcal{L}_{\nabla \hat{f}}^*$ being a tensor)

$$\begin{aligned} & -h \int_{\partial\Omega} \langle \tilde{\omega} | \tilde{\omega} \rangle_{\Lambda^p T_\sigma^* \Omega} \left(\frac{\partial \hat{f}}{\partial n} \right) (\sigma) d\sigma \leq -(1 - \delta) h \int_{\partial\Omega} \langle \tilde{\omega} | \tilde{\omega} \rangle_{\Lambda^p T_\sigma^* \Omega} \left(\frac{\partial \hat{f}}{\partial n} \right) (\sigma) d\sigma \\ & \leq (1 - \delta) \left[h^2 \|\tilde{d}\tilde{\omega}\|_{\Lambda^{p+1} L^2}^2 + h^2 \|d^* \tilde{\omega}\|_{\Lambda^{p-1} L^2}^2 + \| |\nabla f| \tilde{\omega} \|_{\Lambda^p L^2}^2 + C_1 h \|\tilde{\omega}\|_{\Lambda^p L^2}^2 \right]. \end{aligned}$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

- If $k > K$,

$$\begin{aligned} \forall \omega \in \Lambda H_{0,\mathbf{n}}^1, \quad \mathcal{D}_{g_0,f,h}(\chi_k \omega) = \mathcal{D}_{g_0,f,h}(\tilde{\omega}) &\geq \frac{\delta}{2} \|\nabla f| \tilde{\omega}\|_{\Lambda^p L^2}^2 \\ &\geq C_V^{-1} \|\chi_k \omega\|_{\Lambda^p L^2}^2. \end{aligned}$$

Case 2) $-\frac{\partial f}{\partial n}(U_0) = |\nabla f(U_0)|$.

In this case we will conclude by applying Proposition 3.16. We recall that $U_0 \in \partial\Omega$ is a critical point of $f|_{\partial\Omega}$ with $\frac{\partial f}{\partial n}(U_0) < 0$ and with index p_0 . Around U_0 , we introduce adapted local coordinates, denoted by $\bar{x} = (\bar{x}', \bar{x}_n)$. This coordinate system is provided by Lemma 3.18 below, applied with $f_1 = f$ and $\alpha = f|_{\partial\Omega \cap \mathcal{V}_0}$. Then the function Φ_+ of Lemma 3.18 is nothing but f and has the form $f(\bar{x}) = -\bar{x}_n + \frac{1}{2}f_-(\bar{x}')$. Moreover, $\bar{\Omega}$ corresponds locally to $\{\bar{x}_n \leq 0\}$.

In order to apply Proposition 3.16, it remains to check that the function f can be extended to \mathbb{R}_-^n , so that it satisfies Assumption 3.9 where U_0 is a critical point of $f|_{\partial\Omega}$.

We recall that we have not specified the choice of \bar{x}' in the boundary. The function $f|_{\partial\Omega \cap \mathcal{V}_0}$ being a Morse function, we can choose in a small neighborhood $\mathcal{V}_0' \subset \partial\Omega$ of $U_0 = (0, \dots, 0)$ Morse coordinates $\bar{x}' = (\bar{x}_1, \dots, \bar{x}_{n-1})$ for f_- which are normal at U_0 for the metric $\sum_{i,j < n} g_{ij}(\bar{x}', 0) d\bar{x}_i d\bar{x}_j$. With these coordinates, f has the form, in a small neighborhood \mathcal{V}_0'' of 0:

$$f(\bar{x}) = -\bar{x}_n + \sum_{j=1}^{n-1} \lambda_j \bar{x}_j^2 + f(U_0). \quad (3.3.1)$$

We choose χ_k such that $\text{supp } \chi_k \subset \mathcal{V}_0''$.

Choosing a cut-off $\chi^{n-1} \in C_0^\infty(\mathbb{R}^{n-1})$, $\chi^{n-1} = 1$ near $\text{supp } \chi_k \cap \partial\Omega$, f is extended to \mathbb{R}_-^n by:

$$\tilde{f}(\bar{x}) = -\bar{x}_n + \left[\chi^{n-1}(\bar{x}') + \frac{1 - \chi^{n-1}(\bar{x}')}{|\bar{x}'|} \right] \left[\sum_{j=1}^{n-1} \lambda_j \bar{x}_j^2 \right] + f(U_0). \quad (3.3.2)$$

Moreover, choosing another cut-off $\chi^n \in C_0^\infty(\overline{\mathbb{R}_-^n})$, $\chi^n = 1$ near $\text{supp } \chi_k$, we extend g_0 to \mathbb{R}_-^n by:

$$\tilde{g} = \chi^n g_0 + (1 - \chi^n) g_e, \quad (3.3.3)$$

where g_e is the Euclidian metric on \mathbb{R}_-^n .

With these coordinates, the quantity $\mathcal{D}_{\bar{g}, \bar{f}, h}(\chi_k \omega) = \mathcal{D}_{g_0, f, h}(\chi_k \omega)$ attains the form discussed in Proposition 3.16.

We can now discuss the lower bound of $\mathcal{D}_{\bar{g}, \bar{f}, h}(\chi_k \omega)$, depending on the localization by the cut-off χ_k , such that $\text{supp } \chi_k \cap \partial\Omega \neq \emptyset$.

- If $k \leq K$, the origin of the coordinate system is $U_0 = U_k$. If U_k is not a critical point of $f|_{\partial\Omega}$ with index $p_k = p$ and $\frac{\partial f}{\partial n}(U_k) < 0$, then

$$\forall \omega \in \Lambda^p H_{0, \mathbf{n}}^1, \quad \mathcal{D}_{\bar{g}, \bar{f}, h}(\chi_k \omega) \geq C^{-1} h \|\chi_k \omega\|_{\Lambda^p L^2, g}^2.$$

- If $k \leq K$ and U_k is a critical point of $f|_{\partial\Omega}$ with index $p_k = p$ and $\frac{\partial f}{\partial n}(U_k) < 0$, then according to Proposition 3.16-iii) there exists a fixed 1-dimensional space $F_k^{(p)}$ such that the inequality,

$$\forall \omega \in \Lambda^p H_{0, \mathbf{n}}^1, \quad \mathcal{D}_{\bar{g}, \bar{f}, h}(\chi_k \omega) \leq C^{-1} h^{6/5} \|\chi_k \omega\|_{\Lambda^p L^2, g}^2$$

implies:

$$\text{dist}(\chi_k \omega, F_k^{(p)}) \leq C h^{1/10} \|\chi_k \omega\|_{\Lambda^p L^2, g}.$$

We now introduce the set A_p of indices k , $1 \leq k \leq K$, such that

- either U_k is a critical point of f with index p ,
- or U_k is a critical point of $f|_{\partial\Omega}$ with index p such that $\frac{\partial f}{\partial n}(U_k) < 0$.

For $\omega \in \Lambda^p H_{0, \mathbf{n}}^1(\Omega)$ with $\|\omega\|_{\Lambda^p L^2, g} = 1$, we get

$$\left(\mathcal{D}_{g_0, f, h}(\omega) \leq C^{-1} h^{6/5} \right) \Rightarrow \left(\text{dist} \left(\omega, \sum_{k \in A_p} F_k^{(p)} \right) \leq C h^{1/10} \right).$$

Hence the dimension of the spectral subspace,

$$F^{(p)} = \text{Ran} 1_{[0, h^{3/2})}(\Delta_{f, h}^{N, (p)}) \subset \text{Ran} 1_{[0, ch^{6/5})}(\Delta_{f, h}^{N, (p)}),$$

is at most $\#A_p = m_p^{\bar{\Omega}}$.

We next verify that $\dim F^{(p)} \geq \#A_p = m_p^{\bar{\Omega}}$. According to the Min-Max principle, it suffices to find an orthonormal set of p -forms $\omega_k^h \in \Lambda^p H_{0, \mathbf{n}}^1(\Omega)$, $k \in A_p$, such that

$$\mathcal{D}_{g_0, f, h}(\omega_k^h) = o(h^{3/2}).$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Indeed it is enough to take a truncated element of the kernel of the local model for $\Delta_{f,h}^{N,(p)}$ around U_k , $k \in A_p$. We give the details for the case $U_k \in \partial\Omega$.

Take two cut-off $\chi_{1,k} \in C_0^\infty(\overline{\mathbb{R}^n_-})$, $\chi_{1,k} = 1$ near 0 (with $\text{supp } \chi_{1,k} \subset \text{supp } \chi_k$) and $\chi_{2,k}$ such that $\chi_{1,k}^2 + \chi_{2,k}^2 = 1$. With the same coordinate system as above, we write on \mathbb{R}^n_- , using the IMS localization formula (3.2.6) and Proposition 3.16-iv),

$$\mathcal{D}_{\tilde{g}_k, \tilde{f}_k, h}(\omega) \geq \mathcal{D}_{\tilde{g}_k, \tilde{f}_k, h}(\chi_{1,k}\omega) + C^{-1} \|\chi_{2,k}\omega\|^2 - Ch^2 \sum_{i=1,2} \|\|\nabla \chi_{i,k}|\omega\|^2,$$

where \tilde{g}_k and \tilde{f}_k are defined on \mathbb{R}^n_- according to the previous construction and coincide with g_0 and f in a neighborhood of $\text{supp } \chi_k$. According to Proposition 3.16, there exists $\eta_k^h \in \Lambda^p H_{0,\mathbf{n}}^1(\mathbb{R}^n_-)$ in the domain of the associated Witten Laplacian, such that $\mathcal{D}_{\tilde{g}_k, \tilde{f}_k, h}(\eta_k^h) = 0$. By taking $\omega_k^h = \|\|\chi_{1,k} \eta_k^h\|^{-1} \chi_{1,k} \eta_k^h$, we obtain the existence of $h_0 > 0$, C' and C'' such that, for $h \in (0, h_0]$:

$$\|\|\chi_{2,k} \eta_k^h\|^2 \leq C'h^2 \|\|\eta_k^h\|^2,$$

and, consequently,

$$\mathcal{D}_{g_0, f, h}(\omega_k^h) \leq C'h^2 \frac{\|\|\eta_k^h\|^2}{\|\|\chi_{1,k} \eta_k^h\|^2} \leq C''h^2.$$

□

The following lemma, which provides in different situations the suitable coordinate systems, simply makes use of the standard solution to Hamilton-Jacobi equations in the non characteristic case. It is proved in [Lep2] (this is indeed also Lemma 3.4 of [Lep2]).

LEMMA 3.18. — 1) Let be $f_1 \in C^\infty(\overline{\Omega}, \mathbb{R})$ and $U_0 \in \partial\Omega$ a critical point of $f_1|_{\partial\Omega}$ with $\frac{\partial f_1}{\partial n}(U_0) \neq 0$.

Assume furthermore $\alpha \in C^\infty(\partial\Omega, \mathbb{R})$ be a local solution to $|\nabla_T \alpha|^2 = |\nabla_T f_1|^2$ around U_0 .

Then there exists a neighborhood \mathcal{V}_0 of U_0 in $\overline{\Omega}$ such that the eikonal equation:

$$|\nabla \Phi_\pm|^2 = |\nabla f_1|^2$$

(on the boundary, it means $|\partial_n \Phi_\pm|^2 + |\nabla_T \Phi_\pm|^2 = |\partial_n f_1|^2 + |\nabla_T f_1|^2$)

with the boundary conditions

$$\Phi_{\pm}|_{\partial\Omega\cap\nu_0} = \alpha \quad , \quad \partial_n\Phi_{\pm}|_{\partial\Omega\cap\nu_0} = \pm \frac{\partial f_1}{\partial n}|_{\partial\Omega\cap\nu_0}$$

admits a unique local smooth real-valued solution.

2) There exists local coordinates $(x_1, \dots, x_n) = (x', x_n)$ in a neighborhood of U_0 in $\overline{\Omega}$ with $(x', x_n)(U_0) = 0$ where the function Φ_{\pm} and the metric g_0 have the form:

$$\Phi_{\pm} = \mp x_n + \alpha(x') \quad \text{and} \quad g_0 = g_{nn}(x) dx_n^2 + \sum_{i,j=1}^{n-1} g_{ij}(x) dx_i dx_j .$$

Moreover, the boundary $\partial\Omega$ is locally defined by $\{x_n = 0\}$ and Ω corresponds to $\left\{ \operatorname{sgn}\left(\frac{\partial f_1}{\partial n}(U_0)\right) x_n > 0 \right\}$.

Remark 3.19. — Lemma 3.18 will be used with various functions f_1 and α and will provide several coordinate systems:

- We have already introduced the coordinate system $\bar{x} = (\bar{x}, \bar{x}_n)$ associated with $f_1 = f$ and $\alpha = f|_{\partial\Omega}$.
- The coordinate system denoted simply by $x = (x', x_n)$ will be associated with $f_1 = f$ and $\alpha = \varphi$, where φ is the Agmon distance along the boundary. This system will be used to give the simple form $\Phi = \Phi_+ = -x_n + \varphi(x')$ to the Agmon distance Φ , solving $|\nabla\Phi|^2 = |\nabla f|^2$ with the boundary condition $\partial_n\Phi = \partial_n f$. Agmon distances are specified in Section 4 below.
- Finally the coordinate system $\tilde{x} = (\tilde{x}', \tilde{x}_n)$ will be associated with $f_1 = (f + \Phi)$ and $\alpha = f|_{\partial\Omega} + \varphi$ and will be used in the final application of the Laplace method.

4. Accurate WKB analysis near the boundary for $\Delta_{f,h}^{(1)}$

4.1. Introduction

We work here under Assumption 3.1. Like in [HeNi], we have shown that for $0 \leq p < n$, some quasimodes of $\Delta_{f,h}^{N,(p)}$ being near the spectral subspace in $1_{[0, h^{\frac{3}{2}})}(\Delta_{f,h}^{N,(p)})$ are localized near the boundary $\partial\Omega$ and more precisely near critical points of $f|_{\partial\Omega}$ with index p such that $\frac{\partial f}{\partial n} < 0$. In the boundaryless case ([HeKlNi]) and in the case with tangential Dirichlet

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

boundary conditions ([HeNi]), the WKB analysis done in [HeSj4] and in [HeNi] says that the small eigenvalues are of order $\mathcal{O}(e^{-C/h})$ and provides an accurate approximate basis of $\text{Ran}1_{[0,h^{3/2})}(\Delta_{f,h}^{(p)})$.

In order to get a similar result, we need an accurate WKB analysis at the boundary, and like in [HeNi], we restrict our attention on the case $p = 1$ because our motivation is to analyze the Witten Laplacian on 0-forms.

For an accurate comparison between eigenvectors and WKB quasimodes near a critical point U_1 of $f|_{\partial\Omega}$ with index 1 and $\frac{\partial f}{\partial n}(U_1) < 0$, we introduce another self-adjoint realization of $\Delta_{f,h}^{(1)}$ in a neighborhood $\Omega_{U_1,\rho}$ with mixed boundary conditions: Neumann boundary conditions on $\partial\Omega_{U_1,\rho} \cap \partial\Omega$ and full Dirichlet boundary conditions on $\partial\Omega_{U_1,\rho} \setminus \partial\Omega$.

4.2. Local WKB construction

Take U_1 a critical point of $f|_{\partial\Omega}$ with index 1 such that $\frac{\partial f}{\partial n}(U_1) < 0$. According to Definition 3.1 and Proposition 3.2 of [Lep2], there exists a local coordinate system $(\underline{x}_1, \dots, \underline{x}_n) = (\underline{x}', \underline{x}_n)$ which satisfies the following properties:

- i) $d\underline{x}_1, \dots, d\underline{x}_n$ is an orthonormal basis of $T_{U_1}^*(\overline{\Omega})$ positively oriented.
- ii) The boundary $\partial\Omega$ corresponds locally to $\underline{x}_n = 0$ and the interior Ω to $\underline{x}_n < 0$.
- iii) $\frac{\partial}{\partial \underline{x}_n}|_{\partial\Omega} = \vec{n}$, the outgoing normal at the boundary. Moreover, $\frac{\partial}{\partial \underline{x}_n}$ is unitary and normal to $\{\underline{x}_n = \text{Constant}\}$.

Moreover, the choice of the coordinates $(\underline{x}_1, \dots, \underline{x}_{n-1})$ (centered at U_1 such that $d\underline{x}_1, \dots, d\underline{x}_n$ is an orthonormal basis of $T_{U_1}^*(\overline{\Omega})$) in the boundary is arbitrary.

Let φ be the Agmon distance to U_1 on the boundary (i.e. associated with the metric $|\nabla_{\underline{x}'} f(\underline{x}', 0)|^2 d\underline{x}'^2$). Recall that φ satisfies

$$|\nabla_T f|^2 = |\nabla \varphi|^2$$

on the boundary and that φ is smooth near U_1 (see [HeSj1]). Apply now the first point of Lemma 3.18 with $f_1 = f$ and $\alpha = \varphi$ and denote by Φ the function Φ_+ of the lemma (Φ is the Agmon distance to U_1 , i.e. associated with the metric $|\nabla_{\underline{x}'} f(\underline{x}')|^2 d\underline{x}'^2$). Hence the following equalities are locally satisfied:

$$|\partial_n \Phi|^2 + |\nabla_T \Phi|^2 = |\nabla \Phi|^2 = |\nabla f|^2,$$

D. Le Peutrec

$$\begin{aligned}\Phi|_{\partial\Omega} &= \varphi, \\ \partial_n \Phi|_{\partial\Omega} &= \frac{\partial f}{\partial n}|_{\partial\Omega}.\end{aligned}$$

According to [HeSj4] pp. 279–280, there exist Morse coordinates (v_1, \dots, v_{n-1}) for $f|_{\Omega}$ centered at U_1 and such that $dv_1(U_1), \dots, dv_{n-1}(U_1)$, $\vec{n}_{U_1}^*$ is orthonormal and positively oriented. With these coordinates

$$f(v, 0) = \frac{\lambda_1}{2}v_1^2 + \dots + \frac{\lambda_{n-1}}{2}v_{n-1}^2 + f(U_1) \quad (4.2.1)$$

and

$$\varphi(v) = \frac{|\lambda_1|}{2}v_1^2 + \dots + \frac{|\lambda_{n-1}|}{2}v_{n-1}^2, \quad (4.2.2)$$

with $\lambda_1 < 0$.

Moreover, (x_1, \dots, x_{n-1}) can be chosen equal to (v_1, \dots, v_{n-1}) in the boundary. Hence, Theorem 1.1 of [Lep2] (about the Neumann case) implies the following proposition:

PROPOSITION 4.1. — *Consider around U_1 the above system of coordinates $\underline{x} = (\underline{x}', \underline{x}_n)$ which satisfies (4.2.1)(4.2.2) with $\lambda_1 < 0$. There exists locally, in a neighborhood of $\underline{x} = 0$, a C^∞ solution u_1^{wkb} to*

$$\Delta_{f,h}^{(1)} u_1^{wkb} = e^{-\frac{\Phi}{h}} \mathcal{O}(h^\infty) \quad (4.2.3)$$

$$\mathbf{n}u_1^{wkb} = 0 \text{ on } \partial\Omega \quad (4.2.4)$$

$$\mathbf{n}d_{f,h}u_1^{wkb} = 0 \text{ on } \partial\Omega, \quad (4.2.5)$$

where u_1^{wkb} has the form:

$$\begin{aligned}u_1^{wkb} &= a(\underline{x}, h)e^{-\frac{\Phi}{h}}, \\ \text{with } a(\underline{x}, h) &\sim \sum_k a^k(\underline{x})h^k \text{ and } a^0(0) = d\underline{x}_1.\end{aligned}$$

4.3. Another local Neumann realization of $\Delta_{f,h}^{(1)}$

Let U_1 be a critical point of $f|_{\partial\Omega}$ with index 1 and $\frac{\partial f}{\partial n}(U_1) < 0$ and let introduce a new system of local coordinates.

We apply Lemma 3.18 with $f_1 = f$ and $\alpha = \varphi$, the Agmon distance to U_1 on the boundary. The function Φ_+ of the lemma is then Φ , the Agmon

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

distance to U_1 and we have the existence of local coordinates (x', x_n) around U_1 where Φ and the metric g_0 have the form:

$$\Phi = -x_n + \varphi(x') \quad \text{and} \quad g_0 = g_{nn}(x) dx_n^2 + \sum_{i,j=1}^{n-1} g_{ij}(x) dx_i dx_j .$$

Moreover, the boundary $\partial\Omega$ is locally defined by $\{x_n = 0\}$ and Ω corresponds to $\{x_n < 0\}$.

We work now with the local coordinate system defined above and $x \mapsto |x|$ is the Euclidean norm in these coordinates.

As in [HeNi], we consider the domain, for $\rho > 0$,

$$\Omega_{U_1, \rho} = \left\{ |x - (0, 1)|^2 < \rho^2 + 1, x_n < 0 \right\} ,$$

which has the shape of a thin lens stuck on $\partial\Omega$ with radius ρ and thickness $\mathcal{O}(\rho^2)$. Its boundary splits into

$$\Gamma_D := \partial\Omega_{U_1, \rho} \cap \Omega = \left\{ |x - (0, 1)|^2 = \rho^2 + 1, x_n \leq 0 \right\}$$

and

$$\Gamma_{ND} := \partial\Omega_{U_1, \rho} \cap \partial\Omega = \{|x'| < \rho, x_n = 0\} .$$

On this domain, we introduce the functional space

$$\Lambda^1 H_{0;0,\mathbf{n}}^1(\Omega_{U_1, \rho}) = \{u \in \Lambda^1 H^1(\Omega_{U_1, \rho}); \mathbf{n}u|_{\Gamma_{ND}} = 0, u|_{\Gamma_D} = 0\} .$$

The Friedrichs extension associated with the quadratic form:

$$\Lambda^1 H_{0;0,\mathbf{n}}^1(\Omega_{U_1, \rho}) \ni \omega \mapsto \mathcal{D}_{g,f,h}^N(\omega) = \|d_{f,h}\omega\|^2 + \|d_{f,h}^*\omega\|^2 ,$$

is denoted by $\Delta_{f,h}^{N,D,(1)}$. The domain of $\Delta_{f,h}^{N,D,(1)}$ can be embedded in $\Lambda^1 H^2(\Omega_{U_1, \rho'})$ for any $0 < \rho' < \rho$.

An element $\omega \in D(\Delta_{f,h}^{N,D,(1)})$ satisfies indeed:

$$\langle \Delta_{f,h}^{N,D,(1)} \omega \mid \eta \rangle = \langle d_{f,h}\omega \mid d_{f,h}\eta \rangle + \langle d_{f,h}^*\omega \mid d_{f,h}^*\eta \rangle = \mathcal{D}_{g,f,h}(\omega, \eta) ,$$

for all $\eta \in \Lambda^1 H_{0;0,\mathbf{n}}^1$, taking $\omega = \eta \equiv 0$ outside $\Omega_{U_1, \rho}$. By testing with $\eta \in \mathcal{C}_0^\infty(\Omega_{U_1, \rho})$, this gives $\Delta_{f,h}\omega \in \Lambda^1 L^2(\Omega_{U_1, \rho})$ and therefore ω admits a second trace on Γ_{ND} thanks to the Green formula. By testing with any $\eta \in \mathcal{C}_{0;0,\mathbf{n}}^\infty(\Omega_{U_1, \rho})$, we get:

$$\mathbf{n}d_{f,h}\omega|_{\Gamma_{ND}} = 0 .$$

Along Γ_{ND} , ω solves an elliptic boundary value problem $\Delta_{f,h}^{(1)}\omega \in \Lambda^1 L^2$, $\mathbf{n}\omega = 0$, $\mathbf{n}d_{f,h}\omega = 0$, which provides the H^2 regularity in $\Omega_{U_{1,\rho'}}$ for any $\rho' < \rho$.

We now prove:

PROPOSITION 4.2. — *For $\rho > 0$ small enough, there exist $h_\rho > 0$ and $C_\rho > 0$, such that the self-adjoint operator $\Delta_{f,h}^{N,D,(1)}$ satisfies the following properties:*

a) *For $h \in (0, h_\rho]$, the spectral projection $1_{[0, h^{3/2})}(\Delta_{f,h}^{N,D,(1)})$ has rank 1.*

b) *Any family of L^2 -normalized eigenvectors $(u^h)_{h \in (0, h_\rho]}$ of $\Delta_{f,h}^{N,D,(1)}$ such that the corresponding eigenvalue $E(h)$ is $\mathcal{O}(h)$, satisfies*

$$\forall \rho' < \rho, \forall \alpha \in \mathbb{N}^n, \exists N_\alpha \in \mathbb{N}, \exists C_{\alpha, \rho'} > 0 \text{ such that, } \forall x \in \Omega_{U_{1,\rho'}}, \quad (4.3.1)$$

$$\left| \partial_x^\alpha u^h(x) \right| \leq C_{\alpha, \rho'} h^{-N_\alpha} \exp\left(-\frac{\Phi(x)}{h}\right).$$

c) *There exists $\varepsilon_\rho > 0$ such that the first eigenvalue $E_1(h)$ of $\Delta_{f,h}^{N,D,(1)}$ satisfies*

$$E_1(h) = \mathcal{O}(e^{-\varepsilon_\rho/h}).$$

d) *If u_1^h denotes the eigenvector of $\Delta_{f,h}^{N,D,(1)}$ associated with eigenvalue $E_1(h)$ and normalized by the condition $\mathbf{t}u_1^h(0) = \mathbf{t}u_1^{wkb}(0)$, then*

$$\forall \rho' < \rho, \forall \alpha \in \mathbb{N}^n, \forall N \in \mathbb{N}, \exists C_{N, \alpha, \rho'} > 0 \text{ such that, } \forall x \in \Omega_{U_{1,\rho'}}, \quad (4.3.2)$$

$$\left| \partial_x^\alpha (u_1^h - u_1^{wkb})(x) \right| \leq C_{N, \alpha, \rho'} h^N \exp\left(-\frac{\Phi(x)}{h}\right).$$

Once this is proved, one easily gets rough exponentially small upper bounds for the $m_{\ell}^{\overline{\Omega}}$ first eigenvalues of $\Delta_{f,h}^{N,(\ell)}$ ($\ell \in \{0, 1\}$) on Ω , by constructing quasimodes suitably localized near each of the critical points.

The next subsections are devoted to the proof of Proposition 4.2. A fundamental ingredient for the proof is a variant of the integration by parts formula of Lemma 2.6.

LEMMA 4.3. — *Let $\rho > 0$ and let ψ be a real-valued Lipschitz function on $\overline{\Omega_{U_{1,\rho}}}$. The relation*

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

$$\begin{aligned} \operatorname{Re} \mathcal{D}_{g,f,h}^N(\omega, e^{2\frac{\psi}{h}}\omega) &= h^2 \left\| de^{\frac{\psi}{h}}\omega \right\|_{\Lambda^2 L^2}^2 + h^2 \left\| d^* e^{\frac{\psi}{h}}\omega \right\|_{\Lambda^0 L^2}^2 \\ &\quad + \langle (|\nabla f|^2 - |\nabla\psi|^2 + h\mathcal{L}_{\nabla f} + h\mathcal{L}_{\nabla f}^*) e^{\frac{\psi}{h}}\omega \mid e^{\frac{\psi}{h}}\omega \rangle_{\Lambda^1 L^2} \\ &\quad + h \int_{\Gamma_{ND}} \langle \omega \mid \omega \rangle_{\Lambda^1 T_\sigma^* \Omega} e^{2\frac{\psi(\sigma)}{h}} \left(\frac{\partial f}{\partial n} \right) (\sigma) d\sigma \end{aligned} \tag{4.3.3}$$

holds for any $\omega \in \Lambda^1 H_{0,0,\mathbf{n}}^1(\Omega_{U_{1,\rho}})$.

Moreover, when $\omega \in D(\Delta_{f,h}^{N,D,(1)})$, the left-hand side equals $\operatorname{Re} \langle e^{2\frac{\psi}{h}} \Delta_{f,h}^{(1)} \omega \mid \omega \rangle$.

Proof. — For ω in $\Lambda^1 H_{0,0,\mathbf{n}}^1(\Omega_{U_{1,\rho}})$, we have $\tilde{\omega} := e^{2\frac{\psi}{h}}\omega$ in $\Lambda^1 H_{0,0,\mathbf{n}}^1(\Omega_{U_{1,\rho}})$ and the same computations as the ones done in [HeNi] to prove Lemma 4.3 lead to:

$$\begin{aligned} \mathcal{D}_{g,f,h}^N(\omega, e^{2\frac{\psi}{h}}\omega) &= \mathcal{D}_{g,f,h}^N(\tilde{\omega}, \tilde{\omega}) - \langle |\nabla\psi|^2 \tilde{\omega} \mid \tilde{\omega} \rangle \\ &\quad - \langle d\psi \wedge \tilde{\omega} \mid d_{f,h}\tilde{\omega} \rangle + \langle d_{f,h}\tilde{\omega} \mid d\psi \wedge \tilde{\omega} \rangle \\ &\quad + \langle \mathbf{i}_{\nabla\psi}\tilde{\omega} \mid d_{f,h}^*\tilde{\omega} \rangle - \langle d_{f,h}^*\tilde{\omega} \mid \mathbf{i}_{\nabla\psi}\tilde{\omega} \rangle. \end{aligned}$$

By taking the real part, we obtain:

$$\operatorname{Re} \mathcal{D}_{g,f,h}^N(\omega, e^{2\frac{\psi}{h}}\omega) = \mathcal{D}_{g,f,h}^N(\tilde{\omega}, \tilde{\omega}) - \langle |\nabla\psi|^2 \tilde{\omega} \mid \tilde{\omega} \rangle.$$

We conclude by applying Lemma 2.6. \square

4.4. Exponential decay of eigenvectors of $\Delta_{f,h}^{N,D,(1)}$

As in [HeNi], the pointwise estimate, $\partial_x^\alpha u^h(x) = \mathcal{O}(h^{-N\alpha} e^{-\frac{\Phi(x)}{h}})$, which is stated in Proposition 4.2-b), will be proved in several steps. We will first consider H^1 -estimates and deduce afterwards higher order estimates from elliptic regularity.

Even for H^1 -estimates we need two steps: we prove first the exponential decay along the boundary Γ_{ND} by applying Lemma 4.3 with the function ψ similar to φ introduced above ; then the exponential decay in the interior of $\Omega_{U_{1,\rho}}$ is obtained with ψ similar to Φ once the boundary term is well controlled.

Proof of a) and b) in Proposition 4.2. —

Statement a)

Actually it is a simple comparison with the full half-space problem via Min-Max principle as we did for Theorem 3.5. Any $\omega \in \Lambda^1 H_{0,0,\mathbf{n}}^1(\Omega_{U_{1,\rho}})$ can indeed be viewed as an element of $\Lambda^1 H_{0,\mathbf{n}}^1(\mathbb{R}_-^n)$ by setting $\omega = 0$ on $\mathbb{R}_-^n \setminus \Omega_{U_{1,\rho}}$.

Statement b)

Let $u^h \in D(\Delta_{f,h}^{N,D,(1)})$ satisfy

$$\Delta_{f,h}^{(1)} u^h = E(h)u^h, \quad E(h) = \mathcal{O}(h), \quad \|u^h\| = 1.$$

We will use the notation

$$\tilde{u}^h = e^{\frac{\psi^h}{h}} u^h.$$

The integration by parts formula (4.3.3) will be applied with $\psi = \psi^h$ where ψ^h will be similar to φ or similar to Φ .

Let us recall

$$|\nabla f|^2 = |\nabla \Phi|^2, \quad \frac{\partial f}{\partial n} = \frac{\partial \Phi}{\partial n} \quad \text{and} \quad \Phi(x', x_n) = -x_n + \varphi(x'), \quad (4.4.1)$$

where $x' = 0$ is a local minimum for φ with $\varphi(0) = 0$. Moreover we have $\nabla x_n \cdot \nabla \varphi(x') = 0$ so that:

$$|\nabla f|^2 = |\nabla \Phi|^2 = |\nabla x_n|^2 + |\nabla \varphi|^2. \quad (4.4.2)$$

We will first show the decay along the boundary before we propagate the decay in the normal direction inside Ω (see [HeSj5] and [HeNi] for references).

Step 1: Decay along Γ_{ND} .

We take:

$$\psi^h(x', x_n) = \begin{cases} \varphi(x') - Ch \log \frac{\varphi(x')}{h}, & \text{if } \varphi(x') > Ch \\ \varphi(x') - Ch \log C, & \text{if } \varphi(x') \leq Ch, \end{cases}$$

where the constant $C > 1$ will be fixed later.

We associate the sets:

$$\Omega_-^h = \{x = (x', x_n) \in \Omega_{U_{1,\rho}}; \varphi(x') < Ch\},$$

and

$$\Omega_+^h = \{x = (x', x_n) \in \Omega_{U_{1,\rho}}; \varphi(x') > Ch\}.$$

The condition $E(h) = \mathcal{O}(h)$ the formula (4.3.3), (4.4.1) and (4.4.2) imply the existence of $C_1 > 0$ such that:

$$\begin{aligned} C_1 h \|\tilde{u}^h\|_{\Lambda^1 L^2(\Omega_-^h)}^2 &\geq \|hd\tilde{u}^h\|_{\Lambda^2 L^2}^2 + \|hd^* \tilde{u}^h\|_{\Lambda^0 L^2}^2 + \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 L^2} \\ &- h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma + \langle (|\nabla \varphi|^2 - |\nabla \psi^h|^2) \tilde{u}^h | \tilde{u}^h \rangle \\ &- C_1 h \langle 1_{\Omega_+^h}(x) \tilde{u}^h | \tilde{u}^h \rangle, \end{aligned} \quad (4.4.3)$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

with C_1 determined by f and the upper bound of $E(h)$.

Furthermore,

$$\nabla\psi^h = \nabla\varphi - 1_{\Omega_+^h}(x) \frac{Ch\nabla\varphi}{\varphi},$$

so we have:

$$|\nabla\psi^h|^2 = |\nabla\varphi|^2 + 1_{\Omega_+^h}(x) \left(-2Ch \frac{|\nabla\varphi|^2}{\varphi} + C^2h^2 \frac{|\nabla\varphi|^2}{\varphi^2} \right).$$

Consequently,

$$\begin{aligned} C_1h \|\tilde{u}^h\|_{\Lambda^1L^2(\Omega_-^h)}^2 &\geq \|hd\tilde{u}^h\|_{\Lambda^2L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0L^2}^2 + \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1L^2} \\ &\quad - h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1T_\sigma^*\Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma \\ &\quad + \langle \left[|\nabla\varphi|^2 \left(\frac{2Ch}{\varphi} - \frac{C^2h^2}{\varphi^2} \right) - C_1h \right] 1_{\Omega_+^h}(x) \tilde{u}^h | \tilde{u}^h \rangle. \end{aligned}$$

For $x \in \Omega_+^h$,

$$\frac{2Ch}{\varphi} - \frac{C^2h^2}{\varphi^2} \geq \frac{Ch}{\varphi} \quad (\text{since } 2a - a^2 \geq a \quad \forall a \in [0, 1])$$

then, φ being a positive Morse function, there exists $C_2 > 0$ which is determined by φ such that, for all $x \in \Omega_+^h$,

$$C_2 \geq \frac{|\nabla\varphi(x')|^2}{\varphi(x')} \geq C_2^{-1}$$

and we get:

$$\begin{aligned} C_1h \|\tilde{u}^h\|_{\Lambda^1L^2(\Omega_-^h)}^2 &\geq \|hd\tilde{u}^h\|_{\Lambda^2L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0L^2}^2 \\ &\quad + \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1L^2} - h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1T_\sigma^*\Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma \\ &\quad + (CC_2^{-1} - C_1) h \langle 1_{\Omega_+^h}(x) \tilde{u}^h | \tilde{u}^h \rangle. \quad (4.4.4) \end{aligned}$$

Since $\partial_n f(U_1) = \partial_n x_n(U_1) \neq 0$, we can choose ρ small enough such that:

$$C_3 \geq |\nabla x_n|^2 \geq C_3^{-1} \quad \text{on } \Omega_{U_1, \rho},$$

where C_3 is a strictly positive constant.

Hence we get, by adding the term $(CC_2^{-1} - C_1)h\langle 1_{\Omega_-^h}(x)\tilde{u}^h | \tilde{u}^h \rangle$ to (4.4.4):

$$\begin{aligned} CC_2^{-1}h \|\tilde{u}^h\|_{\Lambda^1 L^2(\Omega_-^h)}^2 &\geq \|hd\tilde{u}^h\|_{\Lambda^2 L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0 L^2}^2 \\ &\quad + (1 + 2\delta(C)h) \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 L^2} \\ &\quad - h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma, \end{aligned}$$

where $\delta(C) = \frac{1}{2}C_3^{-1}(CC_2^{-1} - C_1)$ is positive and large for C large enough.

At least, we have on Ω_-^h by the definitions:

$$|\tilde{u}^h| \leq e^C |u^h| \text{ a.e.}$$

and the condition $\|u^h\| = 1$ leads to:

$$\begin{aligned} \tilde{\delta}(C)h &\geq \|hd\tilde{u}^h\|_{\Lambda^2 L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0 L^2}^2 + (1 + 2\delta(C)h) \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 L^2} \\ &\quad - h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma, \end{aligned} \tag{4.4.5}$$

where $\tilde{\delta}(C) = e^{2C}CC_2^{-1}$.

We now apply (4.3.3) to \tilde{u}^h with $\psi = 0$, f and h replaced respectively by $-x_n$ and $\frac{h}{1+\delta(C)h}$, in order to get,

$$\begin{aligned} (1 + \delta(C)h)^{-1} \|hd\tilde{u}^h\|_{\Lambda^2 L^2}^2 &+ (1 + \delta(C)h)^{-1} \|hd^*\tilde{u}^h\|_{\Lambda^0 L^2}^2 \\ &+ (1 + \delta(C)h) \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle - h \int_{\Gamma_{ND}} \langle \tilde{u}^h | \tilde{u}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma \\ &\quad + hC_4 \|\tilde{u}^h\|_{\Lambda^1 L^2}^2 \geq 0, \end{aligned} \tag{4.4.6}$$

with $C_4 > 0$ independent of C .

The difference (4.4.5)–(4.4.6) yields:

$$\begin{aligned} \frac{\delta(C)h^3}{1 + \delta(C)h} \left[\|d\tilde{u}^h\|_{\Lambda^2 L^2}^2 + \|d^*\tilde{u}^h\|_{\Lambda^0 L^2}^2 \right] &- hC_4 \|\tilde{u}^h\|_{\Lambda^1 L^2}^2 \\ &+ \delta(C)h \langle |\nabla x_n|^2 \tilde{u}^h | \tilde{u}^h \rangle \leq \tilde{\delta}(C)h. \end{aligned}$$

We choose $C > 1$ large enough such that $\delta(C)C_3^{-1} - C_4 > 0$.

This leads, after choosing $h_0 > 0$ small enough, to the existence of a constant $C_5 > 0$ such that, for all $h \in (0, h_0]$,

$$C_5 h \geq h^3 \|\tilde{u}^h\|_{\Lambda^1 H^1}^2.$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Since $\psi^h \geq \varphi + \tilde{C}h \log h$ (for all $\tilde{C} > C$), we have proved the existence of $N_0 > 0$ such that:

$$\left\| e^{\frac{\varphi}{h}} u^h \right\|_{\Lambda^1 H^1} \leq C_6 h^{-N_0}. \quad (4.4.7)$$

Remember that $\varphi \geq 0$ vanishes only at $x' = 0$. Using the trace theorem, this also leads to:

$$\left\| e^{\frac{\varphi}{h}} u^h \Big|_{\Gamma_{ND}} \right\|_{\Lambda^1 H^{1/2}(\Gamma_{ND})} \leq C_7 h^{-N_0}. \quad (4.4.8)$$

Step 2: Normal decay inside Ω .

We follow a very similar approach by working with the function Φ .

We take:

$$\psi^h(x', x_n) = \begin{cases} \Phi(x) - Ch \log \frac{\Phi(x)}{h}, & \text{if } \Phi(x) > Ch \\ \Phi(x) - Ch \log C, & \text{if } \Phi(x) \leq Ch, \end{cases}$$

where the constant $C > 1$ will be fixed later.

We associate the sets:

$$\Omega_-^h = \{x = (x', x_n) \in \Omega_{U_1, \rho}; \Phi(x) < Ch\}$$

and

$$\Omega_+^h = \{x = (x', x_n) \in \Omega_{U_1, \rho}; \Phi(x) > Ch\}.$$

The formula (4.3.3) is used like in Step 1, with $\tilde{u}^h = e^{\frac{\psi^h}{h}} u^h$ and $E(h) = \mathcal{O}(h)$. The difference comes from the fact that the boundary term is already estimated with (4.4.8).

We have indeed on the boundary $x_n = 0$ the inequality: $e^{\frac{\psi^h}{h}} \leq e^{\frac{\varphi}{h}}$, due to the relation $\Phi|_{x_n=0} = \varphi$.

From (4.3.3) used like in Step 1 (see (4.4.3)) we get the existence of $C_1 > 0$ such that:

$$\begin{aligned} C_1 h \left\| \tilde{u}^h \right\|_{\Lambda^1 L^2(\Omega_-^h)}^2 + C_1 h \left\| e^{\frac{\varphi}{h}} u \right\|_{H^{1/2}(\Gamma_{ND}; \Lambda^1 T^* \Omega_{U_1, \rho})}^2 &\geq \left\| h d \tilde{u}^h \right\|_{\Lambda^2 L^2}^2 \\ + \left\| h d^* \tilde{u}^h \right\|_{\Lambda^0 L^2}^2 + \langle (|\nabla f|^2 - |\nabla \psi^h|^2) \tilde{u}^h \mid \tilde{u}^h \rangle - C_1 h \langle 1_{\Omega_+^h}(x) \tilde{u}^h \mid \tilde{u}^h \rangle. \end{aligned}$$

Moreover, from (4.4.8) and the inequality

$$|\tilde{u}^h(x)| \leq e^C |u^h(x)| \quad \text{a.e. in } \Omega_-^h,$$

we get, for any $C > 1$, the existence of $\tilde{\delta}(C) > 0$ such that the following estimate is satisfied:

$$\begin{aligned} \tilde{\delta}(C)h^{1-2N_0} &\geq C_1h \|\tilde{u}^h\|_{\Lambda^1L^2(\Omega^h_+)}^2 + C_1h \left\| e^{\frac{\Phi}{h}}u \right\|_{H^{1/2}(\Gamma_{ND};\Lambda^1T^*\Omega_{U_1,\rho})}^2 \\ &\geq \|hd\tilde{u}^h\|_{\Lambda^2L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0L^2}^2 + \langle (|\nabla f|^2 - |\nabla\psi^h|^2)\tilde{u}^h | \tilde{u}^h \rangle \\ &\quad - C_1h \langle 1_{\Omega^h_+}(x)\tilde{u}^h | \tilde{u}^h \rangle. \end{aligned} \quad (4.4.9)$$

Since $|\nabla f|^2 = |\nabla\Phi|^2$ and Φ is a positive function without critical points, we can use the same computations as the ones done in Step 1 with φ replaced by Φ to get:

$$\begin{aligned} |\nabla f|^2 - |\nabla\psi^h|^2 &= 1_{\Omega^h_+}(x) \left(2Ch \frac{|\nabla\Phi|^2}{\Phi} - C^2h^2 \frac{|\nabla\Phi|^2}{\Phi^2} \right) \\ &\geq \frac{Ch|\nabla\Phi|^2}{\Phi} \geq C_2^{-1}Ch, \end{aligned}$$

with $C_2 > 0$ independent of C .

We take $C \geq 2C_1C_2$. By adding the estimated term $(C_2^{-1}C - C_1)h \langle 1_{\Omega^h_+}(x)\tilde{u}^h | \tilde{u}^h \rangle$ to (4.4.9) we get:

$$\tilde{\delta}_2(C)h^{1-2N_0} \geq \|hd\tilde{u}^h\|_{\Lambda^2L^2}^2 + \|hd^*\tilde{u}^h\|_{\Lambda^0L^2}^2 + (C_2^{-1}C - C_1)h \|\tilde{u}^h\|_{\Lambda^1L^2},$$

which gives, by analogy with Step 1, the existence of $C_3 > 0$ and $N_1 > 0$ such that:

$$\left\| e^{\frac{\Phi}{h}}u \right\|_{\Lambda^1H^1(\Omega_{U_1,\rho})} \leq C_3 h^{-N_1}. \quad (4.4.10)$$

Step 3: Elliptic regularity.

We now set $\tilde{u}^h = e^{\frac{\Phi}{h}}u^h$. For $\rho' < \rho$, we take a cut-off $\chi \in \mathcal{C}^\infty(\Omega_{U_1,\rho})$ with compact support in $\Omega_{U_1,\rho} \cup \Gamma_{ND}$ and such that $\chi = 1$ on a neighborhood of $\Omega_{U_1,\rho'}$. The form $v^h = \chi\tilde{u}^h$ satisfies the boundary value problem:

$$\begin{cases} v^h - \Delta v^h = r_0^h & \text{in } \mathbb{R}^n, \\ \mathbf{n}v^h = 0 \text{ and } \mathbf{n}dv^h = r_1^h & \text{on } \{x_n = 0\}, \end{cases}$$

$$\text{with } \|r_0^h\|_{\Lambda^1L^2(\mathbb{R}^n_-)} = \mathcal{O}(h^{-N_1}) \text{ and } \|r_1^h\|_{\Lambda^2H^{1/2}(\mathbb{R}^{n-1})} = \mathcal{O}(h^{-N_1}).$$

This implies, by [Sch], the existence of $N_2 > 0$ such that:

$$\|v^h\|_{\Lambda^1H^2} = \mathcal{O}(h^{-N_2}).$$

We conclude by induction for any finite decreasing sequence $(\rho_k)_{0 \leq k \leq K}$ with $\rho_K > \rho'$ and associated cut-offs χ_k , with $\chi_k = 1$ in a neighborhood of Ω_{U_1,ρ_k} and $\text{supp } \chi_k \subset \{\chi_{k-1} = 1\}$, using the Sobolev injections. \square

4.5. Small eigenvalues are exponentially small

We now check that the eigenvalue $E_1(h)$ of $\Delta_{f,h}^{N,D,(1)}$ lying in $[0, h^{3/2})$ is actually of order $\mathcal{O}(e^{-\varepsilon_\rho/h})$ for some $\varepsilon_\rho > 0$. We prove this by comparison with the half-space problem as it is done in [Lep2] at the end of the proof of Theorem 1.1.

Proof of Proposition 4.2-c. — Again we introduce in a neighborhood of U_1 , the coordinate system $\bar{x} = (\bar{x}', \bar{x}_n)$ leading to (3.3.1). The function f and the metric g_0 are extended according to (3.3.2) and (3.3.3) so that Proposition 3.16 can be applied. Consequently, the half-space Witten Laplacian, $\Delta_{\bar{f},h}^{N,(1)}$, has a one dimension kernel and its second eigenvalue is larger than $Ch^{6/5}$.

Let u^h be a normalized eigenvector of $\Delta_{f,h}^{N,D,(1)}$ associated with the first eigenvalue $E_1(h)$, which belongs to the interval $(0, h^{3/2}]$. Let $\chi \in \mathcal{C}^\infty(\overline{\Omega_{U_1,\rho}})$ be a cut-off function with compact support in $\Omega_{U_1,\rho} \cup \Gamma_{ND}$ and such that $\chi = 1$ in a neighborhood of 0 with $\frac{\partial \chi}{\partial n} \Big|_{\partial \Omega} \equiv 0$.

The form $v^h = \chi u^h \in \Lambda^1 H^2(\mathbb{R}_-^n)$ belongs to the domain of $\Delta_{\bar{f},h}^{N,(1)}$, i.e. $\mathbf{n}v^h = \mathbf{n}d_{\bar{f},h} v^h = 0$. Moreover, v^h satisfies

$$(\Delta_{\bar{f},h}^{(1)} - E_1(h))v^h = -h^2[\Delta, \chi]u^h \text{ in } \mathbb{R}_-^n$$

and the 1-form $r^h = -h^2[\Delta, \chi]u^h$ vanishes in a neighborhood \mathcal{V}_1 of $\bar{x} = 0$. Due to the exponential decay of u^h stated in Proposition 4.2-b), there exist C and N_0 , such that r^h also satisfies

$$|r^h(\bar{x})| \leq Ch^{-N_0} \left[\sum_{1 \leq |\beta| \leq 2} |\partial_{\bar{x}}^\beta \chi(\bar{x})| \right] e^{-\frac{\Phi(\bar{x})}{h}} \leq e^{-\frac{c\chi}{h}}.$$

With $\|v^h\|_{\Lambda^1 L^2} = 1 + \mathcal{O}(e^{-c/h})$, $\|r^h\|_{\Lambda^1 L^2} = \mathcal{O}(e^{-c/h})$ and the a priori estimate $E_1(h) = \mathcal{O}(h^{3/2})$, the spectral theorem implies $|E_1(h) - 0| = \mathcal{O}(e^{-c/h})$ like in the proof of Theorem 1.1 given in [Lep2] (p. 245). \square

4.6. Accurate comparison with the WKB solution

We now compare the eigenvector associated with an exponentially small eigenvalue with its WKB approximation. We adapt the method presented in [Hel2, HeSj1] and in [HeNi] by following the same strategy as in Subsection 4.4. The H^1 -estimates are done in two steps with ψ^h similar to φ

and then with ψ^h similar to Φ . Finally the elliptic regularity is used for the \mathcal{C}^∞ -estimates.

Proof of Proposition 4.2-d). — Let $u_1^h \in D(\Delta_{f,h}^{N,D,(1)})$ be an eigenvector associated with the first eigenvalue $E_1(h)$ of $\Delta_{f,h}^{N,D,(1)}$:

$$\Delta_{f,h}^{N,D,(1)} u_1^h = E_1(h) u_1^h, \quad \|u_1^h\| = 1.$$

According to Proposition 4.2-c), we know that $E_1(h) = \mathcal{O}(e^{-\frac{\varepsilon\rho}{h}})$, with $\varepsilon_\rho > 0$, while the second eigenvalue of $\Delta_{f,h}^{N,D,(1)}$ is larger than $h^{3/2}$.

By taking $\rho > 0$ small enough, the WKB approximation u_1^{wkb} presented in Subsection 4.2 satisfies

$$\begin{cases} \Delta_{f,h}^{(1)} u_1^{wkb} = \mathcal{O}(h^\infty) e^{-\frac{\Phi(x)}{h}} & \text{in } \Omega_{U_1,\rho}, \\ \mathbf{n}u_1^{wkb}|_{\Gamma_{ND}} = 0, \\ \mathbf{n}d_{f,h}u_1^{wkb}|_{\Gamma_{ND}} = 0, \end{cases}$$

and there exists $c > 0$, such that for any $\rho' > 0$, we have

$$\|u_1^{wkb}\|_{\Lambda^1 L^2(\Omega_{U_1,\rho'})} \sim ch^{\frac{n+1}{4}}$$

(see indeed further the proof of Proposition 5.19).

Let us choose the cut-off function $\chi \in \mathcal{C}^\infty(\overline{\Omega_{U_1,\rho}})$ to be supported in $\Omega_{U_1,\rho/2} \cup \Gamma_{ND}$ and to satisfy $\chi = 1$ on $\Omega_{U_1,\rho'}$ with $0 < \rho' < \rho/2$, $\frac{\partial \chi}{\partial n}|_{\partial \Omega} \equiv 0$. Later, we will take $\rho' > 0$ small enough, so that χ can be taken in the form

$$\chi(x', x_n) = \chi_1(x') \chi_n(x_n).$$

From

$$(\Delta_{f,h}^{N,D,(1)} - E_1(h)) \chi u_1^{wkb} = \mathcal{O}(h^\infty)$$

and the comparison result of [Hel2] (see Proposition 4.1.1), the real constant factor $c(h)$ in the truncated WKB approximation $v_1^{wkb} = c(h) \chi u_1^{wkb}$ can be chosen so that

$$\|v_1^{wkb} - u_1^h\|_{\Lambda^1 L^2} = \mathcal{O}(h^\infty).$$

But then $\langle \Delta_{f,h}^{N,D,(1)} (v_1^{wkb} - u_1^h) | v_1^{wkb} - u_1^h \rangle = \mathcal{O}(h^\infty)$, which implies

$$\|v_1^{wkb} - u_1^h\|_{\Lambda^1 H^1} = \mathcal{O}(h^\infty)$$

and, due to the exponential decay of u_1^h and u_1^{wkb} ,

$$\|\chi(u_1^h - c(h)u_1^{wkb})\|_{\Lambda^1 H^1} = \mathcal{O}(h^\infty).$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Set

$$w^h = \chi(u_1^h - c(h)u_1^{wkb}) .$$

The 1-form w^h satisfies in $\Omega_{U_1, \rho}$

$$\begin{aligned} (\Delta_{f,h}^{(1)} - E_1(h))w^h &= \chi(x)(\Delta_{f,h}^{(1)} - E_1(h))(u_1^h - c(h)u_1^{wkb}) \\ &\quad + [\Delta_{f,h}^{(1)}, \chi](u_1^h - c(h)u_1^{wkb}) \\ &= \tilde{r}^h e^{-\frac{\Phi(x)}{h}} + r^h , \end{aligned} \tag{4.6.1}$$

where \tilde{r}^h and r^h satisfy, according to Proposition 4.2-b),

$$\tilde{r}^h = \mathcal{O}(h^\infty), \quad \text{supp } r^h \subset \text{supp } \nabla \chi \quad \text{and} \quad r^h = \mathcal{O}(h^{-N_0})e^{-\frac{\Phi(x)}{h}} .$$

The last estimate can be done for any \mathcal{C}^{k_0} -norm, with $k_0 \in \mathbb{N}$.

On the boundary $\partial\Omega_{U_1, \rho} = \Gamma_{ND} \cup \Gamma_D$, we have simply

$$\begin{aligned} \mathbf{n}w^h|_{\Gamma_{ND}} &= 0, \quad w^h|_{\Gamma_D} = 0, \\ \text{and} \quad \mathbf{n}d_{f,h}w^h|_{\Gamma_{ND}} &= 0 . \end{aligned}$$

With the different of choices for ψ^h given below, we will use the notation

$$\tilde{w}^h = e^{\frac{\psi^h}{h}} w^h .$$

The 1-forms w and \tilde{w} belong to $\Lambda^1 H^2(\Omega_{U_1, \rho})$ and their supports do not meet Γ_D . Hence the integration by parts given in formula (2.2.7) can be used in addition to (4.3.3).

Step 1: Comparison along Γ_{ND} .

Like in the proof of Proposition 4.2-b) presented in Subsection 4.4, we introduce the sets

$$\begin{aligned} \Omega_-^h &= \{x = (x', x_n) \in \Omega_{U_1, \rho} ; \quad \varphi(x') < Ch\} , \\ \text{and} \quad \Omega_+^h &= \{x = (x', x_n) \in \Omega_{U_1, \rho} ; \quad \varphi(x') > Ch\} . \end{aligned}$$

For any $N \in \mathbb{N}$, we take:

$$\varphi_N^h(x') = \min \{ \varphi^h(x') + Nh \log h^{-1}, \psi(x') \} ,$$

$$\text{where} \quad \varphi^h(x') = \begin{cases} \varphi(x') - Ch \log \frac{\varphi(x')}{h} , & \text{if } \varphi(x') > Ch \\ \varphi(x') - Ch \log C , & \text{if } \varphi(x') \leq Ch , \end{cases}$$

$$\text{and} \quad \psi(x') = \min \{ \varphi^h(y') + (1 - \varepsilon)|\varphi(x') - \varphi(y')|, y' \in \text{supp } \nabla \chi_1 \} .$$

We recall that $|\varphi(x') - \varphi(y')|$ is the Agmon distance between x' and y' along the boundary, $d_{Ag}^{\partial\Omega}(x', y')$ (i.e. associated with the metric $|\nabla_{x'} f(x', 0)|^2 dx'^2$), and that $\varphi(x') = d_{Ag}^{\partial\Omega}(x', 0)$. Moreover, due to the triangular inequality,

$$|\psi(x') - \psi(y')| \leq (1 - \varepsilon)|\varphi(x') - \varphi(y')| = (1 - \varepsilon)d_{Ag}^{\partial\Omega}(x', y'), \quad (4.6.2)$$

which implies, owing to the properties of the Agmon distance (see e.g. [DiSj] p. 53):

$$\text{for almost all } x', \quad |\nabla\psi(x')| \leq (1 - \varepsilon)|\nabla_{x'} f(x', 0)| = (1 - \varepsilon)|\nabla\varphi(x')|. \quad (4.6.3)$$

Let us also recall that the cut-off χ writes $\chi(x', x_n) = \chi_1(x')\chi_n(x_n)$. The constant $C \geq 1$ will be fixed at the end like in the proof of Proposition 4.2-b). The constants $\rho' \in (0, \rho/2)$ and $\varepsilon > 0$ are chosen so that, for $h \in (0, h_{N,\rho',\varepsilon})$,

$$\varphi_N^h(x') = \varphi^h(x') + Nh \log h^{-1} \text{ in } \Omega_{U_1,\rho'}. \quad (4.6.4)$$

Consequently, φ being the Agmon distance on the boundary,

$$\varphi_N^h(x') = \varphi^h(x') + Nh \log h^{-1} = \varphi(x') - Ch \log C + Nh \log h^{-1} \text{ on } \Omega_{\pm}^h. \quad (4.6.5)$$

Note furthermore the inequalities:

$$\begin{aligned} \varphi_N^h(x') &\leq \varphi(x') + Nh \log h^{-1} \text{ in } \Omega_{U_1,\rho} \\ \varphi_N^h(x') &\leq \varphi(x') \leq \Phi(x), \text{ if } x' \in \text{supp } \nabla\chi_1, \\ \text{and } \varphi_N^h(x') &\leq \varphi(x') + Nh \log h^{-1} \leq \Phi(x), \text{ if } x_n \in \text{supp } \chi'_n. \end{aligned}$$

In particular, we have for $h \in (0, h_{N,\rho',\varepsilon})$,

$$\varphi_N^h(x') \leq \Phi(x), \text{ for } x \in \text{supp } \nabla\chi,$$

which implies

$$\left\| e^{\frac{\varphi_N^h}{h}} r^h \right\|_{\Lambda^1 L^2} = \mathcal{O}_N(h^{-N_0}).$$

We apply the integration by parts formula (4.3.3), where the left-hand side is computed with (2.2.7), and we obtain for the form $\tilde{w}^h = e^{\frac{\varphi_N^h}{h}} w^h$, by analogy with the proof of Proposition 4.2-b), using (4.6.1) and $E_1(h) = \mathcal{O}(h^\infty) = \mathcal{O}(h)$:

$$\begin{aligned} C_1 h \|\tilde{w}^h\|_{\Lambda^1 L^2(\Omega_{\pm}^h)}^2 &+ \left\| \tilde{r}^h + e^{\frac{\varphi_N^h}{h}} r^h \right\|_{\Lambda^1 L^2} \|\tilde{w}^h\|_{\Lambda^1 L^2} \geq \|hd\tilde{w}^h\|_{\Lambda^2 L^2}^2 \\ &+ \|hd^* \tilde{w}^h\|_{\Lambda^0 L^2}^2 + \langle |\nabla x_n|^2 \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 L^2} \\ &+ h \int_{\Gamma_{ND}} \langle \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma + \langle (|\nabla\varphi|^2 - |\nabla\varphi_N^h|^2) \tilde{w}^h | \tilde{w}^h \rangle \\ &\quad - C_1 h \langle 1_{\Omega_{\pm}^h}(x) \tilde{w}^h | \tilde{w}^h \rangle, \end{aligned}$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

where the constant $C_1 > 0$ is determined by f and $\tilde{r}^h = \mathcal{O}(h^\infty)$.

In Ω_-^h the weight $e^{\frac{\varphi_N^h}{h}}$ is bounded by $C_2(C)h^{-N}$ and this provides

$$\|\tilde{w}^h\|_{\Lambda^1 L^2(\Omega_-^h)} \leq C_2(C)h^{-N} \|w^h\|_{\Lambda^1 L^2(\Omega_-^h)} \leq C_3(C, N),$$

due to $\|w^h\|_{\Lambda^1 H^1} = \mathcal{O}(h^\infty)$.

We obtain:

$$\begin{aligned} \tilde{\delta}(C, N)(h^{-N_0} \|\tilde{w}^h\|_{\Lambda^1 H^1} + 1) &\geq \|hd\tilde{w}^h\|_{\Lambda^2 L^2}^2 + \|hd^*\tilde{w}^h\|_{\Lambda^0 L^2}^2 \\ &+ \langle |\nabla x_n|^2 \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 L^2} + h \int_{\Gamma_{ND}} \langle \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma \\ &+ \langle (|\nabla \varphi|^2 - |\nabla \varphi_N^h|^2) \tilde{w}^h | \tilde{w}^h \rangle - C_1 h \langle 1_{\Omega_+^h}(x) \tilde{w}^h | \tilde{w}^h \rangle. \end{aligned}$$

In Ω_-^h , $|\nabla \varphi|^2 = |\nabla \varphi_N^h|^2$, using (4.6.5).

In Ω_+^h , the point x fulfills almost surely one of the two possibilities:

- **Either** $\nabla \varphi_N^h = \nabla \psi$, and we get, owing to (4.6.3),

$$|\nabla \varphi|^2 - |\nabla \varphi_N^h|^2 \geq (2\varepsilon - \varepsilon^2) |\nabla \varphi|^2 \geq \delta_{\rho, \varepsilon},$$

where the last lower bound is due to the fact that $\varphi_N(x) = \psi(x)$ cannot occur in a neighborhood of $x' = 0$ for $\varepsilon > 0$ small enough and $h \in (0, h_{N, \rho', \varepsilon})$;

- **or** $\nabla \varphi_N^h = \nabla \varphi(1 - \frac{Ch}{\varphi})$.

So we get, similarly to the proof of Proposition 4.2-b), for C big enough and $h \in (0, h_{N, \rho', \varepsilon}]$, with $h_{N, \rho', \varepsilon} > 0$ small enough:

$$\begin{aligned} \tilde{\delta}_2(C, N)(h^{-N_0} \|\tilde{w}^h\|_{\Lambda^1 H^1} + 1) &\geq \\ &\|hd\tilde{w}^h\|_{\Lambda^2 L^2}^2 + \|hd^*\tilde{w}^h\|_{\Lambda^0 L^2}^2 + (1 + 2\delta(C)h) \langle |\nabla x_n|^2 \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 L^2} \\ &+ h \int_{\Gamma_{ND}} \langle \tilde{w}^h | \tilde{w}^h \rangle_{\Lambda^1 T_\sigma^* \Omega} \left(\frac{\partial x_n}{\partial n} \right) (\sigma) d\sigma. \end{aligned}$$

After treating the r.h.s. like in the proof of Proposition 4.2-b)-Step 1, we obtain, for a constant $N_0 > 0$,

$$\|\tilde{w}^h\|_{\Lambda^1 H^1(\Omega_{U_1, \rho})} \leq C_4 h^{-N_0}.$$

Our choice of (ε, ρ') implies

$$\forall x \in \Omega_{U_1, \rho'}, \quad \varphi_N^h(x') \geq \varphi(x') + Nh \log h^{-1} + \tilde{C}h \log h .$$

We have proved the existence of N_1 and ρ'_0 , such that, for any $N \in \mathbb{N}$ and $\rho' \in (0, \rho'_0]$, there exists $h_{N, \rho'} > 0$ and $C_{N, \rho'} > 0$, such that:

$$\left\| e^{\frac{\varphi}{h}} (u_1^h - c(h)u_1^{wkb}) \right\|_{\Lambda^1 H^1(\Omega_{U_1, \rho'})} \leq C_{N, \rho'} h^{N-N_1}$$

holds for any $h \in (0, h_{N, \rho'})$.

This last estimate and $\Phi|_{\Gamma_{ND}} = \varphi$ imply

$$\left\| e^{\frac{\Phi}{h}} (u_1^h - c(h)u_1^{wkb}) \right\|_{\Lambda^1 H^{1/2}(\Omega_{U_1, \rho'} \cap \Gamma_{ND})} = \mathcal{O}(h^\infty) .$$

Step 2: Comparison in the normal direction.

After replacing ρ' by ρ , Step 1 provides the estimate

$$\left\| e^{\frac{\varphi}{h}} (u_1^h - c(h)u_1^{wkb}) \right\|_{\Lambda^1 H^1} = \mathcal{O}(h^\infty) . \quad (4.6.6)$$

We work in $\Omega_{U_1, \rho}$ with the above estimate and $\rho' \in (0, \rho/2)$ will be taken again small enough.

In order to get the interior estimate with the weight $e^{\frac{\Phi}{h}}$, we modify the previous analysis like in the proof of Proposition 4.2-b). The sets Ω_{\pm}^h are now given by

$$\begin{aligned} \Omega_-^h &= \{x = (x', x_n) \in \Omega_{U_1, \rho} ; \Phi(x) < Ch\} , \\ \text{and} \quad \Omega_+^h &= \{x = (x', x_n) \in \Omega_{U_1, \rho} ; \Phi(x) > Ch\} . \end{aligned}$$

The function φ_N^h , $N \in \mathbb{N}$, is given by

$$\begin{aligned} \varphi_N^h(x) &= \min \{ \varphi^h(x) + Nh \log h^{-1}, \psi(x) \} , \\ \text{with} \quad \varphi^h(x) &= \begin{cases} \Phi(x) - Ch \log \frac{\Phi(x)}{h} , & \text{if } \Phi(x) > Ch , \\ \Phi(x) - Ch \log C , & \text{if } \Phi(x) \leq Ch , \end{cases} \\ \text{and} \quad \psi(x) &= \min \{ \varphi^h(y) + (1 - \varepsilon)d_{Ag}(x, y), y \in \text{supp } \nabla \chi \} . \end{aligned}$$

We recall that the Agmon distance $d_{Ag}(x, y)$ is the distance between x and y for the metric $|\nabla f|^2 dx^2$ and that $\Phi(x) = d_{Ag}(x, U_1)$.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Moreover, equivalently to (4.6.3) in Step 1, the following inequality is satisfied:

$$\text{for almost all } x, \quad |\nabla\psi(x)| \leq (1 - \varepsilon) |\nabla f(x)|. \quad (4.6.7)$$

Furthermore, the constant $C \geq 1$ will again be fixed in the end like in the proof of Proposition 4.2-b), while the constants $\rho' \in (0, \rho/2)$ and $\varepsilon > 0$ are chosen so that:

$$\varphi_N^h(x) = \varphi^h(x) + Nh \log h^{-1} \text{ in } \Omega_{U_1, \rho'}.$$

Again, this implies:

$$\varphi_N^h(x) = \varphi^h(x) + Nh \log h^{-1} \quad \text{on } \Omega_-^h$$

Now we have the inequalities

$$\begin{aligned} \varphi_N^h(x) &\leq \Phi(x) + Nh \log h^{-1} \quad \text{in } \Omega_{U_1, \rho} \\ \text{and} \quad \varphi_N^h(x) &\leq \Phi(x) \quad \text{in } \text{supp } \nabla \chi. \end{aligned}$$

Hence the estimate

$$\left\| e^{\frac{\varphi_N^h}{h}} r^h \right\|_{\Lambda^1 L^2} = \mathcal{O}(h^{-N_0})$$

is still valid.

Inequality (4.6.6) implies that the L^2 -norm of the trace of \tilde{w}^h on Γ_{ND} is $\mathcal{O}(h^\infty)$ and we have the following estimate:

$$\|\tilde{w}^h\|_{\Lambda^1 L^2(\Omega_-^h)} \leq C_2(C) h^{-N} \|w^h\|_{\Lambda^1 L^2(\Omega_-^h)} \leq C_3(C, N).$$

With these estimates, the integration by parts formula (4.3.3) and (2.2.7) lead to:

$$\begin{aligned} \tilde{\delta}(C, N)(h^{-N_0} \|\tilde{w}^h\|_{\Lambda^1 L^2} + 1) &\geq \|hd\tilde{w}^h\|_{\Lambda^2 L^2}^2 + \|hd^* \tilde{w}^h\|_{\Lambda^0 L^2}^2 \\ &\quad + \langle (|\nabla f|^2 - |\nabla \varphi_N^h|^2 - C_1 h) 1_{\Omega_+^h}(x) \tilde{w}^h | \tilde{w}^h \rangle. \end{aligned}$$

Finally, for almost all $x \in \Omega_+^h$ we have:

- **Either:** $\nabla \varphi_N^h(x) = \nabla \psi(x)$

and, owing to (4.6.7),

$$|\nabla f|^2 - |\nabla \varphi_N^h|^2 = (2\varepsilon - \varepsilon^2) |\nabla f|^2 \geq \delta_{\rho, \varepsilon} > 0;$$

- **or:** $\nabla \varphi_N^h(x) = \nabla \varphi^h(x)$

and we get like in the proof of Proposition 4.2-b)

$$|\nabla f|^2 - |\nabla \varphi^h|^2 \geq C_4 C h.$$

By taking C big enough, we get that $\|e^{\frac{\varphi_N^h}{h}} w^h\| = \mathcal{O}(h^{-N_0})$ for some $N_0 > 0$.

Like in Step 1, this leads to

$$\left\| e^{\frac{\Phi}{h}} (u_1^h - c(h)u_1^{wkb}) \right\|_{\Lambda^1 H^1(\Omega_{U_1, \rho'})} = \mathcal{O}(h^\infty),$$

for $\rho' \in (0, \rho/2)$ small enough.

Step 3: The estimates in higher order Sobolev spaces are done like in the proof of Proposition 4.2-b) by a bootstrap argument after writing a boundary value problem for $\chi(u_1^h - c(h)u_1^{wkb})$ in \mathbb{R}_-^n . \square

5. Labelling of local minima and construction of the quasimodes

5.1. Preliminaries

Here we adapt to our case with Neumann boundary condition the method of selecting the proper critical points with index 1 which was used in [HeKlNi] and in [HeNi]. We recall that the intuition for getting the good labelling of local minima, which is useful even to state properly the assumptions and results, comes from the probabilistic approach. The local minima have to be labelled according to the decreasing order of exit times. We refer to [BoGaKl], [BoEcGaKl] and [FrWe] for details.

Note that a similar strategy has independently been considered in [CoPaYc] for the spectral analysis on Markov processes on graphs.

The existence of such a labelling is an assumption which is generically satisfied. After this, it is possible to construct accurately quasimodes leading, with the help of the Witten complex structure, to accurate asymptotic expansions of the low lying eigenvalues.

5.2. Generalized critical points and local structure of the level sets of a Morse function

We recall that we work here on a compact connected oriented Riemannian manifold $\bar{\Omega} = \Omega \cup \partial\Omega$ with boundary and that the function f satisfies Assumption 3.1. According to our preliminary results on the Witten Laplacian $\Delta_{f,h}^N$ in Theorem 3.5, we introduce the following definition of generalized critical points with index p .

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

DEFINITION 5.1. — A point $U \in \overline{\Omega}$ will be called a generalized critical point of f with index p if:

- either $U \in \Omega$ and U is a critical point of f with index p ,
- or $U \in \partial\Omega$ and U is a critical point with index p of $f|_{\partial\Omega}$ such that $\frac{\partial f}{\partial \vec{n}}(U) < 0$ (\vec{n} being the outgoing normal vector).

Remark 5.2. — In particular, for $p = 0$, we get that the generalized minima are simply the local minima.

The set of generalized critical points with index p is denoted by $\mathcal{U}^{(p)}$. We recall that we want to analyze the Witten Laplacian on 0-forms so we restrict our attention to the cases $p = 0$ and $p = 1$. From now on, we will use the notation:

$$m_p = \#\mathcal{U}^{(p)} \quad \text{for } p = 0, 1 \quad (5.2.1)$$

instead of $m_p^{\overline{\Omega}}$.

Finally it is convenient to call \mathcal{U} the union of all critical points of f and $f|_{\partial\Omega}$.

Before labelling the local minima, let us recall a few remarks coming from the local analysis of a Morse function which satisfies Assumption 3.1 (we refer to [Mill], [HeKlNi], and [HeNi]).

Local structure of the level sets of a Morse function.

In order to analyze the local situation near a point x_0 of $\overline{\Omega}$, let us introduce:

$$A_f^<(x_0) := \{x \in \overline{\Omega}; f(x) < f(x_0)\} \cap B_{x_0},$$

where B_{x_0} is a ball centered at x_0 . Similarly, we can introduce

$$A_f^{\leq}(x_0) := \{x \in \overline{\Omega}; f(x) \leq f(x_0)\} \cap B_{x_0}.$$

Interior points:

First we observe that, near a non critical point $x_0 \in \Omega$ of f , one can find B_{x_0} and a set of local coordinates such that

$$A_f^<(x_0) = \{y_1 < 0\} \cap B_{x_0}.$$

Secondly, if x_0 is a critical point with index p , then there exists a ball B_{x_0} around x_0 and a set of local coordinates centered at x_0 such that

$$A_f^<(x_0) = \left\{ -\sum_{\ell=1}^p y_\ell^2 + \sum_{\ell=p+1}^n y_\ell^2 < 0 \right\} \cap B_{x_0} ,$$

and

$$A_f^{\leq}(x_0) = \left\{ -\sum_{\ell=1}^p y_\ell^2 + \sum_{\ell=p+1}^n y_\ell^2 \leq 0 \right\} \cap B_{x_0} .$$

We now observe that

1. When $p = 0$ (local minimum), $A_f^<(x_0)$ is empty and $A_f^{\leq}(x_0)$ is reduced to $\{x_0\}$.
2. When $p = 1$, $A_f^<(x_0)$ has two connected components and x_0 belongs to the closure of each of the two components. This property will be crucial in the discussion.
3. When $p \geq 2$, $A_f^<(x_0)$ is (arcwise) connected.

Points on the boundary:

If x_0 belongs to $\partial\Omega$, Assumption 3.1 leads to two cases:

First case.

If x_0 is not a critical point of $f|_{\partial\Omega}$, then the hypersurfaces $\{x \mid f(x) = f(x_0)\}$ and $\partial\Omega$ intersect transversally in a neighborhood of x_0 . Hence there is a ball B_{x_0} around x_0 and a set of local coordinates such that

$$A_f^<(x_0) = \{y_1 < 0, y_n \leq 0\} \cap B_{x_0} ,$$

and

$$A_f^{\leq}(x_0) = \{y_1 \leq 0, y_n \leq 0\} \cap B_{x_0} ,$$

with $\Omega \cap B_{x_0} = \{y_n < 0\} \cap B_{x_0}$.

Second case.

If x_0 is a critical point of $f|_{\partial\Omega}$ with index p and with $\pm \frac{\partial f}{\partial n}(x_0) > 0$, there are local coordinates $(y_1, \dots, y_{n-1}, y_n)$, constructed from the second point of Lemma 3.18, such that (y_1, \dots, y_{n-1}) are Morse coordinates for $f|_{\partial\Omega}$ and such that

$$A_f^<(x_0) = \left\{ \pm y_n - \sum_{i=1}^p y_i^2 + \sum_{i=p+1}^{n-1} y_i^2 < 0 , y_n \leq 0 \right\} \cap B_{x_0} ,$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

and

$$A_f^{\leq}(x_0) = \left\{ \pm y_n - \sum_{i=1}^p y_i^2 + \sum_{i=p+1}^{n-1} y_i^2 \leq 0, y_n \leq 0 \right\} \cap B_{x_0}.$$

These local models allow to see that

1. If x_0 is a local minimum of $f|_{\partial\Omega}$ such that $\frac{\partial f}{\partial n}(x_0) < 0$, then $A_f^<(x_0) = \emptyset$ and $A_f^{\leq}(x_0) = \{x_0\}$.
2. If x_0 is a local minimum of $f|_{\partial\Omega}$ such that $\frac{\partial f}{\partial n}(x_0) > 0$, then $A_f^<(x_0) \cap \partial\Omega = \emptyset$ and $A_f^{\leq}(x_0) \cap \partial\Omega = \{x_0\}$. Moreover, $A_f^<(x_0)$ is connected.
3. If $p = 1$ and $\frac{\partial f}{\partial n}(x_0) < 0$ (i.e. if $x_0 \in \mathcal{U}^{(1)} \cap \partial\Omega$), $A_f^<(x_0)$ has two connected components with a non-empty intersection with $\partial\Omega$ and x_0 belongs to the closure of each of the two components. Again, this property will be crucial in the discussion.
4. In all other cases, $A_f^{\leq}(x_0)$ is connected with a non-empty intersection with $\partial\Omega$.

5.3. Labelling of local minima and first consequence

Remember our main Assumption 1.1:

The function f has $\#\mathcal{U}$ distinct critical values and the quantities $f(U^{(1)}) - f(U^{(0)})$, with $U^{(1)} \in \mathcal{U}^{(1)}$ and $U^{(0)} \in \mathcal{U}^{(0)}$ are distinct.

DEFINITION 5.3. — *For $\lambda \in \mathbb{R}$, we define $H^0(\{f < \lambda\})$ as the number of connected components of the level set $L(\lambda) = f^{-1}((-\infty, \lambda))$.*

Due to local structure of the level sets of a Morse function and to Assumption 1.1, the function $H^0(\{f < \lambda\})$ of $\lambda \in \mathbb{R}$ is a step function which satisfies, with λ decreasing from $+\infty$:

- $H^0(\{f < \lambda\})$ decreases by 1 around every $\lambda = f(U^{(0)})$ with $U^{(0)} \in \mathcal{U}^{(0)}$.
- wherever $H^0(\{f < \lambda\})$ increases by 1, it is around a $\lambda = f(U^{(1)})$ with $U^{(1)} \in \mathcal{U}^{(1)}$.
- $H^0(\{f < \lambda\})$ is locally constant away from those points.

Remark 5.4. — $\overline{\Omega}$ is connected and compact so $H^0(\{f < \lambda\})$ equals respectively 1 or 0 for $\lambda \geq \lambda_f$ or $\lambda \leq -\lambda_f$ for some $\lambda_f > 0$.

Consequently, the previous discussion implies that the number of critical values of f with index 1 where $H^0(\{f < \lambda\})$ increases (by 1) is equal to $m_0 - 1$ and so that $m_1 + 1 \geq m_0$.

We now label the local minima of f as follow:

1. We set $U_1^{(0)} = \min_{x \in \overline{\Omega}} f$, $z_1 = \infty$, $f(z_1) = z_1 = \infty$ and we consider $H^0(\{f < \lambda\})$ for λ decreasing from $f(z_1) = +\infty$.
2. When $U_k^{(0)}$ and z_k are defined for $k = 1, \dots, K - 1$, decrease λ from $f(z_{K-1})$ until $H^0(\{f < \lambda\})$ increases by 1. Denote by λ_K this value.
3. By Assumption 1.1 and by the previous discussion, there exists a unique point in $\mathcal{U}^{(1)}$, that we denote by z_K , satisfying $f(z_K) = \lambda_K$. Then we denote by $U_K^{(0)}$ the global minimum of the new connected component.
4. We iterate 2. and 3. until all the local minima have been considered.
5. At least we permute the k 's to make the sequence $\left(f(z_k) - f(U_k^{(0)})\right)_{k \in \{1, \dots, m_0\}}$ strictly decreasing, which is possible by Assumption 1.1.

DEFINITION 5.5 (*The map j*). — *If the generalized critical points with index 1 are numbered $U_j^{(1)}$, $j = 1, \dots, m_1$, we set $U_1^{(1)} = z_1 = \infty$ and we define the application $k \mapsto j(k)$ from $\{1, \dots, m_0\}$ to $\{0, 1, \dots, m_1\}$ by:*

$$\begin{cases} j(1) = 0 \text{ and } U_{j(1)}^{(1)} = z_1 \\ \forall k \geq 2, U_{j(k)}^{(1)} = z_k . \end{cases}$$

DEFINITION 5.6. — *For $k \in \{1, \dots, m_0\}$, we denote by E_k the connected component of $U_k^{(0)}$ in*

$$f^{-1}((-\infty, f(U_{j(k)}^{(1)})) \setminus \{U_{j(k)}^{(1)}\}) .$$

Remark 5.7. — By the previous construction, $U_k^{(0)}$ is the global minimum of E_k .

PROPOSITION 5.8. — *Under Assumption 1.1, the following properties are satisfied:*

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

a) The sequence $\left(f(U_{j(k)}^{(1)}) - f(U_k^{(0)})\right)_{k \in \{1, \dots, m_0\}}$ is strictly decreasing.

b) $E_1 = \overline{\Omega}$ is compact and for any $k > 1$ the set E_k is a relatively compact subset of $f^{-1}((-\infty, f(U_{j(k)}^{(1)})))$ satisfying $\overline{E_k} = E_k \cup \left\{U_{j(k)}^{(1)}\right\}$.

c) For any $(k, j) \in \{1, \dots, m_0\} \times \{0, 1, \dots, m_1\}$, the relation $U_j^{(1)} \in E_k$ implies:

either $(j = j(k') \text{ for some } k' > k) \text{ or } j \notin j(\{1, \dots, m_0\})$.

d) For any $k \neq k' \in \{1, \dots, m_0\}$, the relation $U_{k'}^{(0)} \in E_k$ implies:

$$\left(k' > k \text{ and } f(U_{k'}^{(0)}) > f(U_k^{(0)})\right).$$

e) The application $j : \{1, \dots, m_0\} \rightarrow \{0, 1, \dots, m_1\}$ is injective.

Proof. — By Assumption 1.1 and by construction, the points **a)**, **b)** and **e)** are obvious.

c) Assume now $U_{j(k')}^{(1)} \in E_k$.

Since $U_{j(k)}^{(1)} \notin E_k$, one has $k \neq k'$. Moreover, by definition of E_k and by Assumption 1.1, we have the inequality $f(U_{j(k')}^{(1)}) < f(U_{j(k)}^{(1)})$ which implies that $E_{k'}$ is contained in E_k , by connectedness of E_k and $E_{k'}$.

Consequently, $U_{k'}^{(0)} \in E_k$ and by Assumption 1.1, $f(U_{k'}^{(0)}) > f(U_k^{(0)})$ (because $U_k^{(0)}$ is the global minimum of f on $\overline{E_k}$) which yields:

$$f(U_{j(k')}^{(1)}) - f(U_{k'}^{(0)}) < f(U_{j(k)}^{(1)}) - f(U_k^{(0)})$$

and the point **a)** gives $k' > k$.

d) Assume $U_{k'}^{(0)} \in E_k$ for $k \neq k'$.

Again one has $f(U_{k'}^{(0)}) > f(U_k^{(0)})$ which implies $k' \neq 1$ (then $U_{j(k')}^{(1)} \in \overline{\Omega}$) and there are two possible cases:

$$U_{j(k')}^{(1)} \in \overline{E_k} \text{ or } U_{j(k')}^{(1)} \notin \overline{E_k}.$$

In the second case, let us look at $E_{k'}$. $E_{k'}$ is connected and $U_{k'}^{(0)}$ is the global minimum of f on $E_{k'}$. Moreover, $U_{k'}^{(0)} \in E_{k'} \cap E_k$ and $U_{j(k')}^{(1)} \in \overline{E_{k'}} \setminus \overline{E_k}$ imply, by connectedness, that $\partial E_k \cap E_{k'} \neq \emptyset$.

$\overline{E_k}$ is then contained in $E_{k'}$ and $U_k^{(0)} \in E_{k'}$, which cannot occur.

Consequently, $U_{j(k')}^{(1)} \in \overline{E_k}$ and the points **b)** and **c)** imply $k' > k$. \square

5.4. Construction of the quasimodes

Like in [HeKlNi] and in [HeNi], we associate with every $U_k^{(0)}$, $k \in \{1, \dots, m_0\}$, a quasimode for $\Delta_{f,h}^{N,(0)}$ which is approximately supported in E_k , while the quasimodes for $\Delta_{f,h}^{N,(1)}$ will be supported in the balls $B(U_j^{(1)}, 2\varepsilon_1)$, for $j \in \{1, \dots, m_1\}$. A ball $B(U, \rho)$, with $U \in \overline{\Omega}$ and $\rho > 0$, is a geodesic ball and the geodesic distance is denoted by d_Ω . The parameter $\varepsilon_1 > 0$ is fixed so that:

- $d_\Omega(U, U') \geq 10 \varepsilon_1$ for $U, U' \in \mathcal{U}$, $U \neq U'$.
- For all $U \in \mathcal{U}$ and all $k \in \{1, \dots, m_0\}$, $U \notin \overline{E_k}$ implies

$$d_\Omega(U, E_k) \geq 10 \varepsilon_1 .$$

- The construction of the WKB approximation of Subsection 4.6 is possible in the ball $B(U_j^{(1)}, 2\varepsilon_1)$. If $U_j^{(1)}$ is a boundary point, this means the introduction of the coordinates (x', x_n) used in Section 4.3 and the existence of Φ . Recall that in these coordinates, Φ and g_0 have the form:

$$\Phi = -x_n + \varphi(x') \text{ and } g_0 = g_{nn}(x) dx_n^2 + \sum_{i,j=1}^{n-1} g_{ij}(x) dx_i dx_j . \quad (5.4.1)$$

The parameter $\varepsilon_1 > 0$ will be kept fixed, while we need another parameter $\varepsilon \in (0, \varepsilon_0)$ which will be fixed in the final step of the proof.

Like in [HeNi], the construction presented in [HeKlNi] has to be adapted when $U_{j(k)}^{(1)} \in \partial\Omega$ or $U_k^{(0)} \in \partial\Omega$ (recall that in [HeNi], the case $U_k^{(0)} \in \partial\Omega$ did not occur) and we focus on these changes.

However, note that in [HeNi] the set $\overline{E_k}$ intersected $\partial\Omega$ at most at one point ($\overline{E_k} \cap \partial\Omega \subset \{U_{j(k)}^{(1)}\}$). It is not the case here and we cannot use the same construction when $U_{j(k)}^{(1)} \in \partial\Omega$.

For every $k \in \{1, \dots, m_0\}$ and $\varepsilon > 0$, we introduce the set:

$$\tilde{\Omega}_k(\varepsilon, \delta) = \left\{ x \in \overline{\Omega}, d_\Omega \left(x, \overline{E_k} \setminus B(U_{j(k)}^{(1)}, \varepsilon) \right) < \delta \right\} \cup B(U_{j(k)}^{(1)}, \varepsilon) ,$$

with $\delta \in (0, \delta_\varepsilon)$, $\delta_\varepsilon > 0$ small enough.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

The cut-off function $\tilde{\chi}_{k,\varepsilon} \in \mathcal{C}_0^\infty(\overline{\Omega})$, $0 \leq \tilde{\chi}_{k,\varepsilon} \leq 1$ is chosen so that:

$$\text{supp } \tilde{\chi}_{k,\varepsilon} \subset \tilde{\Omega}_k(\varepsilon, \delta_\varepsilon) \quad \text{and} \quad \tilde{\chi}_{k,\varepsilon}|_{\tilde{\Omega}_k(\varepsilon, \delta_\varepsilon/2) \setminus B(U_{j(k)}^{(1)}, \varepsilon)} = 1 .$$

Around $U_{j(k)}^{(1)}$, the cut-off function $\tilde{\chi}_{k,\varepsilon}$ is chosen (more accurately below when $U_{j(k)}^{(1)} \in \partial\Omega$) so that $U_{j(k)}^{(1)} \notin \text{supp } \tilde{\chi}_{k,\varepsilon}$ and

$$\forall x \in B(U_{j(k)}^{(1)}, \varepsilon) , \left(\tilde{\chi}_{k,\varepsilon}(x) \neq 0 , \text{ and } f(x) < f(U_{j(k)}^{(1)}) \right) \Rightarrow x \in E_k . \quad (5.4.2)$$

Remark 5.9. — The cut-off functions $\tilde{\chi}_{k,\varepsilon}$ are used in the construction of quasimodes for $\Delta_{f,h}^{N,(0)}$.

Moreover, in the case $k = 1$, we have by construction $\tilde{\chi}_{k,\varepsilon} \equiv 1$ in $\overline{\Omega}$ because $U_{j(1)}^{(1)} \notin \overline{\Omega}$. This case provides directly the eigenvector $\|e^{-f(x)/h}\|^{-1} e^{-f(x)/h}$ (of $\Delta_{f,h}^{N,(0)}$) with the eigenvalue 0.

Like in [HeKlNi] and in [HeNi], we deduce from Proposition 5.8 the following properties for $\tilde{\chi}_{k,\varepsilon}$.

PROPOSITION 5.10. — *By taking $\delta = \delta_\varepsilon$ with $\varepsilon \in (0, \varepsilon_0]$, $0 < \varepsilon_0 \leq \varepsilon_1$ small enough, the cut-off functions $\tilde{\chi}_{k,\varepsilon}$ ($k \in \{1, \dots, m_0\}$) satisfy the following properties:*

- a) *If x belongs to $\text{supp } \tilde{\chi}_{k,\varepsilon}$ and $f(x) < f(U_{j(k)}^{(1)})$, then $x \in E_k$.*
- b) *There exist $C > 0$ and, for any $\varepsilon \in (0, \varepsilon_0]$, a constant $K_\varepsilon > 0$, such that, for $x \in \text{supp } \nabla \tilde{\chi}_{k,\varepsilon}$,*

$$\begin{aligned} \text{either } & x \notin B(U_{j(k)}^{(1)}, \varepsilon) \text{ and } f(U_{j(k)}^{(1)}) + K_\varepsilon^{-1} \leq f(x) \leq f(U_{j(k)}^{(1)}) + K_\varepsilon , \\ \text{or } & x \in B(U_{j(k)}^{(1)}, \varepsilon) \text{ and } \left| f(x) - f(U_{j(k)}^{(1)}) \right| \leq C\varepsilon . \end{aligned}$$

- c) *For any $U \in \mathcal{U}$, $U \neq U_{j(k)}^{(1)}$, the distance $d_\Omega(U, \text{supp } \nabla \tilde{\chi}_{k,\varepsilon})$ is bounded from below by $3\varepsilon_1 > 0$. If in addition $U \in \text{supp } \tilde{\chi}_{k,\varepsilon}$, then $U \in E_k$.*
- d) *If, for some $k' \in \{1, \dots, m_0\}$, $U_{k'}^{(0)}$ belongs to $\text{supp } \tilde{\chi}_{k,\varepsilon}$, then $k' \geq k$ and*

$$f(U_{k'}^{(0)}) > f(U_k^{(0)}), \quad f(U_{j(k')}^{(1)}) \leq f(U_{j(k)}^{(1)}), \quad \text{if } k \neq k' .$$

e) For any $j \in \{1, \dots, m_1\}$, such that $U_j^{(1)} \in \text{supp } \tilde{\chi}_{k,\varepsilon}$,

either $j \notin j(\{1, \dots, m_0\})$,

or $j = j(k')$, for some $k' \geq k$ and $U_{k'}^{(0)} \in \text{supp } \tilde{\chi}_{k,\varepsilon}$.

The quasimodes for $\Delta_{f,h}^{N,(1)}$ associated with the $U_j^{(1)} \in \Omega$ are constructed like in [HeKlNi] and in [HeNi] (and rely on the approximation by the Dirichlet problem in small balls $B(U_j^{(1)}, 2\varepsilon_1)$). We will not recall the complete construction here.

In the same spirit as in [HeNi], the quasimodes associated with the $U_j^{(1)} \in \partial\Omega$ will rely on the approximation by the Neumann realization associated with the neighborhood $\Omega_{U_j^{(1)},\rho}$ ($\rho > 0$ small enough) which was studied in Subsection 4.6.

Once $\rho > 0$ is fixed uniformly for all $U_j^{(1)} \in \partial\Omega$, the parameter $\varepsilon_1 > 0$ is reduced so that $B(U_j^{(1)}, 2\varepsilon_1) \subset \Omega_{U_1,\rho}$ for all $U_j^{(1)} \in \partial\Omega$.

For all $j \in \{1, \dots, m_1\}$, u_j denotes a normalized eigenvector associated with the first (exponentially small) eigenvalue of this Dirichlet or Neumann realization. The cut-off function $\theta_j \in C_0^\infty(B(U_j^{(1)}, 2\varepsilon_1))$ is taken such that $\theta_j = 1$ on $B(U_j^{(1)}, \varepsilon_1)$ and $\frac{\partial\theta_j}{\partial n}\Big|_{\partial\Omega} \equiv 0$ for boundary points $U_j^{(1)} \in \partial\Omega$.

Note that the function $\tilde{\chi}_{k,\varepsilon}$ depends on $\varepsilon \in (0, \varepsilon_0]$, while θ_j is kept fixed like $\varepsilon_1 > 0$.

DEFINITION 5.11. — For cut-off $\chi_{k,\varepsilon}$ satisfying the properties of Proposition 5.10 like $\tilde{\chi}_{k,\varepsilon}$, introduce the following quasimodes.

For any $k \in \{1, \dots, m_0\}$, the (ε, h) -dependent function $\psi_k^{(0)}$ is defined by

$$\psi_k^{(0)}(x) = \left\| \chi_{k,\varepsilon}(x) e^{-(f(x)-f(U_k^{(0)}))/h} \right\|^{-1} \chi_{k,\varepsilon}(x) e^{-(f(x)-f(U_k^{(0)}))/h}.$$

For any $j \in \{1, \dots, m_1\}$, the h -dependent 1-form $\psi_j^{(1)}$ is defined by

$$\psi_j^{(1)}(x) = \left(\|\theta_j u_j\|^{-1} \right) \theta_j(x) u_j(x).$$

We set $\lambda_1^{\text{app}}(\varepsilon, h) = 0$, and for any $k \in \{2, \dots, m_0\}$:

$$\lambda_k^{\text{app}}(\varepsilon, h) = \left| \left\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \right\rangle \right|^2.$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Remark 5.12. —

a) In the case $U_{j^{(k)}}^{(1)} \in \Omega$, $\chi_{k,\varepsilon}$ is $\tilde{\chi}_{k,\varepsilon}$ with additional properties (see [HeKlNi] for details) and we will still denote it here by $\tilde{\chi}_{k,\varepsilon}$. In the case $U_{j^{(k)}}^{(1)} \in \partial\Omega$, the real choice of $\chi_{k,\varepsilon}$ will be fixed further (see Definition 5.20). Moreover, $\chi_{k,\varepsilon}$ also satisfies the properties of Proposition 5.10.

b) For the sake of conciseness, we omit the (ε, h) - and h -dependence in the notations $\psi_k^{(0)}$ and $\psi_j^{(1)}$.

c) We will show in the next section that the $\lambda_k^{app}(\varepsilon, h)$'s are approximated values of the small eigenvalues of $\Delta_{f,h}^{N,(0)}$.

By Remark 5.9, this definition is coherent for $k = 1$ and $\psi_1^{(0)}$ is the normalized eigenvector associate with the eigenvalue 0.

d) Due to the condition $\frac{\partial\theta_j}{\partial n} \Big|_{\partial\Omega} \equiv 0$, $\psi_j^{(1)}$ belongs to $D(\Delta_{f,h}^{N,(1)})$ and this, even if $U_j^{(1)}$ belongs to $\partial\Omega$.

5.5. Quasimodal estimates

We end this section by reviewing the quasimodal estimates which are derived from Propositions 5.8 and 5.10. The asymptotic expansion of the quantity $\langle \psi_{j^{(k)}}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \rangle$ has also be done in [HeKlNi] when $U_k^{(0)}$ and $U_{j^{(k)}}^{(1)} \in \Omega$ are interior points. Like in [HeNi], we will simply complete this analysis by establishing the asymptotic expansion of $\langle \psi_{j^{(k)}}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \rangle$, when $U_k^{(0)}$ or $U_{j^{(k)}}^{(1)}$ is in $\partial\Omega$.

Remark 5.13. — In this subsection, we make computations with different coordinate systems $v = (v_1, \dots, v_n)$ (around $U = U_k^{(0)}$ or $U = U_{j^{(k)}}^{(1)}$) all given given by Lemma 3.18.

Looking at the proof of Lemma 3.18 given in [Lep2], notice that the coordinates (v_1, \dots, v_{n-1}) in the boundary can be chosen freely. Moreover, according to [HeSj4] pp. 279-280, they can be chosen such that $dv_1(U), \dots, dv_{n-1}(U), \vec{n}_U^*$ is orthonormal and positively oriented and

$$f(v, 0) = \frac{\lambda_1}{2} v_1^2 + \dots + \frac{\lambda_{n-1}}{2} v_{n-1}^2 + f(U) \text{ and } \varphi(v) = \frac{|\lambda_1|}{2} v_1^2 + \dots + \frac{|\lambda_{n-1}|}{2} v_{n-1}^2, \quad (5.5.1)$$

with $\lambda_1 < 0$ when $U = U_j^{(1)}$. Hence all the coordinate systems around $U \in \partial\Omega$ will coincide on $\partial\Omega$ while they may differ in Ω according to the case when a normal form is used for f , Φ or $f + \Phi$ in Ω .

Remind that the parameter $\varepsilon_1 > 0$ is fixed, while ε_0 and $\varepsilon \in (0, \varepsilon_0]$ may have to be adapted during the proof. **We shall denote by α a generic positive constant which is independent of $\varepsilon \in (0, \varepsilon_0]$.**

Introduce the following notation which will be very useful:

DEFINITION 5.14. — *The notation $g(h) = \mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}})$ means that, for all $\varepsilon \in (0, \varepsilon_0]$, there exists a constant $C_\varepsilon > 0$ such that:*

$$\forall h \in (0, h_0], \quad |g(h)| \leq C_\varepsilon e^{-\frac{\alpha}{h}}.$$

From Proposition 5.8-d) and the good localization of $\nabla \chi_{k,\varepsilon}$, we deduce the following estimates for $\psi_k^{(0)}$.

PROPOSITION 5.15. — *The system of (ε, h) -dependent functions $(\psi_k^{(0)})_{k \in \{1, \dots, m_0\}}$ of Definition 5.11 is almost orthogonal with*

$$\left(\langle \psi_k^{(0)} \mid \psi_{k'}^{(0)} \rangle \right)_{k, k' \in \{1, \dots, m_0\}} = \text{Id}_{\mathbb{C}^{m_0}} + \mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}}),$$

and there exists $\alpha > 0$ and, for any $\varepsilon \in (0, \varepsilon_0]$, $C(\varepsilon)$ and $h_0(\varepsilon)$ such that, for any $h \in (0, h_0(\varepsilon)]$,

$$\langle \Delta_{f,h}^{N,(0)} \psi_k^{(0)} \mid \psi_k^{(0)} \rangle = \left\| d_{f,h}^{(0)} \psi_k^{(0)} \right\|^2 \leq C(\varepsilon) e^{-2 \frac{f(U_j^{(1)}) - f(U_k^{(0)}) - \alpha \varepsilon}{h}}.$$

COROLLARY 5.16. — *There exists $\varepsilon_0 > 0$ and $\alpha > 0$ such that, for any choice of ε in $(0, \varepsilon_0]$ and for all $k \in \{1, \dots, m_0\}$, the (ε, h) -dependent quasi-modes $\psi_k^{(0)}$ satisfy the estimate*

$$\langle \Delta_{f,h}^{N,(0)} \psi_k^{(0)} \mid \psi_k^{(0)} \rangle = \mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}}).$$

The exponential decay of the first eigenvector u_j , associated with an exponentially small eigenvalue, of the Dirichlet realization of $\Delta_{f,h}^{(1)}$ around $U_j^{(1)}$, provides the following estimates for $\psi_j^{(1)}$. We refer the reader to [HeKINi] or [HeSj4] for $U_j^{(1)} \in \Omega$ and to Subsection 4.6 for $U_j^{(1)} \in \partial\Omega$.

PROPOSITION 5.17. — *The system of h -dependent 1-forms, $(\psi_j^{(1)})_{j \in \{1, \dots, m_1\}}$ given in Definition 5.11 is orthonormal and there exists $\alpha > 0$ independent of ε such that*

$$\langle \Delta_{f,h}^{N,(1)} \psi_j^{(1)} \mid \psi_j^{(1)} \rangle = \mathcal{O}(e^{-\frac{\alpha}{h}}),$$

for all $j \in \{1, \dots, m_1\}$.

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Let us now compute some asymptotic expansions.

PROPOSITION 5.18. — For k in $\{2, \dots, m_0\}$ and x in $\overline{\Omega}$,

$$\psi_k^{(0)}(x) = \gamma_k(h)(1 + a_k(h))\chi_{k,\varepsilon}(x)e^{-\frac{f(x)-f(U_k^{(0)})}{h}},$$

where $\gamma_k(h)$ is defined in Definition 1.2 and $a_k(h) \sim \sum_{\ell=1}^{\infty} a_{k,\ell}h^\ell$.

Proof. — In the case $U_k^{(0)} \in \Omega$, we refer the reader to [HeKINi].

If $U_k^{(0)} \in \partial\Omega$, we use again, in a neighborhood of $U_k^{(0)}$, the coordinate system (\bar{x}', \bar{x}_n) introduced in the second part of the Section 3.3 (with $\bar{x}(U_k^{(0)}) = 0$). In this coordinate system, f and g_0 equal:

$$f(\bar{x}) = -\bar{x}_n + f|_{\partial\Omega}(\bar{x}') = -\bar{x}_n + f(U_k^{(0)}) + \varphi(\bar{x}'), \quad (5.5.2)$$

$$g_0 = g_{nn}(\bar{x}) d\bar{x}_n^2 + \sum_{i,j=1}^{n-1} g_{ij}(\bar{x}) d\bar{x}_i d\bar{x}_j, \quad (5.5.3)$$

where $\varphi = f|_{\partial\Omega} - f(U_k^{(0)})$ is the Agmon distance to $U_k^{(0)}$ on the boundary. We denote by $V_{g_0}(d\bar{x})$ the normalized volume form:

$$V_{g_0}(d\bar{x}) = (\det G_0(\bar{x}))^{1/2} d\bar{x}' \wedge d\bar{x}_n =: \nu(\bar{x}', \bar{x}_n) d\bar{x}' \wedge d\bar{x}_n.$$

From (5.5.2),

$$d\bar{x}_n(U_k^{(0)}) = -\frac{\partial f}{\partial n}(U_k^{(0)}) \vec{n}_{U_k^{(0)}}^* \text{ and } \nu(0,0) = \left(-\frac{\partial f}{\partial n}(U_k^{(0)})\right)^{-1}. \quad (5.5.4)$$

For some constants $\eta > 0$ and $\delta_\eta > 0$,

$$\begin{aligned} \left\| \chi_{k,\varepsilon} e^{-\frac{f(x)-f(U_k^{(0)})}{h}} \right\|^2 &= \int_{\overline{\Omega}} \chi_{k,\varepsilon}^2 e^{-2\frac{f(x)-f(U_k^{(0)})}{h}} V_{g_0}(dx) \\ &= \int_{B(0,\eta)} e^{2\frac{\bar{x}_n}{h}} e^{-2\frac{\varphi(\bar{x}')}{h}} \nu(\bar{x}', \bar{x}_n) d\bar{x}' \wedge d\bar{x}_n + \mathcal{O}(e^{-\frac{\delta_\eta}{h}}). \end{aligned}$$

According to (5.5.1),

$$\begin{aligned} \left\| \chi_{k,\varepsilon} e^{-\frac{f(x)-f(U_k^{(0)})}{h}} \right\|^2 &= \int_{B(0,\eta)} e^{2\frac{\bar{x}_n}{h}} e^{-\frac{|\lambda_1|\bar{x}_1^2 + \dots + |\lambda_{n-1}|\bar{x}_{n-1}^2}{h}} \nu(\bar{x}', \bar{x}_n) d\bar{x} \wedge d\bar{x}_n \\ &\quad + \mathcal{O}(e^{-\frac{\delta_\eta}{h}}). \end{aligned}$$

By expanding $\nu(\bar{x}', \bar{x}_n)$ to a Taylor Series of arbitrary order $k \in \mathbb{N}^*$, we can separate the variables \bar{x}' and \bar{x}_n in the last integral term.

Hence, using the Laplace Method for each term, we obtain an asymptotic expansion of arbitrary order of $\left\| \chi_{k,\varepsilon} e^{-\frac{f(x) - f(U_k^{(0)})}{h}} \right\|^2$.

Moreover, from (5.5.4), the first term is:

$$\left(-\frac{\partial f}{\partial n}(U_k^{(0)}) \right)^{-1} \frac{h}{2} \frac{(\pi h)^{\frac{n-1}{2}}}{\left| \det \text{Hess } f|_{\partial\Omega}(U_k^{(0)}) \right|^{\frac{1}{2}}} = (\gamma_k(h))^{-2}.$$

□

PROPOSITION 5.19. — In $B(U_{j(k)}^{(1)}, \varepsilon_1)$, choose the coordinate system x which satisfies (5.4.1) and (5.5.1) with $\lambda_1 < 0$. For k in $\{2, \dots, m_0\}$, the equality

$$\psi_{j(k)}^{(1)}(x) = \delta_{j(k)}(h) b(x, h) e^{-\frac{\Phi(x)}{h}},$$

holds up to a phase factor, when $\delta_{j(k)}(h)$ is defined according to Definition 1.2, $b(x, h) \sim \sum_{\ell=0}^{\infty} b_{k,\ell}(x) h^\ell$, $b_{k,\ell}(x) = \sum_{i=1}^n b_{k,\ell}^i(x) dx_i$, and $b_{k,0}^i(0) = \delta_{1i}$.

Proof. — In Section 4, we found a WKB approximation u_1^{wkb} of an eigenvector u_1^h such that,

$$\begin{aligned} e^{\frac{\Phi(x)}{h}} u_1^{wkb} &= \sum_{i=1}^n a_i^0(x) dx_i + h a^1(x, h), \\ a_i^0(0) &= \delta_{1i}, \quad a^1(x, h) \sim \sum_{\ell} h^\ell a_\ell(x), \end{aligned}$$

and

$$\forall x \in B(U_{j(k)}^{(1)}, 2\varepsilon_1), \quad e^{\frac{\Phi(x)}{h}} \left| \partial_x^\alpha (u_1^h(x) - u_1^{wkb}(x)) \right| \leq C_{\alpha,N} h^N.$$

The WKB approximation u_1^{wkb} was initially constructed in another coordinate system $(\underline{x}_1, \dots, \underline{x}_n)$. Remark 5.13 recalls that the tangential coordinates $\underline{x}_1, \dots, \underline{x}_{n-1}$ and x_1, \dots, x_{n-1} can coincide in $\partial\Omega$ with different deformations as entering into Ω .

The normalized eigenvector that we take here is

$$u_{j(k)} = \frac{u_1^h}{\|u_1^h\|}.$$

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

Let us first compute accurately:

$$\|u_1^h\| = \|\theta_{j(k)}u_1^h\| + \mathcal{O}(h^\infty) = \|\theta_{j(k)}u_1^{wkb}\| + \mathcal{O}(h^\infty).$$

Moreover,

$$\|\theta_{j(k)}u_1^{wkb}\|^2 = \int \theta_{j(k)}(x)^2 \langle a(x, h) | a(x, h) \rangle e^{-\frac{2\Phi(x)}{h}} V_{g_0}(dx),$$

where the integral is over $x_n \leq 0$. Note furthermore that,

$$dx_n(U_{j(k)}^{(1)}) = -\frac{\partial\Phi}{\partial n}(U_{j(k)}^{(1)}) \vec{n}_{U_{j(k)}^{(1)}}^* = -\frac{\partial f}{\partial n}(U_{j(k)}^{(1)}) \vec{n}_{U_{j(k)}^{(1)}}^*.$$

Proceeding like in the proof of Proposition 5.18, we obtain, using the Laplace method, a full asymptotic expansion of $\|\theta_{j(k)}u_1^{wkb}\|^2$. The first term is given by the first term of

$$\int \theta_{j(k)}(x)^2 \langle a^0(x) | a^0(x) \rangle e^{\frac{2x_n}{h}} e^{-2\frac{\varphi(x')}{h}} V_{g_0}(dx),$$

and from $\langle a^0(x) | a^0(x) \rangle(0) = 1$, we conclude like in the proof of Proposition 5.18. \square

Before stating the next result, let us specify the choice of $\chi_{k,\varepsilon}$ when $U_{j(k)}^{(1)} \in \partial\Omega$. We assume $\varepsilon \in (0, \varepsilon_0)$, with $0 < \varepsilon < \frac{\varepsilon_1}{10}$. We introduce locally near $U_{j(k)}^{(1)}$ a new coordinate system $(\tilde{x}_1, \dots, \tilde{x}_n)$ by application of Lemma 3.18 with $f_1 = f + \Phi$ and $\alpha = (f + \Phi)|_{\partial\Omega}$.

Hence, we can write in $B(U_{j(k)}^{(1)}, 2\varepsilon_1)$, choosing ε_1 small enough:

$$(f + \Phi)(\tilde{x}) = -\tilde{x}_n + (f + \Phi)|_{\partial\Omega}(\tilde{x}') = -\tilde{x}_n + f(\tilde{x}', 0) + \varphi(\tilde{x}')$$

with an arbitrary choice of \tilde{x}' in the boundary.

Remark moreover that in this case,

$$d\tilde{x}_n(U_{j(k)}^{(1)}) = -2\frac{\partial f}{\partial n}(U_{j(k)}^{(1)}) \vec{n}_{U_{j(k)}^{(1)}}^*.$$

We choose the coordinate system \tilde{x}' in the boundary like it was chosen in the boundaryless case (see [HeSj4][HeKlNi]) according to the geometry of stable and unstable manifolds in order to write $(f + \Phi)|_{\partial\Omega}$ as a function of $n - 2$ coordinates:

$$(f + \Phi)|_{\partial\Omega}(\tilde{x}') = f(\tilde{x}', 0) + \varphi(\tilde{x}') = (f + \Phi)|_{\partial\Omega}(\tilde{x}_2, \dots, \tilde{x}_{n-1}). \quad (5.5.5)$$

DEFINITION 5.20. — For any $k \in 1, \dots, m_0$ we define the cut-off $\chi_{k,\varepsilon}$ by:

- If $U_{j(k)}^{(1)} \in \Omega$, $\chi_{k,\varepsilon} = \tilde{\chi}_{k,\varepsilon}$.
- If $U_{j(k)}^{(1)} \in \partial\Omega$, we first construct near $\partial\Omega \cap E_k$ the cut-off $\chi_{k,\varepsilon}^{\partial\Omega}$ like it was constructed in the boundaryless case (see [HeKlNi] pp. 26-29).

Then, choosing a cut-off

$$\chi_n(\tilde{x}_n) \in \mathcal{C}_0^\infty(\mathbb{R}_-), \quad \chi_n = 1 \text{ on } (-\delta_\varepsilon, 0]$$

we take for $\chi_{k,\varepsilon}$:

$$\chi_{k,\varepsilon}(\tilde{x}) = \chi_n(\tilde{x}_n)\chi_{k,\varepsilon}^{\partial\Omega} + (1 - \chi_n(\tilde{x}_n))\tilde{\chi}_{k,\varepsilon}.$$

Note that $\chi_{k,\varepsilon}$, for δ_ε small enough, satisfies the same properties as $\tilde{\chi}_{k,\varepsilon}$ in Proposition 5.10 and we make that choice. Moreover, according to [HeKlNi] p. 28, in a neighborhood of $\{\tilde{x}_1 = 0\} \cap \partial\Omega$, the cut-off $\chi_{k,\varepsilon}$ only depends on \tilde{x}_1 : $\chi_{k,\varepsilon} = \chi_{k,\varepsilon}(\tilde{x}_1)$.

PROPOSITION 5.21. — There exist ε_0 and sequences $(c_{k,m})_{m \in \mathbb{N}^*}$, such that the (ε, h) -dependent and h -dependent quasimodes $\psi_k^{(0)}$ and $\psi_j^{(1)}$ ($(k, j) \in \{1, \dots, m_0\} \times \{1, \dots, m_1\}$ and $\varepsilon \in (0, \varepsilon_0)$) satisfy:

$$|\langle \psi_j^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \rangle| = 0 \quad \text{if } j \neq j(k),$$

$$|\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \rangle| = \gamma_k(h) \delta_{j(k)}(h) \theta_{j(k)}(h) e^{-\frac{f(U_{j(k)}^{(1)}) - f(U_k^{(0)})}{h}} (1 + hc_k^1(h))$$

where $\gamma_k(h)$, $\delta_{j(k)}(h)$, and $\theta_{j(k)}(h)$ are defined in Definition 1.2 and $c_k(h) \sim \sum_{\ell=0}^\infty c_{k,\ell} h^\ell$.

Proof. — The first statement for $j \neq j(k)$ is a consequence of our choice of $\varepsilon_1 > 0$ and $\chi_{k,\varepsilon}$ which gives according to Proposition 5.10-c) $\text{supp } \psi_j^{(1)} \cap \text{supp } \nabla \chi_{k,\varepsilon} = \emptyset$. We conclude with $d_{f,h}^{(0)} \psi_k^{(0)} = C_{\varepsilon,h} (d^{(0)} \chi_{k,\varepsilon}) e^{-f/h}$.

The second case was completely treated in [HeKlNi] when $U_{j(k)}^{(1)} \in \Omega$ and $U_k^{(0)} \in \Omega$. Moreover, in the case when $U_{j(k)}^{(1)} \in \Omega$ and $U_k^{(0)} \in \partial\Omega$, the proof done in [HeKlNi] remains valid if we take the convenient $\gamma_k(h)$.

Show now the cases when $U_{j(k)}^{(1)} \in \partial\Omega$ and $U_k^{(0)} \in \Omega \cup \partial\Omega$ by adapting the proofs done in [HeKlNi] and [HeNi].

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

From Proposition 5.18, Proposition 5.19, and

$$d_{f,h}^{(0)} \left(\chi_{k,\varepsilon} e^{-\frac{f(x)}{h}} \right) = e^{-\frac{f(x)}{h}} h d^{(0)} \chi_{k,\varepsilon},$$

we obtain the existence, for any $\varepsilon > 0$, of $\sigma_\varepsilon > 0$ such that

$$\begin{aligned} \left\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \right\rangle &= h \gamma_k(h) \delta_{j(k)}(h) \\ &\times \int_{B(U_{j(k)}^{(1)}, \varepsilon)} \langle b(x, h) \mid d\chi_{k,\varepsilon} \rangle(x) e^{-\frac{(\Phi(x)+f(x)-f(U_k^{(0)}))}{h}} V_{g_0}(dx) \\ &+ \mathcal{O}_\varepsilon \left(e^{-\frac{f(U_{j(k)}^{(1)})-f(U_k^{(0)})+\sigma_\varepsilon}{h}} \right), \end{aligned}$$

with $b(x, h)$ defined in Proposition 5.19.

Using the coordinate system \tilde{x} , with the choice of $\chi_{k,\varepsilon}$,

$$\begin{aligned} \left\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \right\rangle &= h \gamma_k(h) \delta_{j(k)}(h) \\ &\times \int_{\widetilde{B(U_{j(k)}^{(1)}, \varepsilon)}} \langle b(\tilde{x}, h) \mid d\chi_{k,\varepsilon} \rangle(\tilde{x}) e^{-\frac{-\tilde{x}_n + \varphi(\tilde{x}') + f(\tilde{x}', 0) - f(U_k^{(0)})}{h}} V_{g_0}(d\tilde{x}) \\ &+ \mathcal{O}_\varepsilon \left(e^{-\frac{f(U_{j(k)}^{(1)})-f(U_k^{(0)})+\sigma_\varepsilon}{h}} \right) \\ &= r(h) \int_{C_\varepsilon} \langle b(\tilde{x}, h) \mid d\tilde{x}_1 \rangle \chi'_{k,\varepsilon}(\tilde{x}_1) e^{\frac{\tilde{x}_n - \varphi(\tilde{x}') - (f(\tilde{x}', 0) - f(U_{j(k)}^{(1)}))}{h}} V_{g_0}(d\tilde{x}) \\ &+ \mathcal{O}_\varepsilon \left(e^{-\frac{f(U_{j(k)}^{(1)})-f(U_k^{(0)})+\sigma'_\varepsilon}{h}} \right), \end{aligned}$$

where

$$r(h) = h \gamma_k(h) \delta_{j(k)}(h) e^{\frac{f(U_k^{(0)})-f(U_{j(k)}^{(1)})}{h}}$$

and C_ε is a cylinder $|\tilde{x}'| < c_\varepsilon$, $-c_\varepsilon < \tilde{x}_n < 0$. Expanding $\langle b(\tilde{x}, h) \mid d\tilde{x}_1 \rangle$ to a Taylor Series (of arbitrary order), we can obtain, using the Laplace method, an asymptotic expansion (of arbitrary order) for $\left\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \right\rangle$.

Moreover, the first term in the expansion of $\langle b(\tilde{x}, h) \mid d\tilde{x}_1 \rangle$ equals at $\tilde{x} = 0$, $\langle b_{k,0}(\tilde{x}) \mid d\tilde{x}_1 \rangle(0) = 1$. After recalling (5.5.5) which says that the

exponent $f(\tilde{x}', 0) + \varphi(x')$ does not depend on \tilde{x}_1 , the first term of the wanted expression is then given by

$$r(h) \int e^{\frac{\tilde{x}_n}{h}} d\tilde{x}_n \int e^{-\frac{\varphi(\tilde{x}') + (f(\tilde{x}', 0) - f(U_{j(k)}^{(1)}))}{h}} d\tilde{x}_2 \dots d\tilde{x}_{n-1} \int \chi'_{k,\varepsilon}(\tilde{x}_1) d\tilde{x}_1.$$

Using the Laplace method and

$$\int_{\mathbb{R}} \chi'_{k,\varepsilon}(x_1) dx_1 = -1,$$

we find

$$r(h) \frac{h}{2 \frac{\partial f}{\partial n}(U_{j(k)}^{(1)})} \frac{|\widehat{\lambda}_1^{\partial\Omega}(U_{j(k)}^{(1)})|^{\frac{1}{2}}}{|\det \text{Hess } f|_{\partial\Omega}(U_{j(k)}^{(1)})|^{\frac{1}{2}}} (\pi h)^{\frac{n-2}{2}}.$$

□

6. Final proof

6.1. Main result

Recall first some notations.

The generalized critical points with index 0, $\{U_k^{(0)}, k \in \{1, \dots, m_0\}\}$, are labelled according to Subsection 5.3 and the generalized critical points with index 1, $\{U_{j(k)}^{(1)}, k \in \{2, \dots, m_0\}\}$, are those introduced in Definition 5.5.

Moreover, the quantity $\lambda_k^{app}(\varepsilon, h)$ introduced in Definition 5.11 is associated with the quasimodes $\psi_k^{(0)}$ and $\psi_{j(k)}^{(1)}$:

$$\lambda_k^{app}(\varepsilon, h) = \left| \left\langle \psi_{j(k)}^{(1)} \mid d_{f,h}^{(0)} \psi_k^{(0)} \right\rangle \right|^2.$$

At a generalized critical point U with index i ($i \in \{0, 1\}$), the Hessians $\text{Hess } f(U)$ or $\text{Hess } f|_{\partial\Omega}$ are computed in orthonormal coordinates for the metric g_0 , while considering only the tangential coordinates $x' = (x_1, \dots, x_{n-1})$ for the second case.

At least, for a generalized critical point $U \in W$ with index 1 for $W = \Omega$ or $W = \partial\Omega$, $\widehat{\lambda}_1^W(U)$ denotes the negative eigenvalue of $\text{Hess } f|_W(U)$.

With these notations, we have the following theorem, which implies Theorem 1.3:

Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian

THEOREM 6.1. — *Under Assumptions 3.1 and 1.1, the first eigenvalue $\lambda_1(h)$ of $\Delta_{f,h}^{N,(0)}$ is 0 and its $m_0 - 1$ first non zero eigenvalues $\lambda_2(h), \dots, \lambda_{m_0}(h)$ admit the following asymptotic expansion. There exist $\varepsilon_0 > 0$ and $\alpha > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0]$,*

$$\forall k \in \{2, \dots, m_0\}, \quad \lambda_k(h) = \lambda_k^{app}(\varepsilon, h) \left(1 + \mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}})\right).$$

Recall also that, from Proposition 5.21, for any $\varepsilon \in (0, \varepsilon_0]$,

$$\lambda_k^{app}(\varepsilon, h) = \gamma_k^2(h) \delta_{j(k)}^2(h) \theta_{j(k)}^2(h) e^{-2 \frac{f(U_{j(k)}^{(1)}) - f(U_k^{(0)})}{h}} \left(1 + hc_k^1(h)\right)$$

where $\gamma_k(h)$, $\delta_{j(k)}(h)$, and $\theta_{j(k)}(h)$ are defined in Definition 1.2 and $c_k^1(h)$ admits a complete expansion: $c_k^1(h) \sim \sum_{m=0}^{\infty} h^m c_{k,m}$.

6.2. Finite dimensional reduction and final proof

Set first, for $\ell \in \{0, 1\}$:

$$\forall i \in \{1, \dots, m_\ell\}, \quad v_i^{(\ell)} = 1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(\ell)}) \psi_i^{(\ell)}, \quad (6.2.1)$$

where the $\psi_i^{(\ell)}$ are the (ε, h) - and h - dependent quasimodes introduced in Definition 5.11.

Remark 6.2. — Note that here again we omit the (ε, h) -dependence (resp. h -dependence) of the functions $v_k^{(0)}$ (resp. 1-forms $v_j^{(1)}$) in the notation.

Recall furthermore the definition of the space $F^{(\ell)}$ given in introduction ($\ell \in \{0, 1\}$),

$$F^{(\ell)} = \text{Ran } 1_{[0, h^{\frac{3}{2}})}(\Delta_{f,h}^{(\ell)}),$$

which has dimension m_ℓ according to Theorem 3.5.

According to Lemma 2.11, Corollary 5.16 (for $\ell = 0$) and Proposition 5.17 (for $\ell = 1$), $\left\| 1_{[h^{3/2}, +\infty)}(\Delta_{f,h}^{N,(\ell)}) \psi_i^{(\ell)} \right\|$ is estimated from above by $\mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}})$, which implies the two following propositions:

PROPOSITION 6.3. — *For $\ell \in \{0, 1\}$, the ℓ -forms $(v_i^{(\ell)})_{i \in \{1, \dots, m_\ell\}}$ satisfy:*

$$\left\| v_i^{(\ell)} - \psi_i^{(\ell)} \right\| = \mathcal{O}_\varepsilon(e^{-\frac{\alpha}{h}})$$

for some $\alpha > 0$ independent of $\varepsilon \in (0, \varepsilon_0]$.

PROPOSITION 6.4. — For $\ell \in \{0, 1\}$, the system $(v_i^{(\ell)})_{i \in \{1, \dots, m_\ell\}}$ is a basis of $F^{(\ell)}$ satisfying:

$$V^{(\ell)} := \left(\langle v_i^{(\ell)} | v_{i'}^{(\ell)} \rangle \right)_{i, i' \in \{1, \dots, m_\ell\}} = \text{Id}_{\mathbb{C}^{m_\ell}} + \mathcal{O}_\varepsilon(e^{-\frac{\alpha'}{h}}),$$

for some $\alpha > 0$ independent of $\varepsilon \in (0, \varepsilon_0]$.

Finally, we can also establish:

PROPOSITION 6.5. — There exist $\varepsilon'_0 > 0$ and $\alpha' > 0$ such that, for all $\varepsilon \in (0, \varepsilon'_0]$, the estimates

$$\left| \langle v_j^{(1)} | d_{f,h}^{(0)} v_k^{(0)} \rangle \right| \leq C_\varepsilon e^{-\frac{f(U_{j(k)}^{(1)}) - f(U_k^{(0)}) + \alpha'}{h}}, \quad \text{if } j \neq j(k),$$

and

$$\langle v_{j(k)}^{(1)} | d_{f,h}^{(0)} v_k^{(0)} \rangle = \langle \psi_{j(k)}^{(1)} | d_{f,h}^{(0)} \psi_k^{(0)} \rangle \left(1 + \mathcal{O}_\varepsilon(e^{-\frac{\alpha'}{h}}) \right),$$

hold for all $(k, j) \in \{1, \dots, m_0\} \times \{1, \dots, m_1\}$.

Proof. — Remark first, $1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(1)})$ being a spectral projector and using Corollary 2.10:

$$\begin{aligned} \langle v_j^{(1)} | d_{f,h}^{(0)} v_k^{(0)} \rangle &= \langle 1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(1)}) v_j^{(1)} | d_{f,h}^{(0)} 1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(0)}) \psi_k^{(0)} \rangle \\ &= \langle 1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(1)}) v_j^{(1)} | 1_{[0, h^{3/2})}(\Delta_{f,h}^{N,(1)}) d_{f,h}^{(0)} \psi_k^{(0)} \rangle = \langle v_j^{(1)} | d_{f,h}^{(0)} \psi_k^{(0)} \rangle. \end{aligned}$$

The end of the proof is a straightforward consequence of Proposition 5.15, which gives

$$\left\| d_{f,h}^{(0)} \psi_k^{(0)} \right\| \leq C_\varepsilon e^{-\frac{f(U_{j(k)}^{(1)}) - f(U_k^{(0)}) - \alpha'' \varepsilon}{h}},$$

Propositions 5.21 and 6.3. \square

Proof of Theorem 6.1. — By Propositions 6.4 and 6.5, the bases $(v_i^{(\ell)})_{i \in \{1, \dots, m_\ell\}}$ of $F^{(\ell)}$, for $\ell \in \{0, 1\}$, satisfy Assumptions 2.1 and 2.2 of [Lep1]. Theorem 2.3 of [Lep1] then implies Theorem 6.1 (which immediately implies Theorem 1.3).

Remark 6.6. — The conditions of [Lep1] are not exactly satisfied here because the one to one map j should act from $\{1, \dots, m_0\}$ to $\{1, \dots, m_1\}$, with $\dim F^{(i)} = m_i$.

We can easily reduce the study to this last case, by setting:

$$\overline{m}_0 = m_0 \quad , \quad \overline{m}_1 = m_1 + 1 ,$$

and,

$$\overline{F^{(0)}} = F^{(0)} \quad , \quad \overline{F^{(1)}} = F^{(1)} \oplus^\perp \mathcal{C}v_{m_1+1}^{(1)} .$$

Setting in addition $j(1) = m_1 + 1$ instead of $j(1) = 0$, the conditions of [Lep1] are fulfilled.

Note furthermore that the decreasing sequence $(\alpha_k)_{k \in \{1, \dots, \overline{m}_0\}}$ of [Lep1] is then here $\left(f(U_{j(k)}^{(1)}) - f(U_k^{(0)}) \right)_{k \in \{1, \dots, \overline{m}_0\}}$ whose first term is by definition $+\infty$.

□

Acknowledgement. — The author would like to thank T. Jecko and F. Nier for profitable discussions.

Bibliography

- [Bis] BISMUT (J.M.). — The Witten complex and the degenerate Morse inequalities. *J. Differ. Geom.* 23, p. 207-240 (1986).
- [BoEcGaKl] BOVIER (A.), ECKHOFF (M.), GAYRARD (V.), and KLEIN (M.). — Metastability in reversible diffusion processes I: Sharp asymptotics for capacities and exit times. *JEMS* 6 (4), p. 399-424 (2004).
- [BoGaKl] BOVIER (A.), GAYRARD (V.), and KLEIN (M.). — Metastability in reversible diffusion processes II: Precise asymptotics for small eigenvalues. *JEMS* 7 (1), p. 69-99 (2004).
- [Bur] BURGHELEA (D.). — Lectures on Witten-Helffer-Sjöstrand theory. *Gen. Math.* 5, p. 85-99 (1997).
- [ChLi] CHANG (K.C.) and LIU (J.). — A cohomology complex for manifolds with boundary. *Topological Methods in Non Linear Analysis*, Vol. 5, p. 325-340 (1995).
- [CyFrKiSi] CYCON (H.L.), FROESE (R.G.), KIRSCH (W.), and SIMON (B.). — Schrödinger operators with application to quantum mechanics and global geometry. *Text and Monographs in Physics*, Springer Verlag, 2nd corrected printing (2008).
- [CoPaYc] COLIN DE VERDIÈRE (Y.), PAN (Y.), and YCART (B.). — Singular limits of Schrödinger operators and Markov processes. *J. Operator Theory* 41, No. 1, p. 151-173 (1999).
- [DiSj] DIMASSI (M.) and SJÖSTRAND (J.). — Spectral Asymptotics in the semiclassical limit. *London Mathematical Society, Lecture Note Series* 268, Cambridge University Press (1999).
- [Duf] DUFF (G.F.D.). — Differential forms in manifolds with boundary. *Ann. of Math.* 56, p. 115-127 (1952).

- [DuSp] DUFF (G.F.D.) and SPENCER (D.C.). — Harmonic tensors on Riemannian manifolds with boundary. *Ann. of Math.* 56, p. 128-156 (1952).
- [FrWe] FREIDLIN (M.I.) and WENTZELL (A.D.). — Random perturbations of dynamical systems. Transl. from the Russian by Joseph Szuacs. 2nd ed. *Grundlehren der Mathematischen Wissenschaften*, 260, New York (1998).
- [GaHuLa] GALLOT (S.), HULIN (D.), and LAFONTAINE (J.) *Riemannian Geometry*. Universitext, 2nd Edition, Springer Verlag (1993).
- [Gil] GILKEY (P.B.). — Invariance theory, the heat equation, and the Atiyah-Singer index theorem. *Mathematics Lecture Series*, 11, Publish or Perish, Wilmington (1984).
- [Gol] GOLDBERG (S.I.). — *Curvature and Homology*. Dover books in Mathematics, 3rd edition (1998).
- [Gue] GUÉRINI (P.) Prescription du spectre du Laplacien de Hodge-de Rham. *Annales de l'ENS*, Vol. 37 (2), p. 270-303 (2004).
- [Hel1] HELFFER (B.). — Etude du Laplacien de Witten associé à une fonction de Morse dégénérée. *Publications de l'université de Nantes, Séminaire EDP* 1987-88.
- [Hel2] HELFFER (B.). — Introduction to the semi-classical Analysis for the Schrödinger operator and applications. *Lecture Notes in Mathematics* 1336, Springer Verlag (1988).
- [Hel3] HELFFER (B.). — Semi-classical analysis, Witten Laplacians and statistical mechanics. *World Scientific* (2002).
- [Hen] HENNIART (G.) . — Les inégalités de Morse (d'après E. Witten). *Seminar Bourbaki*, Vol. 1983/84, *Astérisque* No. 121-122, p. 43-61 (1985).
- [HeKINi] HELFFER (B.), KLEIN (M.), and NIER (F.). — Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. *Matematica Contemporanea*, 26, p. 41-85 (2004).
- [HeNi] HELFFER (B.) and NIER (F.). — Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach: the case with boundary. *Mémoire* 105, *Société Mathématique de France* (2006).
- [HeSj1] HELFFER (B.) and SJÖSTRAND (J.). — Multiple wells in the semi-classical limit I. *Comm. Partial Differential Equations* 9 (4), p. 337-408 (1984).
- [HeSj2] HELFFER (B.) and SJÖSTRAND (J.). — Puits multiples en limite semi-classique II -Interaction moléculaire-Symétries-Perturbations. *Ann. Inst. H. Poincaré Phys. Théor.* 42 (2), p. 127-212 (1985).
- [HeSj4] HELFFER (B.) and SJÖSTRAND (J.). — Puits multiples en limite semi-classique IV -Etude du complexe de Witten -. *Comm. Partial Differential Equations* 10 (3), p. 245-340 (1985).
- [HeSj5] HELFFER (B.) and SJÖSTRAND (J.). — Puits multiples en limite semi-classique V - Etude des minipuits-. *Current topics in partial differential equations*, p. 133-186, Kinokuniya, Tokyo (1986).
- [HoKuSt] HOLLEY (R.), KUSUOKA (S.), and STROOCK (D.). — Asymptotics of the spectral gap with applications to the theory of simulated annealing. *J. Funct. Anal.* 83 (2), p. 333-347 (1989).
- [Kol] KOLOKOLTSOV (V.N.). — Semi-classical analysis for diffusions and stochastic processes. *Lecture Notes in Mathematics* 1724, Springer Verlag (2000).
- [KoMa] KOLOKOLTSOV (V.N.), and MAKAROV (K.). — Asymptotic spectral analysis of a small diffusion operator and the life times of the corresponding diffusion process. *Russian J. Math. Phys.* 4 (3), p. 341-360 (1996).

- [KoPrSh] KOLDAN (N.), PROKHORENKOV (I.), and SHUBIN (M.). — Semiclassical Asymptotics on Manifolds with Boundary. Preprint (2008). <http://arxiv.org/abs/0803.2502v1>
- [Lau] LAUDENBACH (F.). — Topologie différentielle. Cours de Majeure de l'Ecole Polytechnique (1993).
- [Lep1] LE PEUTREC (D.). — Small singular values of an extracted matrix of a Witten complex. *Cubo, A Mathematical Journal*, Vol. 11 (4), p. 49-57 (2009).
- [Lep2] LE PEUTREC (D.). — Local WKB construction for Witten Laplacians on manifolds with boundary. *Analysis & PDE*, Vol. 3, No. 3, p. 227-260 (2010).
- [Mic] MICLO (L.). — Comportement de spectres d'opérateurs à basse température. *Bull. Sci. Math.* 119, p. 529-533 (1995).
- [Mil1] MILNOR (J.W.). — Morse Theory. Princeton University press (1963).
- [Mil2] MILNOR (J.W.). — Lectures on the h -cobordism Theorem. Princeton University press (1965).
- [Nie] NIER (F.). — Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. Journées "Equations aux Dérivées Partielles", Exp No VIII, Ecole Polytechnique (2004).
- [Per] PERSSON (A.). — Bounds for the discrete part of the spectrum of a semi-bounded Schrödinger operator. *Math. Scandinavica* 8, p. 143-153 (1960).
- [Sch] SCHWARZ (G.). — Hodge decomposition. A method for Solving Boundary Value Problems. *Lecture Notes in Mathematics* 1607, Springer Verlag (1995).
- [Sima] SIMADER (C.G.). — Essential self-adjointness of Schrödinger operators bounded from below. *Math. Z.* 159, p. 47-50 (1978).
- [Sim] SIMON (B.). — Semi-classical analysis of low lying eigenvalues, I. Non-degenerate minima: Asymptotic expansions. *Ann. Inst. H. Poincaré, Phys. Théor.* 38, p. 296-307 (1983).
- [Wit] WITTEN (E.). — Supersymmetry and Morse inequalities. *J. Diff. Geom.* 17, p. 661-692 (1982).
- [Zha] ZHANG (W.). — Lectures on Chern-Weil theory and Witten deformations. *Nankai Tracts in Mathematics*, Vol. 4, World Scientific (2002).