

YOSHIHIRO MIZUTA

**On the existence of weighted boundary limits  
of harmonic functions**

*Annales de l'institut Fourier*, tome 40, n° 4 (1990), p. 811-833

[http://www.numdam.org/item?id=AIF\\_1990\\_\\_40\\_4\\_811\\_0](http://www.numdam.org/item?id=AIF_1990__40_4_811_0)

© Annales de l'institut Fourier, 1990, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## ON THE EXISTENCE OF WEIGHTED BOUNDARY LIMITS OF HARMONIC FUNCTIONS

by Yoshihiro MIZUTA

---

### 1. Introduction.

In this paper we are concerned with the existence of boundary limits of functions  $u$  which are harmonic in a bounded open set  $G \subset \mathbb{R}^n$  and satisfy a condition of the form :

$$\int_G \Psi(|\text{grad } u(x)|) \omega(x) dx < \infty,$$

where  $\Psi(r)$  is a nonnegative nondecreasing function on the interval  $[0, \infty)$  and  $\omega$  is a nonnegative measurable function on  $G$ . In case  $G$  is a Lipschitz domain,  $\Psi(r) = r^p$  and  $\omega(x) = \rho(x)^\beta$ , many authors studied the existence of (non) tangential boundary limits; see, for example, Carleson [2], Wallin [10], Murai [7], Cruzeiro [3] and Mizuta [5], [6]. Here  $\rho(x)$  denotes the distance of  $x$  from the boundary  $\partial G$ . In this paper, we assume that  $\Psi$  is of the form  $r^p \psi(r)$ , where  $\psi$  is a nonnegative nondecreasing function on the interval  $[0, \infty)$  such that  $\psi(2r) \leq A_1 \psi(r)$  for any  $r > 0$ , with a positive constant  $A_1$ . In case  $G$  is a Lipschitz domain and  $\omega(x)$  is of the form  $\lambda(\rho(x))$ , where  $\lambda$  is a positive and nondecreasing function on the interval  $(0, \infty)$  such that  $\lambda(2r) \leq A_2 \lambda(r)$  for any  $r > 0$  with a positive constant  $A_2$ , our first aim is to find a positive function  $\kappa(r)$  such that  $[\kappa(\rho(x))]^{-1} u(x)$  tends to zero as  $x$  tends to the boundary  $\partial G$ ; when  $\kappa$  is bounded,  $u$  is shown to be extended to a continuous function on  $G \cup \partial G$ .

---

*Key-words* : Harmonic functions - Tangential boundary limits - Bessel capacity Hausdorff measure.

*A.M.S. Classification* : 31B25.

It is known (see [5]) that if  $u$  is a harmonic function on the unit ball  $B$  satisfying

$$\int_B |\text{grad } u(x)|^p (1 - |x|^2)^\beta dx < \infty, \quad \beta \geq p - n,$$

then  $u(x)$  has a finite limit as  $x \rightarrow \xi$  along  $T_\alpha(\xi, a) = \{x \in B; |x - \xi|^\alpha < a\rho(x)\}$  for any  $a > 0$  and any  $\xi \in \partial G$  except those in a suitable exceptional set, where  $\alpha \geq 1$ . Further it is known that this fact is best possible as to the size of the exceptional sets. We shall show in Theorem 1 that if  $u$  is a harmonic function on  $B$  satisfying the stronger condition:

$$\int_B \Psi_p(|\text{grad } u(x)|)(1 - |x|^2)^{p-n} dx < \infty$$

and if  $\psi$  is of logarithmic type (see condition  $(\psi_1)$  below) and  $\int_0^1 [\psi(t^{-1})]^{-1/(p-1)} t^{-1} dt < \infty$ , then  $u$  is extended to a function which is continuous on  $B \cup \partial B$ .

Next let us consider the case where

$$G = G_\alpha \equiv \{x = (x', x_n) \in R^{n-1} \times R^1; |x'|^\alpha < x_n < 1\}.$$

In case  $\alpha < 1$ ,  $G_\alpha$  is not a Lipschitz domain. However, we will also find a positive function  $\kappa(r)$  such that  $[\kappa(|x|)]^{-1}u(x)$  tends to zero as  $x \rightarrow 0$ ,  $x \in G_\alpha$ ; when  $\kappa$  is bounded,  $u$  is shown to have a finite limit at the origin.

Further, we study the existence of (tangential) boundary limits

$$\lim_{x \rightarrow \xi, x \in T_\alpha(\xi, a, b)} u(x)$$

at  $\xi \in \partial G$  except those in a suitable exceptional set, where  $T_\alpha(\xi, a, b) = \{\xi + \Xi_\xi x; x_n > a|x'| + b|x'|^\alpha\}$  with  $a \geq 0$ ,  $b \geq 0$  and an orthogonal transformation  $\Xi_\xi$ . We note here that if  $G$  is a Lipschitz domain, then for any  $\xi \in \partial G$ , there exist  $a_\xi$ ,  $b_\xi \geq 0$ ,  $r_\xi > 0$  and an orthogonal transformation  $\Xi_\xi$  such that  $T_\alpha(\xi, a_\xi, b_\xi) \cap B(\xi, r_\xi) \subset G$ , where  $B(x, r)$  denotes the open ball with center at  $x$  and radius  $r$ . If  $\alpha = 1$ , then our results will imply the usual angular limit theorem.

## 2. Weighted boundary limits.

Throughout this paper, let  $\psi$  be a nonnegative nondecreasing function on the interval  $(0, \infty)$  satisfying the following condition:

$(\psi_1)$  There exists  $A > 1$  such that  $A^{-1}\psi(r) \leq \psi(r^2) \leq A\psi(r)$  whenever  $r > 0$ .

By condition  $(\psi_1)$ , we see that  $\psi$  satisfies the so-called  $(\Delta_2)$  condition, that is, we can find  $A_1 > 1$  such that

$$(\Delta_2) \quad A_1^{-1}\psi(r) \leq \psi(2r) \leq A_1\psi(r) \quad \text{for any } r > 0.$$

For  $p > 1$ , set  $\Psi_p(r) = r^p\psi(r)$ . Since  $\Psi_p(r) \rightarrow 0$  as  $r \rightarrow 0$ , we may assume that  $\Psi_p(0) = 0$ .

If  $\eta$  is a positive measurable function on the interval  $(0, \infty)$ , then we define

$$\kappa_\eta(r) = \left( \int_r^1 s^{p'(1-n/p)} \eta(s)^{-p'/p} s^{-1} ds \right)^{1/p'},$$

where  $1/p + 1/p' = 1$ .

In this paper, let  $M_1, M_2, \dots$  denote various constants independent of the variables in question. Further, we denote by  $B(x, r)$  the open ball with radius  $r$  and center at  $x$ .

Our first aim is to establish the following result, which gives a generalization of Theorem 1 in [6].

**THEOREM 1.** — *Let  $\lambda$  be a nonnegative monotone function on the interval  $(0, \infty)$  satisfying the  $(\Delta_2)$  condition, and let  $\psi$  be a nonnegative nondecreasing function on the interval  $(0, \infty)$  satisfying condition  $(\psi_1)$ . Set  $\eta(r) = \psi(r^{-1})\lambda(r)$ . Suppose  $u$  is a function harmonic in a bounded Lipschitz domain  $G$  in  $R^n$  and satisfying*

$$(1) \quad \int_G \Psi_p(|\text{grad } u(x)|) \lambda(\rho(x)) dx < \infty.$$

*If  $\kappa_\eta(0) = \infty$ , then  $\lim_{x \rightarrow \partial G} [\kappa_\eta(\rho(x))]^{-1} u(x) = 0$ ; if  $\kappa_\eta(0) < \infty$ , then  $u$  has a finite limit at each boundary point of  $G$ .*

*Remark.* — If  $\lambda(r) = r^{p-n}$  and  $\psi$  satisfies the additional condition :

$$(\psi_2) \quad \int_0^1 [\psi(r^{-1})]^{-1/(p-1)} r^{-1} dr < \infty,$$

then  $\kappa_\eta(0) < \infty$ .

For a proof of Theorem 1, we need the following lemma (see [6], Lemma 1).

LEMMA 1. — Let  $G$  be a bounded Lipschitz domain in  $R^n$ . Then for each  $\xi \in \partial G$ , there exist  $r_\xi > 0$  and  $c_\xi > 0$  with the following properties :

- i) if  $0 < r < r_\xi$ , then there exist  $x_r \in G \cap B(\xi, r)$  and  $\sigma_r > 0$  such that

$$E(x, x_r) = \bigcup_{0 \leq t \leq 1} B(X(t), c_\xi \rho(X(t))) \subset G \cap B(\xi, 2r)$$

whenever  $x \in G \cap B(\xi, \sigma_r)$ , where  $X(t) = (1-t)x + tx_r$ ;

- ii)  $\rho(x) + |x-y| < M_1 \rho(y)$  whenever  $y \in E(x, x_r)$ ;

- iii) if  $u$  is a function harmonic in  $G$ , then

$$|u(x) - u(x_r)| \leq M_2 \int_{E(x, x_r)} |\text{grad } u(y)| \rho(y)^{1-n} dy$$

for any  $x \in G \cap B(\xi, \sigma_r)$ . Here  $M_1$  and  $M_2$  are positive constants independent of  $x$ ,  $r$  and  $u$ .

*Proof of Theorem 1.* — Let  $u$  be as in the theorem, and let  $\xi \in \partial G$ . For a sufficiently small  $r > 0$ , by Lemma 1, we find that

$$|u(x) - u(x_r)| \leq M_1 \int_{E(x, x_r)} |\text{grad } u(y)| \rho(y)^{1-n} dy$$

for any  $x \in G \cap B(\xi, \sigma_r)$ . Let  $0 < \delta < 1$ . By condition  $(\psi_1)$ , we can find a constant  $A_\delta > 1$  such that

$$(2) \quad A_\delta^{-1} \psi(r) \leq \psi(r^\delta) \leq A_\delta \psi(r) \quad \text{whenever } r > 0.$$

Hence, from Hölder's inequality we derive

$$\begin{aligned} |u(x) - u(x_r)| &\leq M_1 \left( \int_{\{y \in E(x, x_r); f(y) > \rho(y)^{-\delta}\}} \rho(y)^{p'(1-n)} \psi(f(y))^{-p'/p} \right. \\ &\quad \times \lambda(\rho(y))^{-p'/p} dy \Big)^{1/p'} F(r) + M_1 \int_{E(x, x_r)} \rho(y)^{1-n-\delta} dy \end{aligned}$$

$$\begin{aligned}
&\leq M_2 \left( \int_0^{3r} (\rho(x) + t)^{p'(1-n/p)-1} [\Psi((\rho(x) + t)^{-1})]^{-p'/p} \right. \\
&\quad \times \lambda(\rho(x) + t)^{-p'/p} dt \Big)^{1/p'} F(r) + M_2 \int_{B(x, 2r)} |x - y|^{1-\delta-n} dy \\
&\leq M_3 \kappa_\eta(\rho(x)) F(r) + M_3 r^{(1-\delta)/n},
\end{aligned}$$

where  $f(y) = |\text{grad } u(y)|$  and  $F(r) = \left( \int_{G \cap B(\xi, 2r)} \Psi_p(f(y)) \lambda(\rho(y)) dy \right)^{1/p}$ .

Consequently, if  $\kappa_\eta(0) = \infty$ , then we obtain

$$\limsup_{x \rightarrow \xi} \kappa_\eta(\rho(x))^{-1} |u(x)| \leq M_3 \left( \int_{G \cap B(\xi, 2r)} \Psi_p(f(y)) \lambda(\rho(y)) dy \right)^{1/p}.$$

Condition (1) implies that the right hand side tends to zero as  $r \rightarrow 0$ , so that the left hand side is equal to zero.

On the other hand, if  $\kappa_\eta(0) < \infty$ , then we see that  $\sup_{x \in G \cap B(\xi, \sigma_r)} |u(x) - u(x_r)|$  tends to zero as  $r \rightarrow 0$ , which implies that  $u(x)$  has a finite limit at  $\xi$ . Thus Theorem 1 is established.

### 3. The case $G = G_\alpha$ with $\alpha < 1$ .

If  $\alpha < 1$ , then  $G_\alpha$  is not a Lipschitz domain. However, we study the existence of boundary limits for  $u$  satisfying condition (1).

For simplicity, set

$$\kappa_{\eta, \alpha}(r) = \left( \int_r^1 s^{p'(1-n/p)} [\eta(s)]^{-p'/p} s^{\alpha-2} ds \right)^{1/p'}$$

and

$$K_{\eta, \alpha}(x) = \kappa_\eta(\rho(x)) + \kappa_{\eta, \alpha}(x_n^{1/\alpha}) \quad \text{for } x = (x', x_n).$$

**THEOREM 2.** — *Let  $\lambda$ ,  $\psi$  and  $\eta$  be as in Theorem 1. Let  $u$  be a function harmonic in  $G_\alpha$  and satisfying condition (1). If  $0 < \alpha < 1$  and  $K_{\eta, \alpha}(x) \rightarrow \infty$  as  $x \rightarrow 0$ , then*

$$\lim_{x \rightarrow 0, x \in G_\alpha} [K_{\eta, \alpha}(x)]^{-1} u(x) = 0;$$

*and if  $K_{\eta, \alpha}(x)$  is bounded, then  $u(x)$  has a finite limit as  $x \rightarrow 0$ ,  $x \in G_\alpha$ .*

*Proof.* — For  $r > 0$ , let  $X(r) = (0, \dots, 0, r)$  and  $B_r = B(X(r), \rho(X(r)))$ . If  $E(x, X(r)) \subset B_r$ , then, in view of Lemma 1, we have

$$|u(x) - u(X(r))| \leq M_1 \int_{B_r} |\text{grad } u(y)| \rho(y)^{1-n} dy.$$

As in the proof of Theorem 1, by use of Hölder's inequality we establish

$$(3) \quad |u(x) - u(X(r))| \leq M_2 \kappa_\eta(\rho(x), 2\rho(X(r))) U(r) + M_2 [m_n(B_r)]^{(1-\delta)/n},$$

where  $0 < \delta < \alpha < 1$ ,  $\kappa_\eta(t, r) = \left( \int_t^r s^{p'(1-n/p)} \eta(s)^{-p'/p} s^{-1} ds \right)^{1/p'}$  and

$$U(r) = \left( \int_{B_r} \Psi_p(|\text{grad } u(y)|) \lambda(\rho(y)) dy \right)^{1/p}.$$

For a large integer  $j (\geq j_0)$ , set  $r_j = M_3 j^{-\alpha/(1-\alpha)}$ , where  $j_0$  and  $M_3 > 0$  are chosen so that  $r_j - r_{j+1} < \rho(X(r_j))$ . Now we define

$$F_j = \{x = (x', x_n) \in G_\alpha; |x_n - r_j| < \rho(X(r_j))\}.$$

We shall show the existence of  $N > 0$  such that the number of  $F_m$  with  $F_m \cap F_j \neq \emptyset$  is at most  $N$  for any  $j$ . Letting  $a$  and  $b$  be positive numbers, we assume that  $r_j - ar_j^{1/\alpha} \leq r_{j+k} + b(r_{j+k})^{1/\alpha}$ . Then

$$j[1 - (j/(j+k))^{\alpha/(1-\alpha)}] \leq M_3^{(1-\alpha)/\alpha} [a + b(j/(j+k))^{1/(1-\alpha)}].$$

Since  $M_4 = \inf_{0 < t < 1} (1 - t^{\alpha/(1-\alpha)})/(1 - t) > 0$ , we derive

$$jk/(j+k) \leq M_5 \quad \text{with} \quad M_5 = [M_3^{(1-\alpha)/\alpha} (a + b)]/M_4,$$

so that

$$k \leq M_5 j/(j - M_5) \quad \text{when} \quad j > M_5.$$

From this fact we can readily find  $N > 0$  with the required property. Thus  $\{F_\ell\}$  is shown to satisfy the above condition.

By (3) we have

$$\begin{aligned} |u(X(r_j)) - u(X(r_{j+k}))| &\leq |u(X(r_j)) - u(X(r_{j+1}))| \\ &\quad + |u(X(r_{j+1})) - u(X(r_{j+2}))| + \dots + |u(X(r_{j+k-1})) - u(X(r_{j+k}))| \\ &\leq M_6 \left( \sum_{\ell=j}^{j+k-1} U(r_\ell)^p \right)^{1/p} \left( \sum_{\ell=j}^{j+k-1} \rho(X(r_\ell))^{p'(1-n/p)} [\eta(\rho(X(r_\ell)))]^{p'} \right)^{1/p'} \\ &\quad + M_2 \sum_{\ell=j}^{\infty} [m_n(B_{r_\ell})]^{(1-\delta)/n}. \end{aligned}$$

We note here that

$$\sum_{\ell=j}^{\infty} [m_n(B_{r_\ell})]^{(1-\delta)/n} \leq M_7 \sum_{\ell=j}^{\infty} \ell^{-(1-\delta)/(1-\alpha)} < \infty$$

since  $\delta < \alpha$ , and, by setting  $\sigma(j) = j^{-1/(1-\alpha)}$  for simplicity,

$$\begin{aligned} \sum_{\ell=j}^{j+k-1} \rho(X(r_\ell))^{p'(1-n/p)} [\eta(\rho(X(r_\ell)))]^{-p'/p} \\ \leq M_8 \sum_{\ell=j}^{j+k-1} [\ell^{-1/(1-\alpha)}]^{p'(1-n/p)} [\eta(\ell^{-1/(1-\alpha)})]^{-p'/p} \\ \leq M_9 \int_j^{j+k} [t^{-1/(1-\alpha)}]^{p'(1-n/p)} [\eta(t^{-1/(1-\alpha)})]^{-p'/p} dt \\ = M_{10} \int_{\sigma(j+k)}^{\sigma(j)} s^{p'(1-n/p)} [\eta(s)]^{-p'/p} s^{\alpha-2} ds \\ \leq M_{10} [\kappa_{\eta,\alpha}(\sigma(j+k))]^{p'} \leq M_{11} [\kappa_{\eta,\alpha}(\rho(X(r_{j+k})))]^{p'}. \end{aligned}$$

First suppose  $K_{\eta,\alpha}(x) \rightarrow \infty$  as  $x \rightarrow 0$ . Then, since  $\{F_\ell\}$  meets mutually at most  $N$  times, we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} [K_{\eta,\alpha}(X(r_{j+k}))]^{-1} |u(X(r_{j+k}))| \\ \leq M_6 [M_{11}]^{1/p'} \left( \int_{\bigcup_{\ell \geq j} F_\ell} \Psi_p(|\text{grad } u(y)|) \lambda(\rho(y)) dy \right)^{1/p} \end{aligned}$$

for any  $j$ . Thus it follows that the left hand side is equal to zero. We also see from (3) that

$$\lim_{r \rightarrow 0} \left[ \sup_{x \in B_r \cap G_\alpha} [K_{\eta,\alpha}(x)]^{-1} |u(x) - u(X(r))| \right] = 0.$$

Since  $B_r$  contains some  $X(r_j)$ , it follows that

$$\lim_{x \rightarrow 0, x \in G_\alpha} [K_{\eta,\alpha}(x)]^{-1} u(x) = 0.$$

If  $K_{\eta,\alpha}(x)$  is bounded, then we see that

$$\limsup_{j \rightarrow \infty} |u(X(r_j)) - u(X(r_k))| = 0$$

and

$$\limsup_{r \downarrow 0} \sup_{x \in B_r} |u(x) - u(X(r))| = 0.$$

These facts imply that  $u$  has a finite limit at the origin.

Here we give a result, which is a generalization of Theorem 2.



PROPOSITION 1. — Let  $\lambda_1$  and  $\lambda_2$  be nonnegative monotone functions on the interval  $(0, \infty)$  satisfying the  $(\Delta_2)$  condition, and let  $\psi$  be a nonnegative nondecreasing function on the interval  $(0, \infty)$  satisfying condition  $(\psi_1)$ . Suppose  $u$  is a function harmonic in  $G_\alpha$  and satisfying

$$\int_{G_\alpha} \Psi_p(|\text{grad } u(x)|) \lambda_1(\rho(x)) \lambda_2(|x|^{1/\alpha}) dx < \infty.$$

Set  $\eta_1(r) = \psi(r^{-1})\lambda_1(r)$ ,  $\eta(r) = \psi(r^{-1})\lambda_1(r)\lambda_2(r)$  and

$$K(x) = \kappa_{\eta_1}(\rho(x))[\lambda_2(x_n^{1/\alpha})]^{-1/p} + \kappa_{\eta, \alpha}(x_n^{1/\alpha}).$$

If  $K(0) (= \lim_{x \rightarrow 0} K(x)) = \infty$ , then  $[K(x)]^{-1}u(x) \rightarrow 0$  as  $x \rightarrow 0$ ,  $x \in G_\alpha$ ;

if  $K(x)$  is bounded, then  $u(x)$  has a finite limit as  $x \rightarrow 0$ ,  $x \in G_\alpha$ .

*Proof.* — As in the proof of Theorem 2, for  $x \in B_r$ , we see that

$$\begin{aligned} |u(x) - u(X(r))| &\leq M_1 r^{1-\delta} + M_1 \kappa_{\eta_1}(\rho(x)) \left( \int_{B_r} \Psi_p(f(y)) \lambda_1(\rho(y)) dy \right)^{1/p} \\ &\leq M_1 r^{1-\delta} + M_2 \kappa_{\eta_1}(\rho(x)) \lambda_2(r^{1/\alpha})^{-1/p} \\ &\quad \times \left( \int_{B_r} \Psi_p(f(y)) \lambda_1(\rho(y)) \lambda_2(|y|^{1/\alpha}) dy \right)^{1/p} \end{aligned}$$

and

$$\begin{aligned} |u(X(r_j)) - u(X(r_{j+k}))| &\leq M_3 j^{-(1-\delta)/(1-\alpha)} + M_3 \kappa_{\eta, \alpha}(\rho(X(r_{j+k}))) \\ &\quad \times \left( \int_{(\Delta_{j+k, j})} \Psi_p(f(y)) \lambda_1(\rho(y)) \lambda_2(|y|^{1/\alpha}) dy \right)^{1/p}, \end{aligned}$$

where  $f(y) = |\text{grad } u(y)|$  and  $\Delta_{k, j} = \bigcup_{l=j}^k B_{r_l}$ . Thus the remaining part of the proof is similar to the proof of Theorem 2.

Next, for  $0 < a < 1$ , let  $G_\alpha(a) = \{x = (x', x_n) \in R^{n-1} \times R^1; 0 < x_n < 1, |x'|^\alpha < ax_n\}$ . Then the following result can be proved similarly.

PROPOSITION 2. — Let  $\lambda$ ,  $\psi$  and  $\eta$  be as in Theorem 1. Let  $u$  be a function harmonic in  $G_\alpha$  and satisfying

$$(4) \quad \int_{G_\alpha} \Psi_p(|\text{grad } u(x)|) \lambda(|x|^{1/\alpha}) dx < \infty.$$

If  $0 < \alpha < 1$  and  $\kappa_{\eta, \alpha}(0) = \infty$ , then

$$\lim_{x \rightarrow 0, x \in G_\alpha(a)} [\kappa_{\eta, \alpha}(\rho(x))]^{-1} u(x) = 0$$

for any  $a$  such that  $0 < a < 1$ ; and if  $\kappa_{\eta, \alpha}(r)$  is bounded, then  $u(x)$  has a finite limit as  $x \rightarrow 0$ ,  $x \in G_\alpha(a)$ , for any  $a$  such that  $0 < a < 1$ .

*Remark.* — Proposition 2 is best possible as to the order of infinity in the following sense: if  $\varepsilon > 0$ ,  $\beta > \alpha p - \alpha - 1$  and  $D$  is the half plane  $\{(x, y); x > 0\}$ , then we can find a harmonic function  $u$  on  $D$  which satisfies condition (4) with  $\lambda(r) = r^\beta$  and

$$(5) \quad \lim_{x \rightarrow 0} x^{-\varepsilon} [\kappa_{\eta, \alpha}(x^{1/\alpha})]^{-1} u(x, 0) = \infty.$$

For this purpose, consider  $u(x, y) = r^{-a} \cos a\theta$ , where  $r = (x^2 + y^2)^{1/2}$  and  $\theta = \tan^{-1}(y/x)$ . Then  $u$  is harmonic in  $D$ . Since  $\lambda(r) = r^\beta$ , we see that

$$M_1 \psi(r^{-1})^{-1/p} r^{-a_0} \leq \kappa_{\eta, \alpha}(r) \leq M_2 \psi(r^{-1})^{-1/p} r^{-a_0}$$

with  $a_0 = (2 - p + \beta)/\alpha p + (1 - \alpha)/\alpha p'$ . If  $0 < a < a_0$ , then

$$\int_{G_\alpha} \Psi_p(|\text{grad } u(z)|) \lambda(\rho(z)) dz < \infty.$$

If  $a$  is taken so large that  $-\varepsilon + a_0 < a < a_0$ , then we see that  $u$  also satisfies (5).

#### 4. Removability of the origin.

In this section we are concerned with the removability of the origin for harmonic functions satisfying condition (1) with  $G = B(0, a) - \{0\}$ ,  $a > 0$ .

**THEOREM 3.** — Let  $\lambda$ ,  $\psi$  and  $\eta$  be as in Theorem 1, and let  $u$  be a function which is harmonic in  $B(0, r_0) - \{0\}$  and satisfies

$$\int_{B(0, r_0) - \{0\}} \Psi_p(|\text{grad } u(x)|) \lambda(|x|) dx < \infty.$$

If  $\limsup_{r \downarrow 0} N(r)^{-1} \kappa_\eta(r) < \infty$ , then  $u$  can be extended to a function harmonic in  $B(0, r_0)$ , where  $N(r) = \log(1/r)$  in case  $n = 2$  and  $N(r) = r^{2-n}$  in case  $n \geq 3$ .

*Proof.* — For  $\varepsilon > 0$  and  $x \in B(0, r_0/2) - \{0\}$ , let  $x_\varepsilon = \varepsilon x/|x|$ . Then Lemma 1 gives

$$|u(x) - u(x_\varepsilon)| \leq M \kappa_\eta(|x|) \left( \int_{B(0, 2\varepsilon)} \Psi_p(|\text{grad } u(y)|) \lambda(|x|) dx \right)^{1/p} + M \int_{B(0, 2\varepsilon)} |y|^{1-\delta-n} dy,$$

where  $0 < \delta < 1$ . Consequently, it follows that  $\lim_{x \rightarrow 0} N(|x|)^{-1} u(x) = 0$ .

Now our result is a consequence of a result in [1], p. 204.

## 5. Limits at infinity.

In this section, we discuss the existence of limits at infinity for harmonic functions on a tube domain  $T_\ell = \{x = (x', x'') \in R^\ell \times R^{n-\ell}; |x''| < 1\}$ . This  $T_\ell$  is not generally obtained, by inversion, from  $G_\alpha$ .

THEOREM 4. — Let  $u$  be a harmonic function on  $T_\ell$  satisfying

$$\int_{T_\ell} \Psi_p(|\text{grad } u(x)|) \rho(x)^{p-n} \lambda(|x|) dx < \infty,$$

where  $\lambda$  is a positive monotone function on  $(0, \infty)$  satisfying the  $(\Delta_2)$  condition. Set

$$\tilde{\Psi}(r) = \left( \int_0^r [\Psi(t^{-1})]^{-p'/p} t^{-1} dt \right)^{1/p'}$$

and

$$\kappa(r) = \left( \int_1^r [\tilde{\Psi}(t) \lambda(t)^{-1/p}]^{p'} dt \right)^{1/p'},$$

$r > 1$ . If  $\kappa(r) \rightarrow \infty$  as  $r \rightarrow \infty$ , then  $[\kappa(|x|)]^{-1} u(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ ,  $x \in T_\ell$ ; and if  $\kappa(r)$  is bounded, then  $u(x)$  has a finite limit at infinity.

For the study of the behavior at infinity, we do not think it necessary to replace  $\rho(x)^{p-n}$  by a more general function  $\lambda_1(\rho(x))$ . The proof of this theorem is similar to the proofs of Theorem 2 and Proposition 1; but we give a proof for the sake of completeness.

*Proof of Theorem 4.* — For  $x \in T_\ell$ , take  $x_0 \in T_\ell$  such that  $E(x, x_0) \subset B(x_0, 1)$ . Then, by Lemma 1, we have

$$|u(x) - u(x_0)| \leq M_1 \int_{E(x, x_0)} f(y) \rho(y)^{1-n} dy,$$

where  $f(y) = |\text{grad } u(y)|$ . Hence Hölder's inequality implies that

$$\begin{aligned} |u(x) - u(x_0)| &\leq M_1 \left( \int_{\{y \in E(x, x_0); f(y) \geq \alpha \rho(y)^{-\delta}\}} \Psi_p(f(y)) \rho(y)^{p-n} dy \right)^{1/p} \\ &\quad \times \left( \int_{\{y \in E(x, x_0); f(y) \geq \alpha \rho(y)^{-\delta}\}} \rho(y)^{p'(1-n)} [\Psi(f(y)) \rho(y)^{p-n}]^{-p'/p} dy \right)^{1/p'} \\ &\quad + \alpha \int_{E(x, x_0)} \rho(y)^{1-n-\delta} dy \\ &\geq M_1 \left( \int_{B(x_0, 1)} \Psi_p(f(y)) \rho(y)^{p-n} dy \right)^{1/p} \\ &\quad \times \left( \int_{E(x, x_0)} [\Psi(\alpha \rho(y)^{-\delta})]^{-p'/p} \rho(y)^{-n} dy \right)^{1/p'} + M_2 \alpha, \end{aligned}$$

where  $\alpha > 0$  and  $0 < \delta < 1$ . If we note that

$$\begin{aligned} \left( \int_{E(x, x_0)} [\Psi(\alpha \rho(y)^{-\delta})]^{-p'/p} \rho(y)^{-n} dy \right)^{1/p'} \\ \leq M_3 \left( \int_0^2 [\Psi(\alpha r^{-\delta})]^{-p'/p} r^{-1} dr \right)^{1/p'} \leq M_4 \Psi(\alpha^{-1}), \end{aligned}$$

then

$$|u(x) - u(x_0)| \leq M_5 \left( \int_{B(x_0, 1)} \Psi_p(f(y)) \rho(y)^{p-n} dy \right)^{1/p} \Psi(\alpha^{-1}) + M_2 \alpha.$$

Taking  $\alpha = |x|^{-2}$ , we have

$$\begin{aligned} |u(x) - u(x_0)| &\leq M_6 \left( \int_{B(x_0, 1)} \Psi_p(f(y)) \rho(y)^{p-n} \lambda(|y|) dy \right)^{1/p} \\ &\quad \times \tilde{\Psi}(|x|) \lambda(|x|)^{-1/p} + M_2 |x|^{-2}. \end{aligned}$$

For  $x = (x', x'')$ , let  $k$  be the nonnegative integer such that  $k \leq |x'| < k+1$ . Put  $x_j = j(x', 0)/|x'|$  for  $j = 0, 1, \dots, k$  and

$x_{k+1} = (x', 0)$ . Then

$$\begin{aligned}
 |u(x) - u(x_{j_0})| &\leq |u(x) - u(x_{k+1})| + |u(x_{k+1}) - u(x_k)| + \dots \\
 &\quad + |u(x_{j_0+1}) - u(x_{j_0})| \\
 &\leq M_6 \left( \int_{\Delta(x, x_{j_0})} \Psi_p(f(y)) \rho(y)^{p-n} \lambda(|y|) dy \right)^{1/p} \\
 &\quad \times \left( \sum_{j=j_0}^{k+1} [\tilde{\Psi}(j) \lambda(j)^{-1/p}]^{p'} \right)^{1/p'} + M_2 \left( \sum_{j=j_0}^{k+1} j^{-2} \right) \\
 &\leq M_7 \left( \int_{\Delta(x, x_{j_0})} \Psi_p(f(y)) \rho(y)^{p-n} \lambda(|y|) dy \right)^{1/p} \kappa(|x|) + M_7 j_0^{-1},
 \end{aligned}$$

where  $\Delta(x, x_{j_0}) = \bigcup_{j_0 \leq j \leq k+1} B(x_j, 1)$ . If  $\kappa(r)$  is not bounded, then it follows that

$$\limsup_{|x'| \rightarrow \infty, x \in T_\ell} [\kappa(|x|)]^{-1} |u(x)| \leq M_7 \left( \int_{T_\ell - B(0, j_0-1)} \Psi_p(f(y)) \rho(y)^{p-n} \lambda(|y|) dy \right)^{1/p}$$

for any  $j_0$ , which implies that the left hand side equals zero.

If  $\kappa(r)$  is bounded, then  $u(x)$  is shown to have a finite limit at infinity.

## 6. Global boundary behavior.

In this section we are concerned with the global existence of tangential boundary limits of harmonic functions  $u$  on  $G$  satisfying (1). Our aim is to give generalizations of the author's results [5], [6]. We consider the sets

$$E_0 = \left\{ \xi \in \partial G ; \int_{G \cap B(\xi, 1)} |\xi - y|^{1-n} |\text{grad } u(y)| dy = \infty \right\}$$

and

$$E_h = \left\{ \xi \in \partial G ; \limsup_{r \downarrow 0} h(r)^{-1} \int_{G \cap B(\xi, r)} \Psi_p(|\text{grad } u(y)|) \lambda(\rho(y)) dy > 0 \right\},$$

where  $h$  is a positive nondecreasing function on the interval  $(0, \infty)$ . From condition (1) it follows that  $H_h(E_h) = 0$ ; moreover, in case  $\lambda(r) = r^\beta$ ,  $B_{1-\beta/p, p}(E_0) = 0$ . Here  $H_h$  denotes the Hausdorff measure with the measure function  $h$  and  $B_{\alpha, p}$  denotes the Bessel capacity of index  $(\alpha, p)$  (see Meyers [4]). As to the size of  $E_0$ , we shall give a precise evaluation in Proposition 3 below, after discussing the  $\Psi_p$  norm inequality of singular integrals.

Further, let  $\varphi$  be a positive nondecreasing function on the interval  $(0, \infty)$  such that  $\lim_{r \downarrow 0} \varphi(r) = 0$ ,  $\varphi(r)/r$  is nondecreasing on  $(0, \infty)$  and  $\varphi(2r) \leq M\varphi(r)$  for any  $r > 0$  with a positive constant  $M$ . For  $a > 0$  and  $\xi \in \partial G$ , set

$$S_\varphi(a) = \{x = (x', x_n) \in R^{n-1} \times R^1; \varphi(|x - \xi|) < ax_n\}$$

and

$$T_\varphi(\xi, a) = \{\xi + \Xi_\xi x; x \in S(a)\}$$

with an orthogonal transformation  $\Xi_\xi$ .

**THEOREM 5.** — *Let  $G$  be a Lipschitz domain in  $R^n$ , and let  $u$  be a harmonic function on  $G$  satisfying condition (1). If  $\xi \in \partial G - E_0 \cup E_h$ ,  $T_\varphi(\xi, a) \subset G$  and  $\kappa_\eta(\rho(x)) \leq M(a) h(|\xi - x|)^{-1/p}$  on  $T_\varphi(\xi, a)$ , with a positive constant  $M(a)$ , then  $u(x)$  has a finite limit as  $x \rightarrow \xi$ ,  $x \in T_\varphi(\xi, a)$ .*

*Proof.* — In view of Lemma 1, we can find  $\{r_j\}$ ,  $\{x_j\}$  and  $c > 0$  (in Lemma 1) with the following properties:

- i)  $0 < r_{j+1} < r_j < 1/j$ .
- ii)  $x_j \in G \cap B(\xi, r_j)$ .
- iii) If  $x \in G \cap B(\xi, r_{j+1})$ , then  $E(x, x_j) \subset G \cap B(\xi, r_j)$ ,  $\rho(x) + |x - y| \leq M_1 \rho(y)$  for any  $y \in E(x, x_j)$  and

$$|u(x) - u(x_j)| \leq M_1 \int_{E(x, x_j)} f(y) \rho(y)^{1-n} dy,$$

where  $f(y) = |\text{grad } u(y)|$ . Hence, as in the proof of Theorem 1, we obtain

$$\begin{aligned} |u(x) - u(x_j)| &\leq M_1 \int_{E(x, x_j) - B(\xi, 2|x - \xi|)} f(y) \rho(y)^{1-n} dy \\ &\quad + M_1 \int_{\{y \in G \cap B(\xi, 2|x - \xi|) : f(y) < \rho(y)^{-\delta}\}} \rho(y)^{1-\delta-n} dy \\ &\quad + M_2 \kappa_\eta(\rho(x)) \left( \int_{G \cap B(\xi, 2|\xi - x|)} \Psi_p(f(y)) \lambda(\rho(y)) dy \right)^{1/p} \\ &\leq M_3(I_1 + I_2 + I_3), \end{aligned}$$

where  $0 < \delta < 1$ . If  $y \in E(x, x_j)$  and  $|y - \xi| \geq 2|x - \xi|$ , then  $\rho(y) \geq M_1^{-1}|x - y| \geq M_1^{-1}(|y - \xi| - |x - \xi|) \geq (2M_1)^{-1}|y - \xi|$ , so that

$$I_1 \leq M_4 \int_{E(x, x_j) - B(\xi, 2|x - \xi|)} f(y) |y - \xi|^{1-n} dy.$$

Moreover,  $I_2 \leq M_5|x - \xi|^{1-\delta}$  and  $\kappa_n(\rho(x)) \leq M(a)h(|x - \xi|)^{-1/p}$  for  $x \in T_\varphi(\xi, a)$  by our assumption. Consequently, if  $\xi \in \partial G - (E_0 \cup E_h)$ , then  $\{u(x_\ell)\}_{\ell \geq j+1}$  is bounded, so that we can find a subsequence  $\{u(x_{j_k})\}$  which converges to a number  $u_0$  as  $k \rightarrow \infty$ . Hence, since

$$\lim_{j \rightarrow \infty} [\limsup_{x \rightarrow \xi, x \in T_\varphi(\xi, a)} |u(x) - u(x_j)|] = 0,$$

it follows that  $u(x) \rightarrow u_0$  as  $x \rightarrow \xi$  along  $T_\varphi(\xi, a)$ .

For  $a, b \geq 0$  and  $\alpha > 1$ , set

$$S_\alpha(a, b) = \{x = (x', x_n); x_n > a|x'| + b|x'|^\alpha\}.$$

If  $G$  is a Lipschitz domain, then, for each  $\xi \in \partial G$  we can find  $a_\xi, b_\xi \geq 0, r_\xi > 0$  and an orthogonal transformation  $\Xi_\xi$  such that

$$\{\xi + \Xi_\xi x; x \in S_\alpha(a_\xi, b_\xi)\} \cap B(\xi, r_\xi) \subset G.$$

For  $b > b_\xi$ , put

$$T_\alpha(\xi, b) = T_\alpha(\xi, \Xi_\xi, b) \equiv \{\xi + \Xi_\xi x; x \in S_\alpha(a_\xi, b)\} \cap B(\xi, r_\xi).$$

**COROLLARY** — Let  $G$  be a Lipschitz domain. For  $\alpha > 1$ , let  $\{T_\alpha(\xi, b); \xi \in \partial G, b > b_\xi\}$  be given as above. If  $u$  is a function which is harmonic in  $G$  and satisfies

$$\int_G \Psi_p(|\text{grad } u(x)|) \rho(x)^\beta dx < \infty$$

for  $\beta > p - n$ , then there exists a set  $E \subset \partial G$  such that

$$\text{i) } H_h(E) = 0 \text{ for } h(r) = \inf_{t \geq r} t^{\alpha(n-p+\beta)} \psi(t^{-1});$$

$$\text{ii) } u(x) \text{ has a finite limit as } x \rightarrow \xi \text{ along } T_\alpha(\xi, b) \text{ whenever } \xi \in \partial G - E \text{ and } b > b_\xi.$$

*Proof.* — First note that for  $\varepsilon > 0, r^\varepsilon \psi(r^{-1}) \geq M_1 s^\varepsilon \psi(s^{-1})$  whenever  $0 < s < r$ , on account of condition  $(\psi_1)$ . Hence, since  $\rho(x) \geq M_1|x - \xi|^\alpha$

for  $x \in T_\alpha(\xi, b)$ ,

$$\begin{aligned} \kappa_\eta(\rho(x)) &\leq \left( \int_{M_1 r^\alpha}^1 [s^{n-p+\beta} \psi(s^{-1})]^{-p'/p} s^{-1} ds \right)^{1/p'} \\ &\leq M_2 [r^{\alpha(n-p+\beta-\delta)} \psi(r^{-1})]^{-1/p} \left( \int_{M_1 r^\alpha}^1 s^{-\delta p'/p-1} ds \right)^{1/p'} \\ &\leq M_3 h(r)^{-1/p}, \end{aligned}$$

where  $0 < \delta < n - p + \beta$  and  $r = |x - \xi|$ . Let  $E = E_0 \cup E_h$  in the notation given in Theorem 5. Since  $B_{1-\beta/p,p}(E_0) = 0$  implies that  $E_0$  has Hausdorff dimension at most  $n - p + \beta$ , on account of [4], Theorem 22. Since  $\alpha > 1$  and  $n - p + \beta > 0$ ,  $\lim_{r \rightarrow 0} h(r)/r^{n-p+\beta} = 0$ , so that we see

that  $H_h(E_0) = 0$ . Hence  $H_h(E) = 0$ , and the Corollary follows from Theorem 5.

*Remark 1.* — In case  $\psi(r) \equiv 1$ ,  $\lambda(r) = r^\beta$  with  $p - n \leq \beta < p - 1$  and  $\varphi(r) = r^\alpha$  with  $\alpha > 1$ , we can take  $h$  so that  $h(r) = r^{\alpha(n-p+\beta)}$  if  $n - p + \beta > 0$  and  $h(r) = [\log(2 + r^{-1})]^{1-p}$  if  $n - p + \beta = 0$ . Hence, Theorem 5 and its Corollary give the usual  $T_\alpha$ -limit theorem (see [5]).

*Remark 2.* — Nagel, Rudin and Shapiro [8] proved the existence of  $T_\alpha$ -limits of harmonic functions represented as Poisson integrals in a half space.

## 7. Singular integrals.

Here we establish the following result.

**THEOREM 6.** — *Let  $f$  be a function on  $R^n$  such that*

$$\int (1 + |y|)^{1-n} |f(y)| dy < \infty$$

*and  $\int \Psi_p(|f(y)| |y_n|^{\beta/p}) dy < \infty$ , where  $-1 < \beta < p - 1$ . If we set*

*$u(x) = \int |x - y|^{1-n} f(y) dy$ , then*

$$\int \Psi_p(|\text{grad } u(x)| |x_n|^{\beta/p}) dx \leq M \int \Psi_p(|f(y)| |y_n|^{\beta/p}) dy$$

*with a positive constant  $M$  independent of  $f$ .*



*Proof.* — Without loss of generality, we may assume that  $f \geq 0$  on  $R^n$ . First we consider the case  $\beta = 0$ . We note, by the well-known fact from the theory of singular integral operators, that

$$\begin{aligned}\lambda(a) &\equiv H_n(\{x; |\operatorname{grad} u(x)| > a\}) \\ &\leq M_1 a^{-1} \int_{\{y; f(y) \geq a/2\}} U(y) dy + M_1 a^{-q} \int_{\{y; f(y) < a/2\}} U(y)^q dy \\ &= M_1 \mu_1(a) + M_1 \mu_2(a),\end{aligned}$$

where  $H_n$  denotes the  $n$ -dimensional Lebesgue measure,  $q > p$  and  $U(y) = |\operatorname{grad} u(y)|$ . Hence we have

$$\begin{aligned}\int \Psi_p(|\operatorname{grad} u(x)|) dx &= \int_0^\infty \lambda(a) d\Psi_p(a) \\ &\leq M_1 \int_0^\infty \mu_1(a) d\Psi_p(a) + M_1 \int_0^\infty \mu_2(a) d\Psi_p(a) \\ &\leq M_1 \int U(y) \left( \int_0^{2f(y)} a^{-1} d\Psi_p(a) \right) dy + M_1 \int U(y)^q \left( \int_{2f(y)}^\infty a^{-q} d\Psi_p(a) \right) dy \\ &\leq M_2 \int \Psi_p(U(y)) dy.\end{aligned}$$

In case  $\beta \neq 0$ , set  $g(y) = |y_n|^{\beta/p} U(y)$  and

$$v(x) = \int |x - y|^{1-n} g(y) dy.$$

For  $j = 1, 2, \dots, n$ , we see that

$$||x_n|^{\beta/p} (\partial/\partial x_j) u(x) - (\partial/\partial x_j) v(x)| \leq M_3 \int K_\beta(x_n, y_n) (P_{|x_n - y_n|} g)(x', x_n) dy_n,$$

where  $K_\beta(x_n, y_n) = |1 - |x_n/y_n|^{\beta/p}|/|x_n - y_n|$  and  $P$  denotes the Poisson kernel in the upper half space  $D = \{x = (x', x_n) \in R^{n-1} \times R^1; x_n > 0\}$ . By [9], Theorem 1, (a) in Chap. III and Theorem 1, (c) in Chap. I, we have for  $q \geq 1$

$$\int [P_t g(x', x_n)]^q dx' \leq M_4 \int g(y', y_n)^q dy'.$$

Hence, by using Minkowski's inequality (cf. [9], Appendix A.1), we establish

$$\begin{aligned} \int \left( \int K_{\beta}(x_n, y_n) (P_{|x_n - y_n|} g)(x', x_n) dy_n \right)^q dx \\ \leq M_4 \int \left( \int K_{\beta}(x_n, y_n) \left( \int g(y', y_n)^q dy' \right)^{1/q} dy_n \right)^q dx_n. \end{aligned}$$

Let  $q_1$  and  $q_2$  be positive numbers such that  $\beta < q_1 - 1$  and  $1 < q_1 < p < q_2$ . Applying Appendix A.3 in Stein's book [9], we see that

$$\begin{aligned} \lambda(a) &\equiv H_n(\{x; ||x_n|^{\beta/p}(\partial/\partial x_j)u(x) - (\partial/\partial x_j)v(x)| > a\}) \\ &\leq M_5(\mu_1(a) + \mu_2(a)), \end{aligned}$$

where

$$\mu_1(a) = a^{-q_1} \int_{\{y; g(y) \geq a/2\}} g(y)^{q_1} dy$$

and

$$\mu_2(a) = a^{-q_2} \int_{\{y; g(y) < a/2\}} g(y)^{q_2} dy.$$

Consequently, by the above considerations, we see that

$$\int \Psi_p(|x_n|^{\beta/p}(\partial/\partial x_j)u(x) - (\partial/\partial x_j)v(x)|) \leq M_6 \int \Psi_p(g(y)) dy.$$

Thus it follows that

$$\int \Psi_p(|x_n|^{\beta/p}(\partial/\partial x_j)u(x)|) dx \leq M_7 \int \Psi_p(g(y)) dy,$$

or

$$\int \Psi_p(|x_n|^{\beta/p} \text{grad } u(x)|) dx \leq M_8 \int \Psi_p(g(y)) dy < \infty.$$

*Remark.* — Consider the functions

$$u_j(x) = \int (x_j - y_j) |x - y|^{-n} f(y) dy.$$

Then the same inequality as in Theorem 6 still holds for each  $u_j$ .

For  $\beta > 0$  and  $E \subset R^n$ , we define

$$C_{\beta, \Psi_p}(E) = \inf \int \Psi_p(f(y)) dy,$$

where the infimum is taken over all nonnegative measurable functions  $f$  on  $R^n$  such that  $\int_{B(x, 1)} |x-y|^{\beta-n} f(y) dy \geq 1$  for every  $x \in E$ .

PROPOSITION 3. — Let  $f$  be a nonnegative measurable function on a Lipschitz domain  $G$  such that  $\int_G \Psi_p(f(y)) \rho(y)^\beta dy < \infty$ , and set  $E = \{\xi \in \partial G; \int_{G \cap B(\xi, 1)} |\xi-y|^{1-n} f(y) dy = \infty\}$ . If  $-1 < \beta < p-1$ , then  $C_{1-\beta/p, \Psi_p}(E) = 0$ .

Proof. — By a change of variables, we may assume that  $G$  is the half space  $D$  and  $f$  vanishes outside some ball  $B(0, N)$ . Let  $u(x) = \int_D |x-y|^{1-n} f(y) dy$  for a nonnegative measurable function  $f$  on  $D$  such that  $\int_D \Psi_p(f(y)) y_n^\beta dy < \infty$ . Here note that

$$\begin{aligned} \int \Psi_p(f(y)) y_n^{\beta/p} dy &\leq \int_{\{y \in D; f(y)^\varepsilon \geq y_n^{\beta/p}\}} \Psi_p(f(y)) y_n^{\beta/p} dy \\ &+ \int_{\{y \in D; f(y)^\varepsilon \leq y_n^{\beta/p}\}} \Psi_p(f(y)) y_n^{\beta/p} dy \\ &\leq \int_D y_n^\beta f(y)^p \Psi(f(y)^{1+\varepsilon}) dy \\ &+ \int_{\{y \in D; f(y) > 0\}} \Psi_p(y_n^{(1+\varepsilon-1)\beta/p}) dy < \infty, \end{aligned}$$

if  $\varepsilon > 0$  and  $\beta(1+\varepsilon^{-1}) > -1$ . Hence, from Theorem 6, it follows that

$$\int \Psi_p(|\text{grad } u(x)| |x_n|^{\beta/p}) dx < \infty. \quad \text{Since } |\text{grad } u(x)| = O(|x|^{-n}) \text{ as } |x| \rightarrow \infty, \text{ we see that } \int_{R^n - B(0, a)} \Psi_p(|\text{grad } u(x)|) |x_n|^\beta dx < \infty \text{ for a}$$

sufficiently large  $a$ . Moreover, we have, by letting  $U(x) = |\text{grad } u(x)|$ ,

$$\begin{aligned} \int_{B(0,a)} \Psi_p(U(x)) |x_n|^\beta dx &\leq \int_{\{x \in B(0,a); U(x) \geq |x_n|^{-(1+\delta-1)\beta/p}\}} \Psi_p(U(x)) |x_n|^\beta dx \\ &\quad + \int_{\{x \in B(0,a); U(x) < |x_n|^{-(1+\delta-1)\beta/p}\}} \Psi_p(U(x)) |x_n|^\beta dx \\ &\leq \int \psi([U(x)|x_n|^{\beta/p}]^{1+\delta}) U(x)^p |x_n|^\beta dx \\ &\quad + \int_{B(0,a)} \Psi_p(|x_n|^{-(1+\delta-1)\beta/p}) |x_n|^\beta dx < \infty, \end{aligned}$$

if  $\delta > 0$  and  $\delta > \beta$ . Thus  $\int \Psi_p(U(x)) |x_n|^\beta dx < \infty$ .

Consider the set

$$E^* = \{x \in \partial D; \int_D |x-y|^{1-\beta/p-n} [U(y)y_n^{\beta/p}] dy = \infty\}.$$

Then, by definition,  $C_{1-\beta/p, \Psi_p}(E^*) = 0$ . If  $\xi \in \partial D - E^*$  and  $a > 0$ , then

$$\int_{\Gamma(\xi, a)} |\xi - y|^{1-n} |\text{grad } u(y)| dy < \infty,$$

where  $\Gamma(\xi, a) = \{x \in D; |x - \xi| < ax_n\}$ . It follows that

$$\int_0^{r_0} |\text{grad } u(\xi + r\theta)| dr < \infty \quad \text{for almost every } \theta \in \partial B(0,1),$$

which implies that  $u(\xi + r\theta)$  has a finite limit for almost every  $\theta \in \partial B(0,1)$ . If  $\xi \in E$ , then  $\liminf_{r \rightarrow 0} u(\xi + rx) \geq u(\xi) = \infty$  for any  $x \in D$  by the lower semicontinuity of potentials. Thus  $\xi \in \partial D - E$ . Hence  $E \subset E^*$ , or  $C_{1-\beta/p, \Psi_p}(E) = 0$ .

## 8. Best possibility.

Here we deal with the best possibility of Theorem 1 as to the order of infinity. Let  $D$  be the upper half space, that is,  $D = \{x = (x', x_n) \in R^{n-1} \times R^1; x_n > 0\}$ .

PROPOSITION 4. — Let  $\lambda$ ,  $\psi$  and  $\eta$  be as in Theorem 1. Suppose  $\kappa_\eta(0) = \infty$  and  $r^\delta \eta(r)^{-1}$  is bounded above on  $(0, 1]$  for some  $\delta > 1 - n$ . If  $a(r)$  is a nonincreasing positive function on the interval  $(0, \infty)$  such that  $\lim_{r \downarrow 0} a(r) = \infty$ , then there exists a nonnegative measurable function  $f$  such that  $f = 0$  outside  $B(0, 1)$ ,

$$\int_{\mathbb{R}^n} \Psi_p(f(y)) \lambda(|y_n|) dy < \infty$$

and

$$\limsup_{r \downarrow 0} a(r) \kappa_\eta(r)^{-1} u(r\xi) = \infty \quad \text{for any } \xi \in D,$$

$$\text{where } u(x) = \int_{\mathbb{R}^{n-D}} (x_n - y_n) |x - y|^{-n} f(y) dy.$$

Remark. — By the Remark after Theorem 6, if  $\lambda(r) = r^\beta$  with  $-1 < \beta < p - 1$ , then

$$\int \Psi_p(|\text{grad } u(x)|) |x_n|^\beta dx < \infty.$$

Proof of Proposition 4. — Let  $\{r_j\}$  be a sequence of positive numbers such that  $r_j < r_{j-1}/2$  and

$$\kappa_\eta(r_j) \leq 2 \left( \int_{r_j}^{r_{j-1}} [s^{n-p} \eta(s)]^{-p'/p} s^{-1} ds \right)^{1/p'}.$$

Further take a sequence  $\{b_j\}$  of positive numbers such that  $\lim_{j \rightarrow \infty} b_j a(r_j) = \infty$  and  $\sum_{j=1}^{\infty} b_j^p < \infty$ . Let  $\Gamma(c)$  be the cone  $S_\phi(c)$  with  $\phi(r) \equiv r$ , and set  $\hat{\Gamma}(c) = \{x \in \mathbb{R}^n; -x \in \Gamma(c)\}$ . Now we define

$$f(y) = b_j \kappa_\eta(r_j)^{-p'/p} [|y|^{n-1} \eta(|y|)]^{-p'/p}$$

if  $y \in \hat{\Gamma}_j \equiv \hat{\Gamma}(1) \cap B(0, r_{j-1}) - B(0, r_j)$  and  $f = 0$  otherwise, and consider the function  $u$  defined as in Proposition 4. If

$$x \in \Gamma(c) \cap B(0, 2r_j) - B(0, r_j),$$

then

$$\begin{aligned} u(x) &\geq M_1 b_j \kappa_\eta(r_j)^{-p'/p} \int_{\hat{\Gamma}_j} |y|^{1-n} [|y|^{n-1} \eta(|y|)]^{-p'/p} dy \\ &\geq M_2 b_j \kappa_\eta(r_j), \end{aligned}$$

so that

$$\lim_{x \rightarrow 0, x \in \Delta(c)} a(|x|) \kappa_\eta(|x|)^{-1} u(x) = \infty$$

with  $\Delta(c) = \bigcup_{j=1}^{\infty} \{x \in \Gamma(c); r_j < |x| < 2r_j\}$ . On the other hand, since  $r^\delta \eta(r)^{-1}$  is bounded above by our assumption,  $f(y) \leq M_3 |y|^{-p'(n-1+\delta)/p}$ , so that  $\psi(f(y)) \leq M_4 \psi(|y|^{-1})$  by (2). Hence we establish

$$\begin{aligned} \int_{R^n} \Psi_p(f(y)) \lambda(|y|) dy &\leq M_5 \sum_{j=1}^{\infty} b_j^p \kappa_\eta(r_j)^{-p'} \int_{r_j}^{\infty} |y|^{p'(1-n)} \eta(|y|)^{1-p'} dy \\ &\leq M_6 \sum_{j=1}^{\infty} b_j^p < \infty. \end{aligned}$$

Thus  $f$  satisfies all the required assertions.

The Corollary to Theorem 5 is best possible as to the size of the exceptional sets, in the following sense.

**PROPOSITION 5.** — *Let  $\psi$ ,  $\lambda$  and  $\eta$  be as in Theorem 1. Let  $\varphi$  be a nonnegative nondecreasing function on  $(0, \infty)$  such that  $\varphi(r) \leq Mr$  for any  $r > 0$ , with a positive constant  $M$ , and set*

$$\varphi^*(r) = \int_{\varphi(r)}^{2Mr} [t^{n-p} \eta(t)]^{-p'/p} t^{-1} dt.$$

*Suppose further that the following assertions hold :*

- i)  $r^{\delta_1} \lambda(r)^{-1}$  is nondecreasing on  $(0, \infty)$  for some  $\delta_1 > 1/p - n$ .
- ii)  $r^{\delta_2} \lambda(r)$  is nondecreasing on  $(0, \infty)$  for some  $\delta_2 < 1$ .
- iii)  $\varphi^*(r) \rightarrow \infty$  as  $r \rightarrow 0$ .
- iv)  $\varphi^*(r) \leq M^* \varphi^*(s)$  whenever  $0 < s < r$ , with a positive constant  $M^*$ .

*We now define  $h(r) = \inf_{s \geq r} [\varphi^*(s)]^{-p/p'}$ . Then, for a compact set  $K \subset \partial D$  such that  $H_h(K) = 0$ , there exists a nonnegative measurable function  $f$  on  $R^n$  such that*

$$\int \Psi_p(f(y)) \lambda(|y_n|) dy < \infty$$

*and  $Uf(x) \equiv \int_{R^{n-D}} (x_n - y_n) |y - y|^{-n} f(y) dy$  does not have a finite limit as  $x \in T_\varphi(\xi, 1) \rightarrow \xi$  at any  $\xi \in K$ , where  $T_\varphi(\xi, 1) \equiv \{x + \xi; x \in S_\varphi(1)\}$ .*

*Proof.* — For the construction of such  $f$ , we take, for each positive integer  $m$ , a finite family  $\{B(x_{j,m}, r_{j,m})\}$  of balls such that  $x_{j,m} \in \partial D$ ,  $r_{j,m} < 1/m$ ,  $\sum_j h(r_{j,m}) < 2^{-m}/m$  and  $\bigcup_j B(x_{j,m}, r_{j,m}) \supset K$ . Setting

$$B_{i,j} = B(x_{i,j}, 2Mr_{i,j}) - B(x_{i,j}, \varphi(r_{i,j})),$$

we define

$$f_{m,j}(y) = m^{1/p} [h(r_{j,m})]^{p'/p} [|x_{j,m} - y|^{n-1} \eta(|x_{j,m} - y|)]^{-p'/p}$$

for  $y \in B_{m,j}$  and  $f_{m,j}(y) = 0$  elsewhere. Consider the function  $f(y) = \sup_{m,j} f_{m,j}(y)$ . Since  $f_{m,j}(y) \leq M_1 |x_{j,m} - y|^{-\gamma}$ , where

$$\gamma = 1/p + p'(n-1+\delta_1)/p > 0,$$

we see that  $\psi(f_{m,j}(y)) \leq M_2 \psi(|x_{j,m} - y|^{-1})$  on account of (2). Since  $r^{\delta_2} \lambda(r)$  is nondecreasing and  $\varphi^*(r) \leq M_3 [h(r)]^{-p'/p}$ , we establish

$$\begin{aligned} \int_{R^{n-D}} \Psi_p(f(y)) \lambda(|y_n|) dy &\leq M_4 \sum_m m \left( \sum_j [h(r_{j,m})]^{p'} \int_{B_{j,m}} |x_{j,m} - y|^{p'(1-n)} \right. \\ &\quad \times [\eta(|x_{j,m} - y|)]^{p'} \psi(|x_{j,m} - y|^{-1}) [|x_{j,m} - y|^{\delta_2} \lambda(|x_{j,m} - y|)] |y_n|^{-\delta_2} dy \Big) \\ &\leq M_5 \sum_m m \left( \sum_j [h(r_{j,m})]^{p'} \varphi^*(r_{j,m}) \right) \\ &\leq M_6 \sum_m m \left( \sum_j h(r_{j,m}) \right) \leq M_6 \sum_m 2^{-m} < \infty. \end{aligned}$$

Further,

$$\begin{aligned} Uf(x) &\geq \int (x_n - y_n) |x - y|^{-n} f_{m,j}(y) dy \\ &\geq M_7 m^{1/p} [h(r_{j,m})]^{p'/p} \int_{\varphi(r_{i,j})}^{2Mr_{i,j}} r^{p'(1-n)} [\eta(r)]^{-p'/p} r^{-1} dr \\ &\geq M_7 m^{1/p} \end{aligned}$$

for any  $x \cap D \cap B(x_{j,m}, \varphi(r_{j,m}))$ . If  $\xi \in K$ , then for each  $m$  there exists  $j(m)$  such that  $\xi \in B(x_{j(m),m}, r_{j(m),m})$ . Since

$$B(x_{j(m),m}, \varphi(r_{j(m),m})) \cap T_\varphi(\xi, 1) \neq \emptyset,$$

it follows that

$$\limsup_{x \rightarrow \xi, x \in T_\varphi(\xi, 1)} Uf(x) = \infty.$$

## BIBLIOGRAPHY

- [1] M. BRELOT, *Élément de la théorie classique du potentiel*, 4<sup>e</sup> édition, Centre de Documentation Universitaire, Paris, 1969.
- [2] L. CARLESON, *Selected problems on exceptional sets*, Van Nostrand, Princeton, 1967.
- [3] A. B. CRUZEIRO, Convergence au bord pour les fonctions harmoniques dans  $R^d$  de la classe de Sobolev  $W_1^d$ , C.R.A.S., Paris, 294 (1982), 71-74.
- [4] N. G. MEYERS, A theory of capacities for potentials in Lebesgue classes, *Math. Scand.*, 26 (1970), 255-292.
- [5] Y. MIZUTA, On the Boundary limits of harmonic functions with gradient in  $L^p$ , *Ann. Inst. Fourier*, 34-1 (1984), 99-109.
- [6] Y. MIZUTA, On the boundary limits of harmonic functions, *Hiroshima Math. J.*, 18 (1988), 207-217.
- [7] T. MURAI, On the behavior of functions with finite weighted Dirichlet integral near the boundary, *Nagoya Math. J.*, 53 (1974), 83-101.
- [8] A. NAGEL, W. RUDIN and J. H. SHAPIRO, Tangential boundary behavior of functions in Dirichlet-type spaces, *Ann. of Math.*, 116 (1982), 331-360.
- [9] E. M. STEIN, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, 1970.
- [10] H. WALLIN, on the existence of boundary values of a class of Beppo Levi functions, *Trans. Amer. Math. Soc.*, 120 (1985), 510-525.

Manuscrit reçu le 3 mai 1989.

Yoshihiro MIZUTA,  
Department of Mathematics  
Faculty of Integrated Arts and Sciences  
Hiroshima University  
Hiroshima 730 (Japon).