

ANNALES
DE
L'INSTITUT FOURIER

Bernhard KRÖN

Green functions on self-similar graphs and bounds for the spectrum of the laplacian

Tome 52, n° 6 (2002), p. 1875-1900.

<http://aif.cedram.org/item?id=AIF_2002__52_6_1875_0>

© Association des Annales de l'institut Fourier, 2002, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

GREEN FUNCTIONS ON SELF-SIMILAR GRAPHS AND BOUNDS FOR THE SPECTRUM OF THE LAPLACIAN

by Bernhard KRÖN (*)

1. Introduction.

Self-similar graphs can be seen as discrete versions of fractals (more precisely: compact, complete metric spaces defined as the fixed set of an iterated system of contractions, see Hutchinson in [15]). The simple random walk is a crucial tool in order to study diffusion on fractals, see Barlow and Kigami [1], Barlow and Perkins [2], Grabner [11], Lindstrøm [23] and many others. In the present paper we study Green functions and the spectrum of the corresponding Markov transition operator P and (equivalently) the Laplacian $\Delta = I - P$, where I is the identity.

There are different methods of spectral analysis on self-similar structures. Barlow and Kigami [1] and Sabot [33] used localized eigenfunctions to study the spectrum and to prove that it is a pure point spectrum. A first heuristic result on explicit spectra is due to Rammal [31] who investigated the spectrum of the Sierpiński graph in the setting of statistical physics. Malozemov and Teplyaev studied the spectral self-similarity of operators in a series of papers. Their functional analytic method is fundamentally different from our probabilistic approach. Malozemov computed the spectrum of the Koch graph in [24] and the spectrum of a self-similar tree in [25]. The spectrum of the Sierpiński graph was computed by Teplyaev in [34]. In [26] they discussed spectral properties of a class of self-similar graphs which can be seen as discrete analogues to finitely ramified fractals such

(*) This work is part of the author's PhD-thesis [20]. The author was supported by the projects Y96-MAT and P14379-MAT of the Austrian Science Fund.

Keywords : Self-similar graphs – Green functions.

Math. classification : 60J10 – 30D05 – 05C50.

that the boundaries of their cells consist of two points. We give a first axiomatic definition of self-similar graphs corresponding to fractals with an arbitrary number of boundary points. With our method we do not only obtain information about the spectrum but we are also able to describe the analytic properties of the Green functions precisely. This axiomatic definition of self-similarity of graphs and the central results of the present paper are part of the author's PhD thesis [20]⁽¹⁾. In a recent preprint [27] Malozemov and Teplyaev constructed a similar but more restricted class of self-similar graphs and they obtained the same bounds for the spectrum as in Theorem 5 of this paper.

Bartholdi and Grigorchuk, see [3] and [4], have computed the spectra of several Schreier graphs of fractal groups of intermediate growth. The reader is also referred to the survey article of Bartholdi, Grigorchuk and Nekrashevych [5].

We introduce a new class of rather general self-similar graphs in Section 2. In this definition there occurs a function ϕ which maps a set of vertices to the set of all vertices. This function is a contraction with respect to the natural graph metric and it can be interpreted as the 'self-similarity function' of the graph. A graph theoretic analogue of the Banach fixed point theorem is proved: Either ϕ fixes exactly one origin vertex, or it contracts towards exactly one 'cell' of the graph. The simple random walk on self-similar graphs is recurrent if the graph has bounded geometry, which means that the set of vertex degrees is bounded, the cells are finite and the numbers of vertices in the boundaries of the cells are bounded. Properties concerning bounded geometry and volume growth of self-similar graphs were studied by the author in [21].

In Section 3 the class of 'symmetrically self-similar' graphs is defined. They correspond to finitely ramified fractals for which the renormalization problem for the simple random walk (see Lindstrøm [23] or Kigami [19]) can be solved by using only one variable. In other words: Starting at a point in the boundary of a cell in a fractal, the transition times for the Brownian motion to hit any other point in this boundary have the same distribution for all boundary points. Our class of graphs contains the graphs studied by Malozemov and Teplyaev but also other well-known graphs such as the extensively studied Sierpiński graph, see Barlow and Perkins [2], Grabner

⁽¹⁾ Results and construction of self-similar graphs were presented at the DMV-conference (Dresden, September 2000), 'Dynamic Odyssey' (Luminy-Marseille, February 2001), 'Fractals' (Graz, June 2001) and 'Random walks and Geometry' (Vienna, June/July 2001).

and Woess [12], Hambly [13] and Jones [17] or the Viček graph which corresponds to the Viček snowflake, see Metz and Hambly [14] and [28]. A finite part of this graph can be seen in Figure 1.




Figure 1. A finite part of the Viček snowflake

We apply a variable substitution technique of generating functions (see Goulden and Jackson [10]) in Section 4. This technique was used by Grabner and Woess in [12] to derive a functional equation for a certain Green function on the Sierpiński graph. Rammal and Toulouse used physical arguments to obtain such a functional equation, see [30] and [32]. Grabner and Woess calculated the asymptotic fluctuation behaviour of the n -step return probabilities. Grabner gave further applications of this substitution method in the context of the analysis of stopping times of Brownian motion on the Sierpiński gasket, see [11]. We state similar functional equations for all Green functions on any symmetrically self-similar graph. Such functional equations reflecting the self-similarity of a graph, a group or a fractal occur often in the literature. We refer for example to the papers mentioned above: Bartholdi, Grigorchuk and Nekrashevych ([3], [4] and [5]) and Malozemov and Teplyaev ([24], [25], [26] and [34]).

In Section 5 the functional equations for the Green functions of the whole graph are extended to a decomposition formula. This formula together with the fixed point theorem of Section 2 implies that any Green function can be written as a finite sum of terms consisting of products and concatenations of finitely many Green functions (of the origin cell and the origin vertex) together with the *return function* f , the *transition function* d and the *inner transition functions* h . The functions f , d and h are rational Green functions of the simple random walk with absorbing boundary on the finite subgraph which is spanned by a cell. This decomposition is one

of the central tools in studying the singularities of the Green functions (see also [22]) and in computing the spectrum of the Laplacian.

Green functions of the simple random walk with absorbing boundaries on finite subgraphs spanned by the so-called ‘ n -cells’ are studied in Section 6. These rational functions can be decomposed into the return function, the transition function and the inner transition functions. The zeroes of the return function turn out to be poles of the inner transition functions.

Preliminary results concerning the dynamics of the transition function d , the analytic continuations of the Green functions and the spectrum of the transition operator are stated in Section 7. We define \mathcal{D} as the set of d -backward iterates of the poles of the return function and the inner transition functions. This is the set of poles of the Green functions studied in Section 6 being contained in $\overline{\mathbb{R}} \setminus (-1, 1)$. The reduction algorithm in Section 5 yields the analytic continuation of all Green functions to the immediate basin of attraction A_0^* of the fixed point $z = 0$ of d except for some points in \mathcal{D} . It will turn out that A_0^* coincides with the Fatou set of d .

In Section 8 we put all these results together and give explicit bounds for the spectrum: the reciprocal spectrum

$$\text{spec}^{-1} P = \{1/\lambda ; \lambda \in \text{spec } P\}$$

of the transition operator P of the simple random walk is the set of real numbers z in $\overline{\mathbb{R}}$ such that the Green functions cannot be continued uniquely analytically to z from both half spheres in $\overline{\mathbb{C}} \setminus \overline{\mathbb{R}}$ and

$$\mathcal{J} \subset \text{spec}^{-1} P \subset \mathcal{J} \cup \mathcal{D} \subset \overline{\mathbb{R}} \setminus (-1, 1).$$

A crucial role is played by the singularity $z = 1$ of the Green functions.

The last Section 9 is devoted to the Julia set \mathcal{J} . The Julia set is either a real Cantor set or it is a real interval in the extended complex plane $\overline{\mathbb{C}}$. If \mathcal{J} is a Cantor set, then the reciprocal spectrum coincides with the set of all singularities of all Green functions. We give the example of the two sided infinite line, where the reciprocal spectrum is $\overline{\mathbb{R}} \setminus (-1, 1)$. In this case the Green function has a branch cut along $\overline{\mathbb{R}} \setminus (-1, 1)$. Here the transition functions are conjugated to the Chebychev polynomials.

2. Self-similar graphs.

Graphs $X = (VX, EX)$ are always assumed to be connected, locally finite, infinite and without loops or multiple edges. If X is undirected, then

each $e = \{x, y\}$ is a set of vertices x and y in VX , if X is directed, then $EX \subset VX \times VX$. A *path of length n* from x to y is an $(n+1)$ -tuple of vertices

$$(x_0 = x, x_1, \dots, x_n = y)$$

such that x_i is adjacent to x_{i+1} for $0 \leq i \leq n - 1$. The distance $d_X(x, y)$ is the length of the shortest path from x to y . We define the *vertex boundary* or *boundary* θC of a set C of vertices in VX as the set of vertices in $VX \setminus C$ which are adjacent to a vertex in C . The *closure* of C is defined as $\bar{C} = C \cup \theta C$. Since we will use the topological closure only for sets of complex numbers, this notion will cause no misunderstandings. The set C is called *connected* if every pair of vertices in C can be connected by a path in X that does not leave C . The set C is *adjacent* to a vertex x if $x \in \theta C$. Two sets of vertices are *adjacent* if their boundaries are not disjoint.

For a set $F \subset VX$ let $\mathcal{C}_X(F)$ denote the set of connected components in $VX \setminus F$. We define a new graph X_F by setting $VX_F = F$ and connecting two elements x and y in VX_F by an edge if and only if there exists a $C \in \mathcal{C}_X(F)$ such that x and y are in the boundary of C . The graph X_F is then called the *reduced graph of X with respect to F*.

DEFINITION 1. — A graph X is called *self-similar with respect to F* if

- (F1) no vertices in F are adjacent in X ,
- (F2) the intersection of the closures of two different components in $\mathcal{C}_X(F)$ contains no more than one vertex and
- (F3) there exists an isomorphism $\psi : X \rightarrow X_F$.

We will also write ϕ instead of ψ^{-1} and F^n instead of $\psi^n F$, F^0 is defined as VX . The components of $\mathcal{C}_X(F^n)$ are called *n-cells*, 1-cells are also just called *cells*. Furthermore we define functions

$$S : \mathcal{C}_X(F) \longrightarrow \mathcal{C}_X(F^2)$$

where $S(C)$ is the 2-cell that contains the cell C and

$$\phi_S : \mathcal{C}_X(F) \longrightarrow \mathcal{C}_X(F)$$

such that $\phi_S C$ is the cell whose boundary is $\phi \theta S(C)$; equivalently: $\phi_S C = \phi(S(C) \cap F)$. A cell C such that $\phi_S C = C$ is called an *origin cell*, a vertex which is a fixed point of ϕ is called an *origin vertex*. If $\phi_S^m C = C$ for some integer m , then we call C a *periodic origin cell with period m*.

Note that the boundary of any cell in X spans a complete graph in X_F .

LEMMA 1. — Let C be an m -cell and D be an n -cell. Then either

$$m = n \text{ and } C = D \quad \text{or} \quad |\theta C \cap \theta D| \leq 1.$$

Proof. — If $m = n$, then this is equivalent to Axiom (F2) for the graph $X_{F^{m-1}} = X_{F^{n-1}}$. Otherwise let us suppose that $m < n$. If there are two different elements x and y in $\theta C \cap \theta D$, then they are adjacent in X_{F^m} . The vertices x and y are elements of F^n which is a subset of F^{m+1} . Since the graph X_{F^m} is self-similar with respect to F^{m+1} , this contradicts Axiom (F2) for X_{F^m} . \square

The following theorem can be seen as a graph theoretic analogue for the Banach fixed point theorem.

THEOREM 1. — Let X be a self-similar graph with respect to F . Then X is also self-similar with respect to F^n . Either there exists exactly one origin cell O and for every cell C there exists an integer n such that

$$\phi_S^n C = O$$

or ϕ has exactly one origin vertex and for any cell C there exists a periodic origin cell O_m with period m and an integer n such that

$$\phi_S^n C = O_m.$$

Proof. — The graphs X and X_F are isomorphic, and X_F is isomorphic to X_F reduced by $\psi(F)$ which implies that X is isomorphic to X_{F^2} . The first statement now follows by induction.

Let x and y be vertices in F with $d_{X_F}(x, y) = n$. Since no vertices in F are adjacent in X , every path in X connecting these vertices must have length at least $2n$. Thus

$$d_X(x, y) \geq 2 \cdot d_X(\phi x, \phi y) = 2 \cdot d_{X_F}(x, y)$$

for all pairs of vertices x, y in F . For any cell C in $\mathcal{C}_X(F)$

$$\begin{aligned} d_X(\theta C, \theta \phi_S C) &\geq d_X(\theta S(C), \theta S(\phi_S C)) \\ &\geq 2 \cdot d_X(\phi \theta S(C), \phi \theta S(\phi_S C)) = 2 \cdot d_X(\theta \phi_S C, \theta \phi_S^2 C) \end{aligned}$$

and the sequence $(d_X(\theta \phi_S^{n-1} C, \theta \phi_S^n C))_{n \in \mathbb{N}}$ must eventually be zero.

If ϕ has fixed points x and y , then

$$d_X(x, y) \geq 2 \cdot d_X(\phi x, \phi y) = 2 \cdot d_X(x, y)$$

and x equals y .

Let C_1 and C_2 be different cells such that $\phi_S C_1 = C_2$ and $d_X(\theta C_1, \theta C_2) = 0$. By Axiom (F2) there is exactly one vertex x in $\theta C_1 \cap \theta C_2$. In the case that C_1 and C_2 are in the same 2-cell, C_2 is an origin-cell. Otherwise we distinguish: If $\phi \theta C_2$ lies in $S(C_2)$, then $\phi(S(C_2) \cap F)$ is the cell $\phi_S C_2$ which contains $\phi \theta C_2$, and $\phi_S C_2$ is an origin cell. If $\phi \theta C_2$ does not lie in $S(C_2)$, then $\theta S(C_2)$ and $\theta \phi_S C_2$ must have non-empty intersection, because $d_X(\theta C_2, \theta \phi_S C_2) = 0$. By Lemma 1 this intersection consists only of the vertex x . Thus

$$x \in \theta C_1 \cap \theta \phi_S C_1 \cap \theta \phi_S^2 C_1,$$

and proceeding by induction we obtain $x \in \theta \phi_S^n C_1$ for every integer n . Since X is locally finite, there must be indices k and m such that $\phi_S^{k+m} C_1 = \phi_S^k C_1$. The closures of all different 2-cells containing one of the cells with x in their boundaries must have only the vertex x in common. The function ϕ maps these 2-cells onto the cells having x in their boundaries and therefore it must fix x .

Suppose there is an origin cell C and no fixed point of ϕ . If θC and $\theta S(C)$ are disjoint, then no cell D which is not in $S(C)$ can be fixed by ϕ_S , because $d_X(\theta \phi_S^n D, \theta \phi_S^n C)$ must be zero for sufficiently large integers n , and C is the only origin cell. If there is a vertex x in $\theta C \cap \theta S(C)$, then by the above arguments

$$x \in \phi_S^n C$$

for any integer n and x must be fixed by ϕ . \square

LEMMA 2. — Let X be self-similar with respect to F and the corresponding isomorphism ϕ and let o be an origin vertex. Then there is an integer k such that the subgraphs \bar{A} which are spanned by $\bar{A} = A \cup \{o\}$ for components A in $\mathcal{C}_X(\{o\})$ are self-similar graphs with respect to

$$F^k \cap \bar{A} \quad \text{and} \quad \phi^k|_{F^k \cap \bar{A}},$$

and they have exactly one origin cell O_A .

Proof. — The fixed point o of ϕ must be contained in the boundary of every periodic origin cell. Since X is locally finite, there exist only finitely many periodic cells C_1, C_2, \dots, C_r with periods m_1, m_2, \dots, m_r . For

$$k = \text{lcm}(m_1, m_2, \dots, m_r)$$

we have $\phi_S^k C_{m_i} = C_{m_i}$, where $i \in \{1, 2, \dots, r\}$, and all cells C_i are origin cells with respect to F^k and ϕ^k . Let $A_{i,n}$ be the closed n -cell which

contains C_i . Then, for $n \geq 2$,

$$\phi^k(A_{i,n} \cap F^k) = A_{i,n-1}$$

and if C_i is the origin cell with respect to F^k which lies in the component A in $\mathcal{C}_X(\{o\})$, then

$$\bigcup_{n=1}^{\infty} A_{i,n} = \bar{A}. \quad \square$$

LEMMA 3. — *Let all cells,*

$$\sup\{|\theta C| ; C \in \mathcal{C}_X(F)\} \quad \text{and} \quad \sup\{\deg_X x ; x \in F\}$$

be finite. Then the simple random walk on X is null recurrent.

Proof. — We apply Nash-Williams' recurrence criterion, see [29], as a special case of Rayleigh's short-cut method which, for example, can be found in the book of Doyle and Snell [8], Chapter 6, or in the book of Woess [35], Theorem 2.19 and Corollary 2.20.

Let o be an origin vertex. By Lemma 2 we can choose F such that all cells C with $o \in \theta C$ are origin cells. Let O_n be the union of the closures of all n -cells C_n with $o \in \theta C_n$. If X has no origin vertex, we define O_n as the closure of the n -cell which contains the (in this case unique) origin cell, see Theorem 1. The sequence

$$(A_n)_{n \in \mathbb{N}}, \quad A_1 = O_1 \quad \text{and} \quad A_{n+1} = O_{n+1} \setminus O_n,$$

is a one-dimensional partition of VX in the following sense:

$$\bigcup_{n=1}^{\infty} A_n = VX,$$

and for any $n \geq 2$ the sets A_n are finite and the vertices in A_n are only adjacent to vertices in A_{n-1} , A_n or A_{n+1} . Since

$$\sup\{|\theta C| ; C \in \mathcal{C}_X(F)\} = \sup\{|\theta C| ; C \in \mathcal{C}_X(F^n)\}$$

for every $n \in \mathbb{N}$ and $\{\deg_X(x) ; x \in F\}$ is bounded, the number a_n of edges connecting two sets A_n and A_{n+1} is also bounded. Thus

$$\sum_{n=1}^{\infty} \frac{1}{a_n} = \infty$$

and, by the Nash-Williams' recurrence criterion, the simple random walk on X is recurrent. Recurrence of the simple random walk on infinite graphs is always null recurrence, see for example [35], Theorem 1.18. \square

3. Symmetrically self-similar graphs.

DEFINITION 2. — Let X be a connected graph which is self-similar with respect to F and let \widehat{C} be the subgraph of X which is spanned by the closure \overline{C} of a cell C . From now on we suppose that

(S1) all cells are finite and for any pair of cells C and D in $\mathcal{C}_X(F)$ there exists a graph isomorphism $\alpha : \widehat{C} \mapsto \widehat{D}$ such that $\alpha\theta C = \theta D$.

The graph X is called simply symmetrically self-similar if

(S2) $\text{Aut}(\widehat{C})$ acts transitively on θC ,

and doubly symmetrically self-similar or just symmetrically self-similar if

(S3) $\text{Aut}(\widehat{C})$ acts doubly transitively on θC , which means that it acts transitively on the set of ordered pairs

$$\{(x, y) ; x, y \in \theta C, x \neq y\},$$

where $g((x, y))$ is defined as $(g(x), g(y))$ for any $g \in \text{Aut}(\widehat{C})$.

Let $\text{cells}_X x$ be the number of cells which are adjacent to x . If X satisfies (S1), we write θ_X for the number of vertices in the boundary θC of some cell C .

If X satisfies Axiom (S1), then the number μ of cells in a 2-cell is independent of the choice of this 2-cell and for any cell C the graph \widehat{C} consists of μ copies of the complete graph K_{θ_X} . This fact and many more details can be found in [21].




Figure 2. The subgraph \widehat{C} for the Viček graph.

Figure 2 shows the subgraph \widehat{C} for the Viček graph. It consists of five copies of the complete graph K_4 . The vertices which are drawn fat belong to F ; they constitute the boundary of C . Note that there are vertices which are only contained in the boundary of one cell, see Figure 1, no matter whether the graph has an origin vertex or an origin cell.

LEMMA 4. — Let X be a self-similar graph satisfying Axiom (S1). Then for any $v \in F$,

$$\text{cells}_X v \cdot (\theta_X - 1) = \deg_{X_F} v.$$

Proof. — Each w in the boundary of a cell C with $v \in \theta C$ which is different from v corresponds to a neighbour of v in X_F . \square

LEMMA 5. — Let X be a simply symmetrically self-similar graph. Then the following conditions are equivalent:

- 1) X has bounded geometry.
- 2) If v is in the boundary of some cell C , then

$$\theta_X - 1 = |\{x \in C ; x \sim v\}|.$$

- 3) $\deg_X v = \deg_{X_F} v$ for all $v \in F$.

Proof. — 1) \Rightarrow 2). Suppose X has bounded geometry but

$$(3.1) \quad \frac{\deg_X v}{\text{cells}_X v} = |\{x \in C ; x \sim v\}| = a(\theta_X - 1)$$

for some $a \neq 1$. By the simple symmetry this number is independent of the choice of v and C . Since X and X_F are isomorphic with respect to ψ , by (3.1) and Lemma 4, we have

$$\deg_{X_F} \psi v = \deg_X v = \text{cells}_X v \cdot a(\theta_X - 1) = a \cdot \deg_{X_F} v$$

and for any positive integer n ,

$$\deg_{X_F} \psi^n v = a^n \cdot \deg_{X_F} v.$$

Thus either $(\deg_{X_F} \psi^n v)_{n \in \mathbb{N}}$ tends to zero, which is impossible, or X_F , and therefore also X , do not have bounded geometry.

- 2) \Rightarrow 3). If condition 2) is satisfied, then for all $v \in F$,

$$\deg_X v = \text{cells}_X v \cdot (\theta_X - 1) = \deg_{X_F} v.$$

3) \Rightarrow 1). The condition $\deg_X v = \deg_{X_F} v$ implies $\deg_X v = \deg_{X_{F^n}} v$ for all vertices v in F^n . By Axiom (F3) we have

$$\deg_X x = \deg_{X_F} \psi x = \deg_{X_{F^n}} \psi^n x$$

for all x in VX . For $x = \phi^n v$ this means

$$\deg_X \phi^n v = \deg_{X_{F^n}} \psi^n \phi^n v = \deg_{X_{F^n}} v = \deg_X v.$$

Because of Axiom (S1) the degrees of the vertices in $VX \setminus F$ are bounded. For each $v \in F$ there is an n such that $\phi^n v$ is either a vertex in $VX \setminus F$ or an origin vertex o . Thus, for any cell C ,

$$\max \{ \deg_X x ; x \in \{o\} \cup C \}$$

is the maximal degree in X . \square

4. A functional equation for Green functions on symmetrically self-similar graphs.

Let $P = (p(x, y))_{x, y \in VX}$ be the matrix of the transition operator of the simple random walk on the space of functions

$$\ell^2(X) = \left\{ f : VX \rightarrow \mathbb{C} ; \sum_{x \in VX} \deg_X x \cdot |f(x)|^2 < \infty \right\}$$

with inner product

$$(f, g) = \sum_{x \in VX} \deg_X x \cdot f(x) \overline{g(x)}.$$

Throughout this article, we consider the compact Riemannian sphere $\overline{\mathbb{C}}$, the closure of the complex numbers \mathbb{C} . The Green function for vertices x and y in VX is defined as the generating function for the n -step transition probabilities $p^{(n)}(x, y)$ from x to y . Let \mathbb{U} denote the complex, open unit disc, then

$$G(x, y | z) = \sum_{n=0}^{\infty} p^{(n)}(x, y) z^n, \quad z \in \mathbb{U}.$$

In matrix notation

$$(G(x, y | z))_{x, y \in VX} = \sum_{n=0}^{\infty} P^n z^n = (I - zP)^{-1},$$

whenever $1/z$ is not in the ℓ^2 -spectrum of P .

Following Grabner and Woess in [12] we use the combinatorics of paths to derive a functional equation for the Green functions. Let X be symmetrically self-similar. We write $\Pi_X(x, y)$ for the set of paths in X from x to y and $\Pi_X^*(x, y)$ for those paths from x to y which meet y only at the end. The *weight* of a path $\pi = (x_0, \dots, x_n)$ is defined as

$$W(\pi | z) = \prod_{i=0}^{n-1} \frac{z}{\deg_X x_i}$$

where $z \in \mathbb{C}$. For a set of paths Π we set $W_X(\Pi | z) = \sum_{\pi \in \Pi} W_X(\pi | z)$. Then we have

$$G(x, y | z) = W_X(\Pi_X(x, y) | z).$$

For vertices v and w in F and a path $\pi = (v=x_0, x_1, \dots, w=x_n)$ in X we define $\tau_j = \tau_j(\pi)$ by $\tau_0 = 0$ and if $j \geq 1$, then

$$\tau_j = \min\{i > \tau_{j-1} ; x_i \in F, x_i \neq x_{\tau_{j-1}}\},$$

where $0 \leq j \leq k$ and $k = k(\pi)$ is the last index such that the last set is nonempty. The *shadow* of π is defined as

$$\sigma(\pi) = (v=x_{\tau_0}, x_{\tau_1}, \dots, w=x_{\tau_k})$$

which is a path in X_F . Now let v and w be vertices in F that are adjacent in X_F . We define

$$\Omega(v) = \sigma^{-1}(v) = \{\pi \in \Pi_X(v, v) ; k(\pi) = 0\}$$

and

$$\Lambda(v, w) = \sigma^{-1}((v, w)) \cap \Pi_X^*(v, w) = \{\pi \in \Pi_X^*(v, w) ; k(\pi) = 1\}.$$

For any transition matrix Q of a finite directed graph we write

$$Q^*(z) = \sum_{n=0}^{\infty} (zQ)^n = (I - zQ)^{-1}, \quad \text{where } z \in \mathbb{U}.$$

The coordinates of $Q^*(z)$ are rational functions in z . They can easily be computed explicitly. Let

$$Q_B = (q_B(x, y))_{x, y \in V\widehat{C}}$$

denote the transition matrix of the simple random walk on \widehat{C} with absorbing boundary $B \subset \theta C$. This means that

$$q_B(x, y) = \begin{cases} 1/\deg_{\widehat{C}} x & \text{if } x \sim y \text{ and } x \notin B, \\ 0 & \text{otherwise.} \end{cases}$$

(Note that some authors set $q_B(x, x) = 1$ for vertices x in the absorbing boundary to get stochastic instead of substochastic transition matrices.) Let v and w be two different vertices in θC . Then we define $d, f : \mathbb{U} \rightarrow \overline{\mathbb{C}}$ by

$$d(z) = (Q_{\theta C \setminus \{v\}}^*(z))_{v, w} \cdot (\theta_X - 1) \quad \text{and} \quad f(z) = (Q_{\theta C \setminus \{v\}}^*(z))_{v, v}.$$

The function d is independent of the choice of v and w because $\text{Aut}(\widehat{C})$ acts doubly transitively on θC and f is independent of v by the simple transitivity of $\text{Aut}(\widehat{C})$ on θC . Now d is the generating function of the probabilities that the simple random walk on \widehat{C} starting in some vertex v in θC hits a vertex in $\theta C \setminus \{v\}$ for the first time after n steps, whereas f describes the n -step return probabilities of the random walk to v without hitting the absorbing boundary $\theta C \setminus \{v\}$. We call d the *transition function* and f the *return function*.

LEMMA 6. — *Let X be a graph which is symmetrically self-similar with respect to F and ϕ , and let v and w be vertices in F . Then*

$$G(v, w | z) = G(\phi v, \phi w | d(z)) \cdot f(z).$$

Proof. — For sets of paths Π_1 and Π_2 let $\Pi_1 \circ \Pi_2$ be the set of all possible concatenations of paths in Π_1 with paths in Π_2 . Then for any path

$$\pi = (v_0 = v, v_1, \dots, v_n = w)$$

in $\Pi_{X_F}(v, w)$ we have

$$\sigma^{-1}(\pi) = \Lambda(v_0, v_1) \circ \Lambda(v_1, v_2) \circ \dots \circ \Lambda(v_{n-1}, v_n) \circ \Omega(v_n).$$

For the weight function this implies

$$W(\sigma^{-1}(\pi) | z) = \prod_{i=0}^{n-1} W(\Lambda(v_i, v_{i+1}) | z) \cdot W(\Omega(v_n) | z).$$

The function d describes the transition from v_i to any vertex in a boundary of any cell which is adjacent to v_i and by Lemma 4

$$W(\Lambda(v_i, v_{i+1}) | z) = \frac{d(z)}{\text{cells}_X v_i \cdot (\theta_X - 1)} = \frac{d(z)}{\deg_{X_F} v_i}$$

and

$$W(\Omega(v_n) | z) = f(z).$$

Thus

$$G(v, w | z) = W(\Pi_X(v, w) | z) = \sum_{\pi \in \Pi_{X_F}(v, w)} W(\sigma^{-1}(\pi) | z)$$

$$\begin{aligned}
&= \sum_{\substack{\pi \in \Pi_{X_F}(v, w) \\ \pi = (v_0 = v, v_2, \dots, v_n = w)}} \prod_{i=0}^{n-1} W(\Lambda(v_i, v_{i+1}) \mid z) \cdot W(\Omega(v_n) \mid z) \\
&= \sum_{\substack{\pi \in \Pi_{X_F}(v, w) \\ \pi = (v_0 = v, v_2, \dots, v_n = w)}} \prod_{i=0}^{n-1} \frac{d(z)}{\deg_{X_F} v_i} \cdot f(z) \\
&= G_{X_F}(v, w \mid d(z)) \cdot f(z) = G(\phi v, \phi w \mid d(z)) f(z),
\end{aligned}$$

where G_{X_F} denotes the Green function on X_F . \square

Remark 1. — The main arguments in the proof of Lemma 6 are:

- (i) The graphs X and X_F are isomorphic. The set of shadows of all paths in X connecting two vertices v and w in F is the set of all paths in X_F connecting v and w .
- (ii) A transition along an edge in EX_F from v_i to v_{i+1} corresponds to all possible paths in X from v_i to v_{i+1} in the sense of the transition function d . These paths starting in v_i may return to v_i arbitrarily often and v_{i+1} is the first vertex they hit in $F \setminus \{v_i\}$. This is a concatenation of generating functions in the sense of Lemma 2.2.22 in Goulden and Jackson's book [10] and we substitute $z \mapsto d(z)$.
- (iii) By the substitution in (ii) we did not consider the possibility of returning again to w , after hitting w for the first time. The concatenation of paths that hit w for the first time with their last vertex and paths that start in w and return to w arbitrarily often corresponds to a product of generating functions with distinct configurations in the sense of Lemma 2.2.14 in [10]. Thus $G(\phi v, \phi w \mid d(z))$ has to be multiplied by $f(z)$.

5. Decomposition of the Green functions.

Let C be a cell of a symmetrically self-similar graph X . For any $x \in C$ and $y \in \bar{C}$ we define

$$h_C(x, y \mid z) = (Q_{\theta C}^*(z))_{x,y}$$

as the (x, y) -coordinate of $Q_{\theta C}^*(z)$. In other words, $h_C(x, y \mid \cdot)$ is the generating function of the n -step transition probabilities of the simple random walk on \hat{C} with absorbing boundary θC starting in x and ending

in y . If $x \in \theta C$ and $y \in C$, then we define

$$\tilde{h}(x, y | z) = \frac{\deg_{\widehat{C}} y}{\deg_{\widehat{C}} x} h(y, x | z).$$

This is the generating function for the probabilities that the simple random walk on \widehat{C} , starting in x , hits y after n steps without returning to x and without hitting any other vertex in θC . These functions h and \tilde{h} are called *inner transition functions*. Axiom (S1) implies that there are only finitely many different inner transition functions. Note that any simple random walk is reversible and that $\deg_X a \cdot G(a, b | z) = \deg_X b \cdot G(b, a | z)$ for any vertices a and b . The cell which contains some given vertex $x \in VX \setminus F$ is denoted by $C(x)$.

THEOREM 2. — Let $x, y \in VX \setminus F$ and $v, w \in F$, then

- (i) $G(v, w | z) = G(\phi v, \phi w | d(z)) \cdot f(z),$
- (ii) $G(x, w | z) = \sum_{v \in \theta C(x)} h_{C(x)}(x, v | z) \cdot G(\phi v, \phi w | d(z)) \cdot f(z),$
 $G(v, y | z) = \sum_{w \in \theta C(y)} G(\phi v, \phi w | d(z)) \cdot f(z) \cdot \tilde{h}_{C(y)}(w, y | z),$
- (iii) $G(x, y | z) = \delta_{C(x), C(y)} \cdot h_{C(x)}(x, y | z)$
 $+ \sum_{\substack{v \in \theta C(x) \\ w \in \theta C(y)}} h_{C(x)}(x, v | z) \cdot G(\phi v, \phi w | d(z))$
 $\quad \quad \quad \cdot f(z) \cdot \tilde{h}_{C(y)}(w, y | z),$

where δ denotes the usual Kronecker symbol.

Proof. — Let x and y be two vertices in $VX \setminus F$ such that $C(x) \neq C(y)$. We decompose paths of the simple random walk on X starting in x and ending in y into

- (a) a path from x to a vertex v whose vertices are all contained in $C(x)$ except for the last vertex $v \in \theta C(x)$,
- (b) a path from v to a vertex $w \in \theta C(y)$ and
- (c) a path from w to y which is contained in $C(y)$ except for its first vertex w .

This implies the product decomposition of (iii):

$$\begin{aligned} G(x, y | z) &= \sum_{\substack{v \in \theta C(x) \\ w \in \theta C(y)}} h_{C(x)}(x, v | z) \cdot G(v, w | z) \cdot \tilde{h}_{C(y)}(w, y | z) \\ &= \sum_{\substack{v \in \theta C(x) \\ w \in \theta C(y)}} h_{C(x)}(x, v | z) \cdot G(\phi v, \phi w | d(z)) \cdot f(z) \cdot \tilde{h}_{C(y)}(w, y | z). \end{aligned}$$

If x and y lie in the same cell $C(x) = C(y)$, then paths from x to y need not contain vertices of $\theta C(x)$ and we get the additional additive term $h_{C(x)}(x, y | z)$.

Let x be a vertex in $VX \setminus F$ and let w be a vertex in F . Then part (c) of the above path decomposition and therefore also the functions $\tilde{h}_{C(y)}(w, y | \cdot)$ are cancelled. This implies (ii).

The identity (i) corresponds to Lemma 6. □

By iterating formula (i) in Theorem 2, we obtain:

COROLLARY 1. — Let v and w be vertices in F^n . Then

$$G(v, w | z) = G(\phi^n v, \phi^n w | d^n(z)) \prod_{k=0}^{n-1} f(d^k(z)).$$

For arbitrary vertices x and y we have

$$G(\psi^n x, \psi^n y | z) = G(x, y | d^n(z)) \prod_{k=0}^{n-1} f(d^k(z)).$$

6. Green functions on n -cells.

Let $\text{poles}(g)$ be the set of poles and let $\text{zeroes}(g)$ be the set of zeroes of a complex function g . For a cell C of a symmetrically self-similar graph X we define

$$\text{poles}(\widehat{C}) = \bigcup_{\substack{x \in C \\ y \in \overline{C}}} \text{poles}(h_C(x, y | \cdot)).$$

Axiom (S1) of course implies that $\text{poles}(\widehat{C})$ does not depend on the choice of the cell C .

LEMMA 7. — *The zeroes of the return function f are poles of the inner transition functions h ,*

$$\text{zeroes}(f) \subset \text{poles}(\widehat{C}).$$

Proof. — We consider the simple random walk on \overline{C} with absorbing boundary $\theta C \setminus \{v\}$ for some given vertex v in θC . Let $\widehat{G}(x, y | \cdot)$ be the corresponding Green function on \widehat{C} for vertices x and y in \overline{C} and we define

$$\widehat{F}(x, y | z) = \sum_{n=0}^{\infty} q^{(n)}(x, y) z^n$$

where $q^{(n)}(x, y)$ is the n -step transition probability starting in x for the first hit in y if $n > 0$, while $q^{(0)}(x, y) = 0$. We have

$$\widehat{F}(v, v | z) = 1 - \frac{1}{\widehat{G}(v, v | z)} = 1 - \frac{1}{f(z)}.$$

A proof for this identity can be found for example in [35], Lemma 1.13 (a). Note that any zero of f is a pole of $\widehat{F}(v, v | \cdot)$. For some fixed positive integer n , let $r^{(n)}(v, x)$ be the n -step transition probability from v to x such that the random walk meets v only at the beginning. If the random walk returns to v for the first time after k steps, then either it returns after k steps, where $2 \leq k \leq n$, or after n transitions it is in a vertex $x \in C$, without hitting v during these first n steps, and then it returns to v (after $k-n$ steps, $1 \leq k-n$) in the sense of the inner transition function $h_C(x, v | \cdot)$. In other words:

$$\widehat{F}(v, v | z) = \sum_{k=0}^n q^{(k)}(v, v) \cdot z^k + \sum_{x \in C} r^{(n)}(v, x) \cdot z^n \cdot h_C(x, v | z).$$

This identity holds for any positive integer n . In particular, any pole of $\widehat{F}(v, v | \cdot)$ is a pole in $\text{poles}(\widehat{C})$ which implies $\text{zeroes}(f) \subset \text{poles}(\widehat{C})$. \square

LEMMA 8. — *Let \widehat{C}_n be the graph spanned by the closure of an n -cell C_n and let v be a vertex in the boundary θC_n . By $G_n^A(x, y | \cdot)$ we denote the Green functions of the simple random walk on \widehat{C}_n with absorbing boundary $\theta C_n \setminus \{v\}$ and by $G_n^B(x, y | \cdot)$ those with absorbing boundary θC_n . Let w be a vertex in $\theta C_n \setminus \{v\}$, and let y be a vertex in $F^{n-1} \cap C_n$. Then*

$$(6.1) \quad (\theta_X - 1) \cdot G_n^A(v, w | z) = d^n(z),$$

$$(6.2) \quad G_n^A(v, v | z) = \prod_{k=0}^{n-1} f(d^k(z)) \quad \text{and}$$

$$(6.3) \quad G_n^B(y, v | z) = h_{\phi^{n-1}(C_n \cap F^{n-1})}(\phi^{n-1}y, \phi^{n-1}v | d^{n-1}(z)).$$

Proof. — For $n = 1$ the identities (6.1), (6.2) and (6.3) are just the definitions of the functions d , f and h ; we recall that $F^0 = VX$. Suppose these identities are true for $n - 1$. Then we use the same path decomposition and the same substitution $z \mapsto d(z)$ as in the proof of Lemma 6, with the difference that in (6.1) and in (6.3) the end point of the random walk is an absorbing vertex. Thus we do not have the possibility of returning again to the end point, which means that we do not multiply $G_{n-1}^A(v, w | z)$ and $G_{n-1}^A(v, v | z)$ with $f(z)$, see Remark 1 (iii). In (6.2) the vertex v is not absorbing and we have to perform this multiplication. \square

7. Analytic continuation of Green functions.

Let X again be a doubly symmetrically self-similar graph. The Julia set of d is denoted by \mathcal{J} .

LEMMA 9. — *The point $z = 0$ is a superattracting fixed point and $z = 1$ is a repelling fixed point of the transition function d . The order of d at $z = 0$ is the diameter of the boundary of a cell.*

Proof. — Let v and w be any two different vertices in F . Then $p^{(0)}(v, w) = 0$. Since, by Axiom (F1), no pair of vertices in the boundary of a cell C is adjacent, the order of d at $z = 0$ is $\text{diam } \theta C$ which is at least 2. Thus $d(0) = 0$ and $d'(0) = 0$.

The function d is probability generating which implies $d(1) = 1$. Starting in a vertex $v \in F$, the derivative $d'(1)$ is the expected number of steps needed to hit any other vertex in F for the first time. Hence $d'(1) > 2$. \square

We call *discrete exceptional set* the set

$$\mathcal{D} = \bigcup_{n=0}^{\infty} d^{-n}(\text{poles}(f) \cup \text{poles}(\widehat{C})).$$

Let A_0 be the basin of attraction of 0. Its immediate basin of attraction (the connected component of A_0 which contains 0) is denoted by A_0^* . Let \mathcal{S} be the set of all singularities of all Green functions.

THEOREM 3. — *There is a unique analytic continuation of any Green function to $A_0^* \setminus \mathcal{S}$, which contains $A_0^* \setminus \mathcal{D}$, and*

$$\text{spec}^{-1}P \subset \overline{\mathbb{C}} \setminus (A_0^* \setminus \mathcal{S}) \subset \overline{\mathbb{C}} \setminus (A_0^* \setminus \overline{\mathcal{D}}).$$

Proof. — Let z be a complex number in A_0^* . Then there is an integer n and a neighbourhood $U(z)$ of z such that $U(z) \subset A_0^*$ and $d^n(U(z))$ is contained in the open unit disc \mathbb{U} . Let x and y be any two vertices in VX . Starting with the formula for $G(x, y | \cdot)$ in Theorem 2 and applying it again to the Green functions $G(\phi v, \phi v | d(\cdot))$, we obtain an expression with Green functions of the form $G(\phi^2 v, \phi^2 w | d^2(\cdot))$. Iterating this formula n times in this way we get a term of the form

$$(7.1) \quad G(x, y | z) = A(z) + \sum_{\text{finite sum}} B(z) \cdot G(\phi^n v, \phi^n w | d^n(z))$$

with a finite sum over Green functions of vertices $\phi^n v$ and $\phi^n w$, v and w are elements of F^n , and rational functions A and B which are analytic in $\overline{\mathbb{C}} \setminus \mathcal{D}$. We obtain the analytic continuation of $G(x, y | \cdot)$ in $U(z)$ if z is an element of $A_0^* \setminus \mathcal{D}$. This also implies $\mathcal{S} \cap A_0^* \subset \mathcal{D}$. If z is a point in $\mathcal{D} \setminus \mathcal{S}$ then again (7.1) yields the analytic continuation of $G(x, y | \cdot)$ in $U(z)$. In this case z is a pole in \mathcal{D} which is cancelled in the above sum in (7.1). Thus all Green functions are analytic in $A_0^* \setminus \mathcal{S}$. Note that the choice of n does only depend on z and not on the vertices x and y . A suitable choice for n is such that $|d^n(z)| < 1$. Uniqueness of the analytic continuation of the Green function at the point z follows from the explicit form of (7.1). In particular, if z is real, then we see that there is an $\varepsilon > 0$ such that for any x and y in VX the function $G(x, y | \cdot)$ is real and analytic in $(z - \varepsilon, z + \varepsilon)$. Thus the inversion formula for the resolvent of a self adjoint operator (see Dunford and Schwarz [9], Theorem X.6, implies that $1/z$ is not in the spectrum of the transition operator P . This method was, for example, also used by Kesten (see [18], Lemma 2.1). The values of the Green functions at z are the coordinates of the inverse operator

$$(G(x, y | z))_{x, y \in VX} = \sum_{n=0}^{\infty} z^n P^n = (I - zP)^{-1}. \quad \square$$

8. Non-polar singularities and the Julia set of the transition function.

Let x and y be vertices of a locally finite graph Y . We consider the simple random walk starting in x . Let $q^{(n)}(x, y)$ be the probability of hitting y for the first time after n -steps. We define

$$F(x, y | z) = \sum_{n=0}^{\infty} q^{(n)}(x, y) z^n.$$

The following fact is in principle well known, though hard to be found explicitly stated in the literature.

LEMMA 10. — *Let the simple random walk on a locally finite, infinite graph X be recurrent. Then for any pair of vertices x and y in VX the point $z = 1$ is a non-polar singularity of the Green function $G(x, y | \cdot)$.*

Proof. — Green functions are defined as power series around $z = 0$ with positive coefficients. Pringsheim's theorem implies that the radius of convergence is a singularity. By recurrence the radius of convergence is 1. Now suppose that $z = 1$ is a pole of order $k \geq 1$ of $G(x, y | \cdot)$. Then, near 1,

$$G(x, y | z) = \frac{1}{(1-z)^k} \tilde{G}(z),$$

where \tilde{G} is analytic in a neighbourhood of 1 and $\tilde{G}(1) \neq 0$. We rewrite

$$G(x, y | z) = \frac{F(x, y | z)}{1 - F(y, y | z)}$$

as

$$\frac{1 - F(y, y | z)}{1 - z} = \frac{(1-z)^{k-1} F(x, y | z)}{\tilde{G}(z)}.$$

Note that recurrence implies $F(x, y | 1) = 1$ for all pairs of vertices x and y . Let $z \mapsto 1^-$ along the real axis. Then we obtain $F'(y, y | 1^-) < \infty$. But this is the expected return time to y , which has to be infinite because recurrence of the simple random walk on infinite, connected graphs is equivalent to null recurrence. \square

We are interested in the spectrum of the Laplacian

$$\Delta = I - P$$

on $\ell_2(X)$, where I denotes the identity. These operators are selfadjoint and

$$\text{spec } \Delta = \{1 - \lambda ; \lambda \in \text{spec } P\},$$

$$\text{spec } P \subset [-1, 1] \quad \text{and} \quad \text{spec } \Delta \subset [0, 2].$$

We recall that \mathcal{S} is the set of all singularities of all Green functions and that $\text{spec}^{-1} P$ is the reciprocal spectrum $\{1/\lambda ; \lambda \in \text{spec } P\}$.

The next lemma follows immediately from the fact that the resolvent is analytic outside the spectrum.

LEMMA 11. — *For any locally finite graph we have*

$$\mathcal{S} \subset \text{spec}^{-1} P.$$

For the rest of this section let X again be a symmetrically self-similar graph with bounded geometry.

LEMMA 12. — *One has*

$$\mathcal{D} \subset \overline{\mathbb{R}} \setminus (-1, 1).$$

Proof. — The identities (6.1) and (6.3) imply that $d^{n-1}(\text{poles}(f))$ and $d^{n-1}(\text{poles}(\widehat{C}))$ are poles of $G_n^A(v, w | \cdot)$ and $G_n^B(y, v | \cdot)$, respectively. Let z_0 be a point in $d^{n-1}(\text{poles}(f))$. Either z_0 is a pole of

$$G_n^A(v, v | \cdot) = \prod_{k=0}^{n-1} f \circ d^k$$

or, if it is cancelled in this product, it is a zero of one of the factors $f \circ d^k$, for an integer k with $0 \leq k \leq n-2$. In other words, $z_0 \in d^{-k}(\text{zeroes}(f))$. Then Lemma 7 implies $z_0 \in d^{-k}(\text{poles}(\widehat{C}))$ and, by the above argument, z_0 is a pole of a Green function $G_{k+1}^B(y, v | \cdot)$. We conclude that all points in $d^{n-1}(\text{poles}(f))$ and $d^{n-1}(\text{poles}(\widehat{C}))$ are poles of Green functions of type G_n^A or G_n^B .

Since all our Green functions are analytic in \mathbb{U} , their singularities lie in $\overline{\mathbb{C}} \setminus \mathbb{U}$. By Lemma 11 the reciprocal values of these singularities are in the corresponding spectrum of a self-adjoint operator. Thus they are real and

$$d^{n-1}(\text{poles}(f)) \cup d^{n-1}(\text{poles}(\widehat{C})) \subset \overline{\mathbb{R}} \setminus (-1, 1),$$

for any positive integer n , which implies

$$\mathcal{D} \subset \overline{\mathbb{R}} \setminus (-1, 1). \quad \square$$

Let us define two complex numbers z_1 and z_2 as equivalent if there are integers m and n such that $d^m(z_1) = d^n(z_2)$. A complex number is called *exceptional* if the corresponding equivalence class is finite. The Julia set of d is denoted by \mathcal{J} , the Fatou set by \mathcal{F} .

THEOREM 4. — *For the reciprocal spectrum we have*

$$\text{spec}^{-1} P = \mathcal{J} \cup \mathcal{S} \subset \mathcal{J} \cup \mathcal{D}.$$

It is the set of points z in $\overline{\mathbb{R}} \setminus (-1, 1)$ such that there is a Green function

which cannot be continued analytically from both half spheres of $\overline{\mathbb{C}} \setminus \overline{\mathbb{R}}$ to z . The super attracting fixed point $z = 0$ is the only attracting fixed point of d , $\mathcal{F} = A_0^* = A_0$ and \mathcal{J} is the set of accumulation points of \mathcal{D} . Either d has no exceptional points or $z = 0$ is the only exceptional point of d . In particular, the discrete exceptional set \mathcal{D} does not contain exceptional points.

Proof. — The immediate basin of attraction A_0^* of $z = 0$ is the component of the Fatou set \mathcal{F} which contains $z = 0$, see Theorem 6.3.1 in the book of Beardon [6]. Suppose there is a point z_0 in ∂A_0^* , which is a subset of \mathcal{J} , and a Green function G_0 which continues analytically from A_0^* to an open neighbourhood $U(z_0)$ of z_0 . The Julia set contains all repelling fixed points, and the backward iterates of any point in \mathcal{J} are dense in \mathcal{J} , see for example [6], Theorem 4.2.7 (ii) or Theorems 1.1 and 1.6 in Chapter III of the book of Carlson and Gamelin [7]. Thus there is a \tilde{z} in $U(z_0)$ and an integer n such that $d^n(\tilde{z}) = 1$. The function d is finitely branched. Corollary 1 implies that the Green function G_0 continues analytically to an open neighbourhood of $z = 1$ except for a finite set of singularities. Now $z = 1$ is an isolated singularity of G_0 which is non-polar (see Lemma 10), therefore it is an essential singularity. Corollary 1 also implies that all d -backward iterates of $z = 1$ are essential singularities of G_0 . These backward iterates are dense in \mathcal{J} in contradiction to the assumption that G_0 continues analytically to an open neighbourhood $U(z_0)$ of z_0 in \mathcal{J} . Since all Green functions are analytic on $\mathbb{C} \setminus \text{spec}^{-1}P$, this implies $\partial A_0^* \subset \text{spec}^{-1}P$. The reciprocal spectrum $\text{spec}^{-1}P$ is real and it follows that $z = 0$ is the only attracting fixed point of d and $A_0^* = A_0 = \mathcal{F}$. This implies $\mathcal{J} = \partial A_0^*$. The set $\mathcal{S} \cap \mathcal{F}$ is contained in $\text{spec}^{-1}P$ and Theorem 3 implies that $\mathcal{F} \setminus \mathcal{S}$ is contained in $\mathbb{C} \setminus \text{spec}^{-1}P$. We conclude, $\text{spec}^{-1}P = \mathcal{J} \cup \mathcal{S}$. Since d is rational of degree at least 2, there are at most two exceptional points (see [6], Theorem 4.1.2). Exceptional points are contained in the Fatou set (see [6], Corollary 4.1.3). Thus $z = 0$ is the only exceptional point, or there are no exceptional points. By Lemma 12, there are no exceptional points in \mathcal{D} and Theorem 4.2.7 (i) in [6] now implies that \mathcal{J} is the set of accumulation points of \mathcal{D} . \square

Combining Theorems 3 and 4, we obtain:

THEOREM 5. — One has

$$\mathcal{J} \subset \text{spec}^{-1}P = \mathcal{J} \cup \mathcal{S} \subset \mathcal{J} \cup \mathcal{D} = \overline{\mathcal{D}} \subset \overline{\mathbb{R}} \setminus (-1, 1).$$

Note that $\text{spec } \Delta = \{1 - \lambda; 1/\lambda \in \text{spec}^{-1}P\}$.

9. The Julia set of the transition function.

From the theory of iteration of rational maps we know that the Julia set of a rational function of degree at least 2 is either connected in the complex sphere $\overline{\mathbb{C}}$ or has uncountably many components and each point in the Julia set is an accumulation point of infinitely many distinct components of \mathcal{J} , see for example [6], Theorem 5.7.1 or [16], Lemma 4. In our case the Julia set is real and we obtain the following theorem:

THEOREM 6. — *Either the Julia set of the transition function of a symmetrically self-similar graph is a real interval in $\overline{\mathbb{C}}$ or it is a Cantor set.*

Example 1. — Let o be an origin vertex of the two sided infinite line. The graphs \widehat{C} can be paths of length n for any integer n . Technical calculations show that for the corresponding transition function d_n we have

$$d_n(z) = \frac{z^n}{\sum_{k=0}^{\lfloor \frac{1}{2}n \rfloor} \binom{n}{2k} (1-z^2)^k},$$

these functions are conjugated to the Chebychev polynomials T_n ,

$$T_n(z) = \cos(n \cos^{-1} z) = \frac{1}{d_n(1/z)},$$

and $\mathcal{J} = \overline{\mathbb{R}} \setminus (-1, 1)$. We consider this Green function as the solution

$$\prod_{k=0}^{\infty} f(d_2^k(z)), \quad \text{for } f(z) = \frac{2}{2-z^2} \quad \text{and} \quad d_2(z) = \frac{z^2}{2-z^2},$$

of the functional equation

$$G(o, o | z) = f(z) G(o, o | d_2(z)),$$

where the graphs \widehat{C} spanned by the closures of the cells are paths of length 2. This solution converges as an infinite product in $\mathbb{C} \setminus \mathcal{J}$ to

$$G(o, o | z) = \frac{1}{\sqrt{1-z^2}}, \quad z \in \mathbb{C} \setminus ((-\infty, 1] \cup [1, \infty)).$$

The fact that the product diverges on \mathcal{J} corresponds to the existence of two different branches of the square root.

THEOREM 7. — *Let \mathcal{J} be the Julia set of the transition function of a symmetrically self-similar graph. Then the reciprocal spectrum $\text{spec}^{-1} P$ coincides with the set of singularities of all Green functions \mathcal{S} .*

Proof. — The branches of the analytic continuations from \mathbb{U} to the half spheres in $\overline{\mathbb{C}} \setminus \overline{\mathbb{R}}$ coincide at the real part of the Fatou set $\mathcal{F} \cap \overline{\mathbb{R}}$ which is dense in \mathcal{J} because \mathcal{J} is supposed to be a Cantor set. None of these branches can be continued locally analytically to any point in \mathcal{J} since this local continuation would then have to coincide with both branches of the Green function, which is impossible since no Green function can be continued analytically to a point in the Julia set. For points in $\mathcal{D} \setminus \mathcal{J}$ we have already proved that they are in the spectrum if and only if they are singularities of a Green function, see Theorem 3 and Lemma 11. \square

For symmetrically self-similar graphs with bounded geometry we conjecture that the Julia set \mathcal{J} is an interval if and only if the cell-graph \widehat{C} is a finite line, otherwise it is a Cantor set. According to Lemma 2, the cell graph \widehat{C} is a finite line if and only if X is a ‘star’ consisting of finitely many one-sided infinite lines which have exactly one vertex in common.

Sabot showed in [33] that the integrated density of states for nested fractals with at least three essential fixed points is completely generated by the so-called Neumann-Dirichlet eigenvalues. In [26] Malozemov and Teplyaev proved similar results for symmetric self-similar graphs where each cell has exactly two boundary points.

Acknowledgements. — The author wants to thank Peter Grabner, Elmar Teufl and Wolfgang Woess for fruitful discussions. Representative for many contributions: the idea of choosing a proper set F in Definition 1 (self-similarity of graphs) is due to Peter Grabner, Elmar Teufl found a right idea for the proof of Lemma 7 and Wolfgang Woess is the author of the proof of Lemma 10.

BIBLIOGRAPHY

- [1] M.T. BARLOW, J. KIGAMI, Localized eigenfunctions of the Laplacian on p.c.f. self-similar sets, *J. London Math. Soc.*, 56-2 (1997), 320–332.
- [2] M.T. BARLOW, E.A. PERKINS, Brownian motion on the Sierpiński gasket, *Prob. Theory Related Fields*, 79-4 (1988), 543–623.
- [3] L. BARTHOLDI, Croissance de groupes agissant sur des arbres, Ph. D. thesis, Université de Genève, 2000.

- [4] L. BARTHOLDI, R.I. GRIGORCHUK, On the spectrum of Hecke type operators related to some fractal groups, *Tr. Mat. Inst. Steklova (Din. Sist., Avtom. i Beskon. Gruppy)*, 231 (2000), 5-45.
- [5] L. BARTHOLDI, R.I. GRIGORCHUK, V. NEKRASHEVYCH, From fractal groups to fractal sets, in ‘Fractals in Graz 2001’, P.J. Grabner and W. Woess, eds., Birkhäuser, 2002.
- [6] A.F. BEARDON, Iteration of rational functions, Springer-Verlag, New York, 1991.
- [7] L. CARLESON, T.W. GAMELIN, Complex dynamics, Springer-Verlag, New York, 1993.
- [8] P.G. DOYLE, J.L. SNELL, Random walks and electric networks, Math. Association of America, Washington, DC, 1984.
- [9] N. DUNFORD, J.T. SCHWARTZ, Linear Operators I-II, Interscience, New York, 1963.
- [10] I.P. GOULDEN, D.M. JACKSON, Combinatorial enumeration, John Wiley & Sons, New York, 1983.
- [11] P.J. GRABNER, Functional iterations and stopping times for Brownian motion on the Sierpiński gasket, *Mathematika*, 44-2 (1997), 374–400.
- [12] P.J. GRABNER, W. WOESS, Functional iterations and periodic oscillations for simple random walk on the Sierpiński graph, *Stochastic Process. Appl.*, 69-1 (1997), 127–138.
- [13] B.M. HAMBLY, On the asymptotics of the eigenvalue counting function for random recursive Sierpinski gaskets, *Prob. Theory Related Fields*, 117-2 (2000), 221–247.
- [14] B.M. HAMBLY, V. METZ, The homogenization problem for the Vicsek set, *Stochastic Process. Appl.*, 76-2 (1998), 167–190.
- [15] J.E. HUTCHINSON, Fractals and self-similarity, *Indiana Univ. Math. J.*, 30-5 (1981), 713–747.
- [16] C. INNINGER, Rational iteration, Dissertation, Universitätsverlag Rudolf Trauner, University of Linz, 2001.
- [17] O.D. JONES, Transition probabilities for the simple random walk on the Sierpiński graph, *Stochastic Process. Appl.*, 61-1 (1996), 45–69.
- [18] H. KESTEN, Symmetric random walks on groups, *Trans. Amer. Math. Soc.*, 1959.
- [19] J. KIGAMI, Harmonic calculus on p.c.f. self-similar sets., *Trans. Amer. Math. Soc.*, 1993.
- [20] B. KRÖN, Spectral and structural theory of infinite graphs, PhD. thesis, Graz University of Technology, 2001.
- [21] B. KRÖN, Growth of self-similar graphs, preprint, 2002.
- [22] B. KRÖN, E. TEUFL, Asymptotics of the transition probabilities of the simple random walk on self-similar graphs, preprint, 2002.
- [23] T. LINDSTRØM, Brownian motion on nested fractals, *Mem. Amer. Math. Soc.*, 83 (1990), 420.
- [24] L. MALOZEMOV, The integrated density of states for the difference Laplacian on the modified Koch graph, *Comm. Math. Phys.*, 156-2 (1993), 387–397.
- [25] L. MALOZEMOV, Random walk and chaos of the spectrum. Solvable model, *Chaos Solitons Fractals*, 5-6 (1995), 895–907.

- [26] L. MALOZEMOV, A. TEPLYAEV, Pure point spectrum of the Laplacians on fractal graphs, *J. Funct. Anal.*, 129-2 (1995), 390–405.
- [27] L. MALOZEMOV, A. TEPLYAEV, Self-similarity, operators and dynamics, preprint, 2001.
- [28] V. METZ, How many diffusions exist on the Vicsek snowflake?, *Acta Appl. Math.*, 32-3 (1993), 227–241.
- [29] C.S.J.A. NASH-WILLIAMS, Random walk and electric currents in networks, *Proc. Cambridge Phil. Soc.*, 55 (1959), 181–194.
- [30] R. RAMMAL, Random walk statistics on fractal structures, *J. Stat. Phys.*, 36-5-6 (1984), 547–560.
- [31] R. RAMMAL, Spectrum of harmonic excitations on fractals, *J. Physique*, 45-2 (1984), 191–206.
- [32] R. RAMMAL, TOULOUSE, Random walks on fractal structures and percolation clusters, *J. Physique – Lettres* 44, 36-L13–L22 (1983).
- [33] C. SABOT, Pure point spectrum for the Laplacian on unbounded nested fractals, *J. Funct. Anal.*, 173-2 (2000), 497–524.
- [34] A. TEPLYAEV., Spectral analysis on infinite Sierpiński gaskets, *J. Funct. Anal.*, 159-2 (1998), 537–567.
- [35] W. WOESS, *Random Walks on Infinite Graphs and Groups*, Cambridge University Press, Cambridge, 2000.

Manuscrit reçu le 8 avril 2002,
accepté le 16 mai 2002.

Bernhard KRÖN,
Erwin Schrödinger Institute (ESI)
Boltzmanngasse 9
1090 Wien (Austria).
bernhard.kroen@univie.ac.at.