



ANNALES

DE

L'INSTITUT FOURIER

Jens HORNBOSTEL & Guido KINGS

On non-commutative twisting in étale and motivic cohomology

Tome 56, n° 4 (2006), p. 1257-1279.

http://aif.cedram.org/item?id=AIF_2006__56_4_1257_0

© Association des Annales de l'institut Fourier, 2006, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

ON NON-COMMUTATIVE TWISTING IN ÉTALE AND MOTIVIC COHOMOLOGY

by Jens HORNBOSTEL & Guido KINGS

ABSTRACT. — This article confirms a consequence of the non-abelian Iwasawa main conjecture. It is proved that under a technical condition the étale cohomology groups $H^1(\mathcal{O}_K[1/S], H^i(\overline{X}, \mathbb{Q}_p(j)))$, where $X \rightarrow \text{Spec } \mathcal{O}_K[1/S]$ is a smooth, projective scheme, are generated by twists of norm compatible units in a tower of number fields associated to $H^i(\overline{X}, \mathbb{Z}_p(j))$. Using the “Bloch-Kato-conjecture” a similar result is proven for motivic cohomology with finite coefficients.

RÉSUMÉ. — Cet article confirme une conséquence de la conjecture principale de la théorie d’Iwasawa non abélienne. On démontre que, sous une condition technique, les groupes de cohomologie étale $H^1(\mathcal{O}_K[1/S], H^i(\overline{X}, \mathbb{Q}_p(j)))$, où $X \rightarrow \text{Spec } \mathcal{O}_K[1/S]$ est un schéma projectif lisse, sont engendrés par des unités tordues compatible par rapport aux normes dans une tour de corps de nombres associés à $H^i(\overline{X}, \mathbb{Z}_p(j))$. On établit un résultat similaire pour la cohomologie motivique à coefficients finis en utilisant la conjecture de Bloch-Kato.

Introduction

One of the most astonishing consequences of the non-abelian Iwasawa main conjecture is the twist invariance of the zeta elements, which implies that all motivic elements should be twists of norm compatible units in (big) towers of number fields. More precisely one expects that for a \mathbb{Z}_p -lattice T in a motive with p -adic realization V the image of the twisting map (see 1.3 below)

$$\left(\varprojlim_n H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}_p(1))\right) \otimes T(j-1) \longrightarrow H^1(\mathcal{O}_K[1/S], T(j))$$

Keywords: Étale cohomology, motivic cohomology, non-commutative Iwasawa-theory.
Math. classification: 11R23, 14G40, 14F42, 11R32.

generates a subgroup of finite index. Here the inverse limit runs over the number fields $K_n := K(T/p^n)$ obtained from K by adjoining the elements T/p^n . Moreover, the image of this map should have a motivic meaning, that is the elements should be in the image of the p -adic regulator from motivic cohomology. (This idea is developed in [10] and builds on ideas of Kato [14]).

The philosophy of twisting with the cyclotomic character originates from work of Iwasawa, Tate and Soulé. This already lead to many interesting results. Here Kato's work [15] on the Birch-Swinnerton-Dyer conjecture is the most spectacular example. Earlier Soulé used this idea in the case of Tate motives in his deep investigations about the connection of K -theory and étale cohomology for number rings [26]. He also pointed the way to applications to CM-elliptic curves [29].

The first goal of this paper is to construct the above twisting map and to show that it has for $j \gg 0$ indeed finite cokernel assuming the very reasonable condition that the Iwasawa μ -invariant of the number field K vanishes. This result supports the non-abelian Iwasawa main conjecture and has the astonishing consequence that for the construction of interesting elements in étale cohomology one is reduced to construct norm compatible systems of units in towers of number fields. The proof of this result relies on certain finiteness results of Coates and Sujatha.

In the second part of the paper we consider the statement that the resulting elements are in the image of the regulator from motivic cohomology. Our results in this direction give a hint that the elements obtained as twists of units are motivic. Using the “Bloch-Kato-conjecture” for all fields (as announced by Voevodsky), we prove that there is a twisting map for motivic cohomology compatible with the one for étale cohomology under the cycle class map. This uses constructions of Levine and Geisser.

The authors like to thank Coates and Sujatha for useful discussions and for making available the results of [3] before their publication. The authors are indebted to Geisser for insisting to use motivic cohomology with finite coefficients instead of K -theory in the formulation of the results in the second part.

1. Non-commutative twisting in étale cohomology

In this section we describe the étale situation. All cohomology groups in this paper are étale cohomology groups unless explicitly labeled otherwise.

1.1. The twisting map in étale cohomology

Let K be a number field with ring of integers \mathcal{O}_K . Fix a prime number $p > 2$ (we assume $p \neq 2$ only to have \mathcal{O}_K cohomological dimension 2) and a finite set of primes S of \mathcal{O}_K , which contains the primes dividing p . As usual let

$$G_S := \text{Gal}(K_S/K)$$

be the Galois group of K_S/K , where K_S is the maximal outside of S unramified extension field in a fixed algebraic closure \bar{K} of K . Let T be a finitely generated \mathbb{Z}_p -module with a continuous G_S -action

$$\rho : G_S \longrightarrow \text{Aut}_{\mathbb{Z}_p}(T).$$

We will consider T also as étale sheaf on $\mathcal{O}_K[1/S]$ (see e.g. [5, p. 640] using the fixed embedding into \bar{K} as base point) and write $V := T \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ for the associated \mathbb{Q}_p -sheaf. The most important example is $T = H^i(\bar{X}, \mathbb{Z}_p(j))$, where $\bar{X} := X \times_{\mathcal{O}_K[1/S]} \bar{K}$ and X is a smooth and proper scheme over $\mathcal{O}_K[1/S]$. Let $\mathcal{G} := \text{im } \rho$ be the image of ρ . If we define finite groups

$$G_n := \text{im}\{\rho_n : G_S \longrightarrow \text{Aut}_{\mathbb{Z}/p^n\mathbb{Z}}(T/p^nT)\}$$

then we also have $\mathcal{G} \cong \varprojlim_n G_n$. Note that \mathcal{G} is a p -adic Lie group. The Iwasawa algebra of \mathcal{G} is by definition the continuous group ring

$$\Lambda(\mathcal{G}) := \varprojlim_n \mathbb{Z}_p[G_n] \cong \varprojlim_n \mathbb{Z}/p^n[G_n].$$

Note that $\Lambda(\mathcal{G})$ is flat over \mathbb{Z}_p . If \mathcal{G} is pro- p then $\Lambda(\mathcal{G})$ is a noetherian local ring (see [17, II.2.2.2 and V.2.2.4]) as \mathcal{G} is by construction p -adic analytic. The action of G_S on $\Lambda(\mathcal{G})$ factors through \mathcal{G} and this acts on $\Lambda(\mathcal{G})$ via $\mathcal{G} \subset \Lambda(\mathcal{G})^*$. We denote by K_∞ the field fixed by the kernel of ρ and by K_n the field fixed by the kernel of ρ_n , so that $K_\infty = \bigcup_n K_n$ and $\text{Gal}(K_\infty/K) \cong \mathcal{G}$. Note that $\mathcal{O}_{K_n}[1/S]$ is finite and étale over $\mathcal{O}_K[1/S]$.

Example 1.1. — To make the above definitions more concrete, consider the following important example. Let E/K be an elliptic curve without complex multiplication and $T_p E := \varprojlim_n E[p^n]$ its Tate-module. We have

$$K_n := K(E[p^n]).$$

It is a well-known result of Serre that the image of the Galois group G_S in $\text{Aut}_{\mathbb{Z}_p}(T_p E)$ has finite index and is equal to $\text{Aut}_{\mathbb{Z}_p}(T_p E)$ for almost all p . If we assume the latter case, we have in the above notation $G_n \cong \text{Gl}_2(\mathbb{Z}/p^n)$ and $\mathcal{G} \cong \text{Gl}_2(\mathbb{Z}_p)$.

The following proposition for the étale cohomology should be well-known. For convenience of the reader and to explain the normalizations of the action in detail, we give the proof in an appendix.

PROPOSITION 1.2 (see Appendix B). — *There are canonical isomorphisms of compact finitely generated $\Lambda(\mathcal{G})$ -modules*

$$R\Gamma(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \otimes_{\mathbb{Z}_p}^{\mathbb{L}} T \xrightarrow{\cong} R\Gamma(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T),$$

which induce a map

$$H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \otimes_{\mathbb{Z}_p} T \longrightarrow H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T),$$

and

$$\varprojlim_n H^i(\mathcal{O}_{K_n}[1/S], T/p^n) \cong \varprojlim_n H^i(\mathcal{O}_{K_n}[1/S], T) \cong H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T)$$

(limit over the corestriction maps). Here the $\Lambda(\mathcal{G})$ -module structure on the left hand side is induced by the action of G_n on $H^i(\mathcal{O}_{K_n}[1/S], T/p^n)$. On the right hand side the $\Lambda(\mathcal{G})$ -action is induced by the action on $\Lambda(\mathcal{G})$ via multiplication with the inverse and T has its natural $\Lambda(\mathcal{G})$ -module structure induced by the action of \mathcal{G} .

Remark. — If one assumes that T is a free \mathbb{Z}_p -module, then one gets of course an isomorphism

$$H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \otimes_{\mathbb{Z}_p} T \xrightarrow{\cong} H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T).$$

The extra generality to allow T to be finitely generated is needed, if one wants to consider $T := H^i(\bar{X}, \mathbb{Z}_p)$ for a smooth, projective variety X over $\mathcal{O}_K[1/S]$.

We can now make our main definition:

DEFINITION 1.3. — *The twisting map*

$$\text{Tw}_T : H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \otimes_{\mathbb{Z}_p} T \longrightarrow H^i(\mathcal{O}_K[1/S], T)$$

is the composition of the map of Proposition 1.2 with the map

$$\epsilon : H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T) \longrightarrow H^i(\mathcal{O}_K[1/S], T)$$

induced by the augmentation $\epsilon : \Lambda(\mathcal{G}) \rightarrow \mathbb{Z}_p$.

Our goal is to show that the twisting map is surjective in certain cases after tensoring with \mathbb{Q}_p . In particular it allows to construct elements in $H^i(\mathcal{O}_K[1/S], T)$ starting from (corestriction or norm compatible) elements in

$$H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \cong \varprojlim_n H^i(\mathcal{O}_{K_n}[1/S], \mathbb{Z}_p).$$

We will apply this in the case where $T = H^r(X \times_{\mathcal{O}_K[1/S]} \bar{K}, \mathbb{Z}_p)$.

1.2. Another description of the twisting map

To make the similarity with the twisting map in motivic cohomology and in p -adic K -theory more apparent, we describe the twisting map at finite level.

Fix an integer $n > 0$. We have by Shapiro’s Lemma

$$H^i(\mathcal{O}_K[1/S], \mathbb{Z}/p^n\mathbb{Z}[G_n]) \cong H^i(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n\mathbb{Z})$$

using the identification as explained in Appendix B. As T/p^nT is a trivial sheaf over $\mathcal{O}_{K_n}[1/S]$, we have $T/p^nT \cong H^0(\mathcal{O}_{K_n}[1/S], T/p^nT)$ and the cup product gives an isomorphism

$$H^i(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n\mathbb{Z}) \otimes T/p^nT \xrightarrow{\cup} H^i(\mathcal{O}_{K_n}[1/S], T/p^nT).$$

Together with the corestriction (= trace map in étale cohomology)

$$H^i(\mathcal{O}_{K_n}[1/S], T/p^nT) \longrightarrow H^i(\mathcal{O}_K[1/S], T/p^nT)$$

we get a map

$$(1.1) \quad H^i(\mathcal{O}_K[1/S], \mathbb{Z}/p^n\mathbb{Z}[G_n]) \otimes T/p^nT \longrightarrow H^i(\mathcal{O}_K[1/S], T/p^nT).$$

Observe that by Mittag-Leffler we have

$$\varprojlim_n H^i(\mathcal{O}_K[1/S], T/p^nT) \cong H^i(\mathcal{O}_K[1/S], T).$$

LEMMA 1.4. — *The inverse limit with respect to the trace map and reduction on the coefficients of the maps (1.1) coincides with the twisting map in Definition 1.3.*

Proof. — Straightforward. □

1.3. Tate twist

Let $K^{\text{cyc}} := \bigcup_n K(\mu_{p^n})$ be the field K with all the p -th power roots of unity μ_{p^∞} adjoined. We will assume that K_∞ contains K^{cyc} . If this is not the case it can be achieved by considering $K_n(\mu_{p^n})$ instead of K_n . Let $\Gamma := \text{Gal}(K^{\text{cyc}}/K)$, then we have a map $\mathcal{G} \rightarrow \Gamma$ and we denote its kernel by \mathcal{H} . This map induces a surjection

$$(1.2) \quad \Lambda(\mathcal{G}) \longrightarrow \Lambda(\Gamma).$$

The cyclotomic character induces an inclusion of Γ in \mathbb{Z}_p^* and the associated free $\mathbb{Z}_p - \Gamma$ -module of rank 1 is denoted by $\mathbb{Z}_p(1)$. As usual let

$$\mathbb{Z}_p(j) := \mathbb{Z}_p(1)^{\otimes j} \quad \text{and} \quad T(j) := T \otimes \mathbb{Z}_p(j).$$

We will consider the following important variant of the twisting map, given by combining Definition 1.3 and Proposition 1.2 for $T = \mathbb{Z}_p(1)$ and $T(j - 1)$:

$$(1.3) \quad H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \otimes_{\mathbb{Z}_p} T(j - 1) \xrightarrow{\text{Tw}_{T(j-1)}} H^i(\mathcal{O}_K[1/S], T(j)).$$

Note that $H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) = \varprojlim_n H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}_p(1))$ and that by Kummer theory we have for any m, n an exact sequence

$$(1.4) \quad 0 \rightarrow \mathcal{O}_{K_n}[1/S]^* \otimes \mathbb{Z}/p^m \rightarrow H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^m(1)) \rightarrow \text{Cl}(\mathcal{O}_{K_n}[1/S])[p^m] \rightarrow 0.$$

Here $\text{Cl}(\mathcal{O}_{K_n}[1/S])[p^m]$ is the p^m -torsion subgroup of the class group of $\mathcal{O}_{K_n}[1/S]$. Taking the limit first over m then over n and using that $\text{Cl}(\mathcal{O}_{K_n}[1/S])$ is finite, we get

$$(1.5) \quad \varprojlim_n \mathcal{O}_{K_n}[1/S]^* \otimes \mathbb{Z}_p \cong \varprojlim_n H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}_p(1)).$$

We have a twisted variant of Lemma 1.4. Namely, the map $\text{Tw}_{T(j-1)}$ of (1.3) is again given by taking cup products

$$H^i(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \otimes_{\mathbb{Z}_p} T/p^n(j - 1)$$

and passing to the limit, using again Proposition 1.2.

1.4. The cokernel of the twisting map

To study the cokernel of the twisting map, we factor the augmentation into $\Lambda(\mathcal{G}) \rightarrow \Lambda(\Gamma) \rightarrow \mathbb{Z}_p$ using (1.2) and get:

$$(1.6) \quad H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \rightarrow H^i(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)).$$

The analysis of the cokernel of the twisting map $\text{Tw}_{T(j-1)}$ will proceed in two steps. The first is to investigate the cokernel of (1.6). The second step treats then the cokernel of the map induced by the augmentation $\Lambda(\Gamma) \rightarrow \mathbb{Z}_p$:

$$(1.7) \quad H^i(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)) \rightarrow H^i(\mathcal{O}_K[1/S], T(j)).$$

LEMMA 1.5. — *There is a spectral sequence*

$$E_2^{r,s} = \text{Tor}_r^{\Lambda(\mathcal{G})} (H^s(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \Lambda(\Gamma)) \implies H^{s-r}(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)).$$

Proof. — The projection formula (see e.g. [36, Exercise 10.8.3]) in the derived category gives

$$R\Gamma(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\mathcal{G})}^{\mathbb{L}} \Lambda(\Gamma) \cong R\Gamma(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)).$$

Taking cohomology gives the desired spectral sequence. □

COROLLARY 1.6. — *There is an exact sequence*

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \otimes_{\mathbb{Z}_p} T(j-1) \xrightarrow{Tw_{T(j-1)}} H^1(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)) \longrightarrow \text{Tor}_1^{\Lambda(\mathcal{G})} (H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \Lambda(\Gamma)).$$

Proof. — As $p > 2$ the cohomological p -dimension of $\mathcal{O}_K[1/S]$ is 2 and the result follows from the spectral sequence and the fact that the twisting map factors through

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \longrightarrow H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\mathcal{G})} \Lambda(\Gamma).$$

□

LEMMA 1.7. — *The canonical isomorphism $\Lambda(\mathcal{G}) \otimes_{\Lambda(\mathcal{H})} \mathbb{Z}_p \cong \Lambda(\Gamma)$ induces isomorphisms for all r :*

$$\text{Tor}_r^{\Lambda(\mathcal{G})} (H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \Lambda(\Gamma)) \cong \text{Tor}_r^{\Lambda(\mathcal{H})} (H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p).$$

In particular, one gets from Corollary 1.6 an exact sequence

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \otimes_{\mathbb{Z}_p} T(j-1) \longrightarrow H^1(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)) \longrightarrow \text{Tor}_1^{\Lambda(\mathcal{H})} (H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p).$$

Proof. — The isomorphism $\Lambda(\mathcal{G}) \otimes_{\Lambda(\mathcal{H})} \mathbb{Z}_p \cong \Lambda(\Gamma)$ can be checked at finite level as \mathbb{Z}_p is a finitely generated $\Lambda(\mathcal{H})$ -module. Then

$$\mathbb{Z}_p[G_n] \otimes_{\mathbb{Z}_p[H_n]} \mathbb{Z}_p \cong \mathbb{Z}_p[G_n/H_n]$$

and the claim is obvious. In particular, for finitely generated $\Lambda(\mathcal{G})$ -modules M is the functor $M \mapsto M \otimes_{\Lambda(\mathcal{G})} \Lambda(\Gamma)$ isomorphic to $M \mapsto M \otimes_{\Lambda(\mathcal{H})} \mathbb{Z}_p$. □

From this lemma and the factorization of the twisting map it is clear that the cokernel of the twisting map is controlled by

$$\mathrm{Tor}_1^{\Lambda(\mathcal{H})}(H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p)$$

and by $\mathrm{Tor}_1^{\Lambda(\Gamma)}(H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p)$. To say something about these groups we need some results of Coates and Sujatha on the finiteness of the H^2 's involved.

1.5. The conjectured ranks of the étale cohomology

For the convenience of the reader, we recall the conjecture of Jannsen [12] about the ranks of the étale cohomology.

Let X be a smooth, projective scheme over $\mathcal{O}_K[1/S]$ and denote by

$$\bar{X} := X \times_{\mathcal{O}_K[1/S]} \bar{K}$$

the base change to the algebraic closure.

CONJECTURE 1.8 (Jannsen). — *For $i + 1 < j$ or $i + 1 > 2j$ one has*

$$H^2(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Q}_p(j))) = 0.$$

As a consequence one obtains (for $p \neq 2$) the following formula for the dimension of the H^1 : for $i + 1 < j$

$$\dim_{\mathbb{Q}_p} H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Q}_p(j))) = \dim_{\mathbb{R}} H^i(X \times_{\mathbb{Q}} \mathbb{C}, \mathbb{R}(j))^+,$$

where “+” denotes the invariants under complex conjugation, which acts on $X \times_{\mathbb{Q}} \mathbb{C}$ and on $\mathbb{R}(j) = (2\pi i)^j \mathbb{R}$.

Moreover in analogy with Beilinson’s conjecture that the regulator from K -theory to Beilinson-Deligne cohomology is an isomorphism for $i + 1 < j$, Jannsen also conjectures that the Soulé regulator

$$r_p : H_{\mathrm{mot}}^{i+1}(X, \mathbb{Z}(j)) \otimes_{\mathbb{Z}} \mathbb{Q}_p \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Q}_p(j)))$$

is an isomorphism for $i + 1 < j$. It is shown in [29] that for p -adic K -theory the above regulator is surjective if $j \gg 0$. This should be compared with the result in 2.6.

1.6. Finiteness conditions for H^2

This section contains only slight modifications of results of Coates and Sujatha [3]. We thank them very much for making these results available to us before their publication. One should also compare this section with the Appendix B in Perrin-Riou [23, Prop. B.2].

Let L^{cyc} (resp. L_∞) be the maximal unramified abelian p -extension of K^{cyc} (resp. K_∞), in which every prime above p splits completely.

PROPOSITION 1.9 (Coates-Sujatha [3]). — Assume that $\mathcal{G} = \text{Gal}(K_\infty/K)$ is a pro- p -group, then the following conditions are equivalent:

- i) $\text{Gal}(L^{\text{cyc}}/K^{\text{cyc}})$ is a finitely generated \mathbb{Z}_p -module;
- ii) $\text{Gal}(L_\infty/K_\infty)$ is a finitely generated $\Lambda(\mathcal{H})$ -module;
- iii) $H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T)$ is a finitely generated \mathbb{Z}_p -module;
- iv) $H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T)$ is a finitely generated $\Lambda(\mathcal{H})$ -module.

In particular, if these equivalent conditions are satisfied, the $\Lambda(\Gamma)$ -module $H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T)$ is torsion, i.e., the weak Leopold conjecture is true.

Remarks. — The first condition is equivalent to the famous $\mu = 0$ conjecture of Iwasawa (see [22, Ch. XI, Thm. 11.3.18]), which is known to be true for K/\mathbb{Q} abelian.

Note also that the statements i) and ii) in the proposition are independent of the Galois representation T .

Proof. — The proof of the proposition can be found in Coates and Sujatha [3] in the case of the Tate module for an elliptic curve. The case for an arbitrary Galois representation T is the same. More precisely, the equivalence i) \Leftrightarrow ii) is Lemma 3.7, i) \Leftrightarrow iii) is Thm. 3.4. in *loc. cit.* To prove iii) \Leftrightarrow iv), we have from the spectral sequence in Lemma 1.5 and the vanishing of étale cohomology for $s > 2$ that

$$H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T) \otimes_{\Lambda(\mathcal{H})} \mathbb{Z}_p \cong H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T)$$

and the claim follows from Nakayama’s lemma. □

Example 1.10. — Let E/\mathbb{Q} be an elliptic curve over \mathbb{Q} and F/\mathbb{Q} be an abelian extension such that $E_{p^\infty}(F) \neq 0$. Then it is easy to see (cf. [3, Cor. 3.6]) that $F(E_p^\infty)/F(\mu_p)$ is a pro- p extension. Thus these elliptic curves provide examples where the above Proposition 1.9 with $K = F(\mu_p)$ applies. More specific examples are:

- $E : y^2 + xy = x^3 - x - 1$ and $F = \mathbb{Q}(\mu_7)$ or
- $E : y^2 + xy = x^3 - 3x - 3$ and $F = \mathbb{Q}(\mu_5)$ (see *loc. cit.* 4.7. and 4.8.).

1.7. Étale cohomology classes as twists of units

Recall that L^{cyc} is the maximal unramified abelian p -extension of K^{cyc} , in which every prime above p splits completely, and that we have an isomorphism $H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \cong \varprojlim_n \mathcal{O}_{K_n}[1/S]^* \otimes \mathbb{Z}_p$ by Proposition 1.2 and (1.5).

THEOREM 1.11. — *Suppose that \mathcal{G} as defined in 1.1 is pro- p and that $\text{Gal}(L^{\text{cyc}}/K^{\text{cyc}})$ is a finitely generated \mathbb{Z}_p -module, then there exists a $J \in \mathbb{N}$ such that for all $j \geq J$ the twisting map*

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \otimes_{\mathbb{Z}_p} T(j-1) \xrightarrow{\text{Tw}_{T(j-1)}} H^1(\mathcal{O}_K[1/S], T(j))$$

has finite cokernel. In particular, for $j \geq J$ all elements in $H^1(\mathcal{O}_K[1/S], V(j))$ (where $V = T \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ as before) are “twists” of norm compatible units in

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) = \varprojlim_n H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}_p(1))$$

with a basis of the free part of $T(j-1)$.

Remark. — a) Note that there is always an open subgroup of \mathcal{G} , which is pro- p . In particular, after a finite field extension of K , one can always assume that this is the case.

b) The choice of the twist 1 in $H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1))$ and hence of the group of norm compatible units instead of any other twist is just for esthetic reasons. For the application to Euler systems and the construction of p -adic L -functions the units are certainly the most interesting case. In particular, we see this theorem as a strong confirmation of the philosophy explained in [10], that all p -adic properties of motives in connection with L -values should be encoded in the associated tower of number fields.

It is an interesting question to investigate $H^2(\mathcal{O}_K[1/S], T(j))$ with the above methods and to compare this with the results by McCallum and Sharifi [19].

Proof. — It follows from Proposition 1.9 that under our conditions

$$\text{Tor}_1^{\Lambda(\mathcal{H})}(H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p)$$

is a finitely generated \mathbb{Z}_p -module. Indeed $H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j))$ is a finitely generated $\Lambda(\mathcal{H})$ -module and $\Lambda(\mathcal{H})$ is noetherian (see [17, II.2.2.2 and V.2.2.4]), as \mathcal{H} is a closed subgroup of \mathcal{G} , hence p -adic analytic and pro- p . Thus the groups

$$\text{Tor}_r^{\Lambda(\mathcal{H})}(H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p)$$

are also finitely generated \mathbb{Z}_p -modules by standard homological algebra. The exact sequence 1.6 implies that the cokernel of

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\mathcal{G})} \Lambda(\Gamma) \longrightarrow H^1(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j))$$

is a $\Lambda(\Gamma)$ -module, say $M(j)$, which is finitely generated as a \mathbb{Z}_p -module, hence torsion as $\Lambda(\Gamma)$ -module. By the classification of torsion $\Lambda(\Gamma)$ -modules, the coinvariants of $M(j) \otimes_{\Lambda(\Gamma)} \mathbb{Z}_p$ are finite for sufficiently big j . We get an exact sequence

$$\begin{aligned} & H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\mathcal{G})} \mathbb{Z}_p \\ & \longrightarrow H^1(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\Gamma)} \mathbb{Z}_p \longrightarrow M(j) \otimes_{\Lambda(\Gamma)} \mathbb{Z}_p \rightarrow 0. \end{aligned}$$

To get the twisting map we have to compose with the first map in the following exact sequence (which is similar to Corollary 1.6)

$$\begin{aligned} & H^1(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)) \otimes_{\Lambda(\Gamma)} \mathbb{Z}_p \longrightarrow H^1(\mathcal{O}_K[1/S], T(j)) \\ & \longrightarrow \mathrm{Tor}_1^{\Lambda(\Gamma)}(H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j)), \mathbb{Z}_p). \end{aligned}$$

By our condition and Proposition 1.9 $H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j))$ is also a finitely generated \mathbb{Z}_p -module and thus $\Lambda(\Gamma)$ -torsion. As Γ is cyclic (\mathcal{G} and hence Γ is pro- p), the $\mathrm{Tor}_1^{\Lambda(\Gamma)}$ term identifies with the Γ -invariants of $H^2(\mathcal{O}_K[1/S], \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} T(j))$. Again for j big enough these are finite. \square

Remark. — In fact, if M is the p -adic realization of a motive (say $M = H^n(X \times_K \bar{K}, \mathbb{Q}_p)$ for X/K smooth, projective), one should expect that the $\Lambda(\Gamma)$ -module

$$\mathrm{Tor}_1^{\Lambda(\mathcal{G})}(H^2(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T(j)), \Lambda(\Gamma)) \otimes_{\Lambda(\Gamma)} \mathbb{Z}_p$$

is finite for all $j \geq n + 1$. Compare this with Jannsen’s Conjecture 1.8 about the vanishing of $H^2(\mathcal{O}_K[1/S], H^n(X \times_K \bar{K}, \mathbb{Q}_p(j)))$ for $j \geq n + 1$.

2. Twisting for motivic cohomology with p -adic coefficients

In this section X will always be a smooth and projective scheme over

$$D = \mathcal{O}_K[1/S].$$

The goal in this section is to study the twisting map in motivic cohomology with finite coefficients. The general assumption is that the “Bloch-Kato-conjecture” for motivic cohomology holds as announced by Voevodsky in [34] (do not confuse this with the Tamagawa number conjecture). This implies, using the Beilinson-Lichtenbaum conjecture, that we have to deal with étale cohomology of X .

2.1. Review of motivic cohomology with finite coefficients over Dedekind domains

For a variety X smooth over a Dedekind ring D , we define motivic cohomology groups as the hypercohomology of Bloch’s cycle complex $\mathbb{Z}(j)$. As usual,

$$\Delta_D^s := \text{Spec} \left(D[t_0, \dots, t_s] / \sum_i t_i - 1 \right)$$

denotes the standard algebraic s -simplex.

For a variety X smooth over a Dedekind ring D , let $z^j(X, i)$ be the free abelian group on closed integral subschemes of codimension j on $X \times_D \Delta_D^i$ which intersect all faces properly. The associated complex of presheaves (with $z^j(X, 2j - i)$ in degree i) is denoted $\mathbb{Z}(j)$, and

$$\mathbb{Z}/n(j) := \mathbb{Z}(j) \otimes^{\mathbb{L}} \mathbb{Z}/n.$$

The complex $\mathbb{Z}(j)$ (and thus also $\mathbb{Z}/n(j)$) is a complex of sheaves for the étale topology [6, Lemma 3.1], and we write $\mathbb{Z}/n(j)_{\text{ét}}$ resp. $\mathbb{Z}/n(j)_{\text{Zar}}$ when considering it as a complex of étale resp. Zariski sheaves.

DEFINITION 2.1 (compare [6, p. 779]). — *The motivic cohomology of X is the hypercohomology*

$$(2.1) \quad H_{\text{mot}}^i(X, \mathbb{Z}/n(j)) := H^i(X, \mathbb{Z}/n(j)_{\text{Zar}}).$$

Calling this motivic cohomology is justified by Voevodsky’s [35] Theorem

$$\text{Hom}_{DM^{\text{eff}}, -(K)}(M(X), \mathbb{Z}(j)[i]) =: H_{\text{mot}}^i(X, \mathbb{Z}(j)) \cong CH^j(X, 2j - i)$$

if $D = K$ is a field. In this case, higher Chow groups are defined by taking just cohomology and not hypercohomology. By [6, Thm. 3.2] both definitions coincide not only over a field but still if the base D is a discrete valuation ring.

Observe [6, Section 3] that H_{mot}^i is covariant for proper maps (with degree shift) and contravariant for flat maps. The latter applies in particular to the structural morphisms $p_n : X_n \rightarrow D_n$.

The étale cycle class

$$\text{cl} : H^i(X, \mathbb{Z}/n(j)_{\text{ét}}) \longrightarrow H^i(X, \mathbb{Z}/n(j))$$

factors through the étale sheafification $\mathbb{Z}/n(j)_{\text{ét}}$ via the map

$$\mathbb{Z}/n(j)_{\text{Zar}} \longrightarrow \mathbf{R}\pi_* \mathbb{Z}/n(j)_{\text{ét}}$$

induced by the morphism of sites $\pi : (Sm/D)_{\text{ét}} \rightarrow (Sm/D)_{\text{Zar}}$.

For us the most important consequence of the Bloch-Kato conjecture is the truth of the Beilinson-Lichtenbaum conjecture:

THEOREM 2.2 (Geisser [6, Thm. 1.2(2)(4)]). — *Assume that X is a smooth scheme over a Dedekind domain D with $n \in D^\times$ and that the “Bloch-Kato-conjecture” holds.*

1) *For all i and j there is an isomorphism*

$$H^i(X, \mathbb{Z}/n(j)_{\text{ét}}) \cong H^i(X, \mathbb{Z}/n(j))$$

of the étale hypercohomology of $\mathbb{Z}/n(j)_{\text{ét}}$ with the étale cohomology.

2) *The étale cycle class map induces isomorphisms for $0 \leq i \leq j$*

$$H^i_{\text{mot}}(X, \mathbb{Z}/n(j)) \cong H^i(X, \mathbb{Z}/n(j))$$

of motivic with étale cohomology.

2.2. The geometric twisting map

We are going to define a geometric twisting map, which will allow to relate our results for étale cohomology with motivic cohomology. The main difficulty is that the cup-product is not compatible with corestriction maps. We use the compatibility of the Hochschild-Serre spectral sequence with cup-product to overcome this and to reduce to an observation due to Soulé.

In this section we consider $X \rightarrow \text{Spec } \mathcal{O}_K[1/S]$ smooth and proper. We denote by

$$\bar{X} := X \times \bar{K}$$

and let

$$(2.2) \quad T := \varprojlim_n T_n, \quad T_n := H^i(\bar{X}, \mathbb{Z}/p^n(j-1)).$$

This is a Galois-module and finitely generated \mathbb{Z}_p -module. As in Section 1.1 we define a tower of number fields K_n , where K_n is now obtained by adjoining $H^i(\bar{X}, \mathbb{Z}/p^n(j-1))$ instead of T/p^nT to K , and a p -adic Lie group $\mathcal{G} := \text{Gal}(K_\infty/K)$. The constructions and results of Section 1 carry over to this case. Let

$$X_n := X \times_{\mathcal{O}_K[1/S]} \mathcal{O}_{K_n}[1/S]$$

and denote by $f_n : X_n \rightarrow \text{Spec } \mathcal{O}_K[1/S]$ the structure map. Denote the kernel of the edge morphism γ by

$$\begin{aligned} & H^{i+1}(X_n, \mathbb{Z}/p^n(j))^0 \\ & := \ker \{ H^{i+1}(X_n, \mathbb{Z}/p^n(j)) \xrightarrow{\gamma} H^0(\mathcal{O}_{K_n}[1/S], H^{i+1}(\bar{X}, \mathbb{Z}/p^n(j))) \}. \end{aligned}$$

LEMMA 2.3. — *The product*

$$\begin{aligned}
 H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \times H^i(X_n, \mathbb{Z}/p^n(j-1)) &\longrightarrow H^{i+1}(X_n, \mathbb{Z}/p^n(j)), \\
 \xi \times \eta &\longmapsto f_n^* \xi \cup \eta
 \end{aligned}$$

factors through $H^{i+1}(X_n, \mathbb{Z}/p^n(j))^0$.

Proof. — The pull-back

$$f_n^* : H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \longrightarrow H^1(X_n, \mathbb{Z}/p^n(1))$$

has image in $H^1(X_n, \mathbb{Z}/p^n(1))^0$ and the result follows from the fact that the Hochschild-Serre spectral sequence

$$E_2^{p,q} = H^p(\mathcal{O}_{K_n}[1/S], H^q(\bar{X}, \mathbb{Z}/p^n(j))) \implies H^{p+q}(X_n, \mathbb{Z}/p^n(j))$$

is compatible with cup-products. □

As $E_2^{1,i} = E_\infty^{1,i}$ we have a surjection,

$$(2.3) \quad H^{i+1}(X_n, \mathbb{Z}/p^n(j))^0 \longrightarrow H^1(\mathcal{O}_{K_n}[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))),$$

which we compose with the corestriction map

$$\begin{aligned}
 (2.4) \quad H^1(\mathcal{O}_{K_n}[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))) \\
 \xrightarrow{\text{cores}} H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))).
 \end{aligned}$$

We compose the cup-product in (2.3) with this composition and get

$$\begin{aligned}
 H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \times H^i(X_n, \mathbb{Z}/p^n(j-1)) \\
 \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))).
 \end{aligned}$$

Using again the compatibility of the spectral sequence with products we see that the right factor of the cup-product has to factor through the edge morphism

$$\begin{aligned}
 (2.5) \quad H^i(X_n, \mathbb{Z}/p^n(j-1)) &\xrightarrow{\gamma} H^0(\mathcal{O}_{K_n}[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j-1))) \\
 &\cong H^i(\bar{X}, \mathbb{Z}/p^n(j-1)),
 \end{aligned}$$

where the last isomorphism results from our definition of K_n (see beginning of this section). We get

$$\begin{aligned}
 (2.6) \quad H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \times H^i(\bar{X}, \mathbb{Z}/p^n(j-1)) \\
 \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))).
 \end{aligned}$$

From the construction it is obvious that:

LEMMA 2.4. — *The above map (2.6) coincides with the twisting map at finite level in equation (1.1).*

Together with the isomorphisms in 2.2 we obtain:

DEFINITION 2.5. — For $0 \leq i \leq j - 1$ the geometric twisting map is the map constructed above together with the isomorphisms in 2.2

$$H^1_{\text{mot}}(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \times H^i_{\text{mot}}(X_n, \mathbb{Z}/p^n(j-1)) \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))).$$

We apply Theorem 1.11 in this situation and get:

THEOREM 2.6. — Under the condition of Theorem 1.11 there is an integer m such that for all $n \geq m$ the cokernel of the geometric twisting map in 2.5 for $0 \leq i \leq j - 1$

$$H^1_{\text{mot}}(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \times H^i_{\text{mot}}(X_n, \mathbb{Z}/p^n(j-1)) \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j)))$$

is annihilated by p^m for all $j \geq J$.

It is an important observation by Soulé that, although the cup-product in general is not compatible with corestriction, the image in (2.5) is compatible. To formulate this properly we need:

DEFINITION 2.7. — A sequence of elements $\alpha_n \in H^r(X_n, \mathbb{Z}/p^n)$ is norm compatible if the image of $\text{cores}(\alpha_n)$ under the restriction of coefficients $\mathbb{Z}/p^n \rightarrow \mathbb{Z}/p^{n-1}$ equals α_{n-1} for all $n \geq 2$. The sequence $\{\alpha_n\}$ is reduction compatible if the reduction modulo p^{n-1} of α_n is the pull-back of α_{n-1} for all $n \geq 2$.

Note that the elements $\{\alpha_n\}$ of any \mathbb{Z}_p -lattice $T \subset H^i(\bar{X}, \mathbb{Q}_p(j))$ are reduction compatible. Soulé proves the following:

LEMMA 2.8. — If the sequence $\{\alpha_n\}$ is norm compatible and $\{\beta_n\}$ is reduction compatible, then $\{\alpha_n \cup \beta_n\}$ is norm compatible.

Proof. — This is just the projection formula, see [29, Lemma 1.4]. □

COROLLARY 2.9. — Taking the projective limit over the corestriction maps in (2.6) gives the twisting map of Definition 1.3:

$$H^1(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})(1)) \otimes H^i(\bar{X}, \mathbb{Z}_p(j-1)) \longrightarrow H^1(\mathcal{O}_K[1/S], H^i(\bar{X}, \mathbb{Z}_p(j))).$$

Remarks. — a) More generally, the above construction is possible for any theory A^* , which is covariant for proper maps, contravariant for flat maps and satisfying the projection formula $f_*(a \cup f^*(b)) = f_*(a) \cup b$ for flat

proper (or at least finite étale) maps f . We explain the case of K -theory with finite coefficients in Appendix A.

b) Soulé applies the above construction to get non-torsion elements in the K -groups of rings of integers or elliptic curves with complex multiplication. In these cases the schemes X_n are base changes of X to the ring $\mathcal{O}_{K_n}[1/S]$, where K_n is defined by adjoining p^n -th roots of unity or p^n -th division points of the elliptic curve. The towers of fields are in these cases abelian. It is shown in the cyclotomic case in [11] and [9] (with another method) and in the case of CM-elliptic curves in [16] that these twisted elements are in fact motivic, i.e., are in the image of motivic cohomology.

2.3. Compatibility of cup products in motivic and étale cohomology

The aim of this technical section is to establish the compatibility of cup products in étale and motivic cohomology. More precisely, we show that the pairing of the previous section lifts to motivic cohomology even without assuming the Bloch-Kato conjecture. The problem is that the cup product for motivic cohomology over Dedekind rings is only defined if one factor consists of equi-dimensional cycles (see Definition 2.11 below). We will show that we have a commutative diagram

(2.7)

$$\begin{CD}
 (D_n^\times \otimes \mathbb{Z}/p^n) \times H_{\text{mot}}^i(X_n, \mathbb{Z}/p^n(j-1)) @>{\cup_{\text{mot}} \circ \phi \times \text{id}}>> H_{\text{mot}}^{i+1}(X_n, \mathbb{Z}/p^n(j)) \\
 @V{\phi \times \text{cl}}VV @VV{\text{cl}}V \\
 H^1(D_n, \mathbb{Z}/p^n(1)) \times H^i(X_n, \mathbb{Z}/p^n(j-1)) @>{\cup}>> H^{i+1}(X_n, \mathbb{Z}/p^n(j)) \\
 @V{1 \times \gamma}VV @VV{\tilde{\gamma}}V \\
 H^1(D_n, \mathbb{Z}/p^n(1)) \times H^i(\bar{X}, \mathbb{Z}/p^n(j-1)) @>{\text{Tw}}>> H^1(D_n, H^i(\bar{X}, \mathbb{Z}/p^n(j)))
 \end{CD}$$

where $D_n := \text{Spec}(\mathcal{O}_{K_n}[1/S])$, γ and $\tilde{\gamma}$ are the edge maps as before, the vertical arrows cl are étale cycle class maps and the cup product \cup_{mot} as well as the map ϕ are defined below. The commutativity of the lower square follows from lemma 1.4 and the compatibility of the Hochschild-Serre spectral sequence with cup products. The commutativity of the upper square of (2.7) is discussed below; this generalizes the classical result for the usual cycle class map (see e.g. [20, Prop. VI.9.5]). Recall that by [30, Cor. 4.3] we

have an isomorphism

$$H_{\text{mot}}^i(\bar{X}, \mathbb{Z}/p^n(j)) \cong H^i(\bar{X}, \mathbb{Z}/p^n(j)).$$

Recall from [18, Section 1.7] that an irreducible scheme $Z \rightarrow D$ is *equi-dimensional* if it is dominant over D . The relative dimension $\dim_D Z$ is then defined to be the dimension of the generic fibre. Now we define the relative higher Chow group complex for our smooth $X \rightarrow D$ as follows: $z^j(X/D, p)$ to be the free abelian group generated by irreducible closed subsets $Z \subset X \times_D \Delta_D^p$, such that for each face F of Δ_D^p the irreducible components Z' of $Z \cap (X \times F)$ are equi-dimensional over D and $\dim_D Z' = \dim_D F + d - j$. Note that we have an inclusion of complexes $z^j(X/D, *) \subset z^j(X, *)$. We define equi-dimensional motivic cohomology $H_{\text{mot}}^i(X/D, \mathbb{Z}(j))$ to be the Zariski hypercohomology of the complex which has in degree i the Zariski sheafification of the presheaf $U \mapsto z^j(U/D, 2j - i)$. To define $H_{\text{mot}}^i(X/D, \mathbb{Z}/p^n(j))$ we use the same complex tensored with $\otimes^{\mathbb{L}} \mathbb{Z}/p^n$.

The units D_n^\times we use for twisting are all equi-dimensional:

LEMMA 2.10. — *The map*

$$\phi : D_n^\times \longrightarrow H_{\text{mot}}^1(D_n, \mathbb{Z}(1))$$

induced by sending $u \neq 1 \in D_n^\times$ to the graph of the rational map

$$\left(\frac{1}{1-u}, \frac{u}{u-1} \right) : \text{Spec } D_n \longrightarrow \Delta_D^1$$

restricted to $D_n - x \mid u(x) = 1$ (i.e. to a cycle in $D_n \times_D \Delta_D^1$) factors through $H_{\text{mot}}^1(D_n/D, \mathbb{Z}(1))$. The induced map

$$D_n^\times \otimes \mathbb{Z}/p^n \longrightarrow H_{\text{mot}}^1(D_n/D, \mathbb{Z}/p^n(1))$$

is injective.

Proof. — In [18, Lemma 11.2] Levine constructs a map $D_n^\times \rightarrow CH^1(D_n, 1)$ using the graph of $(1/(1-u), u/(u-1))$. Together with the natural map

$$CH^1(D_n, 1) \longrightarrow H_{\text{mot}}^1(D_n, \mathbb{Z}(1))$$

this defines ϕ and hence a map

$$D_n^\times \otimes \mathbb{Z}/p^n \longrightarrow H_{\text{mot}}^1(D_n, \mathbb{Z}/p^n(1)).$$

If we compose this with the isomorphism in 2.2, we get a map

$$D_n^\times \otimes \mathbb{Z}/p^n \longrightarrow H_{\text{ét}}^1(D_n, \mathbb{Z}/p^n(1)),$$

which is obviously (reduce to the case of a field) the map induced by the Kummer sequence, hence injective. It remains to show that the map factors through $H_{\text{mot}}^1(D_n/D, \mathbb{Z}/p^n(1))$. As the graph of $(1/(1-u), u/(u-1))$ is an equi-dimensional cycle, this follows from the definition. \square

Now we define the upper horizontal map \cup_{mot} of (2.7).

DEFINITION 2.11. — For $f : X_n \rightarrow \text{Spec}(D_n)$ smooth, we define

$$\cup_{\text{mot}} : H_{\text{mot}}^1(D_n/D, \mathbb{Z}/p^n(1)) \times H_{\text{mot}}^i(X_n, \mathbb{Z}/p^n(j-1)) \longrightarrow H_{\text{mot}}^{i+1}(X_n, \mathbb{Z}/p^n(j))$$

as the composition

$$\begin{aligned} & H_{\text{mot}}^1(D_n/D, \mathbb{Z}/p^n(1)) \times H_{\text{mot}}^i(X_n, \mathbb{Z}/p^n(j-1)) \xrightarrow{(f \times \text{id})^* \circ \cup_{D_n/D, X_n}^{r,s}} \\ & \longrightarrow H_{\text{mot}}^{i+1}(X_n \times_D D_n, \mathbb{Z}/p^n(j)) \xrightarrow{(\text{id}, p_n)^*} H_{\text{mot}}^{i+1}(X_n, \mathbb{Z}/p^n(j)). \end{aligned}$$

Here $\cup_{D_n/D, X}^{r,s} : z^s(D_n/D, *) \otimes z^r(X) \rightarrow z^{r+s}(X \times_D D_n)$ is the exterior product with integral coefficients defined by Levine [18, Section 8]. The product of the complexes of presheaves induces a product of complexes of sheaves and (using Godement resolutions as in [7]) on the hypercohomology groups.

Now we return to the commutativity of (2.7). By definition of the twisting map at finite level in 1.2, it is enough to consider the diagram

$$\begin{array}{ccc} H_{\text{mot}}^1(D_n/D, \mathbb{Z}/p^n(1)) \times H_{\text{mot}}^i(X_n, \mathbb{Z}/p^n(j-1)) & \xrightarrow{\cup_{\text{mot}}} & H_{\text{mot}}^{i+1}(X_n \times_D D_n, \mathbb{Z}/p^n(j)) \\ \downarrow \text{cl} & & \downarrow \text{cl} \\ H^1(D_n, \mathbb{Z}/p^n(1)) \times H^i(X_n, \mathbb{Z}/p^n(j-1)) & \xrightarrow{\cup} & H^{i+1}(X_n \times_D D_n, \mathbb{Z}/p^n(j)). \end{array}$$

As pointed out in [6, p. 789], the proof of [7, Prop. 4.7] for the commutativity of the corresponding diagram of varieties over fields carries over to Dedekind domains. The argument in the proof of [7, Prop. 4.7] that \cup equals the product \cup' of *loc. cit.* constructed in a way compatible with $\cup_{D_n/D, X}$ is still valid over Dedekind domains. Hence the commutativity of (2.7).

Appendix A. Twisting in p -adic K -theory

In this appendix, we will reinterpret our results in terms of p -adic K -theory.

As usual, we can define K -theory with coefficients of the exact category $\text{Vect}(X)$ of vector bundles on X using Quillen’s Q -construction and homotopy groups with finite coefficients:

DEFINITION A.1. — *Let*

$$K_r(X, \mathbb{Z}/q) := \pi_r(\Omega \text{BQVect}(X), \mathbb{Z}/q)$$

for $r > 0$ and $K_0(X, \mathbb{Z}/q) := K_0(X)/q$. Moreover, we set

$$K_r(X, \mathbb{Z}_p) := \varprojlim_n K_r(X, \mathbb{Z}/p^n)$$

and define $K_r(X, \mathbb{Q}_p) := K_r(X, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$.

Here we use that we have maps $\epsilon_n : K_r(X, \mathbb{Z}/p^n) \rightarrow K_r(X, \mathbb{Z}/p^{n-1})$ given by reduction of coefficients. Applying \varprojlim to the short exact sequence

$$0 \rightarrow K_r(X)/p^n \rightarrow K_r(X, \mathbb{Z}/p^n) \rightarrow_{p^n} K_{r-1}(X) \rightarrow 0$$

shows that $rk_{\mathbb{Z}} K_r(X) = rk_{\mathbb{Z}_p} K_r(X, \mathbb{Z}_p)$, provided the groups $K_r(X)$ and $K_{r-1}(X)$ are finitely generated (“Bass conjecture”) as proved if $X = \text{Spec}(\mathcal{O}_K)$ by Quillen [24].

We assume as before that X is smooth over $\mathcal{O}_K[1/S]$, of relative dimension d . Adams operations carry over to finite coefficients and their eigenspaces are denoted by $K(X, \mathbb{Z}/p^n)^{(j)}$ as usual. By [28, Prop. 6] the transfer maps $(f_n)_*$ respect these eigenspace decomposition (the hypothesis of *loc. cit.* is satisfied as the field extension K_n/K_{n-1} is finite).

Thomason constructs an algebraic Bott element $\beta \in K_2(X, \mathbb{Z}/p^n)$ and proves that there is an isomorphism

$$K_*(X, \mathbb{Z}/p^n)[\beta^{-1}] \xrightarrow{\cong} K_*^{\text{ét}}(X, \mathbb{Z}/p^n)$$

[32, Thm. 4.11], that $\phi_j : K_j(X, \mathbb{Z}/p^n) \rightarrow K_j^{\text{ét}}(X, \mathbb{Z}/p^n)$ is an epimorphism if $j \geq N$ and β^N annihilates $\ker(\phi_j)$ for all $j \geq 0$, where

$$N = \frac{2}{3}(d+2)(d+3)(d+4)$$

[33, Cor. 3.6]. Multiplying the short exact (for $j \geq 2N$) sequence

$$\ker(\phi_j) \rightarrow K_j(X, \mathbb{Z}/p^n) \rightarrow K_j^{\text{ét}}(X, \mathbb{Z}/p^n)$$

with β^N and applying the Snake Lemma, we get a splitting $K_{j+2N}(X, \mathbb{Z}/p^n) \rightarrow \ker(\phi_{j+2N})$ and thus étale K -theory is a natural direct summand of K -theory in these degrees. So if $2j - i - 2 \geq \frac{8}{3}(d+2)(d+3)(d+4)$, we obtain a pairing

$$K_1^{\text{ét}}(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n) \times K_{2j-i-2}^{\text{ét}}(X_n, \mathbb{Z}/p^n) \rightarrow K_{2j-i-1}^{\text{ét}}(X_n, \mathbb{Z}/p^n)$$

which is a direct summand of the corresponding pairing in algebraic K -theory with finite coefficients. Concerning the first factor, we even have an isomorphism between K_1 and $K_1^{\text{ét}}$ by [4, Prop. 8.2].

Remark A.2. — For $p = 2$, the bounds for j such that

$$K_j(X, \mathbb{Z}/p^n) \xrightarrow{\cong} K_j^{\text{ét}}(X, \mathbb{Z}/p^n)$$

have been improved by Kahn [13, Thm. 2] provided X is “non-exceptional”. He shows that it is an isomorphism if $j \geq cd_2 X - 1$. As he points out [13, p. 104], these improved bounds will carry over to odd p (without the non-exceptional restriction) assuming the Bloch-Kato conjecture for K holds.

The next step is to observe that the étale Atiyah-Hirzebruch spectral sequence degenerates ($E_2 = E_\infty$) provided $p > \frac{1}{2} cd_p X + 1$ where $cd_p X$ is the p -cohomological dimension of X , which is at most $2d + 3$ (see [1, Exposé X]). Moreover, the Adams filtration on K -theory and the weight filtration on étale cohomology coincide in a certain range [27, Thm. 2], so that the left hand side of the above pairing for $K^{\text{ét}}$ is isomorphic to

$$H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \otimes H^i(\bar{X}, \mathbb{Z}/p^n(j-1))$$

provided $p \geq \frac{1}{2}(j + cd_p X + 3)$. As $H^i(\bar{X}, \mathbb{Z}/p^n(j-1))$ is a trivial $\mathcal{O}_{K_n}[1/S]$ -sheaf, the twist of Definition 1.3 yields an isomorphism

$$\begin{aligned} H^1(\mathcal{O}_{K_n}[1/S], \mathbb{Z}/p^n(1)) \otimes H^i(\bar{X}, \mathbb{Z}/p^m(j-1)) \\ \cong H^1(\mathcal{O}_{K_n}[1/S], H^i(\bar{X}, \mathbb{Z}/p^n(j))). \end{aligned}$$

It is now possible to construct elements having property R and N for algebraic K -theory, and to proceed as in the previous section. The p -adic cycle class map has to be replaced by the p -adic regulator (take the inverse limit of [8, Definition 2.22, Example 1.4 (iii)])

$$r_p : K_{2j-i-1}(X, \mathbb{Z}_p)^{(j)} \longrightarrow H^{i+1}(X, \mathbb{Q}_p(j)).$$

Appendix B. Calculation of the inverse limit of Galois cohomology

Here we give the proof of Proposition 1.2. Let $G := G_S$ and $H' \subset H \subset G$ subgroups defining K_n and K_m , so that $G/H' \cong G_m$ and $G/H \cong G_n$ (hence H/H' is finite).

By Shapiro’s Lemma we have

$$H^i(\mathcal{O}_{K_n}[1/S], T) \cong H^i(\mathcal{O}_K[1/S], \text{Hom}_H(G, T)),$$

where $\text{Hom}_H(G, T)$ denotes the continuous maps $f : G \rightarrow T$ such that $f(hg) = hf(g)$. The group G acts on this from the left via $(gf)(x) := f(xg)$. This action we use to take the cohomology. As $H \subset G$ is a normal subgroup

we have also a G/H -action on $\text{Hom}_H(G, T)$ defined by ${}^g f(x) := gf(g^{-1}x)$. This left action commutes with the G -action and defines the structure of a $\mathbb{Z}[G/H]$ -module on $\text{Hom}_H(G, T)$. The corestriction on the left hand side

$$\text{cores} : H^i(\mathcal{O}_{K_m}[1/S], T) \longrightarrow H^i(\mathcal{O}_{K_n}[1/S], T)$$

is induced on the right hand side by the map

$$\text{tr} : \text{Hom}_{H'}(G, T) \longrightarrow \text{Hom}_H(G, T), \quad f \longmapsto \left\{ g \mapsto \sum_{h \in H/H'} hf(h^{-1}g) \right\}$$

A straightforward calculation shows that this is well-defined. Note that on T/p^n the G -action factors through G_n . Consider the isomorphism

$$\phi : \text{Hom}_H(G, T/p^n) \xrightarrow{\cong} \mathbb{Z}_p[G_n] \otimes_{\mathbb{Z}_p} T/p^n, \quad f \longmapsto \sum_{x \in G_n} (x) \otimes f(x^{-1}).$$

The G -action becomes $gf \mapsto \sum_{x \in G_n} (gx) \otimes f(x^{-1})$ and the G/H -action becomes

$${}^g f \longmapsto \sum_{x \in G_n} (g^{-1}x) \otimes gf(x^{-1}).$$

If we put all this together we obtain that the corestriction is induced by

$$\pi \otimes \text{pr} : \mathbb{Z}_p[G_m] \otimes_{\mathbb{Z}_p} T/p^m \longrightarrow \mathbb{Z}_p[G_n] \otimes_{\mathbb{Z}_p} T/p^n$$

where $\pi : \mathbb{Z}_p[G_m] \rightarrow \mathbb{Z}_p[G_n]$ is the canonical surjection (integration over the fibers) and $\text{pr} : T/p^m \rightarrow T/p^n$ the canonical projection. This proves that

$$\varprojlim_n H^i(\mathcal{O}_{K_n}[1/S], T) \cong H^i(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T).$$

The formulas above imply that the G -action on $\Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T$ is only via the first factor so that

$$R\Gamma(\mathcal{O}_K[1/S], \Lambda(\mathcal{G}) \otimes_{\mathbb{Z}_p} T) \cong R\Gamma(\mathcal{O}_K[1/S], \Lambda(\mathcal{G})) \otimes_{\mathbb{Z}_p}^{\mathbb{L}} T.$$

Here the $\Lambda(\mathcal{G})$ -action on the right hand side is induced by multiplication with the inverse on $\Lambda(\mathcal{G})$ and T has its natural $\Lambda(\mathcal{G})$ -structure induced from its G -action. This proves the proposition.

BIBLIOGRAPHY

- [1] M. ARTIN, A. GROTHENDIECK & J.-L. VERDIER, “Théorie des topos et cohomologie étale des schémas, t. 3”, in *Lect. Notes. Math.*, vol. 305, Springer, 1973.
- [2] S. BLOCH & K. KATO, “L-functions and Tamagawa numbers of motives, “The Grothendieck Festschrift”, Vol. I”, in *Progress in Math.*, vol. 86, Birkhäuser, Boston, 1990, p. 333-400.
- [3] J. COATES & R. SUJATHA, “Fine Selmer groups of elliptic curves over p -adic Lie extensions”, *Math. Annalen* **331** (2005), p. 809-839.

- [4] W. G. DWYER & E. M. FRIEDLANDER, “Algebraic and étale K -theory”, *Trans. Amer. Math. Soc.* **292** (1985), p. 247-280.
- [5] J.-M. FONTAINE & B. PERRIN-RIOU, “Cohomologie galoisienne et valeurs des fonctions L ”, in *Proceedings of Symposia in Pure Mathematics, part I*, vol. 55, 1994, p. 599-706.
- [6] T. GEISSER, “Motivic Cohomology over Dedekind rings”, *Math. Z.* **248** (2004), p. 773-794.
- [7] T. GEISSER & M. LEVINE, “The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky”, *J. reine angew. Math.* **530** (2001), p. 55-103.
- [8] H. GILLET, “Riemann-Roch theorems for higher algebraic K -theory”, *Adv. in Math.* **40** (1981), p. 203-289.
- [9] A. HUBER & G. KINGS, “Degeneration of ℓ -adic Eisenstein classes and of the elliptic polylog”, *Invent. Math.* **135** (1999), p. 545-594.
- [10] ———, “Equivariant Bloch-Kato conjecture and non-abelian Iwasawa main conjecture”, in *Proceedings ICM*, vol. II, 2002, p. 149-162.
- [11] A. HUBER & J. WILDESHAUS, “Classical motivic polylogarithm according to Beilinson and Deligne”, *Doc. Math.* **3** (1998), p. 27-133.
- [12] U. JANNSEN, “On the ℓ -adic cohomology of varieties over number fields and its Galois cohomology”, in *Galois groups over \mathbb{Q}* , Ihara et al. (eds.), MSRI Publication, 1989.
- [13] B. KAHN, “ K -theory of semi-local rings with finite coefficients and étale cohomology”, *K-Theory* **25** (2002), p. 99-138.
- [14] K. KATO, “Iwasawa theory and p -adic Hodge theory”, *Kodai Math. J.* **16** (1993), p. 1-31.
- [15] ———, “ p -adic Hodge theory and values of zeta functions of modular forms. Cohomologies p -adiques et applications arithmétiques III”, in *Astérisque*, vol. 295, Soc. Math. Fr., 2004, p. 117-290.
- [16] G. KINGS, “The Tamagawa number conjecture for CM elliptic curves”, *Invent. Math.* **143** (2001), p. 571-627.
- [17] M. LAZARD, “Groupes analytiques p -adiques”, *Pub. Math. IHÉS* **26** (1965), p. 389-603.
- [18] M. LEVINE, “ K -theory and motivic cohomology of schemes”, <http://www.math.uiuc.edu/K-theory/336>.
- [19] W. G. MCCALLUM & R. T. SHARIFI, “A cup product in the Galois cohomology of number fields”, *Duke Math. J.* **120** (2004), p. 269-310.
- [20] J. S. MILNE, *Étale Cohomology*, Princeton University Press, 1980.
- [21] F. MOREL & V. VOEVODSKY, “ \mathbf{A}^1 -homotopy theory of schemes”, *Pub. Math. IHÉS* **90** (1999), p. 45-143.
- [22] J. NEUKIRCH, A. SCHMIDT & K. WINGBERG, “Cohomology of Number Fields”, in *Grundlehren der math. Wiss.*, vol. 323, Springer, 2000.
- [23] B. PERRIN-RIOU, “ p -adic L -functions and p -adic representations”, in *SMF/AMS Texts and Monographs*, vol. 3, American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2000.
- [24] D. QUILLLEN, *Finite generation of the groups K_i of algebraic integers*, LNM, vol. 341, Springer, 1973, 178-198 pages.
- [25] J.-P. SERRE, *Cohomologie galoisienne*, Lecture Notes in Math., vol. 5^e éd., Springer, 1994, x+181 pages.
- [26] C. SOULÉ, “ K -théorie des anneaux d’entiers de corps de nombres et cohomologie étale”, *Invent. Math.* **55** (1979), p. 251-295.

- [27] ———, “Operations on étale K -theory. Applications”, in *Lecture Notes in Math.*, vol. 966, Springer, 1982, p. 271-303.
- [28] ———, “Opérations en K -théorie algébrique”, *Canad. J. Math.* **37** (1985), p. 488-550.
- [29] ———, “ p -adic K -theory of elliptic curves”, *Duke* **54** (1987), p. 249-269.
- [30] A. A. SUSLIN, “Higher Chow Groups and Étale Cohomology, Cycles, Transfers and Motivic Homology Theories”, in *Annals of Math. Studies*, vol. 143, Princeton University Press, 2000.
- [31] A. A. SUSLIN & V. VOEVODSKY, “Relative cycles and Chow sheaves, Cycles, Transfers and Motivic Homology Theories”, in *Annals of Math. Studies*, vol. 143, Princeton University Press, 2000.
- [32] R. W. THOMASON, “Algebraic K -theory and étale cohomology”, *Ann. Sci. École Norm. Sup.* **13** (1985), p. 437-552.
- [33] ———, “Bott stability in Algebraic K -theory, in “Applications of Algebraic K -theory””, *Contemp. Math.* **55** (1986), p. 389-406.
- [34] V. VOEVODSKY, “Motivic cohomology with \mathbb{Z}/ℓ -coefficients”, <http://www.math.uiuc.edu/K-theory/639>.
- [35] ———, “Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic”, *Int. Math. Res. Not.* (2002), p. 351-355.
- [36] C. WEIBEL, “An introduction to homological algebra”, in *Cambridge Studies in Advanced Mathematics*, vol. 38, Cambridge, 1994.

Manuscrit reçu le 6 janvier 2005,
accepté le 9 septembre 2005.

Jens HORNBOSTEL & Guido KINGS
Universität Regensburg
NWF I, Mathematik
93040 Regensburg (Germany)
jens.hornbostel@mathematik.uni-regensburg.de
guido.kings@mathematik.uni-regensburg.de