

ANNALES DE L'INSTITUT FOURIER

MANUEL VALDIVIA

A hereditary property in locally convex spaces

Annales de l'institut Fourier, tome 21, n° 2 (1971), p. 1-2

http://www.numdam.org/item?id=AIF_1971__21_2_1_0

© Annales de l'institut Fourier, 1971, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A HEREDITARY PROPERTY IN LOCALLY CONVEX SPACES ⁽¹⁾

by Manuel VALDIVIA

J. Dieudonné has given in [1] the two following theorems :

1) *If F is a subspace, of finite codimension, of a barrelled space E , then F is a barrelled space.*

2) *If F is a subspace, of finite codimension, of a bornological space, then F is a bornological space.*

In this paper we give a theorem analogous to the previous ones, but using infrabarrelled spaces instead of barrelled or bornological spaces. So we shall prove the following theorem : If F is a subspace, of finite codimension, of an infrabarrelled space E , then F is an infrabarrelled space.

Let K be the field of real or complex numbers. Let E be a locally convex topological vector space over the field K . If \mathfrak{B} is the family of all the absolutely convex, bounded and closed sets of E , we denote with E_B , $B \in \mathfrak{B}$, the linear hull of E with the seminorm associated to B . Let \mathfrak{C} be the topology on E , so that $E[\mathfrak{C}]$ is the inductive limit of the family $\{E_B : B \in \mathfrak{B}\}$.

THEOREM. — *Let F be a subspace of E , with finite codimension. If U is a closed, bornivorous and absolutely convex set of F , then there exists in E an U' , closed, bornivorous and absolutely convex set, such that $U' \cap F = U$.*

In particular, if E is an infrabarrelled space, then F is also an infrabarrelled space.

Proof. — Clearly, the \mathfrak{C} -topology is finer than the initial one on E . On the other hand, for every bounded set A , there exists a set $B \in \mathfrak{B}$, such that $A \subset B$. Hence A is a bounded

⁽¹⁾ Supported in part by the « Patronato para el Fomento de la Investigación en la Universidad ».

set of E_B , therefore A is a bounded set of $E[\mathcal{C}]$. That is, the bounded sets of E and those of $E[\mathcal{C}]$ are the same.

We denote with $F[\mathcal{C}]$ the subspace F , equipped with the topology induced by \mathcal{C} . Since $E[\mathcal{C}]$ is the inductive limit of seminormed spaces, it is a bornological space and, according to theorem 2), $F[\mathcal{C}]$ is a bornological space. Hence, U is a closed neighborhood of 0 in $F[\mathcal{C}]$.

Clearly, it is sufficient to prove the theorem in the case of F being a vector subspace of E , with codimension one. So that we suppose that F is so.

Two cases are possible :

1° $F[\mathcal{C}]$ being dense in $E[\mathcal{C}]$. Let \bar{U} and \bar{U}^* be the closures of U in E and $E[\mathcal{C}]$ respectively. Since U is a neighborhood of 0 in $F[\mathcal{C}]$, then \bar{U}^* is a neighborhood of 0 in $E[\mathcal{C}]$, hence \bar{U}^* is a bornivorous set in the same space.

Furthermore, $\bar{U} \supset \bar{U}^*$, then \bar{U} is a bornivorous set in E . We can take $U' = \bar{U}$, then U' is a closed, bornivorous and absolutely convex set of E , such that $U' \cap F = U$.

2° $F[\mathcal{C}]$ being closed in $E[\mathcal{C}]$. If $U = \bar{U}$, we take a vector x such that $x \in E$ and $x \notin F$. Let C be the balanced hull of the set $\{x\}$, then $U + C$ is a closed set in E and $U + C$ is a neighborhood of 0 in $E[\mathcal{C}]$, therefore, $U + C$ is bornivorous in E . If we take $U' = U + C$ the theorem is satisfied.

If $U \neq \bar{U}$, \bar{U} is absorbing in E , hence there exists an element $z \in \bar{U}$ such as $z \notin F$. Let D be the balanced hull of $\{z\}$. $U + D$ is a neighbourhood of 0 in $E[\mathcal{C}]$, hence it is bornivorous in E . Furthermore $\bar{U} \supset U$ and $\bar{U} \supset D$, then $2\bar{U} \supset U + D$, hence \bar{U} is bornivorous in E . If we take $\bar{U} = U'$ the theorem is satisfied.

BIBLIOGRAPHY

- [1] J. DIEUDONNÉ, Sur les propriétés de permanence de certaines espaces vectoriels topologiques, *Ann. Soc. Pol. Math.*, **25**, (1952), p. 50-55.

Manuscrit reçu le 15 juillet 1970.

Manuel VALDIVIA,
Facultad de Ciencias,
Paseo Valencia al Mar, 13,
Valencia (España).