

# ANNALES DE L'INSTITUT FOURIER

JEAN ERIK BJÖRK

**On extensions of holomorphic functions  
satisfying a polynomial growth condition on  
algebraic varieties in  $\mathbb{C}^n$**

*Annales de l'institut Fourier*, tome 24, n° 4 (1974), p. 157-165

[http://www.numdam.org/item?id=AIF\\_1974\\_\\_24\\_4\\_157\\_0](http://www.numdam.org/item?id=AIF_1974__24_4_157_0)

© Annales de l'institut Fourier, 1974, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

# ON EXTENSIONS OF HOLOMORPHIC FUNCTIONS SATISFYING A POLYNOMIAL GROWTH CONDITION ON ALGEBRAIC VARIETIES IN $\mathbb{C}^n$

by Jan-Erik BJÖRK

## Introduction.

Let  $\mathbb{C}^n$  be the affine complex  $n$ -space with its coordinates  $z_1, \dots, z_n$ . When  $z = (z_1, \dots, z_n)$  is a point in  $\mathbb{C}^n$  we put  $\|z\| = (|z_1|^2 + \dots + |z_n|^2)^{1/2}$ . If  $V$  is an algebraic variety in  $\mathbb{C}^n$  then  $V$  carries a complex analytic structure. A holomorphic function  $f$  on the analytic space  $V$  has a polynomial growth if there exists an integer  $N(f)$  and a constant  $A$  such that

$$|f(z)| \leq A(1 + \|z\|)^{N(f)} \quad \text{for all } z \text{ in } V.$$

Using  $L^2$ -estimates for the  $\bar{\partial}$ -equation very general results dealing with extensions of holomorphic functions from  $V$  into  $\mathbb{C}^n$  satisfying growth conditions defined by plurisubharmonic functions have been proved in [4, 8, 9]. See also [2, 3, 6]. A very special application of this theory proves that when  $V$  is an algebraic variety in  $\mathbb{C}^n$  then there exists an integer  $\varepsilon(V)$  such that the following is valid:

« If  $f$  is a holomorphic function on  $V$  with a polynomial growth of size  $N(f)$  then there exists a polynomial  $P(z_1, \dots, z_n)$  in  $\mathbb{C}^n$  such that  $P = f$  on  $V$  and the degree of  $P$  is at most  $N(f) + \varepsilon(V)$  ».

In this note some further comments about this result are given. We obtain an estimate of  $\varepsilon(V)$  using certain properties

of  $V$  based upon wellknown concepts in algebraic geometry which are recalled in the preliminary section below. The main result occurs in theorem 2.1.

Finally I wish to say that the material in this note is greatly inspired by the (far more advanced) work in [1]. See also [5] for another work closely related to this note.

## 1. Preliminaries.

The subsequent material is standard and essentially contained in [7]. Let  $P_n$  be the projective  $n$ -space over  $C$ . A point  $\xi$  in  $P_n$  is represented by a non-zero  $(n+1)$ -tuple  $(z_0, \dots, z_n)$  of complex scalars, called a coordinate representation of  $\xi$ . Here  $(z_0, \dots, z_n)$  and  $(\lambda z_0, \dots, \lambda z_n)$  represent the same point in  $P_n$  if  $\lambda$  is a non-zero complex scalar. If  $z = (z_1, \dots, z_n)$  is a point in  $C^n$  we get the point  $\mathcal{J}(z)$  in  $P_n$  whose coordinate representation is given by  $(1, z_1, \dots, z_n)$ . Then  $\mathcal{J}$  gives an imbedding of  $C^n$  into an open subset of the compact metric space  $P_n$  and the complementary set  $H_\infty = P_n \setminus \mathcal{J}(C^n)$  is called the hyperplane at infinity.

*1.a. The projective closure of an algebraic variety.* — If  $V$  is an algebraic variety in  $C^n$  then  $\mathcal{J}(V)$  is a locally closed subset of  $P_n$  and its metric closure becomes a projective subvariety of  $P_n$  which is denoted by  $\bar{V}$ . The set

$$\partial V = H_\infty \cap \bar{V}$$

is called the projective boundary of  $V$ .

A point  $\omega$  in  $H_\infty$  has a coordinate representation of the form  $(0, \omega_1, \dots, \omega_n)$  and  $\omega$  gives rise to the complex line  $L(\omega) = \{z \in C^n : z = (\lambda \omega_1, \dots, \lambda \omega_n) \text{ for some complex scalar } \lambda\}$ . In this way  $H^\infty$  is identified with the set of complex lines in  $C^n$ .

Under this identification we know that  $\partial V$  is the projective variety corresponding to the Zariski cone

$$V_c = \{z \in C^n : P^z(z) = 0 \text{ for every } P \text{ in } I(V)\}.$$

Here  $I(V) = \{P \in C[z] : P = 0 \text{ on } V\}$  and  $P^z$  denotes the leading form of a polynomial  $P$ . That is, if  $d = \deg(P)$

we have  $P = P^x + p$  where  $\deg(p) < d$  and  $P^x$  is homogeneous of degree  $d$ . Finally a point  $w$  in  $H_\infty$  belongs to  $\partial V$  if and only if the complex line  $L(w)$  is contained in the conic algebraic variety  $V_c$ .

**1.b. The Vanishing Theorem.** — Let  $\mathcal{O}$  be the sheaf of holomorphic functions on the compact complex analytic manifold  $P_n$ . Recall that  $P_n$  is covered by  $(n+1)$  many open charts  $U_i = \{\xi \in P_n : \xi \text{ has a coordinate representation of the form } (z_0, \dots, z_{i-1}, 1, z_{i+1}, \dots, z_n)\}$ . Here  $U_0 = \mathcal{J}(C^n)$  and in each intersection  $U_i \cap U_j$  we have the nowhere vanishing holomorphic function  $z_i/z_j$ .

Let  $\mathcal{S}$  be a coherent sheaf of  $\mathcal{O}$ -modules and  $m$  an integer. If  $U$  is an open subset of  $P_n$  then the sections over  $U$  of the « twisted sheaf  $\mathcal{S}(m)$  », i.e. the  $H^0(U, \mathcal{O})$ -module  $H^0(U, \mathcal{S}(m))$  are given as follows:

« An element  $a$  in  $H^0(U, \mathcal{S}(m))$  is presented by an  $(n+1)$ -tuple  $\{a_0, \dots, a_n\}$  where each  $a_i \in H^0(U \cap U_i, \mathcal{S})$  and  $a_i = (z_j/z_i)^m a_j$  holds in  $U \cap U_i \cap U_j$ .

Kodaira's Vanishing Theorem says that if  $\mathcal{S}$  is a coherent sheaf of  $\mathcal{O}$ -modules in  $P_n$  then there is an integer  $\rho(S)$  such that the cohomology groups  $H^q(P_n, \mathcal{S}(m)) = 0$  for all  $q > 0$  and every  $m > \rho(S)$ .

Let now  $\bar{V}$  be the projective variety arising from  $V$  as in 1.1 and let  $J(\bar{V})$  be its associated sheaf of ideals in  $\mathcal{O}$ . Then  $J(\bar{V})$  is a coherent sheaf of  $\mathcal{O}$ -modules and  $\bar{V}$  is a complex analytic space with its structure sheaf  $\mathcal{O}_{\bar{V}} = \mathcal{O}/J(\bar{V})$ .

**DEFINITION 1.b.** — Let  $\rho_1(V)$  be the smallest non-negative integer such that  $H^1(P_n, J(\bar{V})(m)) = 0$  for every  $m > \rho_1(V)$ .

**1.c. Normality of  $V$  at infinity.** — Let again  $V$  be an algebraic variety in  $C^n$  and  $\bar{V}$  its projective closure. Then  $\bar{V}$  is a compact analytic space and  $\partial V$  appears as a compact analytic subspace. Let  $\Gamma$  be the sheaf of continuous and complex-valued functions on  $\bar{V}$  which are holomorphic outside  $\partial V$  and vanish identically on  $\partial V$ . It is wellknown, that  $\Gamma$  is a coherent analytic sheaf on  $\bar{V}$  and  $\Gamma$  contains the subsheaf  $\Gamma_0$  consisting of functions which are holomorphic in  $\bar{V}$  and vanish on  $\partial V$ .

In general  $\Gamma_0$  is a proper subsheaf of  $\Gamma$  and we recall how these two sheaves are related to each other. First we consider a general case.

Let  $X$  be a reduced complex analytic space and let  $Y$  be a hypersurface in  $X$ . So if  $y_0 \in Y$  then we can choose an open neighborhood  $U$  of  $y_0$  in  $X$  and some

$$\varphi \in H^0(U, \mathcal{O}_X)$$

such that  $Y \cap U = \{x \in U : \varphi(x) = 0\}$ . Let now  $f$  be a continuous function on  $U$  which is holomorphic outside  $Y \cap U$  and equal to zero on  $Y \cap U$ . We know that if  $K$  is a compact subset of  $U$  then there exists an integer  $M$ , depending on  $K$ ,  $X$  and  $Y$  only, such that the function  $\varphi^M f$  is holomorphic in a neighborhood of  $K$ . We also know that  $f$  is a so called weakly holomorphic function on  $U$  and hence  $f$  is already holomorphic in  $U$  provided that the analytic space  $X$  is normal at each point in  $Y \cap U$ .

**DEFINITION 1.c.** — *We say that the algebraic variety  $V$  is normal at infinity if each point on  $\partial V$  is a normal point for the projective variety  $\bar{V}$ .*

The previous remarks show that if  $V$  is normal at infinity then  $\Gamma = \Gamma_0$  holds. In general the following result holds, using the compactness of  $\partial V$ .

**LEMMA 1.c.** — *Let  $V$  be an algebraic variety in  $C^n$ . Then there exists an integer  $M(V)$  satisfying the following condition. If  $\{f_0, \dots, f_n\}$  is a global section of the sheaf  $\Gamma(m)$ ,  $m$  an arbitrary integer, and if we put  $\tilde{f}_0 = f_0$  and  $\tilde{f}_i = (z_0/z_i)^{M(V)} f_i$  for every  $i = 1, \dots, n$ , then  $\{\tilde{f}_0, \dots, \tilde{f}_n\}$  is a global section of the sheaf  $\Gamma_0(m + M(V))$ .*

## 2. Estimates of $\varepsilon(V)$ .

Let  $f$  be a holomorphic function on  $V$  with a polynomial growth of size  $N(f)$ . Consider a point  $\xi_0 \in \partial V$  and suppose for example that  $\xi_0 \in U_1$ . Hence  $\xi_0$  has a coordinate representation  $(0, 1, y_2, \dots, y_n)$  and we put  $\Omega = \{\xi \in P_n : \xi \text{ has the coordinate representation } (\varpi_0, 1, y_2 + \varpi_2, \dots, y_n + \varpi_n)$

where every  $|\omega_v| < 1$ . Then  $\Omega$  is an open neighborhood of  $\xi_0$  in  $P_n$  and  $\Omega$  can be identified with the open unit polydisc in the  $(\omega_0, \omega_2, \dots, \omega_n)$ -space. That is, a point  $\omega = (\omega_0, \omega_2, \dots, \omega_n)$  gives the point  $\xi(\omega) = (\omega_0, 1, y_2 + \omega_2, \dots, y_n + \omega_n)$  in  $\Omega$ .

If  $z \in V$  and  $\mathcal{J}(z) \in \Omega$  we have  $\mathcal{J}(z) = (1, z_1, \dots, z_n) = (\omega_0(z), 1, y_2 + \omega_2(z), \dots, y_n + \omega_n(z))$  and it follows that  $\omega_0(z) = 1/z_1$  while  $\omega_j(z) = z_j/z_1 - y_j$  for  $j = 2, \dots, n$ .

We define  $\tilde{f}(\omega_0, \omega_2, \dots, \omega_n) = f(\mathcal{J}^{-1}(\xi(\omega)))$  over the set  $\xi^{-1}(\mathcal{J}(V) \cap \Omega)$  and conclude that there exists a constant  $A'$  such that

(x)  $|\tilde{f}(\omega_0, \omega_2, \dots, \omega_n)| |\omega_0|^{N(f)} \leq A'$  holds in  $\xi^{-1}(\mathcal{J}(V) \cap \Omega)$ .

Now  $\bar{V} \cap \Omega$  is an analytic subset of  $\Omega$  and identifying  $\Omega$  with the open unit polydisc in the  $(\omega_0, \omega_2, \dots, \omega_n)$ -space via the mapping  $\xi$  as above we can deduce from (x) that the function  $g(\omega) = (\omega_0)^{N(f)+1} \tilde{f}(\omega)$  extends continuously from  $\mathcal{J}(V) \cap \Omega$  to  $\bar{V} \cap \Omega$  and that  $g$  vanishes on  $\partial V \cap \Omega$ .

This local consideration holds for every point on  $\partial V$  and we obtain the following global result.

LEMMA 2.1. — *Let  $f$  be as above. If  $1 \leq j \leq n$  and if we put  $f_j(1, z_1, \dots, z_n) = (z_0/z_j)^{N(f)+1} f(1, z_1, \dots, z_n)$  on the set  $U_j \cap \mathcal{J}(V)$ , then  $f_j$  extends to a weakly holomorphic function on  $U_j \cap \bar{V}$  which vanishes on  $U_j \cap \partial V$ . Finally, if we put  $f_0(1, z_1, \dots, z_n) = f(z_1, \dots, z_n)$  over*

$$U_0 \cap \bar{V} = \mathcal{J}(V),$$

*then the collection  $\{f_0, \dots, f_n\}$  defines an element of*

$$H^0(\bar{V}, \Gamma(N(f) + 1)).$$

At this stage we can easily estimate  $\varepsilon(V)$ .

THEOREM 2.1. — *Let  $V$  be an algebraic variety in  $C^n$ . Let  $f$  be a holomorphic function on  $V$  with a polynomial growth of size  $N(f)$ . If  $M(V) + N(f) \geq \rho_1(V)$  then there exists a*

polynomial  $P$ , of degree  $M(V) + N(f) + 1$  at most, such that  $P = f$  on  $V$  and  $P^x = 0$  on  $V_c$ .

*Proof.* — Using lemma 2.1 we get the element  $\{f_0, \dots, f_n\}$  in  $H^0(\bar{V}, \Gamma(N(f) + 1))$  and then lemma 1.c gives the element  $\{\tilde{f}_0, \dots, \tilde{f}_n\}$  in  $H^0(\bar{V}, \Gamma_0(N(f) + M(V) + 1))$ .

Since  $m = M(V) + N(f) + 1 > \rho_1(V)$  it follows that the canonical mapping from  $H^0(P_n, \mathcal{O}(m))$  into  $H^0(\bar{V}, \mathcal{O}_{\bar{V}}(m))$  is surjective.

Since  $\Gamma_0$  is a subsheaf of  $\mathcal{O}_{\bar{V}}$  it follows that  $\{\tilde{f}_0, \dots, \tilde{f}_n\}$  belongs to the canonical image of  $H^0(P_n, \mathcal{O}(m))$ . Since  $\tilde{f}_0(\mathcal{X}(z)) = f(z)$  for every  $z$  in  $V$  while each  $\tilde{f}_j$  vanishes over  $U_j \cap \partial V$  when  $j = 1, \dots, n$ , this means that there exists a polynomial  $P(z_1, \dots, z_n)$ , of degree  $m$  at most, such that  $P = f$  on  $V$  and  $P^x = 0$  on  $V_c$ . Here the last fact follows because  $\partial V$  is the projective variety corresponding to the Zariski cone  $V_c$ .

**COROLLARY 2.1.** — *Let  $V$  be an algebraic variety which is normal at infinity. If  $f$  is a holomorphic function on  $V$  with a polynomial growth  $N(f)$ , then there exists a polynomial  $P$  satisfying  $P = f$  on  $V$  while  $P^x = 0$  on  $V_c$  and*

$$\deg(P) \leq \max(1 + N(f), 1 + \rho_1(V)).$$

### 3. The asymptotic estimate of $\varepsilon(V)$ .

Let again  $V$  be an algebraic variety in  $C^n$  where we assume that every irreducible component of  $V$  has a positive dimension. We have the following wellknown result.

**LEMMA 3.1.** — *Let  $f$  be a non-zero holomorphic function on  $V$  with a polynomial growth. Then there exists a non-negative rational number  $Q(f)$  such that  $\limsup \{\|z\|^{-Q(f)} |f(z)| : z \in V \text{ and } \|z\| \rightarrow +\infty\}$  exists as a finite and positive real number.*

**DEFINITION 3.2.** — *When  $k \geq 0$  is an integer we put  $\text{hol}(V, k) = \{f : f \text{ is a holomorphic function on } V \text{ with a polynomial growth } Q(f) \text{ satisfying } Q(f) < k\}$ . We also put  $\text{Hol}(V, k) = \{f : Q(f) = k\}$ .*

In lemma 2.1 we proved that when  $f \in \text{Hol}(V, k)$  then  $f$  determines an element of  $H^0(\bar{V}, \Gamma(k+1))$ . If  $f \in \text{hol}(V, k)$  we can set

$$g_j(\mathcal{J}(z)) = (z_0/z_j)^k f(\mathcal{J}(z)) \text{ for all } z \text{ in } V \cap \mathcal{J}^{-1}(U_j).$$

The same argument as in the proof of lemma 2.1 shows that every  $g_j$  extends continuously to  $\bar{V} \cap U_j$  and vanishes on  $\partial V \cap U_j$ . It follows that  $\{g_0, \dots, g_n\}$  defines an element of  $H^0(\bar{V}, \Gamma(k))$ .

Conversely, if  $\{g_0, \dots, g_n\} \in H^0(\bar{V}, \Gamma(k))$  and if we put  $f(z) = g_0(\mathcal{J}(z))$  for all  $z$  in  $V$  then it is easily verified that  $f \in \text{hol}(V, k)$ . Finally the density of  $V$  in  $\bar{V}$  implies that the section  $\{g_0, \dots, g_n\}$  is uniquely determined by  $f$ .

Summing up, we get the following inclusions.

LEMMA 3.3. — *If  $k \geq 0$  is an integer then*

$$H^0(\bar{V}, \Gamma(k)) = \text{hol}(V, k) \subset \text{Hol}(V, k) \subset H^0(\bar{V}, \Gamma(k+1)).$$

DEFINITION 3.4. — *Let  $V$  be an algebraic variety in  $C^n$ . We let  $\varepsilon_\infty(V)$ , resp.  $e_\infty(V)$ , be the smallest non-negative integer such that for all sufficiently large integers  $k$  and every  $f$  in  $\text{Hol}(V, k)$ , resp. every  $f$  in  $\text{hol}(V, k)$ , there exists a polynomial  $P$  of degree  $k + \varepsilon_\infty(V)$ , resp. of degree  $k + e_\infty(V)$ , at most, such that  $P = f$  on  $V$  and  $P^x = 0$  on  $V_c$ .*

The following invariant of  $V$  is the asymptotic analogue of the integer  $M(V)$ .

DEFINITION 3.5. — *Let  $M_\infty(V)$  be the smallest integer such that for all sufficiently large integers  $k$  and every  $f$  in*

$$H^0(\bar{V}, \Gamma(k)),$$

*it follows that  $\tilde{f} \in H^0(\bar{V}, \Gamma_0(k + M_\infty(V)))$ , where*

$$\tilde{f} = \{\tilde{f}_0, \dots, \tilde{f}_n\}$$

*and  $\tilde{f}_j = (z_0/z_j)^{M_\infty(V)} f_j$  in  $\bar{V} \cap U_j$ .*

Using lemma 3.3 and the same argument as in the proof of theorem 2.1 we get the result below.



**THEOREM 3.1.** —  $M_\infty(V) = e_\infty(V) \leq \varepsilon_\infty(V) \leq M_\infty(V) + 1$ .

We finish this discussion with a remark about the invariant  $M_\infty(V)$ . Recall first that if  $f = \{f_0, \dots, f_n\} \in H^0(\bar{V}, \Gamma(k))$  for some integer  $k$  and if  $P(z_0, \dots, z_n)$  is a homogenous polynomial of degree  $\nu$ , then we get the element  $f \otimes P$  in  $H^0(\bar{V}, \Gamma(k + \nu))$ , where

$$(f \otimes P)_j = (P/z_j^\nu) f_j \quad \text{in } \bar{V} \cap U_j.$$

This simply describes the structure of the graded

$$\mathbb{C}[z_0, \dots, z_n]\text{-module}$$

$G(\Gamma) = \bigoplus H^0(\bar{V}, \Gamma(k))$ . Since  $\Gamma$  is a coherent analytic sheaf we know that  $G(\Gamma)$  is a finitely generated  $\mathbb{C}[z_0, \dots, z_n]$ -module and hence there is an integer  $\nu(\Gamma)$  such that when  $k > \nu(\Gamma)$  then every element in  $H^0(\bar{V}, \Gamma(k))$  is a linear combination of elements of the form  $f \otimes P$ , where

$$f \in H^0(\bar{V}, \Gamma(\nu(\Gamma)))$$

and  $P$  is a homogenous polynomial of degree  $k - \nu(\Gamma)$ .

There is a similar integer  $\nu(\Gamma_0)$  for the graded module  $G(\Gamma_0)$  arising from the coherent sheaf  $\Gamma_0$ . When  $k$  is an integer we let  $\gamma(k)$  be the smallest integer such that for every  $f$  in  $H^0(\bar{V}, \Gamma(k))$  it follows that

$$\tilde{f} \in H^0(\bar{V}, \Gamma_0(k + \gamma(k))),$$

where  $\tilde{f}_j = (z_0/z_j)^{\gamma(k)} f_j$  and  $j = 0, \dots, n$ .

It is easily seen that  $\gamma(k)$  is a decreasing function of  $k$ , provided that  $k \geq \sup \{\nu(\Gamma), \nu(\Gamma_0)\}$ . Finally

$$M_\infty(V) = \lim_{k \rightarrow +\infty} \gamma(k)$$

and we conclude that there exists an integer  $\gamma(V)$  such that

$$M_\infty(V) = \gamma(k) \quad \text{for all } k \geq \gamma(V).$$

## BIBLIOGRAPHY

- [1] P. A. GRIFFITHS, Function theory of finite order on algebraic varieties, *Journ. of Diff. Geom.*, 6 (1972), 285-306.
- [2] L. EHRENPREIS, A fundamental principle for systems of linear differen-

- tial equations with constant coefficients and some of its applications, *Proc. International Symp. on Linear Spaces*, Jerusalem (1960).
- [3] L. EHRENPREIS, Fourier analysis in several complex variables, *Pure and Appl. Math.*, 17 (1970), Wiley-Intersci. Publ.
  - [4] L. HÖRMANDER,  $L^2$ -estimates and existence theorems for the  $\bar{\partial}$ -operator, *Acta Math.*, 113 (1965), 89-152.
  - [5] R. NARASIMHAN, Cohomology with bounds on complex spaces, *Springer Lecture Notes*, 155 (1970), 141-150.
  - [6] V. P. PALAMODOV, Linear differential operators with constant coefficients, Springer-Verlag, 16 (1970).
  - [7] J.-P. SERRE, Géométrie analytique et géométrie algébrique, *Ann. Inst. Fourier*, 6 (1955), 1-42.
  - [8] H. SKODA,  $d''$ -cohomologie à croissance lente dans  $\mathbb{C}^n$ , *Ann. Sci. de l'École Norm. Sup.*, 4 (1971), 97-121.
  - [9] H. SKODA, Applications des techniques à  $L^2$  à la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids, *Ann. Sci. de l'École Norm. Sup.*, 5 (1972), 545-579.

Manuscrit reçu le 5 octobre 1973,  
accepté par J. Dieudonné.

Jan-Erik BJÖRK,  
Department of Mathematics  
University of Stockholm  
Box 6701  
11385 Stockholm (Sweden).